

## Reliability of Hallux Rigidus Radiographic Grading System

Ryan C. Pate, M.D., John W. Fanning, M.D., Naomi N. Shields, M.D.,  
Alexander C.M. Chong, MSAE, MSME  
University of Kansas School of Medicine-Wichita  
Department of Orthopaedics

### Abstract

**Introduction.** The purpose of this study was to determine the inter- and intra-observer reliability of a clinical radiographic scale for hallux rigidus.

**Methods.** A total of 80 patients were retrospectively selected from the patient population of two foot and ankle orthopaedic surgeons. Each corresponding series of radiographic images (weight-bearing anteroposterior, weight-bearing lateral, and oblique of the foot) was randomized and evaluated. Re-randomization was performed and the corresponding radiograph images re-numbered. Four orthopaedic foot and ankle surgeons graded each patient, and each rater reclassified the re-randomized radiographic images three weeks later.

**Results.** Sixty-one out of 80 patients (76%) were included in this study. For intra-observer reliability, most of the raters showed “excellent” agreement except one rater had a “substantial” agreement. For inter-observer reliability, only 14 out of 61 cases (23%) showed total agreement between the eight readings from the four surgeons, and 11 out of the 14 cases (79%) were grade 3 hallux rigidus. One of the raters had a tendency to grade at a higher grade resulting in poorer agreement. If this rater was excluded, the results demonstrated a “substantial” agreement by using this classification.

**Conclusion.** The hallux rigidus radiographic grading system should be used with caution. Although there is an “excellent” level of intra-observer agreement, there is only “moderate” to “substantial” level of inter-observer reliability.

*KS J Med* 2015;8(4):125-134.

### Introduction

Osteoarthritis of the first metatarsophalangeal (MTP) joint of the foot may cause significant pain, disability, and difficulty wearing footwear. The term, hallux rigidus, is used to describe a condition commonly associated with degenerative arthritis of the first MTP joint with osteophyte formation, which results in a painful joint with reduction in the range of motion, especially dorsiflexion.<sup>1-3</sup> Hallux rigidus is a progressive condition, and may present in early or late stages with varying degrees of stiffness and osteophytic thickening of the joint. Chronic MTP joint inflammation leads to capsular distention and eventually to a loss of capsular and collateral ligament integrity.

Throughout the literature discussing foot and ankle disabilities, there have been multiple classification methods for hallux rigidus that have involved clinical findings,<sup>4</sup> radiographic findings,<sup>5-9</sup> or a combination of both.<sup>10-14</sup> The role of these classification systems is to help a physician to choose an appropriate method of treatment as well as to provide a reasonably precise estimation of the outcome of that treatment.<sup>3,5,8,10,14-16</sup> Some researchers have used these classification systems to compare the results of different studies and treatment procedures.<sup>6,7,9,11-13,17-26</sup>

For these classification systems to be useful, the classification system must produce the same desired results time after time in the hands of any physician or researcher who attempts to use it. Reliable

testing is critical to the orthopaedic literature, including hand conditions,<sup>27-29</sup> radiograph measurements,<sup>30-33</sup> Legg-Calve-Perthes disease grading,<sup>34-35</sup> joint arthroplasty loosening,<sup>36</sup> and fracture classifications.<sup>37-48</sup> Beeson et al.<sup>26</sup> performed an exhaustive literature review on hallux rigidus classification systems, and found a total of 18 different classification systems without any studies to determine the reliability of the systems. Clinical radiographic grading system is the fundamental assessment tool to classify the severity of hallux rigidus among all the different classification systems.<sup>5-14</sup> Giannini et al.<sup>5</sup> and Coughlin et al.<sup>10</sup> presented a reasonable summary of the various radiographic grading systems. To our knowledge, there has not been a study that specifically addressed the reliability of radiographic grading for hallux rigidus. The purpose of this study was to determine the inter- and intra-observer reliability of a clinical radiographic scale for hallux rigidus.

## Methods





**Participants.** A total of 80 patients were selected retrospectively from the patient population of two orthopaedic surgeons, who specialized in foot and ankle surgery, in a mid-western city. The study sample was selected based on three radiographs (weight bearing anterior-posterior (AP), oblique, and lateral) of the patients who were diagnosed with hallux rigidus. Poor quality or inadequate radiographs, or evidence of prior surgery were exclusions in this study. Patients with inter-metatarsal angles of greater than 15 degrees (normal is 9 degrees) or hallux valgus angles greater than 20 degree (normal is 15 degrees) also were excluded.

**Instruments and Procedures.** This study was approved by the Human Subjects Committees as minimal risk and with a

waiver of consent and waiver of HIPAA authorization. Three different hard copy radiograph images that had been used for clinical decision-making for each selected patient were obtained. Radiograph images included views of the hallux from the weight-bearing AP, weight-bearing lateral, and oblique radiographs. These radiographs were de-identified of any patient information, and were enhanced and converted to black and white using Kodak EasyShare software (Version 8.2, Kodak, Rockester, NY). Each corresponding series of radiographic images (weight-bearing AP, weight-bearing lateral, and oblique) was randomized, given a number, and recorded on a CD-ROM disk of images.

The inter- and intra-observer reliability for classifying the hallux rigidus involved adjustment of the proportion of agreement among observers with a correction for the proportion of expected agreement by chance. To evaluate inter-observer variability, four attending orthopaedic surgeons whom were trained in foot and ankle surgery were asked to classify the group of radiographic images independently according to the Giannini-modified Coughlin and Shurnas' classification systems (Table 1). Each attending orthopaedic surgeon was given a packet which contained descriptions and diagrams of Giannini's modification of Coughlin and Shurnas' grading system,<sup>5</sup> a score sheet, a CD-ROM disk of radiographic images, and a return mail envelope. To evaluate intra-observer reliability, two rounds of scoring were conducted for each rater with re-randomization of the radiographic images three weeks later and re-numbering between each round.

Table 1. Grading System for Hallux Rigidus<sup>5</sup> (JBJS License Number: 2002840476064).

<b>Grade</b>		<b>Radiographic Findings</b>
Grade 0		Normal
Grade 1		Dorsal osteophyte is main finding, minimal joint space narrowing, minimal periarticular sclerolosis, minimal flattening of the metatarsal heads with a lateral spur
Grade 2		Dorsal, lateral, and possibly medial osteophytes with a flattened appearance of the metatarsal head, no more than ¼ of dorsal joint space involved on the lateral radiograph, and mild to moderate joint space narrowing and sclerosis, sesamoids usually not involved
Grade 3		Substantial joint space narrowing, periarticular cystic changes, more than ¼ of dorsal joint space involved, sesamoids are enlarged, cystic, and/or irregular

**Statistics.** The inter- and intra-observer reliability for classifying the hallux rigidus was calculated with the use of weighted Kappa coefficients by using the SPSS software (Version 16.0; SPSS Inc., Chicago, IL). According to guidelines described by Landis and Koch,<sup>49</sup> a value of  $\leq 0.2$  indicates “poor” or “slight” agreement, 0.21 to 0.40 is “fair” agreement, 0.41 to 0.6 is “moderate” agreement, 0.61 to 0.8 is “substantial” agreement, and  $> 0.80$  is “excellent” agreement. In addition, the percentage of patients where all four examiners agreed on the grade was determined.

## Results

Of the 80 patients diagnosed with hallux rigidus from the two foot and ankle surgeons’ patient populations, 61 patients (76%) met the required criteria and were included in this study. For intra-observer reliability, most of the attending surgeons showed “excellent” agreement by using the Giannini-modified Coughlin and Shurnas’ classification systems to grade the hallux rigidus of the foot (mean weighted Kappa coefficient:  $0.82 \pm 0.07$ ; range: 0.72 - 0.88; Table 2). These results implied that each rater agreed well with themselves when reading the same radiographs at different time points. Only one of the raters had a “substantial” agreement (weighted Kappa coefficient of 0.72).

Table 2. Intra- and inter-observer reliability.

		Rater 1	Rater 2		Rater 3		Rater 4	
		Reading 2	Reading 1	Reading 2	Reading 1	Reading 2	Reading 1	Reading 2
Rater 1	Reading 1	High*	High	Substantial	Substantial	High	Moderate	Moderate
	Reading 2		Substantial	Substantial	Substantial	High	Moderate	Moderate
Rater 2	Reading 1			High*	Substantial	Substantial	Moderate	Moderate
	Reading 2				Substantial	Substantial	Moderate	Moderate
Rater 3	Reading 1					High*	Moderate	Moderate
	Reading 2						Moderate	Moderate
Rater 4	Reading 1							Substantial*

\*Represents intra-observer reliability

For inter-observer reliability, only 14 out of the 61 cases (23%) showed total agreement between the eight readings from the four surgeons, and 11 out of the 14 cases (79%) were grade 3 hallux rigidus. Figures 1 and 2 illustrate “excellent” agreement cases for Grade 2 and Grade 3 hallux rigidus, respectively. Most of the cases showed “excellent” agreement within one grade difference (53 out of 61, 87%) and the mean weighted Kappa was  $0.64 \pm 0.13$  (range:

0.44 -0.83). Figure 3 shows an example of poor agreement. One of the raters had a tendency to grade the hallux rigidus radiographs at a higher grade than the other three raters, resulting in poorer agreement. If this rater was excluded, the results show a “substantial” agreement by using this classification to grade the hallux rigidus of the foot (mean weighted Kappa coefficient:  $0.76 \pm 0.06$ ; range: 0.68 - 0.83).




Figure 1. Radiographs demonstrating good agreement case for Grade 2 hallux rigidus.




Figure 2. Radiographs demonstrating good agreement case for Grade 3 hallux rigidus.




Figure 3. Radiographs demonstrating poor agreement case for hallux rigidus.

### Discussion

Hallux rigidus is a common form of osteoarthritis in the foot.<sup>50</sup> Radiographic examination, including weight-bearing AP and lateral radiographs, usually finds asymmetric joint narrowing and a flattened metatarsal head. The lateral radiographs usually are the most revealing. With advancement of the disease, more of the joint surface is involved. Subchondral cysts, sclerosis, and bony proliferation at the joint margins occur and the joint narrowing progresses.<sup>19,31,51</sup> With the use of the radiographic grading system, orthopaedic surgeons should be able to provide optimum care to patients who have these common acquired disorders of the foot. The Giannini-modified Coughlin and Shurnas'

classification system, like all other classification systems, is intended to aid clinical decision-making for treatment as well as to provide a reasonably precise estimation of the treatment outcome for hallux rigidus. There are many other hallux rigidus classification systems which are very similar to each other. This study used the Giannini-modified Coughlin and Shurnas' classification system because it is widely referred in studies. However, this classification system relies on radiographic findings, regardless of subjective and clinical findings. To be useful, a classification should have at least moderate rater consistency. The results of this study indicated that this particular grading system should be used with caution, as only 75%

reach “excellent” agreement for intra-observer reliability, and “moderate” to “substantial” agreement for the inter-observer reliability. The practical utility of having a system with only high intra-observer reliability is questionable, and it likely would not provide any help with communications between physicians or researchers regarding the population in their studies.

A major point of concern with a radiograph-only system for hallux rigidus was that radiographs are only a part of the evaluation of a patient with hallux rigidus. Coughlin et al.<sup>10</sup> addressed this concern and included a fourth category for patients with pain in the midrange of motion (a clinical finding) and grade 3 radiographic changes. An ideal study with this subject would have both a radiographic and a clinical exam component which would reproduce the clinician’s experience treating this disorder more closely. The logistics of such a study likely would be difficult.

To achieve optimal results, surgical treatment should be individualized with use of different surgical techniques depending upon the degree of arthritis and other clinical considerations. Non-operative treatment, including modifications of shoe wear, use of a shoe insert, and use of anti-inflammatory medication, should be discussed in detail with the patient in accordance to the degree of symptoms.<sup>10,52</sup> If non-operative measures fail, operative intervention, such as arthrodesis, arthroplasty, cheilectomy, proximal phalanx osteotomy, dorsal closing wedge osteotomy, waterman green, Youngswick, Reverdin green, distal oblique sliding osteotomy, sagittal Z osteotomy, and Drago may be indicated.<sup>53</sup> Cheilectomy, which essentially consists of a debridement arthroplasty of the joint, may be appropriate.<sup>54,55</sup> Once more extensive involvement has occurred, arthrodesis is preferred for younger patients

whereas resection arthroplasty may be more appropriate for elderly patients who have a less active lifestyle.<sup>56</sup> Taranow et al.<sup>57</sup> recently presented a different classification system and surgical algorithm for treatment of the varied manifestations of hallux rigidus. This classification includes radiographic findings, motion restriction, and location of pain to guide appropriate surgical choices better. They also recommended procedures to preserve motion, when present, and address the significance of mid-motion and sesamoid pain.

In this study, there were several limitations. First, this was a pilot study that addresses an area where further research is needed. The sample size was relatively small and patients were only drawn from practices of two local foot and ankle surgeons. As such, only four raters were included in the study and bias of an outlier potentially could affect the inter-observer reliability substantially. Furthermore, each rater only evaluated the hallux rigidus radiographs on two occasions.

This study was limited due to the presence of fewer “normal” radiographs rather than “abnormal” radiographs. This also was a retrospective study evaluating a single radiographic classification system. Further research should include a larger sample size, multiple foot and ankle surgeons, and patients should be followed prospectively to assess the validity of the classification system treatment outcome and establish guidelines that would allow orthopedists to allocate their treatment more efficiently.

## **Conclusion**

This study was the first to evaluate the reliability of any hallux rigidus radiographic grading system. Overall, this hallux rigidus radiographic grading system should be used with caution as the results showed that even

though there is an “excellent” level of intra-observer agreement, but there is only “moderate” to “substantial” level of inter-observer reliability. As is common in many orthopaedic grading systems, the overall reliability of this grading system was not “excellent”, thus they may cause confusion with communication in the literature regarding the treatment of hallux rigidus. Further studies are encouraged and needed to support the conclusion of this study.

### Acknowledgements

The authors thank Paul H. Wooley, Ph.D., for his contributions of editing and comments on this manuscript.

### References

- <sup>1</sup> Mann RA, Coughlin MJ, DuVries HL. Hallux rigidus: A review of the literature and a method of treatment. *Clin Orthop Relat Res* 1979; (142):57-63. PMID: 498649.
- <sup>2</sup> Mann RA. Hallux Rigidus. *AAOS Instructional Course Lecture* 1990; (39):15-21.
- <sup>3</sup> Shereff MJ, Baumhauer JF. Hallux rigidus and osteoarthritis of first metatarsophalangeal joint. *J Bone Joint Surg Am* 1998; 80(6):898-908. PMID: 9655109.
- <sup>4</sup> Bonney G, Macnab I. Hallux valgus and hallux rigidus: A critical survey of operative results. *J Bone Joint Surg Br* 1952; 34-B(3):366-385. PMID: 12999918.
- <sup>5</sup> Giannini S, Ceccarelli F, Faldini C, Bevoni R, Grandi G, Vannini F. What’s new in surgical options for hallux rigidus? *J Bone Joint Surg Am* 2004; 86-A Suppl 2:72-83. PMID: 15691111.
- <sup>6</sup> Coughlin MJ. Rheumatoid forefoot reconstruction: A long-term follow-up study. *J Bone Joint Surg Am* 2000; 82(3):322-341. PMID: 10724225.

- <sup>7</sup> Hattrup SJ, Johnson KA. Subjective results of hallux rigidus following treatment with cheilectomy. *Clin Orthop Relat Res* 1988; (226):182-191. PMID: 3335093.
- <sup>8</sup> Hanft JR, Mason ET, Landsman AS, Kashuk KB. A new radiographic classification for hallux limitus. *J Foot Ankle Surg* 1993; 32(4):397-404. PMID: 8251995.
- <sup>9</sup> Roukis TS, Jacobs PM, Dawson DM, Erdmann BB, Ringstrom JB. A prospective comparison of clinical, radiographic, and intraoperative features of hallux rigidus. *J Foot Ankle Surg* 2002; 41(2):76-95. PMID: 11995839.
- <sup>10</sup> Coughlin MJ, Shurnas PS. Hallux rigidus: Grading and long-term results in operative treatment. *J Bone Joint Surg Am* 2003; 85-A(11):2072-2088. PMID: 14630834.
- <sup>11</sup> Viegas GV. Reconstruction of hallux limitus deformity using a first metatarsal saggittal-Z osteotomy. *J Foot Ankle Surg* 1998; 37(3):204-11; discussion 261-262. PMID: 9638545.
- <sup>12</sup> Kurtz DH, Harrill JC, Kaczander BI, Solomon MG. The Valenti procedure for hallux limitus: A long-term follow-up and analysis. *J Foot Ankle Surg* 1999; 38(2):123-130. PMID: 10334699.
- <sup>13</sup> Lombardi CM, Silhanek AD, Connolly FG, Dennis LN, Keslonsky AJ. First metatarsophalangeal arthrodesis for treatment of hallux rigidus: A retrospective study. *J Foot Ankle Surg* 2001; 40(3):137-143. PMID: 11417595.
- <sup>14</sup> Geldwert JJ, Rock GD, McGrath MP, Mancuso JE. Cheilectomy: Still a useful technique for grade I and grade II hallux limitus/rigidus. *J Foot Surg* 1992; 31(2):154-159. PMID: 1645002.
- <sup>15</sup> Pontell D, Gudas CJ. Retrospective analysis of surgical treatment of hallux rigidus/limitus: Clinical and radiographic follow-up of hinged,

- silastic implant arthroplasty and cheilectomy. *J Foot Surg* 1988; 27(6):503-10. PMID: 3243957.
- <sup>16</sup> Vanore JV, Christensen JC, Kravitz SR, et al. Diagnosis and treatment of first metatarsophalangeal joint disorders. Section 2: Hallux rigidus. *J Foot Ankle Surg*. 2003 May-Jun; 42(3):124-36. PMID: 12815579.
- <sup>17</sup> Blyth MJ, Mackay DC, Kinninmonth AW. Dorsal wedge osteotomy in the treatment of hallux rigidus. *J Foot Ankle Surg* 1998; 37(1):8-10. PMID: 9470110.
- <sup>18</sup> Barca F. Tendon arthroplasty of the first metatarsophalangeal joint in hallux rigidus: Preliminary communication. *Foot Ankle Int* 1997; 18(4):222-228. PMID: 9127112.
- <sup>19</sup> Coughlin MJ, Shurnas PJ. Soft-tissue arthroplasty for hallux rigidus. *Foot Ankle Int* 2003; 24(9):661-672. PMID: 14524514.
- <sup>20</sup> Easley ME, Davis WH, Anderson RB. Intermediate to long-term follow-up of medial-approach dorsal cheilectomy for hallux rigidus. *Foot Ankle Int* 1999; 20(3):147-152. PMID: 10195291.
- <sup>21</sup> Mulier T, Steenwerckx A, Thienport E, Sioen W, Hoore KD, Peeraer L, Dereymaeker G. Results after cheilectomy in athletes with hallux rigidus. *Foot Ankle Int* 1999; 20(4):232-237. PMID: 10229279.
- <sup>22</sup> Roukis TS, Jacobs PM, Dawson DM, Erdmann BB, Ringstrom JB. A prospective comparison of clinical, radiographic, and intraoperative features of hallux rigidus: Short-term follow-up and analysis. *J Foot Ankle Surg* 2002; 41(3):158-165. PMID: 12075903.
- <sup>23</sup> Feltham GT, Hanks SE, Marcus RE. Age-based outcomes of cheilectomy for the treatment of hallux rigidus. *Foot Ankle Int* 2001; 22(3):192-197. PMID: 11310859.
- <sup>24</sup> Pulavarti RS, McVie JL, Tulloch CJ. First metatarsophalangeal joint replacement using the bio-action great toe implant: Intermediate results. *Foot Ankle Int* 2005; 26(12):1033-1037. PMID: 16390635.
- <sup>25</sup> Kilmartin TE. Phalangeal osteotomy versus first metatarsal decompression osteotomy for surgical treatment of hallux rigidus: A prospective study of age-matched and condition-matched patients. *J Foot Ankle Surg* 2005; 44(1):2-12. PMID: 15704077.
- <sup>26</sup> Beeson P, Phillips C, Corr S, Ribbans W. Classification systems for hallux rigidus: A review of the literature. *Foot Ankle Int* 2008; 29(4):407-414. PMID: 18442456.
- <sup>27</sup> Guillemin F, Billot L, Boini S, Gerard N, Odegaard S, Kvien TK. Reproducibility and sensitivity to changes of 5 methods of scoring hand radiographic damage in patients with rheumatoid arthritis. *J Rheumatol* 2005; 32(5):778-786. PMID: 15868609.
- <sup>28</sup> Sodha S, Ring D, Zurakowski D, Jupiter J. Prevalence of osteoarthritis of the trapeziometacarpal joint. *J Bone Joint Surg Am* 2005; 87(12):2614-2618. PMID: 16322609.
- <sup>29</sup> Jafarnia K, Collins ED, Kohl HW 3rd, Bennett JB, Ilahi OA. Reliability of the Lichtman classification of Kienbock's disease. *J Hand Surg Am* 2000; 25(3):529-534. PMID: 10811758.
- <sup>30</sup> Saltzman CL, Brandser EA, Berbaum KS, et al. Reliability of standard foot radiographic measurements. *Foot Ankle Int* 1994; 15(12):661-665. PMID: 7894638.
- <sup>31</sup> Coughlin MJ, Freund E. The reliability of angular measurements in hallux valgus deformities. *Foot Ankle Int* 2001; 22(5):369-379. PMID: 11428754.
- <sup>32</sup> Inan M, Jeong C, Chan G, Mackenzie WG, Glutting J. Analysis of lower



- extremity alignment in achondroplasia: Interobserver reliability and intraobserver reproducibility. *J Pediatr Orthop* 2006; 26(1):75-78. PMID: 16439907.
- <sup>33</sup> Carman DL, Browne RH, Birch JG. Measurement of scoliosis and kyphosis radiographs. Intraobserver and interobserver variation. *J Bone Joint Surg Am* 1990; 72(3):328-333. PMID: 2312528.
- <sup>34</sup> Akgun R, Yazici M, Aksoy MC, Cil A, Alpaslan AM, Tumer Y. The accuracy and reliability of estimation of lateral pillar height in determining the herring grade in Legg-Calve-Perthes disease. *J Pediatr Orthop* 2004; 24(6):651-653. PMID: 15502565.
- <sup>35</sup> Kalenderer O, Agus H, Ozcalabi IT, Ozluk S. The importance of surgeon's experience on intraobserver and interobserver reliability of classifications used for Perthes Disease. *J Pediatr Orthop* 2005; 25(4):460-464. PMID: 15958895.
- <sup>36</sup> Temmerman OP, Raijmakers PG, Berkhof J, et al. Diagnostic accuracy and interobserver variability of plain radiography, subtraction arthrography, nuclear arthrography, and bone scintigraphy in the assessment of aseptic femoral component loosening. *Arch Orthop Trauma Surg* 2006; 126(5):316-323. Epub 2006 Mar 18. PMID: 16547724.
- <sup>37</sup> Andersen DJ, Blair WF, Steyers CM Jr, Adams BD, el-Khoury GY, Brandser EA. Classification of distal radius fractures: An analysis of interobserver reliability and intraobserver reproducibility. *J Hand Surg Am* 1996; 21(4):574-582. PMID: 8842946.
- <sup>38</sup> Bernstein J, Adler LM, Blank JE, Dalsey RM, Williams GR, Iannotti JP. Evaluation of the Neer system of classification of proximal humeral fractures with computerized tomographic scans and plain radiographs. *J Bone Joint Surg Am* 1996; 78(9):1371-1375. PMID: 8816653.
- <sup>39</sup> Andersen E, Jorgensen LG, Heddam LT. Evans' classification of trochanteric fractures: An assessment of the interobserver and intraobserver reliability. *Injury* 1990; 21(6):377-378. PMID: 2276801.
- <sup>40</sup> Walton NP, Harish S, Roberts C, Blundell C. AO or Schatzker? How reliable is classification of tibial plateau fractures? *Arch Orthop Trauma Surg* 2003; 123(8):396-398. Epub 2003 Aug 12. PMID: 14574596.
- <sup>41</sup> Martin J, Marsh JL, Nepola JV, Dirschl DR, Hurwitz S, DeCoster TA. Radiographic fracture assessments: Which ones can we reliably make? *J Orthop Trauma* 2000; 14(6):379-385. PMID: 11001410.
- <sup>42</sup> Lamraski G, Monsaert A, De Maeseneer M, Haentjens P. Reliability and validity of plain radiographs to assess angulation of small finger metacarpal neck fractures: Human cadaveric study. *J Orthop Res* 2006; 24(1):37-45. PMID: 16419967.
- <sup>43</sup> Lauder AJ, Inda DA, Bott AM, Clare MP, Fitzgibbons TC, Mormino MA. Interobserver and intraobserver reliability of two classification systems of intra-articular calcaneal fractures. *Foot Ankle Int* 2006; 27(4):251-255. PMID: 16624214.
- <sup>44</sup> Doornberg JN, van Duijn J, Ring D. Coronoid Fracture height in terrible-triad injuries. *J Hand Surg Am* 2006; 31(5):794-797. PMID: 16713844.
- <sup>45</sup> Bjorgul K, Reikeras O. Low interobserver of radiographic signs predicting healing disturbance in displaced intracapsular fracture of the

- femoral neck. *Acta Orthop Scand* 2002; 73(3):307-310. PMID: 12143978.
- <sup>46</sup> Beimers L, Kreder HJ, Berry GK, Stephen DJG, Schemitsch EH, McKee MD, Jaglal S. Subcapital hip fractures: The Garden classification should be replaced, not collapsed. *Can J Surg* 2002; 45(6):411-414. PMID: 12500914.
- <sup>47</sup> Parsons BO, Klepps SJ, Miller S, Bird J, Gladstone J, Flatow E. Reliability and reproducibility of radiographs of greater tuberosity displacement. *J Bone Joint Surg Am* 2005; 87(1):58-65. PMID: 15634814.
- <sup>48</sup> Beaulé PE, Dorey FJ, Matta JM, Letournel classification for acetabular fractures: Assessment of interobserver and intraobserver reliability. *J Bone Joint Surg Am* 2003; 85-A(9):1704-1709. PMID: 12954828.
- <sup>49</sup> Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977; 33(1):159-174. PMID: 843571.
- <sup>50</sup> Weinfeld SB, Schon LC. Hallux metatarsophalangeal arthritis. *Clin Orthop Relat Res* 1998; 349:9-19. PMID: 9584362.
- <sup>51</sup> Shurnas PS. Hallux rigidus: Etiology, biomechanics, and nonoperative treatment. *Foot Ankle Clin* 2009; 14(1):1-8. PMID: 19232987.
- <sup>52</sup> Grady JF, Axe TM, Zager EJ, Sheldon LA. A retrospective analysis of 772 patients with hallux limitus. *J Am Podiatr Med Assoc* 2002; 92(2):102-108. PMID: 11847262.
- <sup>53</sup> Polzer H, Polzer S, Brumann M, Mutschler W, Regauer M. Hallux rigidus: Joint preserving alternatives to arthrodesis - a review of the literature. *World J Orthop* 2014; 5(1):6-13. PMID: 24649409.
- <sup>54</sup> Keiserman LS, Sammarco VJ, Sammarco GJ. Surgical treatment of the hallux rigidus. *Foot Ankle Clin* 2005; 10(1):75-96. PMID: 15831259.
- <sup>55</sup> Beertema W, Draijer WF, van Os JJ, Pilot P. A retrospective analysis of surgical treatment in patients with symptomatic hallux rigidus: Long-term follow-up. *J Foot Ankle Surg* 2006; 45(4):244-251. PMID: 16818152.
- <sup>56</sup> McNeil DS, Baumhauer JF, Glazebrook MA. Evidence-based analysis of the efficacy for operative treatment of hallux rigidus. *Foot Ankle Int* 2013; 34(1):15-32. PMID: 23386758.
- <sup>57</sup> Taranow WS, Moore JR. Hallux rigidus: A treatment algorithm. *Tech Foot & Ankle* 2012; 11(2):65-73.

*Keywords:* hallus rigidus, radiography, reliability