

2017

Kinematic repeatability analyses of multi-segment foot motion in university-level ballet dancers

Sarah Carter

Edith Cowan University, University of Western Australia, sarah.carter@uwa.edu.au

Nahoko Sato

Luke Hopper

Edith Cowan University, l.hopper@ecu.edu.au

Follow this and additional works at: <http://ro.ecu.edu.au/ecuposters>



Part of the [Dance Commons](#), [Podiatry Commons](#), and the [Sports Medicine Commons](#)

Originally presented as Sarah L. Carter, Nahoko Sato & Luke S. Hopper (2017) Kinematic repeatability of a multi-segment foot model for dance. *Poster presented at realise 2017. Australasian Podiatry Conference, May 24-23, Melbourne.*

Recommended Citation

Carter, Sarah; Sato, Nahoko; and Hopper, Luke, "Kinematic repeatability analyses of multi-segment foot motion in university-level ballet dancers" (2017). *ECU Posters*.
<http://ro.ecu.edu.au/ecuposters/24>

This Book is posted at Research Online.
<http://ro.ecu.edu.au/ecuposters/24>

Kinematic repeatability analyses of multi-segment foot motion in university-level ballet dancers

Sarah L. Carter, MSc^{1,2}, Nahoko Sato, PhD³, Luke Hopper, PhD²

¹ Podiatric Medicine and Surgery, School of Allied Health, The University of Western Australia (UWA), Australia

² Western Australian Academy of Performing Arts (WAAPA), Edith Cowan University (ECU), Australia

³ Nagoya Gakuin University, Seto, Aichi, Japan



INTRODUCTION

- Dancers are susceptible to overuse foot and ankle injuries, however there is a lack of biomechanical studies examining foot mechanics to explain the causality of these injuries.
- Previous 3D multi-segment foot models (3DMFMs) applied in dance have only been assessed using a hindfoot, forefoot and hallux segment.¹⁻³
- Nester et al.⁴ found the largest degree of error found in 3DMFMs was where the navicular, 3 cuneiforms and all the metatarsals are analysed as a single segment.
- Dancers exhibit a large range of motion through this region of the foot⁵ as compared to walking⁶ and may exacerbate these errors.
- Dividing the foot into three segments; hindfoot, midfoot and forefoot may improve the repeatability of 3DMFMs for dance and would expand the current understanding of foot movement in dance, particularly regarding plantar flexion through the whole foot.




Figure 1 (A) Anterior view of the marker placement, (B) lateral view of the marker placement and (C) medial view of the marker placement

AIM

- To determine the intra and inter-assessor repeatability of a modified Rizzoli Foot Model (RFM) marker set-up⁶ (Figure 1) by analysing ballet dancers during flex-point-flex movements and static ballet positions.
- Secondary aims were to develop a reliable marker set-up that would:
1. Not impede joint movement.
 2. Minimise error due to skin movement artefact about the 1st metatarsophalangeal joint, midfoot and forefoot.

METHODS

- Modifications were made to the original RFM⁶.
- Six female university-level ballet dancers (age 18.8 ± 0.8 years) performed the following trials in randomized order, natural stance, turnout plié and stance (Figure 2), and two dynamic trials; turnout rise and flex-pointé-flex movement.
- A twelve-camera motion capture system was used to track fourteen reflective markers and one triad on the following segments: tibia, entire foot, hindfoot, midfoot, forefoot and hallux.
- A repeated-measure design was used with each participant undergoing four data collection sessions; i.e. two sessions were conducted by each researcher over two consecutive days.
- Variability of the 3D segment rotations and planar angles were determined using intra-class correlation coefficients (ICC) for the intra and inter-assessor repeatability.



Figure 2. Anterior view of a dancer in (A) parallel stance, (B) demi-plié, (C) turnout stance and (D) turnout rise.

RESULTS

- Intra and inter-assessor reliability demonstrated excellent (ICC ≥ 0.75) repeatability for the 1st metatarsophalangeal joint in the sagittal plane.
- Intra-assessor reliability demonstrated excellent (ICC ≥ 0.75) repeatability during flex-point-flex across all inter-segmental angles except for the tibia-hindfoot and hindfoot-midfoot frontal planes.
- Inter-assessor repeatability ranged from poor to excellent (0.5 > ICC ≥ 0.75) for the 3D segment rotations.
- The most repeatable measure was the tibia-foot dorsiflexion/plantar flexion articulation whereas the least repeatable measure was the hindfoot-midfoot adduction/abduction articulation.
- The variation found in the inter-assessor results is likely due to inconsistencies in marker placement.

CONCLUSIONS

- This 3D dance specific multi-segment foot model provides insight into which kinematic measures can be reliably used to ascertain in vivo technical errors and/or biomechanical abnormalities in a dancer's foot motion.

ACKNOWLEDGEMENTS

The dancers from the Advanced Diploma of Dance program at the WAAPA at ECU who participated in this study. The financial support of the Australian Postgraduate Award scholarship.

1. Fong Yan A, Hiller C, Sinclair PJ, Smith RM. Kinematic analysis of sautés in barefoot and shod conditions. *J Dance Med Sci.* 2014;18(4):149-58.
2. Fong Yan A, Smith R, Vanwanseele B, Hiller C. Mechanics of jazz shoes and their effect on pointing in child dancers. *J Appl Biomech.* 2012;28(3):242-8.
3. Lee H-H, Lin C-W, Wu H-W, Wu T-C, Lin C-F. Changes in biomechanics and muscle activation in injured ballet dancers during a jump-land task with turnout (Sissonne Fermée). *J Sports Sci.* 2012;30(7):689-97.
4. Nester CJ, Liu AM, Ward E, Howard D, Cocheba J, Derrick T. Error in the description of foot kinematics due to violation of rigid body assumptions. *J Biomech.* 2010;43(4):666-72.
5. Lin C-F, Su F-C, Wu H-W. Ankle biomechanics of ballet dancers in relevé en pointé dance. *Res Sports Med.* 2005;13(1):23-35.
6. Leardini A, Benedetti MG, Berti L, Bettinelli D, Nativo R, Giannini S. Rear-foot, mid-foot and fore-foot motion during the stance phase of gait. *Gait Posture.* 2007;25(3):453-62.

