



You have downloaded a document from  
**RE-BUŚ**  
repository of the University of Silesia in Katowice

**Title:** The Wave Nature of Continuous Gravitational Waves from Microlensing

**Author:** Kai Liao, Marek Biesiada, Xi-Long Fan

**Citation style:** Liao Kai, Biesiada Marek, Fan Xi-Long. (2019). The Wave Nature of Continuous Gravitational Waves from Microlensing. "The Astrophysical Journal" (Vol. 875 (2019), Art. No. 139), doi 10.3847/1538-4357/ab1087



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego



# The Wave Nature of Continuous Gravitational Waves from Microlensing

Kai Liao<sup>1</sup> , Marek Biesiada<sup>2,3</sup> , and Xi-Long Fan<sup>4,5</sup> <sup>1</sup> School of Science, Wuhan University of Technology, Wuhan 430070, People's Republic of China<sup>2</sup> Department of Astronomy, Beijing Normal University, Beijing 100875, People's Republic of China<sup>3</sup> Department of Astrophysics and Cosmology, Institute of Physics, University of Silesia, 75 Pułku Piechoty 1, 41-500 Chorzów, Poland<sup>4</sup> School of Physics and Technology, Wuhan University, Wuhan 430072, People's Republic of China; [fanxilong@outlook.com](mailto:fanxilong@outlook.com)<sup>5</sup> Department of Physics and Mechanical and Electrical Engineering, Hubei University of Education, Wuhan 430205, People's Republic of China*Received 2018 November 22; revised 2019 February 23; accepted 2019 March 15; published 2019 April 24*

## Abstract

Gravitational wave predicted by general relativity is the transverse wave of spatial strain. Several gravitational waveform signals from binary black holes and from a binary neutron star system accompanied by electromagnetic counterparts have been recorded by the advanced Laser Interferometer Gravitational-Wave Observatory (LIGO) and the advanced Virgo. In analogy to light, the spatial fringes of diffraction and interference should also exist as the important features of gravitational waves (GWs). We propose that observational detection of such fringes could be achieved through gravitational lensing of continuous GWs. The lenses would play the role of the diffraction barriers. Considering peculiar motions of the observer, the lens, and the source, the spatial amplitude variation of diffraction or interference fringes should be detectable as an amplitude modulation of a monochromatic gravitational signal.

*Key words:* gravitational lensing: micro – gravitational waves

## 1. Introduction

Historically, the nature of light has been described as either waves or particles (corpuscles). These different views have been fiercely debated and supported by famous experiments. For example, Young's double-slit experiment (Young 1804) and Kirchoff–Fresnel and Fraunhofer's diffraction supported the wave nature of light, while Einstein's photoelectric effect and Compton scattering suggested its corpuscular nature. Eventually, with the development of quantum mechanics, we accepted the wave–particle duality inherent to fundamental constituents of the world. On the other hand, gravitational waves (GWs), predicted by general relativity (GR), are transverse waves of spatial strain, powered by varying in time quadrupole moments of the source mass. Reasoning by analogy, one often conjectures that, like the photon, the graviton should exist in the quantum theory of gravity, as a spin two massless particle traveling at the speed of light. Recent successful detections of GW signals from astrophysical sources confirmed their classical nature. However, in order to gain better empirical understanding of the physical nature of a GW, in particular to better investigate its wave nature, it is essential to design appropriate further experiments or observation strategies. This is the main motivation behind this work.

GW signals from coalescing binary black holes (Abbott et al. 2016a, 2016b, 2017a) and a merger of binary neutron stars (Abbott et al. 2017b) have been detected by the Laser Interferometer Gravitational-Wave Observatory (LIGO) and Virgo collaboration, giving birth to GW astronomy and raising multimessenger astronomy to the new level. Theoretical chirp waveforms have been fitted to the observed signals leading to reliable estimates of physical characteristics of coalescing sources, like their chirp masses or luminosity distances.

However, one should not forget that this happened only in one particular place (at Earth), and it would be rewarding to register spatial fringes of the GW (as was done for light), further confirming its wave nature. We propose an experimental setting to achieve this goal.

Gravitational lensing is another effect predicted by GR (e.g., see a review and a history of gravitational lensing in Sauer 2008; Treu 2010), where the light traveling along null geodesics bends in the vicinity of massive bodies. Strong lensing by galaxies and galaxy clusters, weak lensing by dark matter halos, microlensing by stars, and millilensing by dark matter substructure have been widely used in astrophysics (Zackrisson & Riehm 2010) and cosmology (Bonvin et al. 2017). Similarly, a GW also travels along the null geodesic, and it should also display lensing effects (e.g., Ohanian 1974; Halder et al. 2019). If the wavelength of the GW is much shorter than the lens mass scale (like in the case of light), it behaves according to the geometric optics limit based on Fermat's principle. In such cases, gravitational lensing could affect the accuracy of physical inference made from registered signals (Oguri 2018). The event rates of GW lensing by galaxies were studied by Wang et al. (1996), Sereno et al. (2010), Biesiada et al. (2014), Ding et al. (2015), and Li et al. (2018), leading to a robust prediction that the third generation of GW interferometric detectors would yield 50–100 lensed GW events per year. Strongly lensed GW signals, observed together with their electromagnetic (EM) counterparts, have been demonstrated to enhance our understanding regarding fundamental physics (Collett & Bacon 2017; Fan et al. 2017), astrophysics (Liao et al. 2018), and cosmology (Sereno et al. 2011; Liao et al. 2017; Wei & Wu 2017; Li et al. 2019). It is noteworthy that a very recent work showed that the events GW170809 and GW170814 could come from the same source (Broadhurst et al. 2019), but see Hannuksela et al. (2019)

On the contrary, if the wavelength is much longer than the lens mass scale (i.e., its Schwarzschild radius), then one should use the wave optics limit. In the intermediate regime, wave



Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

superposition effects should be considered (Ohanian 1974; Bliokh & Minakov 1975). The lens acts like a diffraction barrier. Wave effects in gravitational lensing of GWs and their influence on the waveforms of chirp signals from inspiraling binaries were discussed in Nakamura (1998), Takahashi & Nakamura (2003), and Takahashi (2017). Fringes due to GW lensing by compact dark matter (Jung & Shin 2019), stars (Christian et al. 2018), and intermediate-mass black holes (Lai et al. 2018) were claimed to be detectable, while the waveforms of strongly lensed GWs were studied in Dai & Venumadhav (2017). These works focused on the distortion of chirping waveforms as a function of frequency. Therefore, they represent a waveform-dependent approach to observe the wave effects of GWs and manifest themselves as systematic uncertainties in the parameter estimation (Cao 2014). Remarkably, Dai et al. (2018) recently proposed an agnostic detection method of diffraction effects on lensed chirp signals based on dynamic programming, which does not require a detailed model of the lensed waveforms.

We propose a strategy to observe spatial diffraction or interference fringes directly, as the spatial variations of the GW amplitude. We find that this could be achieved through monitoring the lensed source of continuous, periodic GWs for a timescale of 0.1–10 yr due to relative peculiar motions of the strong lensing system components. Throughout this paper, we use natural units of  $c = G = 1$  in all equations; SI units are recovered for the purpose of numerical estimates.

## 2. Wave Optics Description

In order to study the wave effects of GWs due to lensing, we consider GWs propagating in the weak gravitational field of the lens (see a further discussion on strong gravitational fields in Zhang & Fan 2018). Thus, the metric can be written as:

$$g_{\mu\nu} = g_{\mu\nu}^{(L)} + h_{\mu\nu}, \quad (1)$$

where  $g_{\mu\nu}^{(L)}$  is the metric determined by the Newtonian potential  $U$  of the lens, and  $h_{\mu\nu}$  is the GW perturbation. Since the polarization tensor  $e_{\mu\nu}$  is parallel transported along the null geodesic, we consider a scalar wave:

$$h_{\mu\nu} = \phi e_{\mu\nu}, \quad (2)$$

whose propagation equation is determined by

$$\partial_\mu(\sqrt{-g^{(L)}} g^{(L)\mu\nu} \partial_\nu \phi) = 0. \quad (3)$$

In the weak field limit and switching to the frequency ( $f$ ) domain of this scalar wave,  $\tilde{\phi}(f, \mathbf{r})$  (with  $\dot{f} \ll 1$ ), one is able to rewrite the above equation in the form of the Helmholtz equation:

$$(\Delta + 4\pi^2 f^2) \tilde{\phi} = 16\pi^2 f^2 U \tilde{\phi}. \quad (4)$$

Following Takahashi & Nakamura (2003), we define the dimensionless amplification factor as

$$F(f) = \tilde{\phi}^L(f) / \tilde{\phi}(f), \quad (5)$$

where  $\tilde{\phi}^L$  and  $\tilde{\phi}$  are the lensed and unlensed ( $U = 0$ ) amplitudes, respectively.

Diffraction results are as a superposition of all possible waves on the lens plane that have different time delays corresponding to different phases. The amplification factor  $F$

( $f$ ) is given by (Takahashi & Nakamura 2003):

$$F(f) = \frac{D_s R_E^2 (1 + z_l) f}{D_l D_{ls}} \int d^2 \mathbf{x} \exp[2\pi i f t_d(\mathbf{x}, \mathbf{y})], \quad (6)$$

where  $R_E$  is the Einstein radius of the lens, and by  $\mathbf{x} = \boldsymbol{\xi}/R_E$ ,  $\mathbf{y} = \boldsymbol{\eta} D_L / (R_E D_S)$  in which we denote dimensionless positions (normalized by  $R_E$ ) with  $\boldsymbol{\xi}$  being the impact parameter in the lens plane and  $\boldsymbol{\eta}$  being the position vector of the source in the source plane. In this notation, time delay functional reads:

$$t_d(\mathbf{x}, \mathbf{y}) = \frac{D_s R_E^2 (1 + z_l)}{D_l D_{ls}} \left[ \frac{1}{2} |\mathbf{x} - \mathbf{y}|^2 - \psi(\mathbf{x}) + \phi_m(\mathbf{y}) \right], \quad (7)$$

where  $\psi(\mathbf{x})$  is dimensionless deflection potential, and  $\phi_m(\mathbf{y})$  is chosen so that the minimum arrival time is zero. Since there is a lot of compact structure in galaxies (stars, black holes, and compact dark matter clumps), we consider the lens described by a point mass, with the lensing potential of  $\psi(\mathbf{x}) = \ln x$  and  $\phi_m(\mathbf{y}) = 0.5(x_m - y)^2 - \ln x_m$

with  $x_m = 0.5(y + \sqrt{y^2 + 4})$ .

For universality, we define the dimensionless parameter  $w = 8\pi M_{Lz} f$ , which serves as a comparison between the barrier and the wavelength. At this point, we would like to be as general as possible, hence we introduce the redshifted mass of  $M_{Lz} = M(1 + z_l)$ . Of course, for nearby sources, one should neglect cosmological effects (setting  $z_l = z_s = 0$ ).

For the case in the limit of  $w \gg 1$ , Equation (6) simplifies to

$$|F(f)| = \sqrt{|\mu_+| + |\mu_-| + 2|\mu_+ \mu_-|^{1/2} \sin(2\pi f \Delta t_d)}, \quad (8)$$

where  $|\mu_+|$  and  $|\mu_-|$  are the magnifications of the brighter and fainter image in the geometrical optics limit.  $\Delta t_d$  is the time delays between multi-signals. For the point mass lens,

$$\Delta t_d = 4M_{Lz} \left[ \frac{y\sqrt{y^2 + 4}}{2} + \ln \left( \frac{\sqrt{y^2 + 4} + y}{\sqrt{y^2 + 4} - y} \right) \right] \sim 8M_{Lz} y. \quad (9)$$


The last term in Equation (8) is the interference between images. For the light sources whose multiple images are independent or the transient GW sources whose lasting times are smaller than  $\Delta t_d$ , this term vanishes. We will see two separate signals, amplified by  $\mu_{\pm}$ , arriving at different times. For the continuous GWs considered in this work, the term gives the interference modulation.

In the earlier works (Takahashi & Nakamura 2003; Jung & Shin 2019; Christian et al. 2018) mentioned above, the authors studied distortions of the waveform of a chirp signal whose frequency increases with time, assuming a fixed  $y$  since the chirping timescale is very small. In contrast, we consider the sources of continuous GW whose frequencies can be assumed to be constant for a long time.


At last, it is worth stressing that, while the fringes of light are determined by  $|F|^2$  (i.e., the intensity), in the context of GWs, the amplitude  $|F|$  is relevant.

## 3. Peculiar Motions and Detectability of the Fringes

Figure 1 shows the amplification factor as a function of the source position  $y$  expressed as a fraction of the Einstein radius



**Figure 1.** Relative amplitude of a lensed continuous gravitational wave as a function of the source position for the point mass lens. Left: diffraction in wave optics description for different values of the  $w$  parameter. Middle: interference pattern of images in the geometric optics limit of  $w = 300$ . Right: zoomed-in of the interval  $y \in [0.4, 0.5]$ . In the geometric optics limit, amplification of light from lensed images is shown in yellow for comparison.



**Figure 2.** Relation between the GW frequency and the lens mass for different values of  $w$ .

of the lens for  $w = 1, 3, 9, 300$  in the case of point mass, respectively. One can see the diffraction and interference fringes and develop the idea of their detectability.

First of all,  $w$  should not be too small, since the amplitude of diffraction fringes would not be detectable in that case. We propose the  $w > 1$  criterion is reasonable. Recalling the definition of  $w$ , one can see that for the frequencies of interest, e.g., of order of,  $f = 1$  kHz masses of the lenses satisfying  $w > (1, 3, 9, 300)$  criterion would be  $M/M_\odot > (8.1, 24.3, 72.9, 2430)$ . Figure 2 shows the dependence between  $w$ , the lens mass, and the GW frequency. Then, one can see that the amplitude pattern of fringes is damped. However, they are noticeable beyond the Einstein ring radius when  $y < 3$ . This is an interesting new feature of GW diffraction effects, which is different from the standard strong lensing/microlensing case.

At last, in order to see complete diffraction or an interference fringe, the monitoring time should be sufficiently large,  $\Delta t > t_f$ , where  $t_f$  is the fringe timescale, such that the peculiar motions would result in relative spatial change (see more details later). An appropriate timescale for the transit of fringes is  $t_f = t_E/w$ , where  $t_E$  is the Einstein radius crossing time. This was noticed in Naderi et al. (2018) who discussed detectability of primordial black holes by detecting the diffraction patterns in lensed quasars. In a similar way, Mehrabi & Rahvar (2013) studied the wave optics features of gravitational microlensing by a binary lens composed of a planet and a parent star. These papers discussed lensing of light, while Ruffa (1999) formulated an idea of the Milky Way's own supermassive black hole amplifying the continuous wave signal from an

extragalactic rotating neutron star and the resulting diffraction pattern.


We summarize the criteria of detectability of the fringes for a point mass lens:

1.  $w > 1$  to have enough amplification variation;
2.  $y < 3$  to detect fringes before they are damped;
3.  $\Delta t > t_f$  to see a fringe pattern.


Of course, with GW signals, one is not able to observe the fringe patterns on the screen. However, peculiar motions (with respect e.g., to the Cosmic Microwave Background (CMB) frame) of the components of lensing system—the observer,  $\mathbf{v}_{\text{obs}}$ ; the lens,  $\mathbf{v}_l$ ; and the source,  $\mathbf{v}_s$ —can be combined to the effective velocity of the source with respect to the lens (Kayser et al. 1986):

$$\mathbf{v}_{\text{eff}} = \mathbf{v}_s - \frac{1 + z_s}{1 + z_l} \frac{D_s}{D_l} \mathbf{v}_l + \frac{1 + z_s}{1 + z_l} \frac{D_{ls}}{D_l} \mathbf{v}_{\text{obs}}. \quad (10)$$

Using this effective relative velocity in the system,  $t_f = \frac{R_E}{v_{\text{eff}} w}$  provides the timescale of observability of diffraction fringes. As the source position changes with respect to the lens—according to  $y(t) = \sqrt{y_0^2 + \left(\frac{t-t_0}{t_E}\right)^2}$ , where  $t_0$  is the time of the closest encounter occurring at the source position  $y_0$ —the amplitude of a GW from a periodic source gets modulated. This creates opportunity to see the diffraction fringes with a favorable configuration of the source and the lens.



**Figure 3.** Amplitude variation of a monochromatic GW in a time interval corresponding to two Einstein crossing times for a  $40 M_{\odot}$  lens. The closest approach time corresponds to  $t_E$ . Diffraction patterns are shown for three frequencies of the GW:  $f = 200$  Hz, 600 Hz, and 1800 Hz. Panels from left to right correspond to three values of the source position at the closest encounter  $y_0 = 0.1, 0.5,$  and  $1$ , respectively.



**Figure 4.** Amplitude variation of a monochromatic GW in a time interval corresponding to two Einstein crossing times for a  $10^4 M_{\odot}$  lens in the geometric optics limit of  $w = 300$ . The source position at the closest encounter is  $y_0 = 0.5$ . For comparison, the yellow line shows the corresponding amplification of light from unresolved lensed images. Middle and right panels enlarge the intervals of 700 days and 100 days around the closest encounter and away from it, respectively.

Central to our idea, the source of the GW is continuous, with the frequency not changing significantly during the observation time. For LIGO and the third generation of ground-based detectors, the source of periodic GWs is expected to be isolated neutron stars emitting at frequencies of  $f \sim 10\text{--}10^3$  Hz. In the case of galactic sources and lenses, distances can be regarded as Euclidean distances, and the Einstein time reads:

$$t_E \approx 34.7 \text{ days} \sqrt{4 \frac{D_l}{D_s} \left(1 - \frac{D_l}{D_s}\right) \left(\frac{D_s}{8 \text{ kpc}}\right)^{1/2}} \times \left(\frac{M}{M_{\odot}}\right)^{1/2} \left(\frac{v_{\text{eff}}}{200 \text{ km s}^{-1}}\right)^{-1}, \quad (11)$$

which would be 220 days for a  $40 M_{\odot}$  lens. One can see from Figure 2 that, in such a case for a source emitting GWs with a frequency of  $f = 600$  Hz, the  $w$  parameter will be  $w = 3$  and  $t_f = 73.3$  days. This is advantageous because this kind of amplitude modulation can be distinguished from the modulation caused by Earth's rotation or orbital motion. In Figure 3, we show the pattern of amplitude modulation of sources emitting at frequencies of 200 Hz, 600 Hz, and 1800 Hz, lensed by a  $40 M_{\odot}$  point mass. This corresponds to the  $w$  parameter:  $w = 1, 3, 9$ , respectively. In the case of a much more massive lens like a compact dark matter clump or a black hole, one could have  $w \gg 1$  corresponding to the geometric optics limit. Yet, the amplitude of a monochromatic GW can be modulated due to interference between two image signals (see Equation (8)). For example, if the lens has a mass of  $M = 10^4 M_{\odot}$ , Einstein time would be  $t_E = 3300$  days, and

for a  $f = 240$  Hz source, the  $w$  parameter would be  $w = 300$  and  $t_f = 11$  days. We display the results in Figure 4.

#### 4. Observing Feasibility

A question arises: how probable is to observe microlensed monochromatic GWs probing the fringes (as discussed above)? In the context of ground-based detectors, the most realistic source of continuous GW radiation are spinning neutron stars (NSs). By extrapolating the NS birth rate (Narayan & Ostriker 1990) from the number of supernovae required to account for the heavy element abundances in the Milky Way (Arnett et al. 1989), one may expect that the galactic bulge contains about  $10^9$  NSs. Globular clusters are expected to contain  $10^3$  NSs (Grindlay & Bailyn 1988), which could be a target population for GW detectors. Putting aside the problem how many of them would be accessible to the third generation of detectors, as a first order estimate, one may use the classical approach to the microlensing probability used in optical studies. In this section, we reintroduce the dimensioned fundamental constants  $G$  and  $c$  in the formulae.

The optical depth is the probability that a given source falls into the Einstein radius of any lensing star along the line of sight:

$$\tau = \int_0^{D_s} n(D_l) \pi R_E^2 dD_l. \quad (12)$$

Let us recall, at this point, that unlike in the optical microlensing, wave effects discussed by us can be observed up to the source position of  $y_{\text{max}} = 3$  Einstein radii. Therefore, optical depth can be boosted by a factor of  $y_{\text{max}}^2$ . Of course, the optical depth for lensing is scenario-dependent. As a first

scenario, let us consider sources located in the bulge, lensed by stellar mass lenses located in the Galactic disk. An order of magnitude estimate is given in the original papers (Paczynski 1986, 1991). Under assumption of a constant density of lenses along the line of sight, one has:

$$\tau = \frac{1}{2c^2} \frac{GM(<D_s)}{D_s} = \frac{v_{\text{rot}}^2}{2c^2}. \quad (13)$$

With the Milky Way's bulge as a target,  $v_{\text{rot}} \approx 220 \text{ km s}^{-1}$  and  $\tau \approx 2.7 \times 10^{-7}$ . In our case, this should be modified to

$$\tau_{\text{GW}} \sim f_l y_{\text{max}}^2 \times 10^{-7} \sim f_l \times 10^{-6}, \quad (14)$$

where  $f_l$  is the fraction of lens mass larger than a solar mass since a smaller mass will enter the wave optics limit. Of course, more realistic estimates have to consider proper mass distribution of potential lenses (e.g., exponential) and proper models of the stellar systems considered (e.g., double exponential disk, spheroidal bulge, Plummer sphere model of globular clusters, etc.). Probability of GW microlensing events expected in time interval  $\Delta t$  can be assessed as

$$P \sim \tau_{\text{GW}} \frac{\Delta t}{t_f}, \quad (15)$$

where the timescale of GW wave effects is  $t_f = t_E/w$ . Considering the Einstein Telescope (ET) operation time as  $\Delta t = 10 \text{ yr}$ , one has  $P \sim f_l (t_f/1 \text{ month})^{-1} \times 10^{-4}$ . This means that if only a fraction of  $f_s = 10^{-5}$  of the source population of  $n = 10^9$ -bulge NSs would be accessible to the ET, such GW microlensing events are expected to occur. Let us emphasize that, unlike in the optical, one does not face the problem of source resolution in the crowded field or light blending.

Another even more promising scenario is associated with lenses located in rich globular clusters like M22 seen against the Galactic bulge. A typical velocity dispersion in a globular cluster is  $\sigma \sim 10 \text{ km s}^{-1}$ , so the typical transverse velocity of the lens is well approximated by that of a cluster as a whole  $\sim 200 \text{ km s}^{-1}$ . As shown by Paczyński (1994), the optical depth for microlensing in such cases is given by:

$$\tau = \frac{\sigma^2}{c^2} \frac{2\pi}{\phi} \left(1 - \frac{D_l}{D_s}\right) \approx 2.4 \cdot 10^{-5} \left(\frac{\sigma}{10 \text{ km s}^{-1}}\right)^2 \left(\frac{1'}{\phi}\right), \quad (16)$$

where the last equality holds when  $D_l/D_s \ll 1$ , and  $\phi$  denotes the angular distance from the cluster center. In the optical band, this implies that by looking at the target field in the bulge within a few arcminutes of the cluster center, the lensing is dominated by objects associated with the cluster itself. This observation triggered much interest since knowledge of the distance and transversal velocity of lenses belonging to the cluster may result with accurate estimates of their masses. First confirmed microlensing of the bulge star at  $D_s = 8.2 \text{ kpc}$  by a low-mass object in the globular cluster M22 located at  $D_l = 2.6 \text{ kpc}$  has been reported in Sahu et al. (2001) and Pietrukowicz et al. (2012). In the context of our considerations, this scenario predicts the optical depth of

$$\tau_{\text{GW}} \sim f_l y_{\text{max}}^2 (1/\phi) \times 10^{-5} \sim f_l (1/\phi) \times 10^{-4}, \quad (17)$$

where,  $\phi$  is expressed in arcminutes. Probability of GW microlensing events in such a scenario is  $P \sim f_l (1/\phi)(\Delta t/10 \text{ yr})(t_{\text{GW}}/1 \text{ month})^{-1} \times 10^{-2}$ . This is particularly promising scenario since the GW signal from the bulge NS can pass through the very center of the globular cluster with very high probability of being lensed. Moreover, it can be lensed by massive objects expected to reside in the center of the cluster, including the intermediate-mass black holes ( $M \sim 10^2\text{--}10^4 M_\odot$ ).

At last, in Safonova & Stalin (2010), the authors considered microlensing scenario where both the source and the lens belong to the globular cluster. They estimated the optical depth for such events as  $\tau \approx 10^{-3}$ . Considering that globular clusters anchor about  $n = 10^3$  NSs and are much closer than the Galactic bulge, this is also a promising scenario from our perspective.

## 5. Conclusion and Discussions

We proposed an observational strategy to test the wave nature of GWs by monitoring the amplitude modulation of lensed sources of continuous gravitational signals. One of the candidates of continuous GW signals is a spinning neutron star that is slightly deformed from perfect spherical symmetry, which is the target of ground-based detectors. Their detection is fairly difficult, since the signal is strongly modulated by the Earth's rotation and orbital motion. Moreover, this modulation is different for every sky position. Diffraction and interference fringes caused by intervening mass acting as a lens and moving with respect to the source also produce modulation. Luckily, the timescales of these effects are significantly different from timescales involved with the motion of the detector. In the case of monochromatic GWs search, strategies are different from chirping signals of coalescing binaries. Depending on what a priori is known about sources of GWs, searches can be targeted (the known position and GW frequency, e.g., the Crab pulsar), directed (only the sky position known), and can be all-sky (or blind) searches. In the setting discussed by us, only blind searches or directed searches (in case of globular cluster lensing) can be applied. The GW signal coming from a rotating NS is so weak that, in order to detect it in the detector's noise, one has to analyze months-long segments of data. A fully coherent analysis of such amount of data is computationally prohibitive in the case of all-sky searches (Brady et al. 1998; Brady & Creighton 2000). Therefore, different hierarchical two-stage schemes were developed, where in the first stage, shorter segments of data are analyzed coherently, and then in the second stage, the results are combined in an incoherent way. Implementation of the F-statistic (Jaranowski et al. 1998) consists of coherent searches over two-day periods, and is followed by a search for coincidences among the candidates from the two-day segments. The timescales for fringe modulation discussed here are larger, and one can expect that respective amplitude modulation could be detectable with current techniques.

In this work, we suggested plausible configurations comprised of bulge NSs lensed by massive objects in the disk, or lensed by massive objects in the globular clusters seen against the bulge, or globular cluster NSs lensed by massive stars in the same cluster. More detailed elaboration of these scenarios is for a further study. This is worth pursuing since the verification of spatial diffraction fringes of lensed continuous gravitational

sources would strengthen our understanding of GWs and contribute to setting up the quantum gravity and shed light on the nature of the graviton. We are looking forward to seeing this phenomenon observed by second-generation detectors like the advanced LIGO and Virgo and by the third-generation detectors like the Einstein Telescope or space-born detectors like the Laser Interferometer Space Antenna (LISA).

K.L. was supported by the National Natural Science Foundation of China (NSFC) No. 11603015 and the Fundamental Research Funds for the Central Universities (WUT:2018IB012). M.B. was supported by the Key Foreign Expert Program for the Central Universities No. X2018002. X. F. was supported by the NSFC 11673008 and Newton International Fellowship Alumni Follow on Funding.

### ORCID iDs

Kai Liao  <https://orcid.org/0000-0002-4359-5994>  
 Marek Biesiada  <https://orcid.org/0000-0003-1308-7304>  
 Xi-Long Fan  <https://orcid.org/0000-0002-8174-0128>

### References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016a, *PhRvL*, **116**, 061102  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016b, *PhRvL*, **116**, 241103  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017a, *PhRvL*, **118**, 221101  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017b, *PhRvL*, **119**, 161101  
 Arnett, W. D., Schramm, D. N., & Truran, J. W. 1989, *ApJL*, **339**, L25  
 Biesiada, M., Ding, X., Piórkowska, A., & Zhu, Z.-H. 2014, *JCAP*, **10**, 080  
 Bliokh, P. V., & Minakov, A. A. 1975, *Ap&SS*, **34**, L7  
 Bonvin, V., Courbin, F., Suyu, S. H., et al. 2017, *MNRAS*, **465**, 4914  
 Brady, P. R., & Creighton, T. 2000, *PhRvD*, **61**, 082001  
 Brady, P. R., Creighton, T., Cutler, C., & Schutz, B. F. 1998, *PhRvD*, **57**, 2101  
 Broadhurst, T., Diego, J. M., & Smoot, G. F., III 2019, arXiv:1901.03190  
 Cao, Z. 2014, *PhRvD*, **90**, 062003  
 Christian, P., Vitale, S., & Loeb, A. 2018, *PhRvD*, **98**, 103022  
 Collett, T. E., & Bacon, D. 2017, *PhRvL*, **118**, 091101  
 Dai, L., Li, S.-S., Zackay, B., Mao, S., & Lu, Y. 2018, *PhRvD*, **98**, 104029  
 Dai, L., & Venumadhav, T. 2017, arXiv:1702.04724  
 Ding, X., Biesiada, M., & Zhu, Z.-H. 2015, *JCAP*, **12**, 006  
 Fan, X.-L., Liao, K., Biesiada, M., et al. 2017, *PhRvL*, **118**, 091102  
 Grindlay, J. E., & Bailyn, C. D. 1988, *Natur*, **336**, 48  
 Halder, A., Banerjee, S., & Majumdar, D. 2019, arXiv:1902.06903v1  
 Hannuksela, O. A., Haris, K., Ng, K. K. Y., et al. 2019, *ApJL*, **874**, L2  
 Jaranowski, P., Królak, A., & Schutz, B. F. 1998, *PhRvD*, **58**, 063001  
 Jung, S., & Shin, C. S. 2019, *PhRvL*, **122**, 041103  
 Kayser, R., Refsdal, S., & Stabell, R. 1986, *A&A*, **166**, 36  
 Lai, K.-H., Hannuksela, O. A., Herrera-Martin, A., et al. 2018, *PRD*, **98**, 083005  
 Li, S.-S., Mao, S., Zhao, Y., & Lu, Y. 2018, *MNRAS*, **476**, 2220  
 Li, Y., Fan, X., & Gou, L. 2019, *ApJ*, **873**, 37  
 Liao, K., Ding, X., Biesiada, M., Fan, X.-L., & Zhu, Z.-H. 2018, *ApJ*, **867**, 69  
 Liao, K., Fan, X.-L., Ding, X., Biesiada, M., & Zhu, Z.-H. 2017, *NatCo*, **8**, 1148  
 Mehrabi, A., & Rahvar, S. 2013, *MNRAS*, **431**, 1264  
 Naderi, T., Mehrabi, A., & Rahvar, S. 2018, *PhRvD*, **97**, 103507  
 Nakamura, T. T. 1998, *PhRvL*, **80**, 1138  
 Narayan, R., & Ostriker, J. O. 1990, *ApJ*, **352**, 222  
 Oguri, M. 2018, *MNRAS*, **480**, 3842  
 Ohanian, H. C. 1974, *IJTP*, **9**, 425  
 Paczyński, B. 1986, *ApJ*, **304**, 1  
 Paczyński, B. 1991, *ApJL*, **371**, L63  
 Paczyński, B. 1994, *AcA*, **44**, 235  
 Pietrukowicz, P., Minniti, D., Jetzer, Ph., Alonso-García, J., & Udalski, A. 2012, *ApJL*, **744**, L18  
 Ruffa, A. A. 1999, *ApJL*, **517**, L31  
 Safonova, M., & Stalin, C. S. 2010, *NewA*, **15**, 450  
 Sahu, K. C., Casertano, S., Livio, M., et al. 2001, *Natur*, **411**, 1022  
 Sauer, T. 2008, *AHES*, **62**, 1  
 Sereno, M., Bleuler, A., Jetzer, P., et al. 2010, *PhRvL*, **105**, 251101  
 Sereno, M., Jetzer, P., Sesana, A., & Volonteri, M. 2011, *MNRAS*, **415**, 2773  
 Takahashi, R. 2017, *ApJ*, **835**, 103  
 Takahashi, R., & Nakamura, T. 2003, *ApJ*, **595**, 1039  
 Treu, T. 2010, *ARA&A*, **48**, 87  
 Wang, Y., Stebbins, A., & Turner, E. L. 1996, *PhRvL*, **77**, 2875  
 Wei, J.-J., & Wu, X.-F. 2017, *MNRAS*, **472**, 2906  
 Young, T. 1804, *Philos. Trans. R. Soc. London*, **94**, 18040001  
 Zackrisson, E., & Riehm, T. 2010, *AdAst*, **2010**, 478910  
 Zhang, H., & Fan, X. 2018, arXiv:1809.06511