

The essential spectrum of Toeplitz operators on the unit ball

Article

Accepted Version

Hagger, R. (2017) The essential spectrum of Toeplitz operators on the unit ball. *Integral Equations and Operator Theory*, 89 (4). pp. 519-556. ISSN 1420-8989 doi: <https://doi.org/10.1007/s00020-017-2399-1> Available at <http://centaur.reading.ac.uk/84023/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s00020-017-2399-1>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



The Essential Spectrum of Toeplitz Operators on the Unit Ball

Raffael Hagger

July 31, 2018

Abstract

In this paper we study the Fredholm properties of Toeplitz operators acting on weighted Bergman spaces $A_p^p(\mathbb{B}^n)$, where $p \in (1, \infty)$ and $\mathbb{B}^n \subset \mathbb{C}^n$ denotes the n -dimensional open unit ball. Let f be a continuous function on the Euclidean closure of \mathbb{B}^n . It is well-known that then the corresponding Toeplitz operator T_f is Fredholm if and only if f has no zeros on the boundary $\partial\mathbb{B}^n$. As a consequence, the essential spectrum of T_f is given by the boundary values of f . We extend this result to all operators in the algebra generated by Toeplitz operators with bounded symbol (in a sense to be made precise down below). The main ideas are based on the work of Suárez et al. ([17, 24]) and limit operator techniques coming from similar problems on the sequence space $\ell^p(\mathbb{Z})$ ([13, 15, 19] and references therein).

AMS subject classification: Primary: 47B35; Secondary: 32A36, 47A53, 47A10

Keywords: Toeplitz operators, Bergman space, essential spectrum, limit operators

1 Introduction

Consider some measure space (X, μ) and a corresponding L^p -space for some $p \in (1, \infty)$, say. Further assume that there is a bounded projection P onto a closed subspace \mathcal{S} of $L^p(X, \mu)$. If we now decompose a multiplication operator parallel to this projection, we obtain a Toeplitz operator. More precisely, if $f: X \rightarrow \mathbb{C}$ is an essentially bounded function and M_f the corresponding multiplication operator on $L^p(X, \mu)$, the corresponding Toeplitz operator is given by $T_f := PM_f|_{\mathcal{S}}$. Toeplitz operators are one of the prime examples for non-normal operators and are thus extensively studied on various different domains, the most prominent example probably being the Hardy space over the circle. In our case here we are going to consider Toeplitz operators on the space of holomorphic L^p -functions defined on the complex open unit ball \mathbb{B}^n , which is called a Bergman space. Using the variables above, we consider $X = \mathbb{B}^n$ with a weighted Lebesgue measure dv_ν for a weight parameter ν and we take \mathcal{S} to be the closed subspace of holomorphic functions contained in $L_\nu^p := L^p(\mathbb{B}^n, dv_\nu)$, here denoted by A_p^p (see Section 2 for more details).

In this paper we are particularly interested in the Fredholm properties of Toeplitz operators. Recall that in the Hardy space case, a Toeplitz operator T_f with a continuous symbol f is Fredholm if and only if f does not have any zeros. A similar result holds for the Bergman space: A Toeplitz operator T_f with a symbol that can be continuously extended to the (Euclidean) boundary $\partial\mathbb{B}^n$ is Fredholm if and only if f has no zeros on the boundary. This result was first established by Coburn in [9] and then generalized in many different directions by several authors (e.g. [1, 2, 6, 8, 16, 17,

23, 24, 28, 29]). One of the latest improvements ([24, Theorem 10.3], [17, Theorem 5.8]) include the following result: Let $\mathfrak{T}_{2,\nu} \subset \mathcal{L}(A_\nu^2)$ denote the closed subalgebra generated by Toeplitz operators T_f with $f \in L^\infty(\mathbb{B}^n)$. Then $A \in \mathfrak{T}_{2,\nu}$ is Fredholm if and only if all of its limit operators are invertible and their inverses are uniformly bounded. Roughly speaking, limit operators are operators that appear when we shift our operator A to the boundary of the domain (a more accurate definition is given in Section 5). This theorem reminds of a seemingly unrelated result in the Fredholm theory of sequence spaces ℓ^p . There, until a few years ago, one of the main theorems was stated as follows: A band-dominated¹ operator A is Fredholm if and only if all of its limit operators are invertible and their inverses are uniformly bounded (see e.g. [19]). There are a few problems with this characterization. Not only is the uniform boundedness condition difficult to work with, it also prevents us from writing the essential spectrum as the union of spectra of limit operators. As a consequence, many different authors worked out particular examples (see e.g. [14, Chapter 3] for a summary) where the uniform boundedness condition could be dropped. Moreover, as there was no known example where the uniform boundedness condition was actually violated, it was conjectured that this condition was actually redundant. And indeed, this was shown in [15] a few years ago. Now the goal of this paper is to show that the same is the case for Toeplitz operators on the Bergman space:

Theorem A. *Let $\mathfrak{T}_{p,\nu} \subset \mathcal{L}(A_\nu^p)$ denote the closed subalgebra generated by Toeplitz operators with bounded symbol and $A \in \mathfrak{T}_{p,\nu}$. Then the following are equivalent:*

- (i) A is Fredholm,
- (ii) A_x is invertible for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$ and $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x^{-1}\| < \infty$,
- (iii) A_x is invertible for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$.

Here, the A_x denote the limit operators of A and they are indexed over the boundary of a certain compactification \mathfrak{M} of \mathbb{B}^n . In particular, we extend [24, Theorem 10.3] and [17, Theorem 5.8] to the Banach space case $p \neq 2$ and show that the uniform boundedness condition is redundant just like it is in the sequence space case. As a consequence, we get

$$\mathrm{sp}_{\mathrm{ess}}(A) = \bigcup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \mathrm{sp}(A_x)$$

for all operators $A \in \mathfrak{T}_{p,\nu}$.

The paper is organized as follows. In Section 2 we introduce all the necessary notation and some preliminary results. Then we proceed by introducing what, in analogy to the sequence space case, we will call band-dominated operators and show some basic properties in Section 3. In particular, we show that Toeplitz operators are band-dominated. In Section 4 we show a Fredholm criterion for band-dominated operators that will be crucial for the proof of Theorem A. In Section 5 we introduce limit operators and finally show our main theorem. After that, we proceed by showing that a similar result holds for the essential norm of an operator $A \in \mathfrak{T}_{p,\nu}$ in Section 6. However, our result is less complete in this case (compare with the corresponding result on ℓ^p : [13, Theorem 3.2]) and leaves some questions open. Section 7 is devoted to some applications of Theorem A.

Note that similar results are expected to hold for more general domains. Sections 2 to 4 are in fact valid word by word for any bounded symmetric domain $\Omega \subset \mathbb{C}^n$. For future reference we

¹a certain property related to the structure of the corresponding infinite matrix

therefore chose to provide full generality in these sections. However, in Sections 5 and 6 there are some open problems in the most general case, which is part of the reason why we restrict ourselves to $\Omega = \mathbb{B}^n$ in this paper. A more general setting will be the topic of future work. The reader who is only interested in the unit ball (which is quite frankly the topic of this paper) may replace any Ω by \mathbb{B}^n , ignore the specific notions for bounded symmetric domains and use the respective (more explicit) formulas.

2 Notation and preliminary results

Let $\Omega \subset \mathbb{C}^n$ be an irreducible bounded symmetric domain of genus g and type (r, a, b) in its Harish-Chandra realization with corresponding Bergman metric β and Jordan triple determinant h . As we only need a handful of properties of bounded symmetric domains, we do not provide an introduction to these notions here. Instead, we refer to [10, 26] for introductions and just mention the properties we actually need. For $p \in (1, \infty)$ and $\nu > -1$ denote by L_ν^p the usual Lebesgue space of p -integrable functions on (Ω, dv_ν) , where

$$dv_\nu(z) = c_\nu h(z, z)^\nu dv(z),$$

dv is the usual Lebesgue measure restricted to Ω and c_ν is a normalizing constant chosen such that $dv_\nu(\Omega) = 1$. The (unique) geodesic symmetry interchanging 0 and z is denoted by ϕ_z . In particular, these symmetries ϕ_z satisfy

$$\beta(\phi_z(x), \phi_z(y)) = \beta(x, y)$$

for all $x, y \in \Omega$. Moreover, h and dv_ν transform under ϕ_z as follows:

$$\begin{aligned} h(\phi_z(x), \phi_z(y)) &= \frac{h(z, z)h(x, y)}{h(x, z)h(z, y)} \\ dv_\nu(\phi_z(w)) &= \frac{h(z, z)^{\nu+g}}{|h(w, z)|^{2(\nu+g)}} dv_\nu(w) \end{aligned}$$

(see [10] for details).

Note that for the unit ball $\Omega = \mathbb{B}^n$ we have $(r, a, b) = (1, 2, n-1)$ and $g = n+1$. Moreover, the Bergman metric is the usual hyperbolic metric on the unit ball and the Jordan triple determinant is simply given by $h(z, w) = 1 - \langle z, w \rangle$ in this case. In case $n = 1$, ϕ_z is given explicitly by the Möbius transform $w \mapsto \frac{z-w}{1-w\bar{z}}$. We refer to [30] for an explicit description of ϕ_z in higher dimensions.

The (closed) subspace of holomorphic functions contained in L_ν^p is denoted by A_ν^p and called a weighted Bergman space. The set of bounded linear operators between Banach spaces X and Y is denoted by $\mathcal{L}(X, Y)$ and we abbreviate $\mathcal{L}(X) := \mathcal{L}(X, X)$. The set of compact operators in $\mathcal{L}(X)$ will be denoted by $\mathcal{K}(X)$. $A \in \mathcal{L}(X)$ is called Fredholm if it is invertible modulo $\mathcal{K}(X)$, i.e. if there exists $B \in \mathcal{L}(X)$ such that both $AB - I$ and $BA - I$ are compact. Equivalently, A is Fredholm if and only if $\ker A$ and $\text{coker } A$ are both finite-dimensional (Atkinson's theorem). The essential spectrum of an operator A will be denoted by $\text{sp}_{\text{ess}}(A)$ and is given by

$$\text{sp}_{\text{ess}}(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not Fredholm}\}.$$

We will say that a net (sequence, series, etc.) of operators converges $*$ -strongly if the net converges strongly and the net of adjoints converges strongly to the adjoint. The characteristic function of a set M will be denoted by χ_M .

Let $P_\nu \in \mathcal{L}(L_\nu^2)$ be the orthogonal projection onto A_ν^2 , called the Bergman projection. One can show (see e.g. [10] or [26]) that P_ν is given by

$$(P_\nu f)(z) = \int_\Omega f(w)h(z, w)^{-\nu-g} dv_\nu(w).$$

Using the same formula also for $p \neq 2$, we can define for every $f \in L^\infty(\Omega)$ the corresponding Toeplitz operator $T_f := P_\nu M_f|_{A_\nu^p}$, where $(M_f)g = f \cdot g$ for all $g \in L_\nu^p$. T_f then defines a bounded linear operator on A_ν^p with $\|T_f\| \leq \|P_\nu\| \|f\|_\infty$, provided that P_ν is indeed a bounded linear operator on L_ν^p . The function f is called the symbol of T_f and M_f , respectively. The algebra generated by all Toeplitz operators acting on A_ν^p will be denoted by $\mathfrak{T}_{p,\nu}$.

The next proposition provides a sufficient condition for P_ν to be bounded. Note that this is certainly not optimal if $r > 1$ as the case $\alpha = \nu$, $p = 2$ demonstrates. For a more optimal condition in the case $\alpha = \nu$ we refer to [10, Lemma 9] (which is a special case of [4, Theorem II.7]).

Proposition 1. *Let $p(\alpha + 1) > \nu + 1 + \frac{(r-1)a}{2} > p\frac{(r-1)a}{2}$. Then $P_\alpha \in \mathcal{L}(L_\nu^p)$ and $P_\alpha f = f$ for all $f \in A_\nu^p$. In particular, P_α is a bounded projection onto $\text{im}(P_\alpha) = A_\nu^p$.*

Proof. By definition,

$$|(P_\alpha f)(z)| \leq \int_\Omega |h(z, w)|^{-\alpha-g} |f(w)| dv_\alpha(w) = \frac{c_\alpha}{c_\nu} \int_\Omega |h(z, w)|^{-\alpha-g} h(w, w)^{\alpha-\nu} |f(w)| dv_\nu(w).$$

We want to show that the integral operator with kernel $R(z, w) = |h(z, w)|^{-\alpha-g} h(w, w)^{\alpha-\nu}$ is bounded on L_ν^p . To do this, we apply the Schur test with the test function $h(z) := h(z, z)^s$, where $s \in \mathbb{R}$ is to be determined later. We thus need to show that there exists a constant C such that

$$\begin{aligned} \int_\Omega |h(z, w)|^{-\alpha-g} h(w, w)^{\alpha-\nu} h(w, w)^{sq} dv_\nu(w) &= c_\nu \int_\Omega |h(z, w)|^{-\alpha-g} h(w, w)^{sq+\alpha} dv(w) \\ &\leq Ch(z, z)^{sq}, \\ \int_\Omega |h(z, w)|^{-\alpha-g} h(w, w)^{\alpha-\nu} h(z, z)^{sp} dv_\nu(z) &= c_\nu h(w, w)^{\alpha-\nu} \int_\Omega |h(z, w)|^{-\alpha-g} h(z, z)^{sp+\nu} dv(z) \\ &\leq Ch(w, w)^{sp}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$. By [12, Theorem 4.1], the first inequality holds for $s \in (-\frac{\alpha+1}{q}, -\frac{(r-1)a}{2q})$, whereas the second one holds for $s \in (-\frac{\nu+1}{p}, -\frac{\nu}{p} + \frac{\alpha}{p} - \frac{(r-1)a}{2p})$. A simple computation shows that $(-\frac{\alpha+1}{q}, -\frac{(r-1)a}{2q}) \cap (-\frac{\nu+1}{p}, -\frac{\nu}{p} + \frac{\alpha}{p} - \frac{(r-1)a}{2p})$ is non-empty if the inequalities stated in the proposition are assumed (cf. [30, Theorem 2.11] in the case $\Omega = \mathbb{B}^n$). Thus $P_\alpha \in \mathcal{L}(L_\nu^p)$.

As $P_\alpha: L_\alpha^2 \rightarrow A_\alpha^2$ is the orthogonal projection, we get $P_\alpha f = f$ for $f \in A_\alpha^2 \cap A_\nu^p$. As $A_\alpha^2 \cap A_\nu^p$ is dense in A_ν^p (polynomials are dense), this generalizes to all of A_ν^p . Similarly, $P_\alpha f \in A_\nu^p$ for $f \in L_\alpha^2 \cap L_\nu^p$ generalizes to all of L_ν^p . Therefore $P_\alpha: L_\nu^p \rightarrow A_\nu^p$ is a bounded projection as well. \square

For the unit ball \mathbb{B}^n the condition on ν and α simplifies significantly because we have $r = 1$ in this case.

Corollary 2. *Let $\Omega = \mathbb{B}^n$ and $p(\alpha + 1) > \nu + 1 > 0$. Then $P_\alpha \in \mathcal{L}(L_\nu^p)$ and $P_\alpha f = f$ for all $f \in A_\nu^p$. In particular, P_α is a bounded projection onto $\text{im}(P_\alpha) = A_\nu^p$.*

Definition 3. We will call $(\alpha, \nu, p) \in \mathbb{R}^2 \times (1, \infty)$ an admissible triple (for Ω) if the inequalities in Proposition 1 are satisfied.

Note that by Corollary 2, (ν, ν, p) is always admissible if $\Omega = \mathbb{B}^n$. Therefore the Toeplitz algebra $\mathfrak{T}_{p, \nu}$ can be defined for all $\nu > -1$ and $p \in (1, \infty)$ in this case. For a more general bounded symmetric domain Ω we will always assume that ν and p are chosen in such a way that (ν, ν, p) is admissible. Clearly, this assumption also implies that (α, ν, p) will always be admissible provided that $\alpha \geq \nu$. This observation will be crucial later on.

We will also need the following two simple propositions that will be used several times later on. The first one is basically a sloppy version of Jensen's inequality, but sufficient for our purposes.

Proposition 4. Let $n \in \mathbb{N}$, $p \in (1, \infty)$ and $x_1, \dots, x_n \geq 0$. Then

$$\left(\sum_{k=1}^n x_k \right)^p \leq n^p \sum_{k=1}^n x_k^p.$$

Proof. Obviously,

$$\left(\sum_{k=1}^n x_k \right)^p \leq \left(n \max_{k=1, \dots, n} x_k \right)^p = n^p \max_{k=1, \dots, n} x_k^p \leq n^p \sum_{k=1}^n x_k^p.$$

□

Proposition 5. Let $(U_k)_{k \in \mathbb{N}}$ be a sequence of measurable sets in Ω such that every $z \in \Omega$ belongs to at most N of the sets U_k and let $f \in L^p_\nu$. Then

$$\sum_{k=1}^{\infty} \int_{U_k} |f(z)|^p \, dv_\nu(z) \leq N \|f\|^p.$$

Proof. For every $k \in \mathbb{N}$ there is a disjoint decomposition $U_k = A_k^1 \cup \dots \cup A_k^N$ such that the sets $(A_k^i)_{k \in \mathbb{N}}$ are again measurable and pairwise disjoint for every $i \in \{1, \dots, N\}$ (see [24, p. 2195] for details). Thus

$$\sum_{k=1}^{\infty} \int_{U_k} |f(z)|^p \, dv_\nu(z) = \sum_{i=1}^N \sum_{k=1}^{\infty} \int_{A_k^i} |f(z)|^p \, dv_\nu(z) \leq \sum_{i=1}^N \int_{\Omega} |f(z)|^p \, dv_\nu(z) = N \|f\|^p. \quad \square$$

3 Band-dominated operators

In this section we introduce the notion of band-dominated operators. The name is chosen in analogy to the sequence space case $\ell^p(\mathbb{Z})$, where band-dominated operators are in fact norm limits of infinite band matrices, see e.g. [13, 14, 15, 19, 20, 21].

Definition 6. An operator $A \in \mathcal{L}(L^p_\nu)$ is called a band operator if there exists a positive real number ω such that $M_f A M_g = 0$ for all $f, g \in L^\infty(\Omega)$ with $\text{dist}_\beta(\text{supp } f, \text{supp } g) > \omega$. The number

$$\inf \{ \omega \in \mathbb{R} : M_f A M_g = 0 \text{ for all } f, g \in L^\infty(\Omega) \text{ with } \text{dist}_\beta(\text{supp } f, \text{supp } g) > \omega \}$$

is called the band width of A . An operator $A \in \mathcal{L}(L^p_\nu)$ is called band-dominated if it is the norm limit of band operators. The set of band-dominated operators will be denoted by BDO^p_ν .

The definition of band-dominated operators can be extended to operators acting on the Bergman space A_ν^p . If (α, ν, p) is admissible and $Q_\alpha := I - P_\alpha$, we can consider the natural extension $\hat{A} := AP_\alpha + Q_\alpha$ of $A \in \mathcal{L}(A_\nu^p)$. An operator acting on A_ν^p is then called band-dominated if its extension is band-dominated. As will be immediate, this definition does not depend on the chosen extension. In this language the main result of this section reads as follows: $\hat{A} \in \text{BDO}_\nu^p$ for all $A \in \mathfrak{T}_{p,\nu}$, i.e. Toeplitz operators are band-dominated.

Theorem 7. *Let (α, ν, p) be an admissible triple and $A \in \mathfrak{T}_{p,\nu}$. Then $\hat{A} \in \text{BDO}_\nu^p$.*

Before we proceed with the proof of this theorem, we show some equivalent characterizations of band-dominated operators that will prove useful later on. For this we need some cut-off functions to decompose our domain Ω . Let us state the following auxiliary lemma, which is due to Carlsson and Goldfarb [7] (see [5, Theorem 91] for a more explicit version). For the unit ball Suárez constructed an explicit cover in [24, Lemma 3.1].

Lemma 8. *There is a (smallest possible) positive integer N (depending only on the bounded symmetric domain Ω) such that for any $\sigma > 0$ there is a cover of Ω by Borel sets $(B_j)_{j \in \mathbb{N}}$ satisfying*

- (i) *the sets B_j are pairwise disjoint,*
- (ii) *every point of Ω belongs to at most N of the sets $\{z \in \Omega : \text{dist}_\beta(z, B_j) \leq \sigma\}$,*
- (iii) *there is a constant $C(\sigma) > 0$ such that $\text{diam}_\beta(B_j) \leq C(\sigma)$ for every $j \in \mathbb{N}$.*

For every $t \in (0, 1)$ let $(B_{j,t})_{j \in \mathbb{N}}$ be a cover of Ω that satisfies (i) to (iii) in Lemma 8 in the case $\sigma = \frac{1}{t}$ and define

$$\Xi_{j,t,k} := \left\{ z \in \Omega : \text{dist}_\beta(z, B_{j,t}) \leq \frac{k}{3t} \right\}$$

for $j \in \mathbb{N}$, $t \in (0, 1)$ and $k = 1, 2, 3$. We now construct families of uniformly Lipschitz continuous functions (partitions of unity) according to these decompositions. Let $f_{j,t} : \Omega \rightarrow [0, 1]$ be defined by

$$f_{j,t}(z) := \frac{\text{dist}_\beta(z, \Omega \setminus \Xi_{j,t,1})}{\text{dist}_\beta(z, B_{j,t}) + \text{dist}_\beta(z, \Omega \setminus \Xi_{j,t,1})}.$$

Clearly, $\text{supp } f_{j,t} = \Xi_{j,t,1}$ and $f_{j,t}(z) = 1$ for $z \in B_{j,t}$. Moreover, it is easy to see that

$$|f_{j,t}(z) - f_{j,t}(w)| \leq \frac{\beta(z, w)}{\text{dist}_\beta(B_{j,t}, \Omega \setminus \Xi_{j,t,1})} = 3t\beta(z, w).$$

Define $g_t : \Omega \rightarrow \mathbb{R}$ by $g_t(z) := \sum_{j=1}^{\infty} f_{j,t}(z)$ and $\varphi_{j,t} : \Omega \rightarrow [0, 1]$ by $\varphi_{j,t} := \frac{f_{j,t}}{g_t}$. The functions $\varphi_{j,t}$ then satisfy the following properties

- (i) $\sum_{j=1}^{\infty} \varphi_{j,t}(z) = 1$ for all $z \in \Omega$,
- (ii) $\text{supp } \varphi_{j,t} = \Xi_{j,t,1}$ for all $j \in \mathbb{N}$, $t \in (0, 1)$,
- (iii) $|\varphi_{j,t}(z) - \varphi_{j,t}(w)| \leq 6Nt\beta(z, w)$ for all $w, z \in \Omega$, $j \in \mathbb{N}$ and $t \in (0, 1)$.

Similarly, we can define functions $\psi_{j,t} : \Omega \rightarrow [0, 1]$ with the following properties:

- (i) $\psi_{j,t}(z) = 1$ for all $z \in \Xi_{j,t,2}$, $j \in \mathbb{N}$ and $t \in (0, 1)$,
- (ii) $\text{supp } \psi_{j,t} = \Xi_{j,t,3}$ for all $j \in \mathbb{N}$ and $t \in (0, 1)$,
- (iii) $|\psi_{j,t}(z) - \psi_{j,t}(w)| \leq 3t\beta(z, w)$ for all $w, z \in \Omega$, $j \in \mathbb{N}$ and $t \in (0, 1)$.

In particular, we have $\varphi_{j,t}\psi_{j,t} = \varphi_{j,t}$ for all $j \in \mathbb{N}$ and $t \in (0, 1)$.

Proposition 9. *Let $A \in \mathcal{L}(L^p_\nu)$ and $N \in \mathbb{N}$ as in Lemma 8. Moreover, let $\varphi_{j,t}$ and $\psi_{j,t}$ be defined as above. Then the following are equivalent:*

- (i) A is band-dominated,
- (ii) $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{a_{j,t}} A M_{1-b_{j,t}} f\|^p = 0$ for all families of functions $a_{j,t}, b_{j,t}: \Omega \rightarrow [0, 1]$ that satisfy $\lim_{t \rightarrow 0} \inf_{j \in \mathbb{N}} \text{dist}_\beta(\text{supp } a_{j,t}, \text{supp}(1 - b_{j,t})) = \infty$ and for every $t \in (0, 1)$ and $z \in \Omega$ the sets $\{j \in \mathbb{N} : z \in \text{supp } a_{j,t}\}$ and $\{j \in \mathbb{N} : z \in \text{supp } b_{j,t}\}$ contain at most N elements,
- (iii) $\lim_{t \rightarrow 0} \left\| \sum_{j=1}^{\infty} M_{a_{j,t}} A M_{1-b_{j,t}} \right\| = 0$ under the same assumptions as in (ii),
- (iv) $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f\|^p = 0$,
- (v) $\lim_{t \rightarrow 0} \left\| \sum_{j=1}^{\infty} M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} \right\| = 0$.

Proof. Let $A \in \text{BDO}_\nu^p$. Then there is a sequence $(A_n)_{n \in \mathbb{N}}$ of band operators such that $A_n \rightarrow A$. Let $\varepsilon > 0$ and choose n sufficiently large such that $\|A - A_n\| < \varepsilon$. Now choose t sufficiently small such that $\inf_{j \in \mathbb{N}} \text{dist}_\beta(\text{supp } a_{j,t}, \text{supp}(1 - b_{j,t}))$ is larger than the band-width of A_n . This implies that $M_{a_{j,t}} A_n M_{1-b_{j,t}} = 0$ for all $j \in \mathbb{N}$. Therefore

$$\begin{aligned} \sum_{j=1}^{\infty} \|M_{a_{j,t}} A M_{1-b_{j,t}} f\|^p &= \sum_{j=1}^{\infty} \|M_{a_{j,t}} (A - A_n) M_{1-b_{j,t}} f\|^p \\ &\leq 2^p \sum_{j=1}^{\infty} (\|M_{a_{j,t}} (A - A_n) f\|^p + \|M_{a_{j,t}} (A - A_n) M_{b_{j,t}} f\|^p) \\ &\leq 2^{p+1} N \varepsilon^p \|f\|^p \end{aligned}$$

for all $f \in L^p_\nu$ and sufficiently small t by Proposition 5. As ε was arbitrary, (ii) follows.

Now assume

$$\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{a_{j,t}} A M_{1-b_{j,t}} f\|^p = 0.$$

Then

$$\left\| \sum_{j=1}^{\infty} M_{a_{j,t}} A M_{1-b_{j,t}} f \right\|^p = \int_{\Omega} \left| \sum_{j=1}^{\infty} M_{a_{j,t}} A M_{1-b_{j,t}} f \right|^p d\nu$$

$$\begin{aligned}
&\leq \int_{\Omega} N^p \sum_{j=1}^{\infty} |M_{a_j,t} AM_{1-b_j,t} f|^p \, dv_{\nu} \\
&= N^p \sum_{j=1}^{\infty} \int_{\Omega} |M_{a_j,t} AM_{1-b_j,t} f|^p \, dv_{\nu} \\
&= N^p \sum_{j=1}^{\infty} \|M_{a_j,t} AM_{1-b_j,t} f\|^p
\end{aligned}$$

for all $f \in L^p_{\nu}$ and $t \in (0, 1)$ by Proposition 4 (for every $z \in \Omega$ the sum over j in the first line contains at most N non-zero terms). It follows

$$\lim_{t \rightarrow 0} \left\| \sum_{j=1}^{\infty} M_{a_j,t} AM_{1-b_j,t} \right\| = 0.$$

Similarly, (iv) implies (v).

That (ii) implies (iv) and (iii) implies (v) is clear since $\text{dist}_{\beta}(\text{supp } \varphi_{j,t}, \text{supp}(1 - \psi_{j,t})) \geq \frac{1}{3t}$ and every $z \in \Omega$ belongs to at most N of the sets $\text{supp } \varphi_{j,t}$ and $\text{supp } \psi_{j,t}$ by construction (see Lemma 8).

We are thus left with the assertion that (v) implies (i). Define

$$A_n := \sum_{j=1}^{\infty} M_{\varphi_{j,\frac{1}{n}}} AM_{\psi_{j,\frac{1}{n}}}.$$

It is easily seen that this defines a bounded linear operator (see also Lemma 16 below). Moreover, A_n is obviously a band operator of band width at most $C(n) + 2n$, where $C(n)$ is the constant from Lemma 8 (iii). As $\sum_{j=1}^{\infty} M_{\varphi_{j,t}} = I$ for all $t \in (0, 1)$, we obtain

$$\|A - A_n\| = \left\| \sum_{j=1}^{\infty} (M_{\varphi_{j,\frac{1}{n}}} A - M_{\varphi_{j,\frac{1}{n}}} AM_{\psi_{j,\frac{1}{n}}}) \right\| = \left\| \sum_{j=1}^{\infty} M_{\varphi_{j,\frac{1}{n}}} AM_{1-\psi_{j,\frac{1}{n}}} \right\|$$

and this tends to 0 as $n \rightarrow \infty$ by assumption. \square

Corollary 10. *Let $A \in \text{BDO}_{\nu}^p$ and let $a_{j,t}, b_{j,t}: \Omega \rightarrow [0, 1]$ satisfy*

$$\liminf_{t \rightarrow 0} \inf_{j \in \mathbb{N}} \text{dist}_{\beta}(\text{supp } a_{j,t}, \text{supp}(1 - b_{j,t})) = \infty$$

and for every $t \in (0, 1)$ and $z \in \Omega$ the sets $\{j \in \mathbb{N} : z \in \text{supp } a_{j,t}\}$ and $\{j \in \mathbb{N} : z \in \text{supp } b_{j,t}\}$ contain at most N elements. Then

$$\limsup_{t \rightarrow 0} \sup_{j \in \mathbb{N}} \|M_{a_j,t} AM_{1-b_j,t}\| = 0 = \limsup_{t \rightarrow 0} \sup_{j \in \mathbb{N}} \|M_{1-b_j,t} AM_{a_j,t}\|.$$

Proof. As

$$\|M_{a_j,t} AM_{1-b_j,t}\|^p = \sup_{\|f\|=1} \|M_{a_j,t} AM_{1-b_j,t} f\|^p \leq \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{a_j,t} AM_{1-b_j,t} f\|^p$$

the first limit follows directly from Proposition 9. To show the second limit we may either repeat the first part of the proof of Proposition 9 with the reversed ordering or just observe that $A \in \text{BDO}_V^p$ is equivalent to $A^* \in \text{BDO}_V^q$ for $\frac{1}{p} + \frac{1}{q} = 1$. \square

The following characterization will also prove itself useful. The proof is quite similar to the proof of Theorem 2.1.6 in [20]. We use the standard notation $[A, B] := AB - BA$ for the commutator of two operators A and B .

Proposition 11. *Let $A \in \mathcal{L}(L_V^p)$. Then A is band-dominated if and only if*

$$\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[A, M_{\varphi_{j,t}}]f\|^p = 0. \quad (3.1)$$

Moreover, $\lim_{t \rightarrow 0} \sup_{j \in \mathbb{N}} \|[A, M_{\varphi_{j,t}}]\| = 0$ in this case.

We divide the proof in two parts. In the first part we deal with band operators only and show a little bit more than we need here. This will come in handy later on.

Lemma 12. *Let $\omega > 0$ and let $a_{j,t}: \Omega \rightarrow [0, 1]$ be measurable functions for $j \in \mathbb{N}$ and $t \in (0, 1)$. If $\liminf_{t \rightarrow 0} \inf_{j \in \mathbb{N}} \text{dist}_\beta(a_{j,t}^{-1}(U), a_{j,t}^{-1}(V)) \rightarrow \infty$ for all sets $U, V \subset [0, 1]$ with $\text{dist}(U, V) > 0$, then for every $\varepsilon > 0$ there exists a $t_0 > 0$ such that for all $t < t_0$ and all band operators of band width at most ω the estimate*

$$\sup_{j \in \mathbb{N}} \|[A, M_{a_{j,t}}]\| \leq 3 \|A\| \varepsilon$$

holds.

Proof. Let $A \in \mathcal{L}(L_V^p)$ be a band operator of band width at most ω , fix $\varepsilon > 0$ and set $m := \lceil \frac{1}{\varepsilon} \rceil$. For $k = 1, \dots, m$ we define the following sets:

$$U_{k,t}^j := \{z \in \Omega : a_{j,t}(z) \geq k\varepsilon\} \quad \text{and} \quad V_{k,t}^j := \left\{z \in \Omega : a_{j,t}(z) \geq \left(k - \frac{1}{2}\right)\varepsilon\right\}.$$

Moreover, we set

$$a_{j,t}^U := \varepsilon \sum_{k=1}^m \chi_{U_{k,t}^j} \quad \text{and} \quad a_{j,t}^V := \varepsilon \sum_{k=1}^m \chi_{V_{k,t}^j}.$$

Clearly, for every $z \in \Omega$, $t \in (0, 1)$ and $j \in \mathbb{N}$ either $a_{j,t}(z) < \varepsilon$ or $a_{j,t}(z) \in [l\varepsilon, (l+1)\varepsilon)$ for some $l \in \{1, \dots, m\}$. In either case this implies $|a_{j,t}(z) - a_{j,t}^U(z)| < \varepsilon$ and hence $\sup_{j \in \mathbb{N}} \|a_{j,t} - a_{j,t}^U\|_\infty < \varepsilon$.

Similarly, we obtain $\sup_{j \in \mathbb{N}} \|a_{j,t} - a_{j,t}^V\|_\infty < \varepsilon$. This of course implies

$$\sup_{j \in \mathbb{N}} \|M_{a_{j,t}} - M_{a_{j,t}^U}\| < \varepsilon \quad \text{and} \quad \sup_{j \in \mathbb{N}} \|M_{a_{j,t}} - M_{a_{j,t}^V}\| < \varepsilon.$$

It follows

$$\sup_{j \in \mathbb{N}} \|AM_{a_{j,t}} - M_{a_{j,t}}A\| \leq \sup_{j \in \mathbb{N}} \left(\|A(M_{a_{j,t}} - M_{a_{j,t}^V})\| + \|(AM_{a_{j,t}^V} - M_{a_{j,t}^V}A)\| + \|(M_{a_{j,t}^U} - M_{a_{j,t}})A\| \right)$$

$$\begin{aligned}
&\leq 2 \|A\| \varepsilon + \sup_{j \in \mathbb{N}} \left\| (AM_{a_{j,t}^V} - M_{a_{j,t}^U} A) \right\| \\
&= 2 \|A\| \varepsilon + \varepsilon \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m (AM_{\chi_{V_{k,t}^j}} - M_{\chi_{U_{k,t}^j}} A) \right\| \\
&\leq 2 \|A\| \varepsilon + \varepsilon \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{\Omega \setminus U_{k,t}^j}} AM_{\chi_{V_{k,t}^j}} \right\| + \varepsilon \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{U_{k,t}^j}} AM_{\chi_{\Omega \setminus V_{k,t}^j}} \right\|. \tag{3.2}
\end{aligned}$$

Since $U_{k,t}^j = a_{j,t}^{-1}([k\varepsilon, 1])$ and $\Omega \setminus V_{k,t}^j = a_{j,t}^{-1}([0, (k - \frac{1}{2})\varepsilon])$, we obtain

$$\inf_{j \in \mathbb{N}} \text{dist}_\beta(U_{k,t}^j, \Omega \setminus V_{k,t}^j) \rightarrow \infty$$

for all k as $t \rightarrow 0$ by assumption. Choose t sufficiently small such that $\text{dist}_\beta(U_{k,t}^j, \Omega \setminus V_{k,t}^j) > \omega$ for all $j \in \mathbb{N}$. As A is a band operator of band width at most ω , the third term in (3.2) vanishes.

Similarly, setting $U_{0,t}^j := \Omega$ and $V_{m+1,t}^j := \emptyset$, we get

$$\begin{aligned}
\sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{\Omega \setminus U_{k,t}^j}} AM_{\chi_{V_{k,t}^j}} \right\| &\leq \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{\Omega \setminus U_{k,t}^j}} AM_{\chi_{V_{k+1,t}^j}} \right\| + \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{\Omega \setminus U_{k-1,t}^j}} AM_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} \right\| \\
&\quad + \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{U_{k-1,t}^j} \setminus U_{k,t}^j} AM_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} \right\| \\
&= \sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{U_{k-1,t}^j} \setminus U_{k,t}^j} AM_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} \right\|.
\end{aligned}$$

As the sets $U_{k-1,t}^j \setminus U_{k,t}^j$ are pairwise disjoint for $k = 1, \dots, m$, this can be further estimated as

$$\begin{aligned}
\sup_{j \in \mathbb{N}} \left\| \sum_{k=1}^m M_{\chi_{U_{k-1,t}^j} \setminus U_{k,t}^j} AM_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} f \right\|^p &= \sup_{j \in \mathbb{N}} \sum_{k=1}^m \left\| M_{\chi_{U_{k-1,t}^j} \setminus U_{k,t}^j} AM_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} f \right\|^p \\
&\leq \|A\|^p \sup_{j \in \mathbb{N}} \sum_{k=1}^m \left\| M_{\chi_{V_{k,t}^j} \setminus V_{k+1,t}^j} f \right\|^p \\
&= \|A\|^p \sup_{j \in \mathbb{N}} \left\| M_{\chi_{V_{1,t}^j}} f \right\|^p \\
&\leq \|A\|^p \|f\|^p
\end{aligned}$$

for all $f \in L_v^p$. Thus

$$\sup_{j \in \mathbb{N}} \left\| (AM_{a_{j,t}} - M_{a_{j,t}} A) \right\| \leq 2 \|A\| \varepsilon + \|A\| \varepsilon = 3 \|A\| \varepsilon \quad \square.$$

Now we can prove Proposition 11.

Proof of Proposition 11. Let $A \in \text{BDO}_v^p$ and fix $\varepsilon > 0$. Then there is a sequence of band operators $(A_n)_{n \in \mathbb{N}}$ such that $A_n \rightarrow A$. Choose n sufficiently large such that $\|A - A_n\| < \varepsilon$. Now observe that

the functions $\varphi_{j,t}$ satisfy the assumption in Lemma 12. Indeed, let $U, V \subset [0, 1]$ with $\text{dist}(U, V) > 0$ and $w_{j,t} \in \varphi_{j,t}^{-1}(U)$, $z_{j,t} \in \varphi_{j,t}^{-1}(V)$. Then

$$\beta(z_{j,t}, w_{j,t}) \geq \frac{1}{6Nt} |\varphi_{j,t}(z_{j,t}) - \varphi_{j,t}(w_{j,t})| \geq \frac{1}{6Nt} \text{dist}(U, V) \rightarrow \infty$$

as $t \rightarrow 0$. Thus there is a $t > 0$ such that

$$\sup_{j \in \mathbb{N}} \|[A, M_{\varphi_{j,t}}]\| \leq \sup_{j \in \mathbb{N}} \|[A_n, M_{\varphi_{j,t}}]\| + \sup_{j \in \mathbb{N}} \|[A - A_n, M_{\varphi_{j,t}}]\| \leq 3 \|A_n\| \varepsilon + 2\varepsilon \leq 3(\|A\| + \varepsilon)\varepsilon + 2\varepsilon$$

by Lemma 12. As ε was arbitrary, this implies

$$\limsup_{t \rightarrow 0} \sup_{j \in \mathbb{N}} \|[A, M_{\varphi_{j,t}}]\| = 0.$$

Using

$$\begin{aligned} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[A, M_{\varphi_{j,t}}]f\|^p &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|[A, M_{\varphi_{j,t}}]M_{\psi_{j,t}}f\|^p + \|[A, M_{\varphi_{j,t}}]M_{1-\psi_{j,t}}f\|^p) \\ &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|[A, M_{\varphi_{j,t}}]\|^p \|M_{\psi_{j,t}}f\|^p + \|M_{\varphi_{j,t}}AM_{1-\psi_{j,t}}f\|^p) \\ &\leq 2^p N \sup_{j \in \mathbb{N}} \|[A, M_{\varphi_{j,t}}]\|^p + \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}}AM_{1-\psi_{j,t}}f\|^p, \end{aligned}$$

and Proposition 9, we obtain

$$\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[A, M_{\varphi_{j,t}}]f\|^p = 0$$

as claimed.

Conversely, assume that (3.1) holds. Clearly, this implies $\limsup_{t \rightarrow 0} \sup_{j \in \mathbb{N}} \|[A, M_{\varphi_{j,t}}]\| = 0$ as well (cf. proof of Corollary 10). We can thus proceed as above to obtain

$$\begin{aligned} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}}AM_{1-\psi_{j,t}}f\|^p &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|[M_{\varphi_{j,t}}, A]f\|^p + \|[M_{\varphi_{j,t}}, A]M_{\psi_{j,t}}f\|^p) \\ &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|[M_{\varphi_{j,t}}, A]f\|^p + \|[M_{\varphi_{j,t}}, A]\|^p \|M_{\psi_{j,t}}f\|^p) \\ &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[M_{\varphi_{j,t}}, A]f\|^p + 2^p N \sup_{j \in \mathbb{N}} \|[M_{\varphi_{j,t}}, A]\|^p. \end{aligned}$$

and by assumption, this tends to 0 as $t \rightarrow 0$. Thus $A \in \text{BDO}_\nu^p$ by Proposition 9. \square

Here are some algebraic properties of BDO_ν^p :

Proposition 13. BDO_ν^p has the following properties:

- (i) It holds $M_f \in \text{BDO}_\nu^p$ for all $f \in L^\infty(\Omega)$.
- (ii) BDO_ν^p is a closed subalgebra of $\mathcal{L}(L_\nu^p)$.
- (iii) If $A \in \text{BDO}_\nu^p$ is Fredholm, then every regularizer B of A is again in BDO_ν^p . In particular, BDO_ν^p is inverse closed.
- (iv) BDO_ν^p contains $\mathcal{K}(L_\nu^p)$ as a closed two-sided ideal.
- (v) It holds $A \in \text{BDO}_\nu^p \iff A^* \in \text{BDO}_\nu^q$ for $\frac{1}{p} = \frac{1}{q} = 1$. In particular, BDO_ν^2 is a C^* -algebra.

Proof. (i), (ii) and (v) are easy to see.

(iii): Set $\tilde{\varphi}_{j,t} := \varphi_{j,t} - \varphi_{j,t}(0)$ for all $j \in \mathbb{N}$ and $t \in (0, 1)$. If A is Fredholm, there exist a regularizer $B \in \mathcal{L}(L_\nu^p)$ and compact operators $K_1, K_2 \in \mathcal{K}(L_\nu^p)$ such that $AB = I + K_1$ and $BA = I + K_2$. This implies

$$[B, M_{\varphi_{j,t}}] = [B, M_{\tilde{\varphi}_{j,t}}] = B[M_{\tilde{\varphi}_{j,t}}, A]B - BM_{\tilde{\varphi}_{j,t}}K_1 + K_2M_{\tilde{\varphi}_{j,t}}B.$$

Thus

$$\sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[B, M_{\varphi_{j,t}}]f\|^p \leq 3^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|B[M_{\tilde{\varphi}_{j,t}}, A]Bf\|^p + \|BM_{\tilde{\varphi}_{j,t}}K_1f\|^p + \|K_2M_{\tilde{\varphi}_{j,t}}Bf\|^p). \quad (3.3)$$

As A is band-dominated, the first term tends to 0 as $t \rightarrow 0$ by Proposition 11. To estimate the other two terms, define $D(0, R) := \{z \in \Omega : \beta(0, z) < R\}$ and $D(0, R)^c := \Omega \setminus D(0, R)$ for $R > 0$. Since K_1 and K_2 are compact, $\|M_{\chi_{D(0, R)^c}}K_1\|$ and $\|K_2M_{\chi_{D(0, R)^c}}\|$ tend to 0 as $R \rightarrow \infty$. Moreover, as for fixed t the origin is contained in at most N of the sets $\Xi_{j,t,3} = \{z \in \Omega : \text{dist}_\beta(z, B_{j,t}) \leq \frac{1}{t}\}$ and $\text{supp } \varphi_{j,t} = \Xi_{j,t,1} = \{z \in \Omega : \text{dist}_\beta(z, B_{j,t}) \leq \frac{1}{3t}\}$, we get that $\tilde{\varphi}_{j,t}$ vanishes on $D(0, \frac{2}{3\sqrt{t}}) \subset D(0, \frac{2}{3t})$ for all but at most N integers j . W.l.o.g. we may assume that these integers are $j = 1, \dots, N$. For $j = 1, \dots, N$ and $z \in D(0, \frac{2}{3\sqrt{t}})$ it holds $|\tilde{\varphi}_{j,t}(z)| \leq 6Nt\frac{2}{3\sqrt{t}} = 4N\sqrt{t}$ by property (iii) of $\varphi_{j,t}$. Therefore

$$\begin{aligned} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|BM_{\tilde{\varphi}_{j,t}}K_1f\|^p &\leq 2^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} (\|BM_{\tilde{\varphi}_{j,t}}M_{\chi_{D(0, R)^c}}K_1f\|^p + \|BM_{\tilde{\varphi}_{j,t}}M_{\chi_{D(0, R)}}K_1f\|^p) \\ &\leq 2^p \|B\|^p \|M_{\chi_{D(0, R)^c}}K_1\|^p \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\tilde{\varphi}_{j,t}}f\|^p \\ &\quad + 2^p N \|B\|^p (4N\sqrt{t})^p \|K_1\|^p \end{aligned}$$

for $t < \frac{4}{9R^2}$. As

$$\sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\tilde{\varphi}_{j,t}}f\|^p = \sup_{\|f\|=1} \left(\sum_{j=1}^N \|M_{\tilde{\varphi}_{j,t}}f\|^p + \sum_{j=N+1}^{\infty} \|M_{\varphi_{j,t}}f\|^p \right)$$

is bounded by $2N$, this implies $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|BM_{\tilde{\varphi}_{j,t}}K_1f\|^p = 0$. Similarly, we obtain the equality

$\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|K_2M_{\tilde{\varphi}_{j,t}}Bf\|^p = 0$. Plugging these observations into (3.3), we conclude

$$\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[B, M_{\varphi_{j,t}}]f\|^p = 0$$

and hence $B \in \text{BDO}_\nu^p$ by Proposition 11.

(iv) The argument in (iii) shows $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\tilde{\varphi}_{j,t}}Kf\|^p = \lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|KM_{\tilde{\varphi}_{j,t}}f\|^p = 0$ for $K \in \mathcal{K}(L_\nu^p)$. Thus the assertion follows again by Proposition 11. \square

To prove Theorem 7, we will only need one more auxiliary lemma. A similar result for the unit ball was shown in [17, Lemma 3.4].

Lemma 14. *Let (α, ν, p) be an admissible triple and for every j let $a_j, b_j: \Omega \rightarrow [0, 1]$ be measurable functions. If*

- (i) *there exists a $\sigma \geq 0$ such that $\text{dist}_\beta(\text{supp } a_j, \text{supp}(1 - b_j)) \geq \sigma$ for all $j \in \mathbb{N}$,*
- (ii) *there is an integer N such that every $z \in \Omega$ belongs to at most N of the sets $\text{supp } a_j$ and to at most N of the sets $\text{supp } b_j$,*

then there is a function $\sigma \mapsto \beta_{p,\alpha,\nu}(\sigma)$ (depending only on p, α and ν) converging to 0 as $\sigma \rightarrow \infty$ such that

$$\left\| \sum_{j=1}^{\infty} M_{a_j} P_\alpha M_{1-b_j} \right\| \leq N \beta_{p,\alpha,\nu}(\sigma).$$

In other words,

$$\left\| \sum_{j=1}^{\infty} M_{a_j} P_\alpha M_{1-b_j} \right\| \rightarrow 0$$

as $\inf_{j \in \mathbb{N}} \text{dist}_\beta(\text{supp } a_j, \text{supp}(1 - b_j)) \rightarrow \infty$.

Proof. Let us consider the case where every $z \in \Omega$ belongs to at most 1 of the sets $\text{supp } a_j$ first. Define

$$\Phi(z, w) := \sum_{j=1}^{\infty} \chi_{\text{supp } a_j}(z) \chi_{\text{supp}(1-b_j)}(w) |h(z, w)|^{-\alpha-g}.$$

Then

$$\begin{aligned} \left| \sum_{j=1}^{\infty} (M_{a_j} P_\alpha M_{1-b_j} f)(z) \right| &= \left| \sum_{j=1}^{\infty} a_j(z) \int_{\Omega} (1 - b_j(w)) f(w) h(z, w)^{-\alpha-g} \, dv_\alpha(w) \right| \\ &\leq \frac{c_\alpha}{c_\nu} \int_{\Omega} \Phi(z, w) h(w, w)^{\alpha-\nu} |f(w)| \, dv_\nu(w). \end{aligned}$$

As in the proof of Proposition 1, we want to apply Schur's test with $h(z) := h(z, z)^s$, where $s \in \mathbb{R}$ has to be determined later. We thus need to show that there exist two constants C_1 and C_2 such that

$$(I1) := \int_{\Omega} \Phi(z, w) h(w, w)^{\alpha-\nu} h(w, w)^{sq} dv_{\nu}(w) \leq C_1 h(z, z)^{sq}$$

for (almost) every $z \in \Omega$ and

$$(I2) := \int_{\Omega} \Phi(z, w) h(w, w)^{\alpha-\nu} h(z, z)^{sp} dv_{\nu}(z) \leq C_2 h(w, w)^{sp}$$

for (almost) every $w \in \Omega$.

So let $z \in \Omega$. We may assume that $z \in \text{supp } a_j$ for some $j \in \mathbb{N}$, otherwise the left-hand side is just 0. Now choose $s \in (-\frac{\alpha+1}{q}, -\frac{(r-1)a}{2q}) \cap (-\frac{\nu+1}{p}, -\frac{\nu}{p} + \frac{\alpha}{p} - \frac{(r-1)a}{2p})$ as in the proof of Proposition 1 and $q_0 > 1$ sufficiently small such that $(sq + \alpha)q_0 > -1$ and $(sq + g)q_0 - g < -\frac{(r-1)\alpha}{2}$. Moreover, let $D(w, r) := \{z \in \Omega : \beta(w, z) < r\}$ for midpoints w and radii r . Since we assumed $N = 1$, there is only one term contributing to $\Phi(z, w)$ and so

$$\begin{aligned} (I1) &= c_{\nu} \int_{\Omega} \Phi(z, w) h(w, w)^{sq+\alpha} dv(w) \\ &= c_{\nu} \int_{\text{supp}(1-b_j)} |h(z, w)|^{-\alpha-g} h(w, w)^{sq+\alpha} dv(w) \\ &\leq c_{\nu} \int_{\Omega \setminus D(z, \sigma)} |h(z, w)|^{-\alpha-g} h(w, w)^{sq+\alpha} dv(w) \\ &= c_{\nu} \int_{\Omega \setminus D(0, \sigma)} |h(\phi_z(0), \phi_z(u))|^{-\alpha-g} h(\phi_z(u), \phi_z(u))^{sq+\alpha} d\phi_z(u) \\ &= c_{\nu} \int_{\Omega \setminus D(0, \sigma)} \left| \frac{h(z, z)}{h(z, u)} \right|^{-\alpha-g} \left(\frac{h(z, z)h(u, u)}{|h(z, u)|^2} \right)^{sq+\alpha} \frac{h(z, z)^g}{|h(z, u)|^{2g}} dv(u) \\ &= c_{\nu} h(z, z)^{sq} \int_{\Omega \setminus D(0, \sigma)} |h(z, u)|^{-2sq-\alpha-g} h(u, u)^{sq+\alpha} dv(u) \\ &\leq c_{\nu} h(z, z)^{sq} |\Omega \setminus D(0, \sigma)|^{1/p_0} \left(\int_{\Omega} |h(z, u)|^{-(2sq+\alpha+g)q_0} h(u, u)^{(sq+\alpha)q_0} dv(u) \right)^{1/q_0} \\ &\leq c_{\nu} h(z, z)^{sq} |\Omega \setminus D(0, \sigma)|^{1/p_0} C^{1/q_0}, \end{aligned}$$

where C is some constant (coming from the Rudin-Forelli estimates [12, Theorem 4.1]) and $\frac{1}{p_0} + \frac{1}{q_0} = 1$ as usual.

Now let $w \in \Omega$. We obtain

$$\begin{aligned} (I2) &= c_{\nu} h(w, w)^{\alpha-\nu} \int_{\Omega} \Phi(z, w) h(z, z)^{sp+\nu} dv(z) \\ &\leq c_{\nu} h(w, w)^{\alpha-\nu} \int_{\Omega} \sum_{j=1}^{\infty} \chi_{\text{supp } a_j}(z) |h(z, w)|^{-\alpha-g} h(z, z)^{sp+\nu} dv(z) \\ &\leq c_{\nu} h(w, w)^{\alpha-\nu} \int_{\Omega} |h(z, w)|^{-\alpha-g} h(z, z)^{sp+\nu} dv(z) \end{aligned}$$

$$\leq C_2 h(w, w)^{sp}$$

for $s \in (-\frac{\nu+1}{p}, -\frac{\nu}{p} + \frac{\alpha}{p} - \frac{(r-1)a}{2p})$ as in Proposition 1. Thus, by Schur's test, we have the following norm estimate:

$$\left\| \sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j} \right\| \leq \left(c_{\nu} |\Omega \setminus D(0, \sigma)|^{1/p_0} C^{1/q_0} \right)^{1/q} C_2^{1/p} =: \beta_{p, \alpha, \nu}(\sigma)$$

with $\beta_{p, \alpha, \nu}(\sigma) \rightarrow 0$ as $\sigma \rightarrow \infty$. This proves the estimate in the case $N = 1$.

Now let us consider the case $N > 1$. As in the proof of Proposition 5, there is a disjoint decomposition $\text{supp } a_j = A_j^1 \cup \dots \cup A_j^N$ such that the sets $(A_j^i)_{j \in \mathbb{N}}$ are again measurable and pairwise disjoint for every $i \in \{1, \dots, N\}$. It follows

$$\begin{aligned} \left\| \sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j} \right\| &= \left\| \sum_{j=1}^{\infty} \sum_{i=1}^N M_{a_j \chi_{A_j^i}} P_{\alpha} M_{1-b_j} \right\| \\ &\leq \sum_{i=1}^N \left\| \sum_{j=1}^{\infty} M_{a_j \chi_{A_j^i}} P_{\alpha} M_{1-b_j} \right\| \\ &\leq N \beta_{p, \alpha, \nu}(\sigma). \end{aligned} \quad \square$$

Proof of Theorem 7. Combining Lemma 14 and Proposition 9, we get that P_{α} is band-dominated. By Proposition 13 the set BDO_{ν}^p is a Banach algebra that contains all multiplication operators. It thus contains all operators of the form $T_f P_{\alpha} + Q_{\alpha} = P_{\nu} M_f P_{\alpha} + Q_{\alpha}$ and therefore all operators of the form $A P_{\alpha} + Q_{\alpha}$ with $A \in \mathfrak{T}_{p, \nu}$. \square

4 A Fredholm criterion for band-dominated operators

In this rather short section we show a Fredholm criterion for band-dominated operators. First, we need another auxiliary proposition that is of course well-known. For completeness we include a short proof.

Proposition 15. *Let (α, ν, p) be an admissible triple and let $D \subset \Omega$ be a compact set. Then the operators $P_{\alpha} M_{\chi_D}$ and $M_{\chi_D} P_{\alpha} : L_{\nu}^p \rightarrow L_{\nu}^p$ are compact.*

Proof. By definition,

$$(P_{\alpha} M_{\chi_D} f)(z) = c_{\alpha} \int_{\Omega} \chi_D(w) f(w) h(z, w)^{-\alpha-g} h(w, w)^{\alpha} dv(w)$$

for all $f \in L_{\nu}^p$. As D is compact, $\chi_D(w) h(z, w)^{-\alpha-g} h(w, w)^{\alpha}$ is uniformly bounded (shown for example in the proof of [11, Proposition 3]). This implies that $P_{\alpha} M_{\chi_{B_R}}$ is compact by the Hille-Tamarkin theorem (see e.g. [27, Theorem 41.6]).

Similarly,

$$(M_{\chi_D} P_{\alpha} f)(z) = c_{\alpha} \int_{\Omega} \chi_D(z) f(w) h(z, w)^{-\alpha-g} h(w, w)^{\alpha} dv(w)$$

and thus $M_{\chi_D} P_{\alpha}$ is compact by the same argument. \square

We will also need the following lemma, which is a small modification of [19, Proposition 13].

Lemma 16. *For $j \in \mathbb{N}$ let $a_j, b_j: \Omega \rightarrow [0, 1]$ be measurable functions and $A_j \in \mathcal{L}(L_\nu^p)$ so that the sequence $(A_j)_{j \in \mathbb{N}}$ is uniformly bounded. If there is an integer N such that every $z \in \Omega$ belongs to at most N of the sets $\text{supp } a_j$ and to at most N of the sets $\text{supp } b_j$, then the series $\sum_{j=1}^{\infty} M_{a_j} A_j M_{b_j}$ converges $*$ -strongly and*

$$\left\| \sum_{j=1}^{\infty} M_{a_j} A_j M_{b_j} \right\| \leq N^2 \sup_{j \in \mathbb{N}} \|A_j\|.$$

Moreover,

$$\sum_{j=1}^{\infty} \|M_{a_j} A_j M_{b_j} f\|^p \leq N \sup_{j \in \mathbb{N}} \|A_j\|^p \|f\|^p$$

for all $f \in L_\nu^p$.

Proof. Since every $z \in \Omega$ belongs to at most N of the sets $\text{supp } a_j$, it follows

$$\begin{aligned} \left\| \sum_{j=1}^{\infty} M_{a_j} A_j M_{b_j} f \right\|^p &= \int_{\Omega} \left| \sum_{j=1}^{\infty} M_{a_j} A_j M_{b_j} f \right|^p dv_\nu \\ &\leq \int_{\Omega} N^p \sum_{j=1}^{\infty} |M_{a_j} A_j M_{b_j} f|^p dv_\nu \\ &= N^p \sum_{j=1}^{\infty} \int_{\Omega} |M_{a_j} A_j M_{b_j} f|^p dv_\nu \\ &= N^p \sum_{j=1}^{\infty} \|M_{a_j} A_j M_{b_j} f\|^p \end{aligned}$$

for all $f \in L_\nu^p$ by Proposition 4 (for every $z \in \Omega$ the sum over j in the first line contains at most N non-zero terms). Using Proposition 5, we also get $\sum_{j=1}^{\infty} \|M_{b_j} f\|^p \leq N \|f\|^p$. We thus obtain

$$\left\| \sum_{j=1}^{\infty} M_{a_j} A_j M_{b_j} f \right\|^p \leq N^p \sum_{j=1}^{\infty} \|M_{a_j} A_j M_{b_j} f\|^p \leq N^p \sum_{j=1}^{\infty} \|A_j\|^p \|M_{b_j} f\|^p \leq N^{p+1} \sup_{j \in \mathbb{N}} \|A_j\|^p \|f\|^p$$

for all $f \in L_\nu^p$. This yields both inequalities and the $*$ -strong convergence follows easily as well. \square

Proposition 17. *Let (α, ν, p) be an admissible triple, $A \in \text{BDO}_\nu^p$ satisfy $[A, P_\alpha] = 0$ and $\psi_{j,t}$ as above. If there is a constant $M > 0$ such that for every $t \in (0, 1)$ there is a $j_0 \in \mathbb{N}$ such that for all $j \geq j_0$ there are operators $B_{j,t}, C_{j,t} \in \mathcal{L}(L_\nu^p)$ with $\|B_{j,t}\|, \|C_{j,t}\| \leq M$ and*

$$B_{j,t} A M_{\psi_{j,t}} = M_{\psi_{j,t}} = M_{\psi_{j,t}} A C_{j,t},$$

then $A|_{A_\nu^p} \in \mathcal{L}(A_\nu^p)$ is Fredholm and $\|(A|_{A_\nu^p} + \mathcal{K}(A_\nu^p))^{-1}\| \leq 2 \min\{\|P_\nu\|, \|P_\alpha\|\} N^2 M$.

Proof. Let the functions $\varphi_{j,t}$ be as above and $\varepsilon > 0$. Then by Lemma 16, the series

$$B_t := \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}}$$

converges strongly with $\|B_t\| \leq N^2 M$. Multiplying by A , we obtain

$$B_t A = \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}} A M_{\psi_{j,t}} + \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}} A M_{1-\psi_{j,t}}, \quad (4.1)$$

where the strong convergence of the two series on the right-hand side is again guaranteed by Lemma 16. As every $z \in \Omega$ belongs to at most N of the sets $\text{supp } \psi_{j,t}$, the same computation as in the proof of Lemma 16 yields

$$\begin{aligned} \left\| \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f \right\|^p &\leq N^p \sum_{j=j_0}^{\infty} \left\| M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f \right\|^p \\ &\leq N^p M^p \sum_{j=j_0}^{\infty} \left\| M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f \right\|^p \end{aligned}$$

for every $f \in L^p_\nu$. Therefore the second term in (4.1) tends to 0 by Proposition 9. For the first term we further compute

$$\sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} M_{\varphi_{j,t}} A M_{\psi_{j,t}} = \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} A M_{\varphi_{j,t}} M_{\psi_{j,t}} + \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} [M_{\varphi_{j,t}}, A] M_{\psi_{j,t}},$$

where the latter term tends to 0 by Lemma 16 and Proposition 11. Furthermore, we have

$$\sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} A M_{\varphi_{j,t}} M_{\psi_{j,t}} = \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} B_{j,t} A M_{\psi_{j,t}} M_{\varphi_{j,t}} = \sum_{j=j_0}^{\infty} M_{\psi_{j,t}} M_{\psi_{j,t}} M_{\varphi_{j,t}} = \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}}.$$

Combining all these estimates, we conclude

$$\lim_{t \rightarrow 0} \left\| B_t A - \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} \right\| = 0.$$

In particular, we have

$$\lim_{t \rightarrow 0} \left\| P_\nu B_t A|_{A_\nu^p} - \sum_{j=j_0}^{\infty} P_\nu M_{\varphi_{j,t}}|_{A_\nu^p} \right\| \leq \lim_{t \rightarrow 0} \|P_\nu\| \left\| B_t A - \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} \right\| = 0.$$

Now as the functions $\varphi_{j,t}$ have compact support, the operators $P_\nu M_{\varphi_{j,t}}|_{A_\nu^p}$ are compact by Proposition 15 and hence $\sum_{j=j_0}^{\infty} P_\nu M_{\varphi_{j,t}}|_{A_\nu^p} \in I + \mathcal{K}(A_\nu^p)$ for every $t > 0$. We deduce that $P_\nu B_t A|_{A_\nu^p} + \mathcal{K}(A_\nu^p)$

converges to $I + \mathcal{K}(A_\nu^p)$. By a Neumann series argument, this implies that there exists a $B \in \mathcal{L}(A_\nu^p)$ such that $BA|_{A_\nu^p} \in I + \mathcal{K}(A_\nu^p)$ and

$$\|B + \mathcal{K}(A_\nu^p)\| \leq 2\|P_\nu\| \|B_t\| \leq 2\|P_\nu\| N^2 M.$$

As $A^* \in \text{BDO}_\nu^q$, we can apply the above to A^* to obtain

$$\lim_{t \rightarrow 0} \left\| AC_t - \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} \right\| = \lim_{t \rightarrow 0} \left\| C_t^* A^* - \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} \right\| = 0$$

for

$$C_t := \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} C_{j,t} M_{\psi_{j,t}}.$$

This implies

$$\lim_{t \rightarrow 0} \left\| AP_\alpha C_t|_{A_\nu^p} - \sum_{j=j_0}^{\infty} P_\alpha M_{\varphi_{j,t}}|_{A_\nu^p} \right\| \leq \lim_{t \rightarrow 0} \|P_\alpha\| \left\| AC_t - \sum_{j=j_0}^{\infty} M_{\varphi_{j,t}} \right\| = 0$$

because $[A, P_\alpha] = 0$. Now we can precede as above to obtain an operator $C \in \mathcal{L}(A_\nu^p)$ with

$$\|C + \mathcal{K}(A_\nu^p)\| \leq 2\|P_\alpha\| N^2 M$$

such that $A|_{A_\nu^p} C \in I + \mathcal{K}(A_\nu^p)$. □

5 Limit operators (unit ball)

From this section onwards we focus on the case of the unit ball $\Omega = \mathbb{B}^n$. The corresponding results are expected to hold for general bounded symmetric domains as well, but need some more preparations. These are postponed to future work.

As we expect the Fredholm information to be located at the boundary, we consider the following shift operators². Let $U_z^p: L_\nu^p \rightarrow L_\nu^p$ be defined by

$$(U_z^p f)(w) = f(\phi_z(w)) \frac{(1 - |z|^2)^{\frac{\nu+n+1}{p}}}{(1 - \langle w, z \rangle)^{\frac{2(\nu+n+1)}{p}}}.$$

Using the standard identities mentioned in Section 2, one obtains that U_z^p is a surjective isometry with $(U_z^p)^2 = I$. In particular, $(U_z^q)^*$ is also an isometry. Moreover, it holds $U_z^p(A_\nu^p) \subseteq A_\nu^p$. However, note that $(U_z^q)^*(A_\nu^p) \not\subseteq A_\nu^p$ in general so that we have to distinguish between $(U_z^q)^*|_{A_\nu^p}$ and $(U_z^q|_{A_\nu^p})^* = P_\nu(U_z^q)^*|_{A_\nu^p}$.

If $A \in \mathfrak{T}_{p,\nu}$ and (z_γ) is a net in \mathbb{B}^n converging to $x \in \mathfrak{M}$, the maximal ideal space of $\text{BUC}(\mathbb{B}^n)$ considered as a compactification of \mathbb{B}^n (see [17, Section 4] for a discussion), then $U_{z_\gamma}^p A(U_{z_\gamma}^q|_{A_\nu^p})^*$ converges $*$ -strongly in $\mathcal{L}(A_\nu^p)$ (see [17, Proposition 4.11]) and the limit is denoted by A_x . If x is

²Strictly speaking, they are rather reflections than shifts, but they serve the purpose of “shifting” operators to the boundary as $z \rightarrow \partial\mathbb{B}^n$.

located at the boundary $\mathfrak{M} \setminus \mathbb{B}^n$, we will call the operator A_x a limit operator of A , which is in accordance with the sequence space case ([13, 14, 15, 19, 20, 21, 22] and the references therein). Note that $\mathcal{K}(A_\nu^p) \subset \mathfrak{T}_{p,\nu}$ and $K_x = 0$ for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$ and $K \in \mathcal{K}(A_\nu^p)$ by [17, Proposition 4.12, Theorem 5.5].

The set of all limit operators $\{A_x : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ is sometimes called the operator spectrum of A because it shares some properties with the usual spectrum, e.g. some kind of compactness (see Proposition 18 below). Note that the operator A_x does not depend on the net (z_γ) converging to $x \in \mathfrak{M}$ (but of course on the element $x \in \mathfrak{M}$). Let $b_z : \Omega \rightarrow \mathbb{C}$ be given by

$$b_z(w) := \frac{(1 - \langle z, w \rangle)^{(\nu+n+1)(1/q-1/p)}}{(1 - \langle w, z \rangle)^{(\nu+n+1)(1/q-1/p)}}.$$

Then $T_{b_z} = (U_z^q|_{A_\nu^q})^* U_z^p|_{A_\nu^p}$ is invertible with $T_{b_z}^{-1} = U_z^p(U_z^q|_{A_\nu^q})^*$ for all $z \in \mathbb{B}^n$. Moreover, as $z_\gamma \rightarrow x$ the net $(T_{b_{z_\gamma}})$ converges $*$ -strongly to another Toeplitz operator, which is denoted by T_{b_x} . T_{b_x} is again invertible and $T_{b_{z_\gamma}}^{-1} \rightarrow T_{b_x}^{-1}$ (see [17, Lemma 4.10]). As we will need it frequently, let us fix the strong continuity in a proposition.

Proposition 18. *For all $A \in \mathfrak{T}_{p,\nu}$ the two maps $A_\bullet : \mathfrak{M} \rightarrow \mathcal{L}(A_\nu^p)$, $x \mapsto A_x$ and $T_{b_\bullet} : \mathfrak{M} \rightarrow \mathcal{L}(A_\nu^p)$, $x \mapsto T_{b_x}$ are bounded and continuous w.r.t. the strong operator topology. In particular, the two sets $\{A_x T_{b_x} : x \in \mathfrak{M}\}$ and $\{A_x T_{b_x} : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ are strongly compact.*

Proof. This follows directly from [17, Proposition 4.11]. \square

So the aim now is to shift a Toeplitz operator A to the boundary to obtain limit operators A_x and then retrieve information about A via Proposition 17. Here is our first step:

Proposition 19. *Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)(n + 1) + \frac{2\nu}{p}$, $A \in \mathfrak{T}_{p,\nu}$ and let (z_γ) be a net in \mathbb{B}^n converging to $x \in \mathfrak{M} \setminus \mathbb{B}^n$ such that A_x is invertible. Then for every real-valued $\xi \in L^\infty(\mathbb{B}^n)$ with compact support there is a γ_0 such that for all $\gamma \geq \gamma_0$ there are operators $B_\gamma, C_\gamma \in \mathcal{L}(L_\nu^p)$ with $\|B_\gamma\|, \|C_\gamma\| \leq 2(\|A_x^{-1}\| \|P_\alpha\| + \|Q_\alpha\|)$ and*

$$B_\gamma \hat{A} M_{\xi \circ \phi_{z_\gamma}} = M_{\xi \circ \phi_{z_\gamma}} = M_{\xi \circ \phi_{z_\gamma}} \hat{A} C_\gamma.$$

Proof. First note that $p \leq 2$ implies $\alpha \geq \nu$ and hence (α, ν, p) is admissible (cf. Corollary 2). Setting $h(w, z) := 1 - \langle w, z \rangle$ and $g := n + 1$ and using the standard transformation identities, we observe

$$\begin{aligned} (U_z^p M_\xi U_z^p f)(w) &= \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(w, z)^{\frac{2(\nu+g)}{p}}} (M_\xi U_z^p f)(\phi_z(w)) \\ &= \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(w, z)^{\frac{2(\nu+g)}{p}}} \xi(\phi_z(w)) (U_z^p f)(\phi_z(w)) \\ &= \xi(\phi_z(w)) \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(w, z)^{\frac{2(\nu+g)}{p}}} \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(\phi_z(w), z)^{\frac{2(\nu+g)}{p}}} f(w) \\ &= \xi(\phi_z(w)) \frac{h(z, z)^{\frac{2(\nu+g)}{p}}}{h(w, z)^{\frac{2(\nu+g)}{p}}} \frac{h(w, z)^{\frac{2(\nu+g)}{p}}}{h(z, z)^{\frac{2(\nu+g)}{p}}} f(w) \\ &= \xi(\phi_z(w)) f(w) \end{aligned}$$

so that

$$U_z^p M_\xi U_z^p = M_{\xi \circ \phi_z}. \quad (5.1)$$

The special value we chose for α also ensures that $P_\alpha U_z^p = U_z^p P_\alpha$ for all $z \in \mathbb{B}^n$. Indeed,

$$\begin{aligned} (P_\alpha U_z^p f)(x) &= \int_{\mathbb{B}^n} f(\phi_z(w)) \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(w, z)^{\frac{2(\nu+g)}{p}}} h(x, w)^{-\alpha-g} dv_\alpha(w) \\ &= \int_{\mathbb{B}^n} f(y) \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(\phi_z(y), z)^{\frac{2(\nu+g)}{p}}} h(x, \phi_z(y))^{-\alpha-g} \frac{h(z, z)^{\alpha+g}}{|h(y, z)|^{2(\alpha+g)}} dv_\alpha(y) \\ &= \int_{\mathbb{B}^n} f(y) \frac{h(z, z)^{\frac{\nu+g}{p}} h(y, z)^{\frac{2(\nu+g)}{p}}}{h(z, z)^{\frac{2(\nu+g)}{p}}} \frac{h(x, z)^{-\alpha-g} h(\phi_z(x), y)^{-\alpha-g}}{h(z, y)^{-\alpha-g}} \frac{h(z, z)^{\alpha+g}}{|h(y, z)|^{2(\alpha+g)}} dv_\alpha(y) \\ &= \frac{h(z, z)^{-\frac{\nu+g}{p} + \alpha + g}}{h(x, z)^{\alpha+g}} \int_{\mathbb{B}^n} f(y) h(y, z)^{\frac{2(\nu+g)}{p} - \alpha - g} h(\phi_z(x), y)^{-\alpha-g} dv_\alpha(y) \\ &= \frac{h(z, z)^{\frac{\nu+g}{p}}}{h(x, z)^{\frac{2(\nu+g)}{p}}} \int_{\mathbb{B}^n} f(y) h(\phi_z(x), y)^{-\alpha-g} dv_\alpha(y) \\ &= (U_z^p P_\alpha f)(x). \end{aligned} \quad (5.2)$$

Let $B_R := \{z \in \mathbb{B}^n : |z| \leq R\}$, where $R < 1$ is chosen sufficiently large such that $\text{supp } \xi \subseteq B_R$. By Proposition 18, $U_{z_\gamma}^p A U_{z_\gamma}^p = U_{z_\gamma}^p A (U_{z_\gamma}^q |_{A^q})^* (U_{z_\gamma}^q |_{A^q})^* U_{z_\gamma}^p = A_{z_\gamma} T_{b_\gamma}$ converges $*$ -strongly to $A_x T_{b_x}$. Moreover, the operator $P_\alpha M_{\chi_{B_R}}$ is compact by Proposition 15. Combining these facts and using Equation (5.2), we get

$$\left\| \left(U_{z_\gamma}^p (A P_\alpha + Q_\alpha) U_{z_\gamma}^p - (A_x T_{b_x} P_\alpha + Q_\alpha) \right) M_{\chi_{B_R}} \right\| = \left\| \left(U_{z_\gamma}^p A U_{z_\gamma}^p - A_x T_{b_x} \right) P_\alpha M_{\chi_{B_R}} \right\| \rightarrow 0$$

as $z_\gamma \rightarrow x$. $A_x T_{b_x} P_\alpha + Q_\alpha$ is invertible with

$$(A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} = T_{b_x}^{-1} A_x^{-1} P_\alpha + Q_\alpha.$$

This implies that there exists a γ_0 such that

$$R_\gamma := (A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} \left(U_{z_\gamma}^p (A P_\alpha + Q_\alpha) U_{z_\gamma}^p - (A_x T_{b_x} P_\alpha + Q_\alpha) \right) M_{\chi_{B_R}}$$

satisfies $\|R_\gamma\| < \frac{1}{2}$ for all $\gamma \geq \gamma_0$. In particular, $I + R_\gamma \in \mathcal{L}(L_v^p)$ is invertible for all $\gamma \geq \gamma_0$.

We then have

$$U_{z_\gamma}^p (A P_\alpha + Q_\alpha) U_{z_\gamma}^p M_{\chi_{B_R}} = (A_x T_{b_x} P_\alpha + Q_\alpha) M_{\chi_{B_R}} + (A_x T_{b_x} P_\alpha + Q_\alpha) R_\gamma$$

and therefore

$$(A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} U_{z_\gamma}^p (A P_\alpha + Q_\alpha) U_{z_\gamma}^p M_\xi = M_\xi + R_\gamma M_\xi = (I + R_\gamma) M_\xi,$$

which implies

$$(I + R_\gamma)^{-1} (A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} U_{z_\gamma}^p (A P_\alpha + Q_\alpha) U_{z_\gamma}^p M_\xi = M_\xi.$$

Applying $U_{z_\gamma}^p$ from both sides and using (5.1) yields

$$U_{z_\gamma}^p (I + R_\gamma)^{-1} (A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} U_{z_\gamma}^p (AP_\alpha + Q_\alpha) M_{\xi \circ \phi_{z_\gamma}} = M_{\xi \circ \phi_{z_\gamma}}$$

and the first assertion follows. For the second assertion note that $M_{\chi_{B_R}} P_\alpha$ is compact as well (see Proposition 15). Thus

$$\left\| M_{\chi_{B_R}} \left(U_{z_\gamma}^p (AP_\alpha + Q_\alpha) U_{z_\gamma}^p - (A_x T_{b_x} P_\alpha + Q_\alpha) \right) \right\| = \left\| M_{\chi_{B_R}} P_\alpha \left(U_{z_\gamma}^p A U_{z_\gamma}^p - A_x T_{b_x} \right) P_\alpha \right\| \rightarrow 0$$

and we obtain

$$M_{\xi \circ \phi_{z_\gamma}} (AP_\alpha + Q_\alpha) U_{z_\gamma}^p (A_x T_{b_x} P_\alpha + Q_\alpha)^{-1} (I + S_\gamma)^{-1} U_{z_\gamma}^p = M_{\xi \circ \phi_{z_\gamma}}$$

for sufficiently large γ and

$$S_\gamma := M_{\chi_{B_R}} \left(U_{z_\gamma}^p (AP_\alpha + Q_\alpha) U_{z_\gamma}^p - (A_x T_{b_x} P_\alpha + Q_\alpha) \right) (A_x T_{b_x} P_\alpha + Q_\alpha)^{-1}. \quad \square$$

Combining Proposition 19 with Proposition 17, we obtain the following theorem.

Theorem 20. *Let $A \in \mathfrak{F}_{p,\nu}$. If A_x is invertible for every $x \in \mathfrak{M} \setminus \mathbb{B}^n$ and $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x^{-1}\| < \infty$, then A is Fredholm.*

Proof. W.l.o.g. we may assume that $p \leq 2$ because otherwise we can just pass to the adjoint, noting that $(A^*)_x = (A_x)^*$ for all $x \in \mathfrak{M}$.

Let $\psi_{j,t}$ be the functions defined above and assume that A is not Fredholm. It is clear that

$$[\hat{A}, P_\alpha] = (AP_\alpha + Q_\alpha)P_\alpha - P_\alpha(AP_\alpha + Q_\alpha) = AP_\alpha - P_\alpha AP_\alpha = 0$$

as $A \in \mathcal{L}(L_\nu^p)$. Thus by Proposition 17, there is a $t \in (0, 1)$ and a strictly increasing sequence $(j_m)_{m \in \mathbb{N}}$ such that

$$B \hat{A} M_{\psi_{j_m, t}} \neq M_{\psi_{j_m, t}} \quad \text{or} \quad M_{\psi_{j_m, t}} \hat{A} B \neq M_{\psi_{j_m, t}}$$

for all $m \in \mathbb{N}$ and all $B \in \mathcal{L}(L_\nu^p)$ with $\|B\| \leq 2 \left(\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x^{-1}\| \|P_\alpha\| + \|Q_\alpha\| \right)$. Taking a suitable subsequence if necessary, we may assume w.l.o.g. that

$$B \hat{A} M_{\psi_{j_m, t}} \neq M_{\psi_{j_m, t}}$$

for all $m \in \mathbb{N}$ (the other case is exactly the same). By Lemma 8, there is a constant C such that $\text{diam}_\beta \text{supp } \psi_{j,t} \leq C$ for all $j \in \mathbb{N}$. We may thus choose a radius R and a sequence of midpoints $(w_m)_{m \in \mathbb{N}}$ with $w_m \rightarrow \partial \mathbb{B}^n$ such that

$$\text{supp } \psi_{j_m, t} \subseteq D(w_m, R) = \{z \in \mathbb{B}^n : \beta(w_m, z) < R\}.$$

As \mathfrak{M} is compact, there exists a subnet (w_{m_γ}) of (w_m) such that $w_{m_\gamma} \rightarrow x$ for some $x \in \mathfrak{M} \setminus \mathbb{B}^n$. Moreover, choosing $\xi = \chi_{D(0, R)}$ in Proposition 19, we obtain a γ_0 such that for all $\gamma \geq \gamma_0$ there is an operator $B_\gamma \in \mathcal{L}(L_\nu^p)$ with $\|B_\gamma\| \leq 2 (\|A_x^{-1}\| \|P_\alpha\| + \|Q_\alpha\|)$ and

$$B_\gamma \hat{A} M_{\chi_{D(w_{m_\gamma}, R)}} = B_\gamma \hat{A} M_{\chi_{D(0, R)} \circ \phi_{w_{m_\gamma}}} = M_{\chi_{D(0, R)} \circ \phi_{w_{m_\gamma}}} = M_{\chi_{D(w_{m_\gamma}, R)}}.$$

Multiplying with $M_{\psi_{j_{m\gamma},t}}$ from the right yields

$$B_\gamma \hat{A} M_{\psi_{j_{m\gamma},t}} = M_{\psi_{j_{m\gamma},t}}$$

for all $\gamma \geq \gamma_0$. This is clearly a contradiction. \square

In the next theorem we show that the converse of Theorem 20 is true as well. In fact, the converse is not limited to Toeplitz operators.

Theorem 21. *Let $A \in \mathcal{L}(A_\nu^p)$ be Fredholm and let (z_γ) be a net in \mathbb{B}^n that tends to $x \in \mathfrak{M} \setminus \mathbb{B}^n$ such that $U_{z_\gamma}^p A(U_{z_\gamma}^q |_{A_\nu^q})^*$ converges $*$ -strongly to $A_x \in \mathcal{L}(A_\nu^p)$. Then A_x is invertible and $\|A_x^{-1}\| \leq \|P_\nu\| \|(A + \mathcal{K}(A_\nu^p))^{-1}\|$. Moreover, if B is a Fredholm regularizer of A , $U_{z_\gamma}^p B(U_{z_\gamma}^q |_{A_\nu^q})^*$ converges $*$ -strongly to $T_{b_x}^{-1} A_x^{-1} T_{b_x}^{-1}$ as $z_\gamma \rightarrow x$.*

Proof. If B is a Fredholm regularizer of A , then $AB - I$ and $BA - I$ are compact and hence $U_{z_\gamma}^p (AB - I)(U_{z_\gamma}^q |_{A_\nu^q})^* \rightarrow 0$ and $U_{z_\gamma}^p (BA - I)(U_{z_\gamma}^q |_{A_\nu^q})^* \rightarrow 0$ $*$ -strongly as $z_\gamma \rightarrow x$ (see [17, Proposition 4.12, Theorem 5.5]). Moreover,

$$\begin{aligned} \|f\| &= \left\| T_{b_{z_\gamma}} T_{b_{z_\gamma}}^{-1} f \right\| \leq \|P_\nu\| \left\| U_{z_\gamma}^p (U_{z_\gamma}^q |_{A_\nu^q})^* f \right\| \\ &\leq \|P_\nu\| \left\| U_{z_\gamma}^p BA(U_{z_\gamma}^q |_{A_\nu^q})^* f \right\| + \|P_\nu\| \left\| U_{z_\gamma}^p (I - BA)(U_{z_\gamma}^q |_{A_\nu^q})^* f \right\| \\ &\leq \|P_\nu\| \|B\| \left\| U_{z_\gamma}^p A(U_{z_\gamma}^q |_{A_\nu^q})^* f \right\| + \|P_\nu\| \left\| U_{z_\gamma}^p (I - BA)(U_{z_\gamma}^q |_{A_\nu^q})^* f \right\| \end{aligned}$$

for every $f \in A_\nu^p$, using that $U_{z_\gamma}^p$ is an isometry. Taking the limit $z_\gamma \rightarrow x$, we obtain $\|f\| \leq \|P_\nu\| \|B\| \|A_x f\|$ for every $f \in A_\nu^p$. This implies that A_x is injective and has a closed range. By the dual argument, we also obtain $\|f\| \leq \|P_\nu\| \|B^*\| \|A_x^* f\|$ for every $f \in A_\nu^q$, which implies that A_x is surjective, hence invertible. Moreover, it shows that $\|A_x^{-1}\| \leq \|P_\nu\| \|B\|$. As this is true for every regularizer B , we obtain $\|A_x^{-1}\| \leq \|P_\nu\| \|(A + \mathcal{K}(A_\nu^p))^{-1}\|$.

Moreover, using $(U_{z_\gamma}^q |_{A_\nu^q})^* T_{b_{z_\gamma}} U_{z_\gamma}^p = I$, we have

$$\begin{aligned} U_{z_\gamma}^p B(U_{z_\gamma}^q |_{A_\nu^q})^* - T_{b_x}^{-1} A_x^{-1} T_{b_x}^{-1} &= U_{z_\gamma}^p B(U_{z_\gamma}^q |_{A_\nu^q})^* T_{b_{z_\gamma}} (A_x - U_{z_\gamma}^p A(U_{z_\gamma}^q |_{A_\nu^q})^*) A_x^{-1} T_{b_{z_\gamma}}^{-1} \\ &\quad + U_{z_\gamma}^p (BA - I)(U_{z_\gamma}^q |_{A_\nu^q})^* A_x^{-1} T_{b_{z_\gamma}}^{-1} \\ &\quad + T_{b_{z_\gamma}}^{-1} A_x^{-1} (T_{b_{z_\gamma}}^{-1} - T_{b_x}^{-1}) + (T_{b_{z_\gamma}}^{-1} - T_{b_x}^{-1}) A_x^{-1} T_{b_x}^{-1} \end{aligned}$$

and all terms on the right-hand side tend $*$ -strongly to 0 as $z_\gamma \rightarrow x$. \square

In particular, we have shown that a Toeplitz operator is Fredholm if and only if all of its limit operators are invertible and their inverses are uniformly bounded. We will state this result in a separate theorem below. But let us first argue why the condition on uniform boundedness is actually redundant. The argument is very similar to the sequence space case, cf. [15].

Let $r_t := \sup_{j \in \mathbb{N}} \text{diam}_\beta \text{supp } \varphi_{j,t}$, where $\varphi_{j,t}$ is defined as usual. By Lemma 8, r_t is finite for every $t \in (0, 1)$. Now, for every $t \in (0, 1)$, $A \in \mathcal{L}(L_\nu^p)$ and every Borel set $F \subseteq \mathbb{B}^n$ we define

$$\nu_t(A|_F) := \inf \{ \|Af\| : f \in L_\nu^p, \|f\| = 1, \text{supp } f \subseteq D(w, r_t) \cap F \text{ for some } w \in \mathbb{B}^n \}$$

and

$$\nu(A|_F) := \inf \{ \|Af\| : f \in L_\nu^p, \|f\| = 1, \text{supp } f \subseteq F \}.$$

Moreover, $\nu(A) := \nu(A|_{\mathbb{B}^n})$.

Proposition 22. For all $A, B \in \mathcal{L}(L_\nu^p)$ and all Borel sets $F \subseteq \mathbb{B}^n$ it holds $|\nu(A|_F) - \nu(B|_F)| \leq \|(A - B)M_{\chi_F}\|$. The same statement also holds if we replace ν by ν_t for some $t \in (0, 1)$.

Proof. We only prove the first claim, but the same proof also works for the second claim. Let $\varepsilon > 0$. Choose $f \in L_\nu^p$ with $\|f\| = 1$, $\text{supp } f \subseteq F$ and $\|Bf\| \leq \nu(B|_F) + \varepsilon$. This implies

$$\nu(A|_F) - \nu(B|_F) - \varepsilon \leq \nu(A|_F) - \|Bf\| \leq \|Af\| - \|Bf\| \leq \|(A - B)f\| \leq \|(A - B)M_{\chi_F}\|.$$

Similarly, $\nu(B|_F) - \nu(A|_F) - \varepsilon \leq \|(A - B)M_{\chi_F}\|$. Since ε was arbitrary, the assertion follows. \square

For $p \leq 2$ and $\alpha = (\frac{2}{p} - 1)(n + 1) + \frac{2\nu}{p}$ we will use the (abuse of) notation $\hat{A}_x := A_x T_{b_x} P_\alpha + Q_\alpha$.

Proposition 23. Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)(n + 1) + \frac{2\nu}{p}$ and $A \in \mathfrak{T}_{p, \nu}$. Then for every $\varepsilon > 0$ there exists a $t \in (0, 1)$ such that for every Borel set $F \subseteq \mathbb{B}^n$ and every operator $B \in \{\hat{A}\} \cup \{\hat{A}_x : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ it holds

$$\nu(B|_F) \leq \nu_t(B|_F) \leq \nu(B|_F) + \varepsilon. \quad (5.3)$$

Proof. The first inequality is clear by definition. For the second inequality we start with a few simple observations. By Theorem 7, \hat{A} is band-dominated. Therefore there is a sequence of band operators $(\hat{A}_n)_{n \in \mathbb{N}}$ that converges to \hat{A} . Choose n sufficiently large such that $\|\hat{A} - \hat{A}_n\| < \frac{\varepsilon}{4}$ and denote the band width of \hat{A}_n by ω . Let $x \in \mathfrak{M}$ and choose a net (z_γ) that converges to x . Then $(U_{z_\gamma} \hat{A}_n U_{z_\gamma})$ is a bounded net and hence there is a subnet of (z_γ) , again denoted by (z_γ) such that $(U_{z_\gamma} \hat{A}_n U_{z_\gamma})$ converges in the weak operator topology as $z_\gamma \rightarrow x$. Let us denote this limit by $(\hat{A}_x)_n$. As $(U_{z_\gamma} \hat{A} U_{z_\gamma}) = (U_{z_\gamma} A U_{z_\gamma}) P_\alpha + Q_\alpha$ converges to $A_x T_{b_x} P_\alpha + Q_\alpha = \hat{A}_x$ in the strong operator topology (see Proposition 18), we obtain that $(U_{z_\gamma} (\hat{A} - \hat{A}_n) U_{z_\gamma})$ converges to $\hat{A}_x - (\hat{A}_x)_n$ in the weak operator topology. This implies

$$\|\hat{A}_x - (\hat{A}_x)_n\| \leq \sup_\gamma \|(U_{z_\gamma} (\hat{A} - \hat{A}_n) U_{z_\gamma})\| = \|\hat{A} - \hat{A}_n\| < \frac{\varepsilon}{4}.$$

Let $f, g \in L^\infty(\mathbb{B}^n)$ with $\text{dist}_\beta(\text{supp } f, \text{supp } g) > \omega$. Then equation (5.1) implies

$$M_f(U_{z_\gamma} \hat{A}_n U_{z_\gamma}) M_g = U_{z_\gamma} M_{f \circ \phi_{z_\gamma}} \hat{A}_n M_{g \circ \phi_{z_\gamma}} U_{z_\gamma} = 0$$

because $\text{dist}_\beta(\text{supp } f \circ \phi_{z_\gamma}, \text{supp } g \circ \phi_{z_\gamma}) = \text{dist}_\beta(\text{supp } f, \text{supp } g) > \omega$. Hence all elements in the net $(U_{z_\gamma} \hat{A}_n U_{z_\gamma})$ have the same band width ω . As $M_f(U_{z_\gamma} \hat{A}_n U_{z_\gamma}) M_g$ converges to $M_f(\hat{A}_x)_n M_g$ in weak operator topology, $(\hat{A}_x)_n$ is also a band operator of band width at most ω .

The strategy now is to prove that there exists a $t \in (0, 1)$ such that for all $F \subseteq \mathbb{B}^n$ and all operators $B \in \{\hat{A}_n\} \cup \{(\hat{A}_x)_n : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ it holds

$$\nu_t(B|_F) \leq \nu(B|_F) + \frac{\varepsilon}{2}$$

and then use Proposition 22. Indeed, suppose that the above is true. Then Proposition 22 implies

$$|\nu(\hat{A}|_F) - \nu(\hat{A}_n|_F)| \leq \|\hat{A} - \hat{A}_n\| < \frac{\varepsilon}{4} \quad \text{and} \quad |\nu(\hat{A}_x|_F) - \nu((\hat{A}_x)_n|_F)| \leq \|\hat{A}_x - (\hat{A}_x)_n\| < \frac{\varepsilon}{4}$$

and

$$|\nu_t(\hat{A}|_F) - \nu_t(\hat{A}_n|_F)| \leq \|\hat{A} - \hat{A}_n\| < \frac{\varepsilon}{4} \quad \text{and} \quad |\nu_t(\hat{A}_x|_F) - \nu_t((\hat{A}_x)_n|_F)| \leq \|\hat{A}_x - (\hat{A}_x)_n\| < \frac{\varepsilon}{4}$$

for all $t \in (0, 1)$ and the proposition follows.

Choose $f \in L^p_\nu$ with $\|f\| = 1$ and $\text{supp } f \subseteq F$ such that

$$\|Bf\| \leq \nu(B|_F) + \frac{\varepsilon}{4}.$$

Let $\varphi_{j,t}$ and $\psi_{j,t}$ be defined as usual. Then by Minkowski's inequality in $\ell^p(\mathbb{N})$, we obtain

$$\begin{aligned} \left(\sum_{j=1}^{\infty} \|BM_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p} &= \left(\sum_{j=1}^{\infty} \|BM_{\varphi_{j,t}^{1/p}} M_{\psi_{j,t}} f\|^p \right)^{1/p} \\ &\leq \left(\sum_{j=1}^{\infty} \|M_{\varphi_{j,t}^{1/p}} Bf\|^p \right)^{1/p} + \left(\sum_{j=1}^{\infty} \|M_{\varphi_{j,t}^{1/p}} BM_{1-\psi_{j,t}} f\|^p \right)^{1/p} \\ &\quad + \left(\sum_{j=1}^{\infty} \|[B, M_{\varphi_{j,t}^{1/p}}] M_{\psi_{j,t}} f\|^p \right)^{1/p}. \end{aligned}$$

The first term is just $\|Bf\|$ (recall that $\sum_{j=1}^{\infty} |\varphi_{j,t}(z)| = 1$ for all $z \in \mathbb{B}^n$, $t \in (0, 1)$). The second term vanishes for $\text{dist}_\beta(\text{supp } \varphi_{j,t}, \text{supp}(1 - \psi_{j,t})) > \omega$ as B has band width ω . The third term can be estimated as

$$\begin{aligned} \left(\sum_{j=1}^{\infty} \|[B, M_{\varphi_{j,t}^{1/p}}] M_{\psi_{j,t}} f\|^p \right)^{1/p} &\leq \sup_{j \in \mathbb{N}} \|[B, M_{\varphi_{j,t}^{1/p}}]\| \left(\sum_{j=1}^{\infty} \|M_{\psi_{j,t}} f\|^p \right)^{1/p} \\ &\leq N^{1/p} \sup_{j \in \mathbb{N}} \|[B, M_{\varphi_{j,t}^{1/p}}]\| \end{aligned} \tag{5.4}$$

by Proposition 5. Observe that the functions $\varphi_{j,t}^{1/p}$ satisfy the assumptions in Lemma 12. Indeed, let $U, V \subset [0, 1]$ with $\text{dist}(U, V) > 0$ and $w_{j,t} \in (\varphi_{j,t}^{1/p})^{-1}(U)$, $z_{j,t} \in (\varphi_{j,t}^{1/p})^{-1}(V)$. Clearly, we have $\text{dist}(U^p, V^p) > 0$ as well and therefore

$$\beta(z_{j,t}, w_{j,t}) \geq \frac{1}{6Nt} |\varphi_{j,t}(z_{j,t}) - \varphi_{j,t}(w_{j,t})| \geq \frac{1}{6Nt} \text{dist}(U^p, V^p) \rightarrow \infty$$

as $t \rightarrow 0$. Lemma 12 thus implies that for every $\delta > 0$ there is a $t > 0$ such that

$$\left(\sum_{j=1}^{\infty} \|[B, M_{\varphi_{j,t}^{1/p}}] M_{\psi_{j,t}} f\|^p \right)^{1/p} \leq \delta \|B\|.$$

As $\|B\| \leq \|\hat{A}\| + \frac{\varepsilon}{4}$ for all $B \in \{\hat{A}_n\} \cup \{(\hat{A}_x)_n : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ by the above, we may choose $\delta > 0$ such that $\delta \|B\| \leq \frac{\varepsilon}{4}$ for all B . Therefore

$$\left(\sum_{j=1}^{\infty} \|BM_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p} \leq \|Bf\| + \frac{\varepsilon}{4} \leq \nu(B|_F) + \frac{\varepsilon}{2} = \left(\nu(B|_F) + \frac{\varepsilon}{2} \right) \left(\sum_{j=1}^{\infty} \|M_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p}.$$

This implies, in particular, that there exists a $j \in \mathbb{N}$ such that

$$\|BM_{\varphi_{j,t}^{1/p}} f\| \leq \left(\nu(B|_F) + \frac{\varepsilon}{2} \right) \|M_{\varphi_{j,t}^{1/p}} f\|$$

for sufficiently small t . Since $\text{supp} \left(M_{\varphi_{j,t}^{1/p}} f \right) \subseteq \text{supp} \varphi_{j,t} \subseteq D(w, r_t)$ for some $w \in \mathbb{B}^n$ by definition, this implies $\nu_t(B|_F) \leq \nu(B|_F) + \frac{\varepsilon}{2}$ for all $B \in \{\hat{A}_n\} \cup \{(\hat{A}_x)_n : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$. As t is chosen independently of F (as it is chosen independently of f) and B , the assertion follows. \square

The next lemma shows some kind of shift invariance of the operator spectrum. This will allow us to shift functions to the right place in the subsequent lemma.

Lemma 24. *Let $p \leq 2$, $\alpha = \left(\frac{2}{p}-1\right)(n+1) + \frac{2\nu}{p}$, $A \in \mathfrak{T}_{p,\nu}$ and $f \in L^p_\nu$ with $\text{supp} f \subseteq D(w, r)$ for some $w \in \mathbb{B}^n$ and $r > 0$. Then for every $x \in \mathfrak{M} \setminus \mathbb{B}^n$ there exist $y \in \mathfrak{M} \setminus \mathbb{B}^n$ and $g \in L^p_\nu$ with $\|g\| = \|f\|$ and $\text{supp} g \subseteq D(0, r)$ such that $\|\hat{A}_x f\| = \|\hat{A}_y g\|$. Moreover, $\nu(\hat{A}_y|_{D(0, r+\beta(0, w))}) \leq \nu(\hat{A}_x|_{D(0, r)})$.*

Proof. A direct computation yields

$$(U_z^p U_w^p f)(\zeta) = (f \circ \phi_w \circ \phi_z)(\zeta) \frac{(1 - |\phi_z(w)|^2)^{(\nu+n+1)/p}}{(1 - \langle \zeta, \phi_z(w) \rangle)^{2(\nu+n+1)/p}} \left(\frac{1 - \langle w, z \rangle}{|1 - \langle w, z \rangle|} \right)^{2(\nu+n+1)/p}$$

for all $w, z \in \mathbb{B}^n$. Using Cartan's theorem, i.e. $\phi_w \circ \phi_z = V \circ \phi_{\phi_z(w)}$ for some unitary map $V: \mathbb{C}^n \rightarrow \mathbb{C}^n$, we get

$$U_z^p U_w^p = \left(\frac{1 - \langle w, z \rangle}{|1 - \langle w, z \rangle|} \right)^{2(\nu+n+1)/p} U_{\phi_z(w)}^p V_*,$$

where $V_* f := f \circ V$ is a composition operator and, by taking inverses, also

$$U_w^p U_z^p = \left(\frac{1 - \langle z, w \rangle}{|1 - \langle w, z \rangle|} \right)^{2(\nu+n+1)/p} V_*^{-1} U_{\phi_z(w)}^p.$$

Choose a net (z_γ) that converges to $x \in \mathfrak{M} \setminus \mathbb{B}^n$. Then

$$\begin{aligned} U_w^p U_{z_\gamma}^p A U_{z_\gamma}^p U_w^p|_{A_\nu^p} &= \left(\frac{1 - \langle z_\gamma, w \rangle}{|1 - \langle w, z_\gamma \rangle|} \right)^{2(\nu+n+1)/p} \left(\frac{1 - \langle w, z_\gamma \rangle}{|1 - \langle w, z_\gamma \rangle|} \right)^{2(\nu+n+1)/p} V_*^{-1} U_{\phi_{z_\gamma}(w)}^p A \\ &\quad \cdot U_{\phi_{z_\gamma}(w)}^p V_*|_{A_\nu^p} \\ &= V_*^{-1} U_{\phi_{z_\gamma}(w)}^p A U_{\phi_{z_\gamma}(w)}^p V_*|_{A_\nu^p}, \end{aligned}$$

where V_* of course depends on w and z_γ . As the unitary group of \mathbb{C}^n is compact, we may assume that V converges to some unitary map \tilde{V} and hence $V_*|_{A_\nu^p}$ converges strongly to $\tilde{V}_*|_{A_\nu^p}$ and $V_*^{-1}|_{A_\nu^p}$ converges strongly to $\tilde{V}_*^{-1}|_{A_\nu^p}$ as $z_\gamma \rightarrow x$. Similarly, using Proposition 18, we may assume that

$$U_{\phi_{z_\gamma}(w)}^p A U_{\phi_{z_\gamma}(w)}^p |_{A_\nu^p} = U_{\phi_{z_\gamma}(w)}^p A (U_{\phi_{z_\gamma}(w)}^q |_{A_\nu^q})^* (U_{\phi_{z_\gamma}(w)}^q |_{A_\nu^q})^* U_{\phi_{z_\gamma}(w)}^p |_{A_\nu^p} = A_{\phi_{z_\gamma}(w)} T_{b_{\phi_{z_\gamma}(w)}}$$

converges strongly to $A_y T_{b_y}$ for some $y \in \mathfrak{M}$. Since $\phi_{z_\gamma}(w) \rightarrow \partial \mathbb{B}^n$ as $z_\gamma \rightarrow \partial \mathbb{B}^n$, it is clear that $y \in \mathfrak{M} \setminus \mathbb{B}^n$. As the limit of a strongly convergent net is unique and $\lim_{z_\gamma \rightarrow x} U_{z_\gamma}^p A U_{z_\gamma}^p = A_x T_{b_x}$, we obtain

$$U_w^p A_x T_{b_x} U_w^p |_{A_\nu^p} = \tilde{V}_*^{-1} A_y T_{b_y} \tilde{V}_* |_{A_\nu^p}.$$

Since P_α commutes with both U_w^p and \tilde{V}_* (the former was shown in the proof of Proposition 19, the latter is immediate from the definition of P_α and \tilde{V}_*), this implies

$$U_w^p \hat{A}_x U_w^p = \tilde{V}_*^{-1} \hat{A}_y \tilde{V}_*.$$

Now let $f \in L_\nu^p$ with $\text{supp } f \subseteq D(w, r)$ for some $w \in \mathbb{B}^n$ and $r > 0$. Set $g := \tilde{V}_* U_w^p f$. Then clearly $\|g\| = \|f\|$ and $\|\hat{A}_x f\| = \|\hat{A}_y g\|$. As $D(0, r)$ is invariant under \tilde{V} , it remains to show that $\text{supp } U_w^p f \subseteq D(0, r)$. But this is clear since $\phi_w(D(0, r)) = D(w, r)$.

For the second assertion consider $f \in L_\nu^p$ with $\text{supp } f \subseteq D(0, r)$. Then $g := \tilde{V}_* U_w^p f$ satisfies $\|g\| = \|f\|$, $\|\hat{A}_x f\| = \|\hat{A}_y g\|$ and $\text{supp } g \subseteq \tilde{V}^{-1}(\phi_w^{-1}(D(0, r))) \subseteq D(0, r + \beta(0, w))$ as above. \square

In the next lemma we show that the infimum $\inf \left\{ \nu(\hat{A}_x) : x \in \mathfrak{M} \setminus \mathbb{B}^n \right\}$ is actually attained by a certain limit operator \hat{A}_y . The proof is very much the same as the proofs of [13, Theorem 3.2] and [15, Theorem 8] and is based on a miraculous procedure invented by Markus Seidel. We therefore refer to [15] for a helpful illustration of the main idea.

Lemma 25. *Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)g + \frac{2\nu}{p}$ and $A \in \mathfrak{T}_{p, \nu}$. Then there exists a $y \in \mathfrak{M} \setminus \mathbb{B}^n$ with*

$$\nu(\hat{A}_y) = \inf \left\{ \nu(\hat{A}_x) : x \in \mathfrak{M} \setminus \mathbb{B}^n \right\}.$$

Proof. Recall $r_t = \sup_{j \in \mathbb{N}} \text{diam}_\beta \text{supp } \varphi_{j, t}$. Proposition 23 yields a sequence $(t_k)_{k \in \mathbb{N}_0}$ with $r_{t_{k+1}} > 2r_{t_k}$ and

$$\nu_{t_k}(B|_F) \leq \nu(B|_F) + 2^{-(k+1)}$$

for all $k \in \mathbb{N}_0$, $F \subseteq \mathbb{B}^n$ and $B \in \{\hat{A}\} \cup \{\hat{A}_x : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$. Choose a sequence $(x_n)_{n \in \mathbb{N}}$ in $\mathfrak{M} \setminus \mathbb{B}^n$ such that

$$\lim_{n \rightarrow \infty} \nu(\hat{A}_{x_n}) = \inf \left\{ \nu(\hat{A}_x) : x \in \mathfrak{M} \setminus \mathbb{B}^n \right\}.$$

Now, for every $n \in \mathbb{N}$ there exists an $f_n^0 \in L_\nu^p$ with $\|f_n^0\| = 1$, $\text{supp } f_n^0$ contained in some $D(w, r_{t_n})$ and

$$\left\| \hat{A}_{x_n} f_n^0 \right\| \leq \nu_{t_n}(\hat{A}_{x_n}) + 2^{-(n+1)} \leq \nu(\hat{A}_{x_n}) + 2^{-n}.$$

By Lemma 24, we can choose a $y_n^0 \in \mathfrak{M} \setminus \mathbb{B}^n$ and a $g_n^0 \in L_\nu^p$ with $\|g_n^0\| = 1$ and $\text{supp } g_n^0 \subseteq D(0, r_{t_n})$ such that

$$\left\| \hat{A}_{y_n^0} g_n^0 \right\| = \left\| \hat{A}_{x_n} f_n^0 \right\| \leq \nu(\hat{A}_{x_n}) + 2^{-n}.$$

Furthermore, for every $n \in \mathbb{N}$ we can find $f_n^1 \in L_\nu^p$ with $\|f_n^1\| = 1$, $\text{supp } f_n^1 \subseteq D(w, r_{t_{n-1}}) \cap D(0, r_{t_n})$ for some $w \in D(0, r_{t_n} + r_{t_{n-1}})$ and

$$\left\| \hat{A}_{y_n^0} f_n^1 \right\| \leq \nu_{t_{n-1}}(\hat{A}_{y_n^0} |_{D(0, r_{t_n})}) + 2^{-n} \leq \nu(\hat{A}_{y_n^0} |_{D(0, r_{t_n})}) + 2^{-n+1}.$$

Thus, using Lemma 24 again, we can choose $y_n^1 \in \mathfrak{M} \setminus \mathbb{B}^n$ and $g_n^1 \in L_\nu^p$ with $\|g_n^1\| = 1$ and $\text{supp } g_n^1 \subseteq D(0, r_{t_{n-1}})$ such that

$$\left\| \hat{A}_{y_n^1} g_n^1 \right\| = \left\| \hat{A}_{y_n^0} f_n^1 \right\| \leq \nu(\hat{A}_{y_n^0} |_{D(0, r_{t_n})}) + 2^{-n+1}.$$

In particular,

$$\nu(\hat{A}_{y_n^1} |_{D(0, r_{t_{n-1}})}) \leq \nu(\hat{A}_{y_n^0} |_{D(0, r_{t_n})}) + 2^{-n+1} \leq \nu(\hat{A}_{x_n}) + 2^{-n+1} + 2^{-n}.$$

Repeating this argument, we obtain $y_n^k \in \mathfrak{M} \setminus \mathbb{B}^n$, $w \in D(0, r_{t_{n-k+1}} + r_{t_{n-k}})$ and $f_n^k \in L_\nu^p$ with $\|f_n^k\| = 1$, $\text{supp } f_n^k \subseteq D(w, r_{t_{n-k}}) \cap D(0, r_{t_{n-k+1}})$ and

$$\left\| \hat{A}_{y_n^{k-1}} f_n^k \right\| \leq \nu_{t_{n-k}}(\hat{A}_{y_n^{k-1}} |_{D(0, r_{t_{n-k+1}})}) + 2^{-n+k-1} \leq \nu(\hat{A}_{y_n^{k-1}} |_{D(0, r_{t_{n-k+1}})}) + 2^{-n+k}$$

for $k = 2, \dots, n$. Moreover, we obtain $g_n^k \in L_\nu^p$ with $\|g_n^k\| = 1$ and $\text{supp } g_n^k \subseteq D(0, r_{t_{n-k}})$ such that

$$\left\| \hat{A}_{y_n^k} g_n^k \right\| = \left\| \hat{A}_{y_n^{k-1}} f_n^k \right\| \leq \nu(\hat{A}_{y_n^{k-1}} |_{D(0, r_{t_{n-k+1}})}) + 2^{-n+k}.$$

In particular,

$$\begin{aligned} \nu(\hat{A}_{y_n^k} |_{D(0, r_{t_{n-k}})}) &\leq \nu(\hat{A}_{y_n^{k-1}} |_{D(0, r_{t_{n-k+1}})}) + 2^{-n+k} \leq \dots \leq \nu(\hat{A}_{y_n^0} |_{D(0, r_{t_n})}) + 2^{-n+k} + \dots + 2^{-n+1} \\ &\leq \nu(\hat{A}_{x_n}) + 2^{-n+k} + \dots + 2^{-n+1} + 2^{-n} \leq \nu(\hat{A}_{x_n}) + 2^{-n+k+1}. \end{aligned}$$

Furthermore, by repeatedly applying the second part of Lemma 24 and collecting all the shifts being made during the process above, we get

$$\begin{aligned} \nu(\hat{A}_{y_n^{n-l}} |_{D(0, r_{t_l})}) &\geq \nu(\hat{A}_{y_n^{n-l+1}} |_{D(0, r_{t_l} + r_{t_{l-1}})}) = \nu(\hat{A}_{y_n^{n-l+1}} |_{D(0, 2r_{t_l} + r_{t_{l-1}})}) \\ &\geq \nu(\hat{A}_{y_n^{n-l+2}} |_{D(0, 2r_{t_l} + r_{t_{l-1}} + r_{t_{l-2}})}) = \nu(\hat{A}_{y_n^{n-l+2}} |_{D(0, 2r_{t_l} + 2r_{t_{l-1}} + r_{t_{l-2}})}) \\ &\geq \dots \geq \nu(\hat{A}_{y_n^n} |_{D(0, 2r_{t_l} + 2r_{t_{l-1}} + 2r_{t_{l-2}} + \dots + 2r_{t_1} + r_{t_0})}) \geq \nu(\hat{A}_{y_n^n} |_{D(0, 4r_{t_l})}) \end{aligned}$$

for fixed $l \leq n$, using $r_{t_{k+1}} > 2r_{t_k}$ for all k .

Now set $y_n := y_n^n$. By Proposition 18, the sequence $(A_{y_n} T_{b_{y_n}})_{n \in \mathbb{N}}$ has a strongly convergent subnet, denoted by $(A_{y_{n_\gamma}} T_{b_{y_{n_\gamma}}})$, that converges to $A_y T_{b_y}$ for some $y \in \mathfrak{M} \setminus \mathbb{B}^n$. Now since $\overline{D(0, 4r_{t_l})}$ is a compact set, $P_\alpha M_{\mathcal{X}_{D(0, 4r_{t_l})}} \in \mathcal{L}(L_\nu^p)$ is a compact operator (see Proposition 15) and therefore

$$\left\| (\hat{A}_{y_{n_\gamma}} - \hat{A}_y) M_{\mathcal{X}_{D(0, 4r_{t_l})}} \right\| = \left\| (A_{y_{n_\gamma}} T_{b_{y_{n_\gamma}}} - A_y T_{b_y}) P_\alpha M_{\mathcal{X}_{D(0, 4r_{t_l})}} \right\| \rightarrow 0.$$

This also implies $\nu(\hat{A}_{y_{n_\gamma}} |_{D(0, 4r_{t_l})}) \rightarrow \nu(\hat{A}_y |_{D(0, 4r_{t_l})})$ by Proposition 22. Thus

$$\begin{aligned} \nu(\hat{A}_y) &\leq \nu(\hat{A}_y |_{D(0, 4r_{t_l})}) = \lim_\gamma \nu(\hat{A}_{y_{n_\gamma}} |_{D(0, 4r_{t_l})}) \leq \lim_\gamma \nu(\hat{A}_{y_{n_\gamma}^{n-l}} |_{D(0, r_{t_l})}) \leq \lim_\gamma \nu(\hat{A}_{x_{n_\gamma}}) + 2^{-l+1} \\ &= \inf \left\{ \nu(\hat{A}_x) : x \in \mathfrak{M} \setminus \mathbb{B}^n \right\} + 2^{-l+1}. \end{aligned}$$

Sending $l \rightarrow \infty$, we obtain $\nu(\hat{A}_y) = \inf \left\{ \nu(\hat{A}_x) : x \in \mathfrak{M} \setminus \mathbb{B}^n \right\}$ as claimed. \square

Now we can summarize this section with the main result of this paper.

Theorem A. *Let $A \in \mathfrak{F}_{p,\nu}$. Then the following are equivalent:*

- (i) A is Fredholm,
- (ii) A_x is invertible and $\|A_x^{-1}\| \leq \|P_\nu\| \|(A + \mathcal{K}(A_\nu^p))^{-1}\|$ for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$,
- (iii) A_x is invertible for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$ and $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x^{-1}\| < \infty$,
- (iv) A_x is invertible for all $x \in \mathfrak{M} \setminus \mathbb{B}^n$,

Proof. That (i) implies (ii) follows from Proposition 18 and Theorem 21, whereas (ii) obviously implies (iii) and (iii) implies (iv). It remains to show that (iv) implies (i). By duality, it suffices to show the case $p \leq 2$. If A_x is invertible, then $(\hat{A}_x)^{-1} = T_{b_x}^{-1} A_x^{-1} P_\alpha + Q_\alpha$, which implies that $(\hat{A}_x)^{-1}$ is also invertible. Now observe that $\nu(B) = \|B^{-1}\|^{-1} > 0$ whenever an operator $B \neq 0$ is invertible (see e.g. [14, Lemma 2.35] for a quick proof). Thus by Lemma 25, $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|\hat{A}_x^{-1}\| < \infty$.

Moreover, if \hat{A}_x is invertible, then $T_{b_x} \hat{A}_x^{-1}|_{A_\nu^p}$ is the inverse of A_x and so $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x^{-1}\| < \infty$.

Therefore (iv) implies (iii). That (iii) implies (i) is Theorem 20. \square

Of course, we get the following corollary for the essential spectrum:

Corollary 26. *Let $A \in \mathfrak{F}_{p,\nu}$. Then*

$$\text{sp}_{\text{ess}}(A) = \bigcup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \text{sp}(A_x).$$

6 Norm estimates (unit ball)

In this section we prove a similar result for the essential norm of an operator $A \in \mathfrak{F}_{p,\nu}$. For the most part this is just a modification of the proofs in Section 5. Recall that $r_t = \sup_{j \in \mathbb{N}} \text{diam}_\beta \text{supp } \varphi_{j,t}$.

For $t \in (0, 1)$, $\alpha \geq \nu$, $A \in \mathcal{L}(A_\nu^p)$ and a Borel set $F \subseteq \mathbb{B}^n$ we define

$$\| \|AP_\alpha|_F \|_t := \sup \{ \|AP_\alpha f\| : f \in L_\nu^p, \|f\| = 1, \text{supp } f \subseteq D(w, r_t) \cap F \text{ for some } w \in \mathbb{B}^n \}$$

and

$$\|AP_\alpha|_F\| := \sup \{ \|AP_\alpha f\| : f \in L_\nu^p, \|f\| = 1, \text{supp } f \subseteq F \}.$$

Proposition 27. *Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)(n + 1) + \frac{2\nu}{p}$ and $A \in \mathfrak{F}_{p,\nu}$. Then for every $\varepsilon > 0$ there exists a $t \in (0, 1)$ such that for all Borel Sets $F \subseteq \mathbb{B}^n$ and all operators B in the set*

$$\{A\} \cup \{A_x T_{b_x} : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$$

it holds

$$\|BP_\alpha|_F\| \geq \| \|BP_\alpha|_F \|_t \geq \|BP_\alpha|_F\| - \varepsilon.$$

Proof. The first inequality is clear by definition. For the second inequality we proceed as in the proof of Proposition 23. Let $A \in \mathfrak{T}_{p,\nu}$ and fix $\varepsilon > 0$. Observe that $AP_\alpha = \hat{A} - Q_\alpha$ is band-dominated by Theorem 7. We may thus choose a band operator A_n of band width ω such that $\|AP_\alpha - A_n\| < \frac{\varepsilon}{4}$. Let $x \in \mathfrak{M}$. Now as in the proof of Proposition 23, there is a net (z_γ) converging to x such that the net $(U_{z_\gamma} A_n U_{z_\gamma})$ converges in weak operator topology. This limit will be denoted by $(A_x)_n$. It follows that $(A_x)_n$ is a band operator of band width at most ω and $\|A_x - (A_x)_n\| < \frac{\varepsilon}{4}$. Let $B \in \{A_n\} \cup \{(A_x)_n : x \in \mathfrak{M} \setminus \mathbb{B}^n\}$ and choose $f \in L_\nu^p$ with $\|f\| = 1$ and $\text{supp } f \subseteq F$ such that

$$\|BP_\alpha f\| \geq \|BP_\alpha|_F\| - \frac{\varepsilon}{4}.$$

We can apply the same reasoning as in the proof of Proposition 23 (just reverse the inequalities and use the reverse triangle inequality) to obtain

$$\left(\sum_{j=1}^{\infty} \|BP_\alpha M_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p} \geq \|BP_\alpha f\| - \frac{\varepsilon}{4}$$

for sufficiently small t . This implies

$$\left(\|BP_\alpha|_F\| - \frac{\varepsilon}{2} \right) \left(\sum_{j=1}^{\infty} \|M_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p} \leq \left(\sum_{j=1}^{\infty} \|BP_\alpha M_{\varphi_{j,t}^{1/p}} f\|^p \right)^{1/p}.$$

Thus there exists a $j \in \mathbb{N}$ such that

$$\left(\|BP_\alpha|_F\| - \frac{\varepsilon}{2} \right) \|M_{\varphi_{j,t}^{1/p}} f\| \leq \|BP_\alpha M_{\varphi_{j,t}^{1/p}} f\|$$

for sufficiently small t . Of course, this implies $\|BP_\alpha|_F\| - \varepsilon \leq \|BP_\alpha|_F\|_t$. As t does not depend on f or B , the result follows as in the proof of Proposition 23. \square

Theorem 28. *Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)(n + 1) + \frac{2\nu}{p}$ and $A \in \mathfrak{T}_{p,\nu}$. Then*

$$\frac{1}{\|P_\alpha\| \|P_\nu\|} \|A + \mathcal{K}(A_\nu^p)\| \leq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\| \leq \|P_\nu\| \|A + \mathcal{K}(A_\nu^p)\|.$$

Proof. Let $x \in \mathfrak{M} \setminus \mathbb{B}^n$, $K \in \mathcal{K}(A_\nu^p)$ and choose a net (z_γ) in \mathbb{B}^n that converges to x . As K is compact, we get $K_x = 0$ by [17, Proposition 4.12, Theorem 5.5]. Banach-Steinhaus thus implies

$$\|A_x\| = \|A_x + K_x\| = \|(A + K)_x\| \leq \sup_\gamma \left\| U_{z_\gamma}^p (A + K) (U_{z_\gamma}^q|_{A_\nu^q})^* \right\| \leq \|P_\nu\| \|A + K\|,$$

where we used that $U_{z_\gamma}^p$ and $(U_{z_\gamma}^q)^*$ are isometries. Since this is true for all $K \in \mathcal{K}(A_\nu^p)$ and $x \in \mathfrak{M} \setminus \mathbb{B}^n$, we get

$$\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\| \leq \|P_\nu\| \|A + \mathcal{K}(A_\nu^p)\|.$$

For the other inequality observe that

$$\|A + \mathcal{K}(A_\nu^p)\| \leq \inf_{K \in \mathcal{K}(L_\nu^p, A_\nu^p)} \|AP_\alpha + K\|.$$

Indeed,

$$\|AP_\alpha + K\| = \sup_{\|f\|=1} \|(AP_\alpha + K)f\| \geq \sup_{\substack{f \in A_\nu^p, \\ \|f\|=1}} \|(AP_\alpha + K)f\| = \sup_{\substack{f \in A_\nu^p, \\ \|f\|=1}} \|(A + K)f\| = \|A + K\|_{A_\nu^p}$$

for every $K \in \mathcal{K}(L_\nu^p, A_\nu^p)$. We will now show

$$\inf_{K \in \mathcal{K}(L_\nu^p, A_\nu^p)} \|AP_\alpha + K\| \leq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\|, \quad (6.1)$$

which obviously implies the desired inequality. So assume that (6.1) is violated, i.e. that there is an $\varepsilon > 0$ such that

$$\inf_{K \in \mathcal{K}(L_\nu^p, A_\nu^p)} \|AP_\alpha + K\| > \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \varepsilon.$$

In particular,

$$\|AP_\alpha|_{\mathbb{B}^n \setminus D(0,s)}\| = \|AP_\alpha M_{1-\chi_{D(0,s)}}\| = \|AP_\alpha - AP_\alpha M_{\chi_{D(0,s)}}\| > \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \varepsilon$$

for all $s > 0$ since $P_\alpha M_{\chi_{D(0,s)}} \in \mathcal{K}(L_\nu^p, A_\nu^p)$ by Proposition 15. Now, by Proposition 27, there is a $t \in (0, 1)$ such that for all $s > 0$ we have

$$\|AP_\alpha|_{\mathbb{B}^n \setminus D(0,s)}\|_t \geq \|AP_\alpha|_{\mathbb{B}^n \setminus D(0,s)}\| - \frac{\varepsilon}{2} > \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \frac{\varepsilon}{2}.$$

In particular, for every $s > 0$ we get a $w_s \in \mathbb{B}^n$ such that

$$\|AP_\alpha M_{\chi_{D(w_s, r_t)}}\| \geq \|AP_\alpha M_{\chi_{D(w_s, r_t)} \setminus D(0,s)}\| > \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \frac{\varepsilon}{2}.$$

It is clear that $w_s \rightarrow \partial \mathbb{B}^n$ as $s \rightarrow \infty$. Since $M_{\chi_{D(w_s, r_t)}} = U_{w_s}^p M_{\chi_{D(0, r_t)}} U_{w_s}^p$ and $P_\alpha U_{w_s}^p = U_{w_s}^p P_\alpha$ (see the proof of Proposition 19), we get

$$\|U_{w_s}^p A U_{w_s}^p P_\alpha M_{\chi_{D(0, r_t)}}\| > \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \frac{\varepsilon}{2}.$$

As \mathfrak{M} is compact, (w_s) has a convergent subnet, denoted again by (w_s) , converging to some $x \in \mathfrak{M} \setminus \mathbb{B}^n$. Thus $U_{w_s}^p A U_{w_s}^p|_{A_\nu^p}$ converges strongly to $A_x T_{b_x}$ and hence

$$\|U_{w_s}^p A U_{w_s}^p P_\alpha M_{\chi_{D(0, r_t)}}\| \rightarrow \|A_x T_{b_x} P_\alpha M_{\chi_{D(0, r_t)}}\|$$

since $P_\alpha M_{\chi_{D(0, r_t)}}$ is compact. This yields

$$\|A_x T_{b_x} P_\alpha M_{\chi_{D(0, r_t)}}\| \geq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\| + \frac{\varepsilon}{2},$$

which is certainly a contradiction. Thus $\inf_{K \in \mathcal{K}(L_\nu^p, A_\nu^p)} \|AP_\alpha + K\| \leq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x T_{b_x} P_\alpha\|$. \square

For $p \geq 2$ we get the following corollary by using the adjoint of P_α instead.

Corollary 29. Let $p \geq 2$, $\alpha = (1 - \frac{2}{p})(n+1) + 2\nu(1 - \frac{1}{p})$ and $A \in \mathfrak{T}_{p,\nu}$. Then

$$\frac{1}{\|P_\alpha^*\| \|P_\nu\|} \|A + \mathcal{K}(A_\nu^p)\| \leq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\| \leq \|P_\nu\| \|A + \mathcal{K}(A_\nu^p)\|.$$

Lemma 30. Let $p \leq 2$, $\alpha = (\frac{2}{p} - 1)(n+1) + \frac{2\nu}{p}$ and $A \in \mathfrak{T}_{p,\nu}$. Then there exists a $y \in \mathfrak{M} \setminus \mathbb{B}^n$ with

$$\|A_y T_{b_y} P_\alpha\| = \sup \{ \|A_x T_{b_x} P_\alpha\| : x \in \mathfrak{M} \setminus \mathbb{B}^n \}.$$

Proof. Replacing all ν by $\|\cdot\|$ and ν_t by $\|\cdot\|_t$ in the proof of Lemma 25 and using Proposition 27 instead of Proposition 23 one easily obtains a proof of Lemma 30 (see also [13, Theorem 3.2]). \square

Let us summarize these results in a final theorem. This may be seen as an analogue of Theorem A and a slight improvement of [17, Theorem 5.2]. Unfortunately this result is far less complete than in the case of the spectrum and thus leaves some questions open: Are $\|A + \mathcal{K}(A_\nu^p)\|$ and $\sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\|$ equal also for $p \neq 2$? And is the supremum also a maximum in case $p \neq 2$?

Theorem 31. Let $A \in \mathfrak{T}_{p,\nu}$ and $\alpha = \left\lfloor \frac{2}{p} - 1 \right\rfloor (\nu + n + 1) + \nu$. Then

$$\frac{1}{\|P_\alpha\| \|P_\nu\|} \|A + \mathcal{K}(A_\nu^p)\| \leq \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\| \leq \|P_\nu\| \|A + \mathcal{K}(A_\nu^p)\|,$$

where the norm of P_α is taken on $L_\nu^{\min\{p,q\}}$ ($\frac{1}{p} + \frac{1}{q} = 1$ as usual). Moreover,

$$\|A + \mathcal{K}(A_\nu^p)\| = \max_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\|$$

if $p = 2$.

Proof. The first statement is a combination of Theorem 28 and Corollary 29. In case $p = 2$ we have $\|P_\alpha\| = \|P_\nu\| = 1$ and therefore

$$\|A + \mathcal{K}(A_\nu^p)\| = \sup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} \|A_x\|$$

(see also [17, Theorem 5.6]). Moreover, as the norms of $A_x T_{b_x} P_\alpha = A_x P_\nu$ and A_x coincide in this case, the second statement follows from Lemma 30. \square

7 Application to symbols of vanishing oscillation

In this section we apply Theorem A to the case of functions of vanishing oscillation. Even though the results obtained in this section are not completely new, it is worth mentioning that they are special cases of Theorem A.

For $z \in \mathbb{B}^n$ and a bounded continuous (BC) function $f: \mathbb{B}^n \rightarrow \mathbb{C}$ we define

$$\text{Osc}_z(f) := \sup \{ |f(z) - f(w)| : w \in \mathbb{B}^n, \beta(z, w) \leq 1 \}$$

and $\text{VO}_\partial(\mathbb{B}^n) := \left\{ f \in \text{BC}(\mathbb{B}^n) : \lim_{|z| \rightarrow 1} \text{Osc}_z(f) = 0 \right\}$. Note that

$$C(\overline{\mathbb{B}^n}) \subset \text{VO}_\partial(\mathbb{B}^n) \subset \text{BUC}(\mathbb{B}^n).$$

Applying Corollary 26 to Toeplitz operators with symbol in $\text{VO}_\partial(\mathbb{B}^n)$, we obtain the following result:

Proposition 32. *Let $f \in \text{VO}_\partial(\mathbb{B}^n)$. Then*

$$\text{sp}_{\text{ess}}(T_f) = f(\partial\mathbb{B}^n),$$

where $f(\partial\mathbb{B}^n)$ denotes the set of limit points of f as $z \rightarrow \partial\mathbb{B}^n$.

In case f is contained in $C(\overline{\mathbb{B}^n})$, $f(\partial\mathbb{B}^n)$ is just the image of $f|_{\partial\mathbb{B}^n}$ and we obtain the classical result mentioned in the introduction.

Proof. Let (z_γ) be a net in \mathbb{B}^n converging to some $x \in \mathfrak{M} \setminus \mathbb{B}^n$ and consider $f \in \text{VO}_\partial(\mathbb{B}^n) \subset \text{BUC}(\mathbb{B}^n)$. Then we have $(f \circ \phi_{z_\gamma})(0) = f(z_\gamma) \rightarrow x(f)$. Moreover, since $\beta(\phi_{z_\gamma}(0), \phi_{z_\gamma}(w)) = \beta(0, w)$, we get

$$|(f \circ \phi_{z_\gamma})(0) - (f \circ \phi_{z_\gamma})(w)| \leq \text{Osc}_{z_\gamma}(f) \rightarrow 0$$

if $\beta(0, w) \leq 1$. Thus $(f \circ \phi_{z_\gamma})(w) \rightarrow x(f)$ uniformly on $\{w \in \mathbb{B}^n : \beta(0, w) \leq 1\}$. By repeating this argument and using that $|\phi_{z_\gamma}(w)| \rightarrow 1$ if $|z_\gamma| \rightarrow 1$, this generalizes to arbitrary compact subsets of \mathbb{B}^n . Therefore $f \circ \phi_{z_\gamma}$ converges uniformly on compact sets to the constant function $x(f)$. Using [17, p. 222], we obtain

$$(T_f)_x = \text{s-}\lim_{z_\gamma \rightarrow x} U_{z_\gamma}^p T_f (U_{z_\gamma}^q |_{A_\nu^q})^* = \text{s-}\lim_{z_\gamma \rightarrow x} T_{b_{z_\gamma}}^{-1} T_{(f \circ \phi_{z_\gamma})_{b_{z_\gamma}}} T_{b_{z_\gamma}}^{-1} = T_{b_x}^{-1} T_{x(f) \cdot b_x} T_{b_x}^{-1} = x(f) T_{b_x}^{-1}.$$

As $T_{b_x}^{-1}$ is always invertible, this implies that $(T_f)_x$ is invertible if and only if $x(f) \neq 0$. Corollary 26 thus simplifies to

$$\text{sp}_{\text{ess}}(T_f) = \bigcup_{x \in \mathfrak{M} \setminus \mathbb{B}^n} x(f) = f(\partial\mathbb{B}^n) \tag{7.1}$$

in this case. □

Let us add two final remarks to this result.

Remark 33. If we introduce $\mathfrak{M}_{\text{VO}_\partial}$ as the maximal ideal space of VO_∂ , we can formulate Proposition 32 like this:

$$\text{sp}_{\text{ess}}(T_f) = \bigcup_{x \in \mathfrak{M}_{\text{VO}_\partial} \setminus \mathbb{B}^n} x(f) =: f(\mathfrak{M}_{\text{VO}_\partial} \setminus \mathbb{B}^n) \tag{7.2}$$

for $f \in \text{VO}_\partial$. This can be seen as follows. Let $\iota: \text{VO}_\partial \rightarrow \text{BUC}(\mathbb{B}^n)$ be the inclusion mapping. By transposition, this induces the continuous map $\pi: \mathfrak{M} \rightarrow \mathfrak{M}_{\text{VO}_\partial}$, $(\pi(x))(f) := x(\iota(f))$. (7.2) thus follows from (7.1).

Remark 34. In [18, 25, 29] similar results were shown for symbols in $\text{VMO}_\partial(\mathbb{B}^n) \cap L^\infty(\mathbb{B}^n)$ (see [3] or [29] for definitions and descriptions). We can also recover this result from Proposition 32: If $f \in \text{VMO}_\partial(\mathbb{B}^n) \cap L^\infty(\mathbb{B}^n)$, then

$$\text{sp}_{\text{ess}}(T_f) = \tilde{f}(\partial\mathbb{B}^n),$$

where \tilde{f} denotes the Berezin transform of f . Indeed, by [3, Theorem B], \tilde{f} is contained in $\text{VO}_\partial(\mathbb{B}^n)$ and $T_{f-\tilde{f}}$ is compact (see also [17, Theorem 5.5]). Thus the assertion follows.

Acknowledgements

I am very grateful to Wolfram Bauer, Marko Lindner and Christian Seifert for their support and useful discussions. I would also like to thank Robert Fulsche for careful reading and pointing out a lot of typos.

References

- [1] S. Axler, J. Conway, G. McDonald, *Toeplitz Operators on Bergman Spaces*, Can. J. Math., Vol. 34, No. 2 (1982), 466-483.
- [2] W. Bauer, L. Coburn, *Toeplitz Operators with Uniformly Continuous Symbols*, Integr. Equ. Oper. Theory 83 (2015), 25-34.
- [3] D. Békollé, C. Berger, L. Coburn, K. Zhu, *BMO in the Bergman metric on bounded symmetric domains*, J. Funct. Anal., Vol. 93 (1990), 310-350.
- [4] D. Békollé, A. Temgoua Kagou, *Reproducing Kernels and L^p -Estimates for Bergman Projections in Siegel Domains of Type II*, Studia Math. 115 (1995), 219-239.
- [5] G. Bell, A. Dranishnikov, *Asymptotic Dimension*, Topology and its Applications, Vol. 155, No. 12 (2018), 1265-1296.
- [6] C. Berger, L. Coburn, *Toeplitz Operators on the Segal-Bargmann Space*, T. Am. Math. Soc., Vol. 301, No. 2 (1987), 813-129.
- [7] G. Carsson, B. Goldfarb, *On Homological Coherence of Discrete Groups*, J. Algebra, Vol. 276, No. 2 (2004), 502-514.
- [8] B. Choe, Y. Lee, *The Essential Spectra of Toeplitz Operators with Symbols in $H^\infty + C$* , Math. Japon., Vol. 45, No. 1 (1997), 57-60.
- [9] L. Coburn, *Singular Integral Operators and Toeplitz Operators on Odd Spheres*, Indiana Univ. Math. J. 23 (1973), 433-439.
- [10] M. Engliš, *Compact Toeplitz Operators via the Berezin Transform on Bounded Symmetric Domains*, Integr. Equ. Oper. Theory 33 (4), 426-455 (1999).
- [11] M. Engliš, *A Mean Value Theorem on Bounded Symmetric Domains*, P. Am. Math. Soc., Vol. 127, No. 11, 3259-3268 (1999).
- [12] J. Faraut, A. Koranyi, *Function Spaces and Reproducing Kernels on Bounded Symmetric Domains*, J. Funct. Anal. 88 (1990), 64-89.
- [13] R. Hagger, M. Lindner, M. Seidel: *Essential Pseudospectra and Essential Norms of Band-Dominated Operators*, J. Math. Anal. Appl., Vol. 437, No. 1 (2016), 255-291.
- [14] M. Lindner, *Infinite Matrices and their Finite Sections*, Birkhäuser Verlag, Basel, Boston, Berlin, 2006.

- [15] M. Lindner, M. Seidel, *An Affirmative Answer to a Core Issue on Limit Operators*, J. Funct. Anal., Vol. 267, No. 3 (2014), 901-917.
- [16] G. McDonald, *Fredholm Properties of a Class of Toeplitz Operators on the Ball*, Indiana Univ. Math. J., Vol. 26, No. 3 (1977), 567-576.
- [17] M. Mitkovski, D. Suárez, B. Wick, *The Essential Norm of Operators on $A_\alpha^p(\mathbb{B}_n)$* , Integr. Equ. Oper. Theory 75 (2013), 197-233.
- [18] A. Perälä, A. Virtanen, *A Note on the Fredholm Properties of Toeplitz Operators on Weighted Bergman Spaces with Matrix-Valued Symbols*, Oper. Matrices, Vol. 5, No. 1 (2011), 97-106.
- [19] V. Rabinovich, S. Roch, B. Silbermann, *Fredholm Theory and Finite Section Method for Band-Dominated Operators*, Integr. Equat. Oper. Th., Vol. 30, No. 4 (1998), 452-495.
- [20] V. Rabinovich, S. Roch, B. Silbermann, *Limit Operators and Their Applications in Operator Theory*, Birkhäuser Verlag, Basel, Boston, Berlin, 2004.
- [21] M. Seidel, *Fredholm Theory for Band-Dominated and Related Operators: a Survey*, Linear Algebra Appl., Vol. 445 (2014), 373-394.
- [22] J. Špakula, R. Willett, *A Metric Approach to Limit Operators*, Trans. Amer. Math. Soc., Vol. 369 (2017), 263-308.
- [23] K. Stroethoff, D. Zheng, *Toeplitz and Hankel Operators on Bergman Spaces*, Trans. Amer. Math. Soc., Vol. 329, No. 2 (1992), 773-794.
- [24] D. Suárez, *The Essential Norm of Operators in the Toeplitz Algebra on $A^p(\mathbb{B}_n)$* , Indiana Univ. Math. J., Vol. 56, No. 5 (2007), 2185-2232.
- [25] J. Taskinen, J. Virtanen, *Toeplitz Operators on Bergman Spaces with Locally Integrable Symbols*, Rev. Mat. Iberoam., Vol. 26, No. 2 (2010), 693-706.
- [26] H. Upmeyer, *Toeplitz Operators and Index Theory in Several Complex Variables*, Operator Theory: Advances and Applications, Vol. 81, Birkhäuser Verlag, Basel, 1996.
- [27] A. Zaanen, *Introduction to Operator Theory in Riesz Spaces*, Springer-Verlag, New York and Heidelberg, 1997.
- [28] X. Zeng, *Toeplitz Operators on Bergman Spaces*, Houston J. Math., Vol. 18, No. 3 (1992), 387-407.
- [29] K. Zhu, *VMO, ESV, and Toeplitz Operators on the Bergman Space*, Trans. Amer. Math. Soc., Vol. 302, No. 2 (1987), 617-646.
- [30] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, Graduate Texts in Mathematics, Vol. 226, Springer-Verlag, New York, 2005.

Raffael Hagger
 Institute of Analysis
 Leibniz Universität Hannover

Welfengarten 1
30167 Hannover
GERMANY
raffael.hagger@math.uni-hannover.de