



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:

Journal of Sports Sciences

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa50479>

Paper:

Birdsey, L., Weston, M., Russell, M., Johnston, M., Cook, C. & Kilduff, L. (2019). Neuromuscular, physiological and perceptual responses to an elite netball tournament. *Journal of Sports Sciences*, 1-6.

<http://dx.doi.org/10.1080/02640414.2019.1625613>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

1 Full title: Neuromuscular, physiological and perceptual responses to an elite
2 netball tournament.

3

4 Running title: Neuromuscular, physiological and perceptual responses to elite
5 netball

6

7 Laurence P. Birdsey^{ab}, Matthew Weston^c, Mark Russell^d, Michael Johnston^e,
8 Christian J. Cook^f and Liam P.Kilduff^{a*}

9 *^aA-STEM College of Engineering, Swansea University, Swansea, UK; ^bSport*

10 *Wales, Cardiff, UK; ^cDepartment of business and law, Teesside University,*

11 *Middlesborough, UK; ^dSchool of social and health sciences, Leeds trinity*

12 *University, UK; ^eBritish Athletics, Birmingham, UK; ^fSchool of sport, health and*

13 *exercise sciences, Bangor University, UK*

14 Corresponding author: Liam P. Kilduff, A-STEM College of engineering,

15 Swansea University Bay Campus, Fabian Way, Crymlyn Burrows, Swansea,

16 SA81 8EN. L.kilduff@swansea.ac.uk

17

18 Word count: 3354

19

20

21

22

23

24 **Neuromuscular, physiological and perceptual responses to an elite**
25 **netball tournament**

26 **Abstract**

27 To examine responses to an International netball tournament, female athletes
28 ($n=11$) played three matches over consecutive days. External (accelerometry) and
29 internal (heart rate; HR, session; sRPE, and differential; dRPE, rating of perceived
30 exertion) load measures quantified match intensity. On match-day mornings, and
31 three days after match three, well-being (brief assessment of mood; BAM+),
32 biochemical (creatinine kinase concentration; CK), neuromuscular (jump height; JH,
33 peak power output; PPO) and endocrine function (salivary cortisol; C, testosterone;
34 T, concentrations) were assessed. External load was similar between matches
35 whereas dRPE and sRPE was greatest for match three. Following match one, CK
36 increased, whereas BAM+, JH, C and T decreased. Following two matches, BAM+,
37 PPO, and T decreased with CK increasing versus baseline. Following consecutive
38 matches, CK (likely moderate; $27.9\% \pm 19.5\%$) and C (possibly moderate; 43.3%
39 $\pm 46.8\%$) increased, whilst BAM+ (possibly moderate; $-20.6\% \pm 24.4\%$) decreased.
40 Three days post-tournament BAM+, T, PPO, and JH decreased. Mid-court elicited
41 higher mean HR (possibly moderate; $3.7\% \pm 3.8\%$), internal and external intensities
42 (possibly very large; $85.7\% \pm 49.6\%$) compared with goal-based positions.
43 Consecutive matches revealed a dose-response relationship for well-being and
44 physiological function; a response evident three days post-tournament.

45 **Keywords:** recovery; monitoring; load; team sport; readiness to train

46 **Introduction**

47 Whilst several studies have reported the movement demands of elite netball in
48 recent years (Bailey, Gastin, Mackey, & Dwyer, 2017; Fox, Spittle, Otago, &
49 Saunders, 2013; Young, Gastin, Sanders, Mackey, & Dwyer, 2016), to date no
50 studies have profiled the physiological responses to elite level tournament match-
51 play. Indeed, only three studies have reported the movement demands of elite
52 netball, one by use of notational analysis (Fox et al., 2013) and two by
53 accelerometry (Bailey et al., 2017; Young et al., 2016). Goal defence (GD),
54 goalkeeper (GK) and goal shooter (GS) positions were reported to perform at the
55 the lowest playing intensities and highest proportions of match time spent in the
56 low-intensity zones when compared to players occupying wing attack (WA), wing
57 defence (WD), centre (C), and goal attack (GA) positions (Young et al., 2016).
58 Additionally, Bailey et al., (2017) reported the accelerometer-based loads
59 associated with typical activities, reporting off-ball guarding to elicit the highest
60 load per instance, whilst jogging accumulated the greatest load across a match.

61

62 At present, a single study has reported the responses to an isolated match reporting
63 a reduction in perception of fatigue and neuromuscular performance immediately
64 and 24 h after an 80 min elite level match, returning to baseline 36 h later (Wood,
65 Kelly, & Gabbett, 2013). Many tournaments require teams to play up to eight
66 matches in 10 days, therefore, the demands are not limited to that of a single match,
67 rather the ability to perform and recover over a series of days. Findings of previous
68 studies reporting the neuromuscular and perceptual recovery profiles (Wood et al.,
69 2013) may be limited by match duration (80 min compared to 60 min for
70 International matches), small sample size ($n=6$) and single match design as opposed

71 to that of a tournament, leading to an underestimation of the responses to
72 tournament match-play. Recent reports of match demands have differed (Fox et al.,
73 2013) to previous reports in elite players (Otago, 1983), as such recent rule changes
74 (January 2016), intended to reduce stoppages and increase the speed and intensity
75 of match-play, may have compromised the application of previous literature
76 regarding the demands and responses to netball match-play. Limited information
77 exists regarding the external loads of professional netball, (Bailey et al., 2017;
78 Young et al., 2016) and no studies have examined the physiological demands and
79 responses to either a single or multiple instances of International-standard netball
80 match-play. A deeper understanding of the movement patterns, coupled with
81 physiological demands, can allow effective training to be prescribed to optimise
82 adaptation and performance, however this information is currently limited (Bailey
83 et al., 2017). Therefore, the purpose of this study was to examine the physiological,
84 neuromuscular, endocrine and perceptual responses to an International netball
85 tournament as well as the physiological demands of International-standard netball.

86 **Methods**

87 This observational study examined the response to a netball tournament performed
88 over three consecutive days. Matches commenced at 19:00, 15:00 and 15:00 h on
89 days one to three, respectively. On the morning of each match (~07:30 h), and three
90 days (approximately 62 h) after the final match (~07:30 h), scores for perceived
91 well-being (adapted brief assessment of mood+; BAM+), and samples of whole
92 blood (Creatine Kinase concentration; CK) and saliva (cortisol; C and testosterone:
93 T concentrations) were collected, and countermovement jump testing performed.
94 Match intensity was quantified using both internal (heart rate telemetry) and
95 external (accelerometry) load metrics. Following the match, players individually

96 recorded session (sRPE: Foster et al., 2001) and differential ratings of perceived
97 exertion (dRPE: Weston, Siegler, Bahnert, McBrien, & Lovell, 2015) using a
98 numerically blinded CR100® scale via an Android tablet. These values were
99 recorded during the cool down period, ~15 min after match-play.

100

101 Eleven female players (age: 25 ± 4 years; mass: 71.8 ± 7.8 kg; height: 1.8 ± 0.1 m)
102 from an International netball team were recruited. Players were assigned according
103 to positions to goal-based ($n=2$, GS and GK) and mid-court ($n=9$, GD, WD, WA,
104 C and GA) groups based on court movement restrictions. This study included an
105 International tournament played at the end of the 2016 domestic season. As such,
106 all players had competed weekly in the British Super League (highest netball league
107 in Britain) and were engaged in full-time training (strength, speed, endurance and
108 netball-specific training sessions four to six times per week) as part of their club's
109 performance preparation program. Five players used no form of hormonal
110 contraceptive and players were requested to self-monitor menstrual cycles and days
111 of contraceptive consumption. Subsequent analyses revealed no bias in hormonal
112 markers as a function of contraceptive use. This study was approved by the Swansea
113 University ethics committee, players were informed of the benefits and risks of the
114 investigation before signing informed consent forms and completing health
115 screening and were made aware that all material would be anonymised. All
116 mandatory health and safety procedures were complied with in completing this
117 research study.

118

119 Players completed BAM+ which is correlated to high-intensity match activity, and
120 is sensitive to physiological responses following elite team sport match-play

121 (Shearer et al., 2017). Using an Android tablet (Iconia One 7 B1-750, Taipei,
122 Taiwan: Acer inc), a series of questions was answered with a 100 mm visual
123 analogue scale anchored with “not at all [0]” and “extremely [100]”. An overall
124 recovery score was generated by subtracting the mean score of negative related
125 items from the mean score of the positively related questions using Equation 1:
126 (Shearer et al., 2017)

127

128 Equation 1: (Alertness + sleep quality + confidence + motivation) /4 - (Anger +
129 confusion + tension + depression + fatigue + muscle soreness)/6.

130

131 For salivary hormone analysis, players were instructed to avoid eating food or
132 drinking fluids other than water after waking to avoid contaminating saliva samples.
133 Prior to breakfast, a two ml sample of saliva was collected via passive drool
134 (Crewther et al., 2013) into sterile containers, with samples subsequently stored at
135 -70°C until assay. After thawing and centrifugation (2000 revolutions·min⁻¹ for 10
136 min), the samples were analysed in duplicate for T and C using commercial kits
137 (Salimetrics, LLC, State College, PA, USA). The minimum detection limit for the
138 testosterone assay was 6.1 pg·ml⁻¹, with interassay coefficient of variation (CV)
139 <10%. The cortisol assay had a detection limit of 0.12 ng·ml⁻¹ with interassay CV
140 <7%. Samples for each player were assayed in the same plate to eliminate inter-
141 assay variability.

142

143 Whole blood CK concentrations were measured via capillary blood (120 µl) being
144 sampled from the fingertip and stored on ice in EDTA prepared collection tubes
145 (Microvette 500, Sarstedt, Numbrecht, Germany) before being centrifuged at 3000

146 revolutions·min⁻¹ for 10 min (Labofuge 400R; Kendro Laboratories,
147 Langensfeld, Germany). Plasma samples were then stored at -70°C before being
148 analysed for CK concentration using commercially available kits (CK-NAC, ABX
149 Diagnostics, Northampton, United Kingdom) on a spectrophotometer (Cobas Mira,
150 ABX Diagnostics, Northampton, United Kingdom). Samples were measured in
151 duplicate (CV=3%) and recorded as a mean.

152

153 A portable force platform with built-in charge amplifier (Kistler type 92866AA,
154 Kistler Instruments Ltd., Farnborough, UK) measured the ground reaction force-
155 time history of countermovement jumps. A sample rate of 1000 Hz was used, and
156 the platform's calibration was confirmed prior to testing. Power (CV=2.4%) and
157 jump height (JH; calculated from takeoff velocity; CV=3.4%) was calculated using
158 previously established procedures (Owen et al., 2014; West et al., 2011) and have
159 been reported to be sensitive to changes following competitive matches (Russell et
160 al., 2015; West et al., 2014). Players performed a standardised warm up before
161 jumping, placed hands on hips throughout the jump, and performed two jumps at
162 each time-point with the best jump taken as the highest peak power output (PPO)
163 and used in subsequent analyses.

164

165 External load was quantified by use of a microtechnology unit (Catapult S5,
166 Catapult, Innovations, Leeds, UK) housing an in-built tri-axial accelerometer
167 sampling at 100 Hz. Players wore a custom-made vest (Catapult Innovations,
168 Leeds, UK) in which units were held in place vertically on the upper back to
169 minimise movement. Data were downloaded using the manufacturer's software
170 (Catapult sprint 5.1, Catapult Innovations, Leeds, UK), analysed for player-load for

171 each quarter, excluding breaks between quarters, with data represented as external
172 load intensity ($\text{AU}\cdot\text{min}^{-1}$). Data was pooled and reported for each position rather
173 than individual players, such that for every match each position would have a single
174 external load intensity for each quarter. Player-load has been reported to be a valid
175 and reliable method (Barrett, Midgley, & Lovell, 2014; Boyd, Ball, & Aughey,
176 2011) of measuring activities performed in team sports movements, with high
177 within and between-device ($\text{CV}\sim 1\%$: Boyd et al., 2011) reliability and has been
178 widely used in team sports (Luteberget & Spencer, 2017; Polgaze, Dawson,
179 Hiscock, & Peeling, 2015) including netball (Chandler, Pinder, Curran, & Gabbett,
180 2014; Young et al., 2016) with detailed calculations described previously (Barrett
181 et al., 2014). Players wore heart rate (HR) monitors (Polar Team System 2, Polar
182 Electro, Warwick, UK) throughout matches, with HR recorded at beat-to-beat
183 intervals. Data was downloaded and analysed for each quarter, excluding breaks
184 between quarters, and only whilst the player was on-court, using the Polar team
185 system software (Polar Team 2, Polar Electro, Warwick, UK). HR data was
186 reported for each player and associated to the position which had been played.

187

188 Following each match, players recorded sRPE along with indices of dRPE
189 including ratings for breathlessness (RPE-B), leg muscle exertion (RPE-L), upper
190 body muscle exertion (RPE-U) and cognitive/technical demands (RPE-T) (Weston
191 et al., 2015). Ratings were provided using a numerically blinded CR100® scale
192 with verbal anchors using a bespoke application on an Android tablet. dRPE
193 provides a detailed quantification of internal load during team sport activities
194 (McLaren, Smith, Spears, & Weston, 2017), is a sensitive marker of match exertion
195 (Weston et al., 2015) and distinguishes between different areas of effort (McLaren

196 et al., 2017; Weston et al., 2015). Players must have performed a minimum of one
197 quarter for sRPE and dRPE to be included in subsequent analyses.

198

199 Data are reported as mean difference \pm 90% confidence limits unless otherwise
200 stated. Visual inspection of the residual plots revealed evidence of
201 heteroscedasticity; therefore, except for sRPE, dRPE, BAM+ and HR, analyses
202 were performed on log transformed data. Separate mixed linear mixed models
203 (SPSS v.24, Armonk, NY: IBM Corp) were used to examine the effect of
204 tournament match-play on measures of physical exertion (external load, HR, sRPE,
205 dRPE) and, thereafter, the effect of playing position on match physical exertion,
206 and, the effects of tournament match-play on the players' neuromuscular,
207 physiological and perceptual responses (PPO, JH, CK, T, C). In these models,
208 match (match 1, match 2, match 3), playing position (mid-court, goal-based) and
209 time (day 1, day 2, day 3, 3 days post), respectively were entered as the fixed effects.
210 In all models, players were included as a random effect with random intercept to
211 account for the dependency that arises from a hierarchical data structure such as
212 ours (i.e., repeated measurements from the same players). From here, a custom-
213 made spreadsheet (Hopkins, 2007) was used to determine magnitude based
214 inferences (Batterham & Hopkins, 2006) for all differences, with inferences based
215 on standardised thresholds for small, moderate, large and very large differences of
216 0.2, 0.6, 1.2 and 2.0 of the pooled between-subject standard deviations (SD)
217 (Hopkins, Marshall, Batterham, & Hanin, 2009). The chance of the difference being
218 substantial or trivial was interpreted using the following scale: 25–75%, possibly;
219 75–95%, likely; 95–99.5%, very likely; >99.5%, most likely (Batterham &
220 Hopkins, 2006). Uncertainty in all estimates is expressed via 90% confidence limits

221 and the magnitude of effects assessed mechanistically, whereby if the confidence
222 limits overlapped the thresholds for the smallest worthwhile positive and negative
223 effects, effects were deemed unclear (Hopkins et al., 2009).

224 **Results**

225 Match data are presented in Table 1. Mean playing time for players across the three
226 matches was 119.8 min (± 48.5 min; \pm SD) and outcomes included two wins and a
227 loss for matches one to three respectively. In response to a single netball match,
228 from day one to day two, CK (likely very large; $72.6\% \pm 26.4\%$) and fatigue (likely
229 small; $56.2\% \pm 46.0\%$) increased, whilst motivation (likely moderate; $-19.5\% \pm$
230 14.3%), BAM+ (likely moderate; $-27.9\% \pm 17.6\%$), sleep quality (possibly
231 moderate; $-16.3\% \pm 15.6\%$), C (likely small; $-27.4\% \pm 23.7\%$), T (possibly small;
232 $-10.8\% \pm 10.8\%$) and JH (possibly small; $-4.0\% \pm 2.5\%$) decreased, with a possible
233 trivial difference for PPO and unclear difference for soreness (Table 2). Following
234 two netball matches, from day one to day three, CK (most likely very large; 120.8%
235 $\pm 33.7\%$), fatigue (possibly large; $146.9\% \pm 46.0\%$) and soreness (possibly
236 moderate; $57.7\% \pm 37.9\%$) increased, whilst BAM+ (likely large; $-42.8 \pm 17.6\%$),
237 motivation (likely moderate; $-20.6\% \pm 14.3\%$), sleep quality (possibly moderate; $-$
238 $30.8\% \pm 15.6\%$), T (possibly small; $-8.7\% \pm 11.0\%$) and PPO (possibly small; $-$
239 $3.3\% \pm 1.7\%$) decreased, with a possible trivial difference for JH and most likely
240 trivial difference for C. Following the performance of two consecutive matches,
241 from day two to three, CK (likely moderate; $27.9\% \pm 19.5\%$), fatigue (likely
242 moderate; $58.1\% \pm 29.5\%$), soreness (possible moderate; $49.6\% \pm 36.0\%$) and C
243 (possibly moderate; $43.3\% \pm 46.8\%$) increased whilst BAM+ (possibly moderate;
244 $-20.6\% \pm 24.4\%$) and sleep quality (possibly moderate; $-17.3\% \pm 18.6\%$) decreased,
245 with an unclear difference for T and motivation, and likely trivial difference for JH

246 and PPO. Three days post-tournament BAM+ (likely very large; $-57.5\% \pm 20.5\%$),
247 sleep quality (likely large; $-38.7\% \pm 18.1\%$), motivation (likely moderate; -24.3%
248 $\pm 16.6\%$), PPO (likely small; $-4.2\% \pm 1.9\%$), JH (possibly small; $-3.9\% \pm 2.8\%$)
249 and T (possibly small; $-10.0\% \pm 12.7\%$) decreased, whilst fatigue increased (very
250 likely moderate; $127.2\% \pm 53.6\%$) compared to day one, with unclear differences
251 for C, CK and soreness.

252

253 *****Table 1 about here*****

254 *****Table 2 about here*****

255

256 Greater mean HR for match one occurred relative to match two (possibly small;
257 $1.2\% \pm 1.2.0\%$). Likely trivial differences were observed for external load intensity
258 and unclear differences for sRPE and dRPE variables. For match three versus one
259 for RPE-B (likely small; $20.1\% \pm 25.4\%$), RPE-L (possibly small; $18.2\% \pm 24.5\%$),
260 RPE-U (possibly small; $18.1\% \pm 22.4\%$) and RPE-T (possibly moderate; $23.2\% \pm$
261 19.8%), greater values were observed. A possible trivial difference existed for
262 external load intensity and unclear differences for sRPE and mean HR. Match three
263 produced greater sRPE (likely small; $21.7\% \pm 27.4\%$), RPE-B (possibly moderate;
264 $32.0\% \pm 26.7\%$), RPE-L (possibly moderate; $30.8\% \pm 25.9\%$), RPE-U (likely small;
265 $30.6\% \pm 23.7\%$), RPE-T (possibly moderate; $27.1\% \pm 20.2\%$) and mean HR
266 (possibly small; $1.1\% \pm 2.0\%$) versus match two. There was a possible trivial
267 difference for external load intensity.

268

269 Overall, mid-court positions performed at a greater external load intensity (possibly
270 very large; $85.7\% \pm 49.6\%$), mean HR (possibly moderate; $3.7\% \pm 3.8\%$) (Table

271 3), and reported higher sRPE (possibly moderate; $40.7\% \pm 40.0\%$), RPE-B (likely
272 moderate; $55.9\% \pm 51.9\%$), RPE-L (possibly large; $79.3\% \pm 48.1\%$), RPE-U
273 (possibly moderate; $47.2\% \pm 54.9\%$) and RPE-T (possibly moderate; $36.9\% \pm$
274 36.7%) compared to goal-based positions (Table 4).

275

276 *****Table 3 about here*****

277 *****Table 4 about here*****

278 **Discussion**

279 The aims of this study were to characterise the physiological, neuromuscular,
280 endocrine and perceptual responses to an International tournament and to identify
281 the position-specific demands of International netball. The primary findings were
282 that the performance of both a single, and multiple matches resulted in a varied
283 recovery profile, with greater perturbations in perceived well-being and
284 physiological function following consecutive matches, and fatigue evident up to
285 three days post-tournament. Additionally, mid-court positions performed at greater
286 internal and external load intensity compared to goal-based positions.

287

288 Across the tournament, CK, reported to be indicative of skeletal muscle damage
289 (Cunniffe et al., 2010), accumulated before returning to baseline thereafter. Whilst
290 there are no reports in netball, investigations in other team sports have reported
291 peak values occurring 24 h post-match, remaining elevated for females for up to 69
292 h (Andersson et al., 2008). Three days post-tournament, CK and perceived soreness
293 had returned to baseline, however neuromuscular performance and T
294 concentrations remained suppressed. This may suggest that neuromuscular
295 performance is impacted by T concentration rather than muscle damage, that CK is

296 not sensitive to detect changes in muscle damage, or that various markers of fatigue
297 collectively interact.

298

299 Following the performance of a single match, T was reduced, and remained reduced
300 until three days post-tournament, whilst C decreased following the first match, then
301 returned and remained at baseline following the second match. Testosterone
302 concentration is associated with enhanced neuromuscular performance (Cook,
303 Kilduff, Crewther, Beaven, & West, 2014), decision making, behaviour, contractile
304 signalling (Crewther, Cook, Cardinale, & Weatherby, 2011), motivation (Cook,
305 Kilduff, & Crewther, 2018) and performance (Crewther et al., 2013). A reduction,
306 as seen in the present study, may have negatively affected one or more of these
307 reported associations, with a resultant impact upon performance. The recovery of
308 C following two matches may suggest a varied anticipatory response with a greater
309 anticipatory rise prior to the first and final match (higher ranked opponent for the
310 final match). However, alternatively the late commencement (19:00 h compared to
311 15:00 h) of match one may have negatively affected post-match processes and
312 recovery. Menstrual phase and hormonal contraceptive use were not controlled for
313 in the present study, however no difference was found in basal T between hormonal
314 contraceptive users and non-users. Additionally, recent reports highlight only a
315 difference in magnitude of T response to a stimulus, rather than the response itself,
316 and no impact upon performance with hormonal contraceptive use (Cook et al.,
317 2018).

318

319 This is the first study to characterise playing demands during an International
320 tournament reporting external load, perceived effort and HR. Internal and external

321 load was greater for mid-court compared to goal-based positions (Table 3). Greater
322 external load intensity for mid-court positions has been previously reported in
323 professional netball (Fox et al., 2013; Young et al., 2016), and is likely due to court
324 movement restrictions resulting in a higher active time (Fox et al., 2013), time spent
325 in high-intensity zones (Young et al., 2016) and type of on and off-ball locomotor
326 and non-locomotor activity (Bailey et al., 2017). Collectively, this suggests that
327 players should not only be conditioned for the position specific movement
328 demands, as previously reported, but also the different physiological and type of
329 effort (as indicated by dRPE) experienced during International match-play. Both
330 sRPE and dRPE can be used by conditioning staff to guide the individualisation of
331 the training stimulus to the positional demands. As markers of fatigue were further
332 reduced following a greater number of consecutive matches, training should aim to
333 replicate these demands to minimise this disturbance, especially when considering
334 that some International tournaments are up to twice as long as in the present study.
335 Unlike perceptual and endocrine responses, neuromuscular performance was not
336 further reduced following consecutive matches. Perceptual markers could therefore
337 be considered as a simple monitoring tool to identify sufficient training load to
338 replicate the fatiguing consequences associated with International netball. Sleep
339 quality was negatively affected following a single, and to a greater extent following
340 consecutive matches, a consideration for coaching and support staff, as sleep has
341 been reported to be vital for recovery (Halson, 2008). Three days post-tournament,
342 when players commenced training, perceived well-being, sleep quality, T
343 concentration and neuromuscular function were reduced, suggesting longer
344 recovery is required than anticipated by conditioning staff.

345 Conclusion

346 This is the first study to report the physiological demands of and responses to an
347 International netball tournament, providing vital information for International
348 coaches and conditioning coaches. Markers of fatigue increased following the
349 performance of a single match, whilst markers of muscle damage and perceived
350 well-being were further affected following consecutive matches. A varied recovery
351 profile was apparent as recovery to baseline of all variables examined did not occur
352 62 h post-tournament. Mid-court positions performed at higher external and internal
353 intensities compared to goal-based positions, an important consideration for
354 conditioning staff in order to individualise training to positional specific demands.

355

356

357

358

359

360

361 Acknowledgements

362 We would like to acknowledge the Welsh netball players, coaching and support staff for
363 their support through this testing. Additionally, to Liam Harper, Christopher Lewis
364 and Charlie Finn for assisting with data collection.

365

366 This research did not receive any specific grant from funding agencies in the public,
367 commercial, or not-for-profit sectors.

368

369 Disclosure of interest

370 The authors report no conflict of interest.

371

372

373

374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396

397 **References**

- 398 Andersson, H., Raastad, T., Nilsson, J., Paulsen, G., Garthe, I., & Kadi, F. (2008).
399 Neuromuscular fatigue and recovery in elite female soccer: Effects of active
400 recovery. *Medicine and Science in Sports and Exercise*, 40(2), 372–380.
401 <https://doi.org/10.1249/mss.0b013e31815b8497>
- 402 Bailey, J. A., Gastin, P. B., Mackey, L., & Dwyer, D. B. (2017). The Player Load
403 Associated With Typical Activities in Elite Netball. *International Journal of*
404 *Sports Physiology and Performance*, 12, 1218–1223.
405 <https://doi.org/10.1123/ijsspp.2016-0378>

- 406 Barrett, S., Midgley, A., & Lovell, R. (2014). PlayerLoadTM: Reliability,
407 convergent validity, and influence of unit position during treadmill running.
408 *International Journal of Sports Physiology and Performance*, 9(6), 945–952.
409 <https://doi.org/10.1123/ijsp.2013-0418>
- 410 Batterham, A. M., & Hopkins, W. G. (2006). Making meaningful inferences about
411 magnitudes. *International Journal of Sport Physiology and Performance*,
412 1(1), 50–57.
- 413 Boyd, L. J., Ball, K., & Aughey, R. J. (2011). The reliability of MinimaxX
414 accelerometers for measuring physical activity in Australian football.
415 *International Journal of Sport Physiology and Performance*, 6, 311–321.
416 <https://doi.org/10.1123/ijsp.6.3.311>
- 417 Chandler, P., Pinder, S., Curran, J., & Gabbett, T. J. (2014). Physical demands of
418 training and competition in collegiate netball players, 28(10), 2732–2737.
419 <https://doi.org/10.1519/JSC.0000000000000486>
- 420 Cook, C. J., Kilduff, L. P., & Crewther, B. T. (2018). Basal and stress-induced
421 salivary testosterone variation across the menstrual cycle and linkage to
422 motivation and muscle power. *Scandinavian Journal of Medicine & Science
423 in Sports*, 28(4), 1345–1353. <https://doi.org/10.1111/sms.13041>
- 424 Cook, C. J., Kilduff, L. P., Crewther, B. T., Beaven, M., & West, D. J. (2014).
425 Morning based strength training improves afternoon physical performance in
426 rugby union players. *Journal of Science and Medicine in Sport*, 17(3), 317–
427 321. <https://doi.org/10.1016/j.jsams.2013.04.016>
- 428 Crewther, B. T., Cook, C., Cardinale, M., & Weatherby, R. (2011). Two emerging
429 concepts for elite athletes cortisol and testosterone. *Review Literature And
430 Arts Of The Americas*, 41(2), 103–123. <https://doi.org/10.2165/11536850->

- 431 000000000-00000
- 432 Crewther, B. T., Sanctuary, C. E., Kilduff, L. P., Carruthers, J. S., Gaviglio, C.
433 M., & Cook, C. J. (2013). The Workout Responses of Salivary-Free
434 Testosterone and Cortisol Concentrations and Their Association With the
435 Subsequent Competition Outcomes in Professional Rugby League. *Journal*
436 *of Strength and Conditioning Research*, 27(2), 471–476.
437 <https://doi.org/10.1519/JSC.0b013e3182577010>
- 438 Cunniffe, B., Hore, A. J., Whitcombe, D. M., Jones, K. P., Baker, J. S., & Davies,
439 B. (2010). Time course of changes in immunoendocrine markers following
440 an international rugby game. *European Journal of Applied Physiology*,
441 108(1), 113–122. <https://doi.org/10.1007/s00421-009-1200-9>
- 442 Foster, C., Florhaug, J., Franklin, J., Gottschall, L., Hrovatin, L., Parker, S., ...
443 Dodge, C. (2001). A new approach to monitoring exercise training. *Journal*
444 *of Strength and Conditioning Research*, 15(1), 109–115.
- 445 Fox, A., Spittle, M., Otago, L., & Saunders, N. (2013). Activity profiles of the
446 Australian female netball team players during international competition:
447 implications for training practice. *Journal of Sports Sciences*, 31(14), 1588–
448 1595. <https://doi.org/10.1080/02640414.2013.792943>
- 449 Halson, S. L. (2008). Nutrition, sleep and recovery. *European Journal of Sport*
450 *Science*, 8(2), 119–126. <https://doi.org/10.1080/17461390801954794>
- 451 Hopkins, W. G. (2007). A spreadsheet for deriving confidence interval,
452 mechanistic inference and clinical inference from a P value. *SportScience*,
453 11, 16+.
- 454 Hopkins, W. G., Marshall, S. W., Batterham, A. M., & Hanin, J. (2009).
455 Progressive statistics for studies in sports medicine and exercise science.

- 456 *Medicine and Science in Sports and Exercise*, 41(1), 3–12.
457 <https://doi.org/10.1249/MSS.0b013e31818cb278>
- 458 Luteberget, L. S., & Spencer, M. (2017). High-intensity events in international
459 women's team handball matches. *International Journal of Sports Physiology*
460 *and Performance*, 12(1), 56–61. <https://doi.org/10.1123/ijsp.2015-0641>
- 461 McLaren, S. J., Smith, A., Spears, I. R., & Weston, M. (2017). A detailed
462 quantification of differential ratings of perceived exertion during team-sport
463 training. *Journal of Science and Medicine in Sport*, 20(3), 290–295.
464 <https://doi.org/10.1016/j.jsams.2016.06.011>
- 465 Otago, L. (1983). A game analysis of the activity patterns of netball players.
466 *Sports Coach*, 7, 24–28.
- 467 Owen, N., Watkins, J., Kilduff, L., Cunningham, D., Bennett, M., & Bevan, H.
468 (2014). Development of a criterion methods to determine peak mechanical
469 power output in a countermovement jump. *Journal of Strength and*
470 *Conditioning Research*, 28(6), 1552–1558.
- 471 Polgaze, T., Dawson, B., Hiscock, D., & Peeling, P. (2015). A comparative
472 analysis of accelerometer and time motion data in elite mens hockey training
473 and competition. *International Journal of Sports Physiology and*
474 *Performance*, 10(4), 446–451. <https://doi.org/10.1123/ijsp.2014-0233>
- 475 Russell, M., Northeast, J., Atkinson, G., Shearer, D. A., Sparkes, W., Cook, C. J.,
476 & Kilduff, L. P. (2015). Between-Match Variability of Peak Power Output
477 and Creatine Kinase Responses to Soccer Match-Play. *Journal of Strength*
478 *and Conditioning Research*, 29(8), 2079–2085.
479 <https://doi.org/10.1519/JSC.0000000000000852>
- 480 Shearer, D. A., Sparkes, W., Northeast, J., Cunningham, D. J., Cook, C. J., &

- 481 Kilduff, L. P. (2017). Measuring recovery: An adapted Brief Assessment of
482 Mood (BAM+) compared to biochemical and power output alterations.
483 *Journal of Science and Medicine in Sport*, *20*(5), 512–517.
484 <https://doi.org/10.1016/j.jsams.2016.09.012>
- 485 West, D. J., Cook, C. J., Stokes, K. A., Atkinson, P., Drawer, S., Bracken, R. M.,
486 & Kilduff, L. P. (2014). Profiling the time-course changes in neuromuscular
487 function and muscle damage over two consecutive tournament stages in elite
488 rugby sevens players. *Journal of Science and Medicine in Sport*, *17*(6), 688–
489 692. <https://doi.org/10.1016/j.jsams.2013.11.003>
- 490 West, D. J., Owen, N. J., Jones, M. R., Bracken, R. M., Cook, C. J., Cunningham,
491 D. J., ... Kilduff, L. P. (2011). Relationships between force-time
492 characteristics of the isometric midthigh pull and dynamic performance in
493 professional rugby league players. *Journal of Strength and Conditioning*
494 *Research*, *25*(11), 3070–3075.
495 <https://doi.org/10.1519/JSC.0b013e318212dcd5>
- 496 Weston, M., Siegler, J., Bahnert, A., McBrien, J., & Lovell, R. (2015). The
497 application of differential ratings of perceived exertion to Australian Football
498 League matches. *Journal of Science and Medicine in Sport*, *18*(6), 704–708.
499 <https://doi.org/10.1016/j.jsams.2014.09.001>
- 500 Wood, A., Kelly, V., & Gabbett, T. J. (2013). Neuromuscular And Perceptual
501 Fatigue Responses To An Elite Level Netball Match. *Journal of Australian*
502 *Strength and Conditioning*, *21*(1), 24–29.
- 503 Young, C. M., Gatin, P. B., Sanders, N., Mackey, L., & Dwyer, D. B. (2016).
504 Player load in elite netball: Match, training, and positional comparisons.
505 *International Journal of Sports Physiology and Performance*, *11*(8), 1074–

506 1079. <https://doi.org/10.1123/ijsp.2015-0156>

507