


Health Behavior Theory and cumulative knowledge regarding health behaviors: are we moving in the right direction?

metadata, citation and similar papers at core.ac.uk

brought to you by  CC BY

provided by Carolina Digital Repository

Seth M. Noar^{1,2} and Rick S. Zimmerman¹

Abstract

Although research on Health Behavior Theory (HBT) is being conducted at a rapid pace, the extent to which the field is truly moving forward in understanding health behavior has been questioned. This issue is examined in the current article. First, we discuss the problems within the HBT literature. Second, we discuss the proliferation of HBT and why theory comparison is essential to this area of research. Finally, we reflect on ways that the field might move forward by suggesting a new agenda for HBT research. It is argued that increased recognition of the similarity of health behavior constructs as well as increased empirical comparisons of theories are essential for true scientific progress in this line of inquiry.

Introduction

It is the goal of many researchers interested in health behavior to understand both the determinants of health behaviors and the process of health behavior change. One key route to an understanding of health behavior has been the development and empirical testing of Health Behavior Theory (HBT). Research in this area has implications including (1) a better understanding of health behavior, and (2) a basis upon which interventions to improve the

public health of individuals and communities can be developed and evaluated.

Although research in this area continues at a rapid pace (Norman and Conner, 1996; Glanz *et al.*, 1997b, 2002; Fisher and Fisher, 2000; DiClemente *et al.*, 2002), the extent to which the field is truly moving forward has been questioned [e.g. (Weinstein, 1993; Zimmerman and Vernberg, 1994; Bandura, 1998; Smedslund, 2000a,b; Noar *et al.*, 2003; Ogden, 2003)]. That is, because we are conducting more research on health behaviors does not necessarily mean that we are adding substantive cumulative knowledge to this area of research. Approximately 10 years ago Neil Weinstein (Weinstein, 1993) quite clearly articulated the problems in this area of research and suggested some potential solutions. Weinstein made the case that we may not be moving forward toward a better understanding of health behavior because of a *lack of empirical comparisons* between the numerous HBTs that exist. He stated [(Weinstein, 1993), p. 324]:

...despite a large empirical literature, there is still no consensus that certain models of health behavior are more accurate than others, that certain variables are more influential than others, or that certain behaviors or situations are understood better than others. In general, researchers have failed to carry out the winnowing process that is necessary for scientific progress.

Not only did Weinstein challenge researchers to carry out more empirical comparisons of such theories, but he laid out guidelines for how one might design studies to do just that. Other researchers have echoed such sentiments for theory comparison research [e.g. (Zimmerman and Vernberg,

¹Department of Communication, University of Kentucky, Lexington, KY 40506-0042, USA

²Correspondence to: S. M. Noar;
E-mail: snoar2@uky.edu

1994; Murray-Johnson *et al.*, 2001; Nigg *et al.*, 2002a; Noar *et al.*, 2003)].

However, it is not clear whether anything has changed since Weinstein's (Weinstein, 1993) article. This leads us to ask the question: what is the best way for the field to move forward? The overriding purpose of the current article is to offer a critique of the current direction of HBT research and to suggest a new agenda of research in this area. We move to accomplish this task in three parts. First, we discuss in detail the problems within this area of research and how these problems may lead to fragmentation rather than cumulative knowledge. Second, we discuss the proliferation of HBT and why theory comparison is essential to this area of research. Finally, we reflect on ways that the field might move forward, including specific suggestions for new studies to be undertaken. It is *not* the purpose of the current article to review all the available evidence regarding what HBTs are most accurate in explaining health behavior. Rather, the purpose is to continue and extend a dialogue on the direction this field might move in—one that has been the topic of articles by theorists in the area [see, e.g. (Cummings *et al.*, 1980; Bandura, 1998; Rosenstock *et al.*, 1988), as well as recent special issues of *Health Education Research* (Connelly, 2002; Glasgow *et al.*, 2002; Nigg *et al.*, 2002b) and *Journal of Health Psychology* (Glanz and Maddock, 2000; Smedslund, 2000a,b)].

What is theory in the first place?

A theory has been defined as [(Glanz *et al.*, 1997a), p. 21]:

A set of interrelated concepts, definitions, and propositions that presents a *systematic* view of events or situations by specifying relations among variables in order to *explain* and *predict* events or situations.

Thus, according to this definition, HBT should describe (1) what variables are most important and (2) how the variables relate or interact, and perhaps

HBT should explain differences across situations, contexts, populations and with regard to different behaviors.

Within the study of health behavior, theories have been proposed at a variety of levels, including the individual, interpersonal, group, organizational and community levels. Further, theories vary in their focus on individual as compared to environmental determinants of behavior and cognitive as compared to affective determinants (Glanz *et al.*, 1997b; Crosby *et al.*, 2002). The primary focus of HBT has been at the individual level [see (Crosby *et al.*, 2002)] and thus this article focuses on individual-level HBTs.

Kegler *et al.* (Kegler *et al.*, 2002) describe a number of ways in which theories are developed, including refining existing theories (e.g. adding new constructs, making theory more parsimonious) as well as formalizing practitioner-based approaches into theory. In addition, Crosby *et al.* (Crosby *et al.*, 2002) describe a trajectory of how theories are developed, modified, and discarded. They state [(Crosby *et al.*, 2002), p. 5]:

Theory development is a dynamic process...as theories become less useful...they are modified or even discarded... As new theories are synthesized and embraced, they too are subject to empirical validation, and if they are found lacking, they are similarly discarded.

Although these may be ideal ways to develop, modify and discard theory, it is not clear that the literature has always followed such systematic methods. Specifically, it is not clear that significant modification of theories takes place very often and we would argue that rarely, if ever, are theories completely discarded. In fact, a recent study found that many HBTs are not falsifiable according to the field's current standards for testing such theories [see (Ogden, 2003)]. Further, when new theories are introduced, it is not always clear why. That is, new theories are often introduced to explain health phenomena when it is not clear that existing theories are inadequate for explaining such phenomena.

The problem: numerous theories, little consensus

Numerous individual-level HBTs exist in the literature. These include the Health Belief Model [HBM (Becker, 1974)], Theory of Reasoned Action [TRA (Ajzen and Fishbein, 1980)] and Theory of Planned Behavior [TPB (Ajzen and Madden, 1986)], Social Cognitive Theory [SCT (Bandura, 1986)], and the Transtheoretical Model [TTM (Prochaska and DiClemente, 1983)]. There are models specific to behavioral areas such as safer sex (Catania *et al.*, 1990; Fisher and Fisher, 1992) and alcohol use (Werch *et al.*, 2000; Jones *et al.*, 2001). Further, Fishbein (Fishbein, 2000) has proposed an integrated theory that combines concepts from several existing theories. In sum, there seems to be no lack of HBTs in the literature [also see (Glanz *et al.*, 1997b, 2002; DiClemente *et al.*, 2002)].

Which of the many theories that exist is most precise in explaining health-related behavior? Glanz *et al.* (Glanz *et al.*, 1997a) found in their review of the literature that the HBM, TRA/TPB, SCT and TTM were among the most widely used theories in the literature. Is one of these theories 'best' in terms of explaining health behavior? Reviews and meta-analyses of the HBM (Becker, 1974; Janz and Becker, 1984; Harrison *et al.*, 1992), TRA and TPB (Ajzen and Fishbein, 1980; Godin and Kok, 1996; Hausenblaus *et al.*, 1997; Albarracin *et al.*, 2001; Hagger *et al.*, 2002), SCT (Bandura, 1986, 1998; Strecher *et al.*, 1986), and TTM (Prochaska *et al.*, 1992, 1994; Rosen, 2000; Burkholder and Evers, 2002; Spencer *et al.*, 2002) all demonstrate support for their respective theories. Given this *apparent* parity in the literature, how do we move toward a better understanding of health behavior? How are researchers supposed to choose the most precise and fruitful theory or theories to apply in their studies? The recent Special Issue of *Health Education Research* highlighted 15 health intervention research projects, and in the articles each researcher was asked to describe how and why a certain theory was chosen to guide one's particular health intervention (Nigg *et al.*, 2002b). Var-

ious reasons were offered, although empirical studies showing the superiority of the chosen theory over other theories was rarely among the reasons.

Problems resulting from a lack of consensus

In addition, each of these theories can be viewed as producing 'mini-literatures', where knowledge accumulates with regard to theories on parallel tracks that inform a particular theory but offer little contribution to cumulative knowledge across theories. This creates several related problems.

First, as Weinstein (Weinstein, 1993) and others have discussed [e.g. (Norman and Conner, 1996; Rimer, 1997; Bandura, 1998; Rosenstock *et al.*, 1988; Institute of Medicine, 2002; Nigg *et al.*, 2002a; Noar *et al.*, 2003)], many of these theories contain constructs that are very similar (or identical), but use different terminology, creating the illusion that they are different. Whether similar constructs are indeed essentially the same is an important question that deserves much research attention. If constructs in different theories are the same, then this adds to our ability to synthesize knowledge *if* we are able to recognize this. Said another way, the *lack of consensus* regarding what to call certain constructs has resulted in a fragmented literature that could be better integrated if a common set of terminology was agreed upon.

In addition, constructs may come from different theoretical and conceptual origins, and have different names, yet when they are measured are essentially the same. Table I presents an example of constructs across different theories that are either similar or exactly the same. For instance, there is likely to be little difference between benefits and barriers, attitudes, positive and negative expectancies, and pros and cons. In fact, in a paper comparing alcohol expectancies and pros and cons, we found that they were quite similar (Noar *et al.*, 2003). On the other hand, in the exercise arena a study demonstrated that attitudes (from the TRA/TPB) and pros and cons are quite different (Jordan *et al.*, 2002). What is clear is that more work is needed in this area, and in addition that both conceptual and empirical points of view should be

Concept	General tenet of the concept <i>'Engaging in the behavior is likely if...'</i>	HBM	TRA	TPB	SCT	TTM
<i>Attitudinal beliefs</i>						
Appraisal of the positive and negative aspects of the behavior and expected outcome of the behavior	the positive aspects outweigh the negative aspects	benefits, barriers/health motive	behavioral beliefs and evaluation of those beliefs (attitudes)	behavioral beliefs and evaluation of those beliefs (attitudes)	outcome expectations/expectancies	pros, cons (decisional balance)
<i>Self-efficacy beliefs/beliefs about control over the behavior</i>						
Belief in one's ability to perform the behavior; confidence	one believes in their ability to perform the behavior	self-efficacy	–	perceived behavioral control	self-efficacy	self-efficacy/temptation
<i>Normative and norm-related beliefs and activities</i>						
Belief that others want you to engage in the behavior (and one's motivation to comply); may include actual support of others	one believes that people important to them want them to engage in the behavior; person has others' support	cues from media, friends (cues to action)	normative beliefs and motivation to comply (subjective norms)	normative beliefs and motivation to comply (subjective norms)	social support	helping relationships (process of change)
Belief that others (e.g. peers) are engaging in the behavior	one believes that other people are engaging in the behavior	–	–	–	social environment/norms; modeling	social liberation (process of change)
Responses to one's behavior that increase or decrease the likelihood one will engage in the behavior; may include reminders	one receives positive reinforcement from others or creates positive reinforcements for themselves	cues from media, friends (cues to action)	– ^a	– ^a	reinforcement	reinforcement management/stimulus control (processes of change)

Table 1. *Continued*

Concept	General tenet of the concept 'Engaging in the behavior is likely if...'	HBM	TRA	TPB	SCT	TTM
<i>Risk-related beliefs and emotional responses</i>						
Belief that one is at risk if one does not engage in the behavior, and that the consequences may be severe; may include actually experiencing negative emotions or symptoms and coping with them	one feels at risk with regard to a negative outcome or disease	perceived susceptibility/severity (perceived threat)	–	–	emotional coping responses/expectancies about environmental cues	dramatic relief (process of change)
<i>Intention/commitment/planning</i>						
Intending or planning to perform the behavior; setting goals or making a commitment to perform the behavior	one has formed strong behavioral intentions to engage in the behavior; one has set realistic goals or made a firm commitment to engage in the behavior	–	behavioral intentions	behavioral intentions	self-control/self-regulation	contemplation/preparation (stages of change); self-liberation (process of change)

Variable names in parentheses indicate that the variable(s) above it are part of that larger category, according to the theory.

^aBoth the TRA and TPB contain normative components that are conceptualized as *beliefs* in reinforcement (normative beliefs), rather than the actual reinforcement itself. It is not clear which conceptualization of these ideas is best for a theoretical framework.

carefully considered when such constructs are compared.

In addition, we as a field should never be *blind* advocates interested only in finding support for particular theories. Rather, we should put theories to the strongest possible tests and when such theories do not stand up to rigorous evaluation, appropriate modifications should be made. As Greenwald *et al.* [(Greenwald *et al.*, 1986), p. 227] state:

Theory obstructs research progress when...the researcher has more faith in the correctness of the theory than in the suitability of the procedures that were used to test it.

Ogden (Ogden, 2003) recently examined a systematic sample of 47 HBT studies and found that when data did not demonstrate support for certain aspects of theories, the authors tended to offer various explanations, *none* of which included that the theory may be incorrect. Ogden concluded that such theories cannot be tested because data are used to support HBT but are rarely if ever used to refute HBT. In fact, though a basic tenet of theory development is that theories should be falsifiable, this work by Ogden strongly suggests that many HBT's are in fact *not* falsifiable. In addition, many have suggested that a number of HBT's are not well specified in terms of the relations between variables [e.g. Fisher and Fisher, 1992; Weinstein, 1993; Rimer, 1997)], making them difficult to test and subsequently verify or falsify.

Further, as already discussed, there are a finite number of questions that HBT should address. Our reading of the literature is that there is *some* consensus regarding which variables are most important to health behavior, though we use the word *some* generously [see Table I (Fishbein *et al.*, 2001; Noar *et al.*, 2004)]. There is much less consensus as to how the variables combine in an equation to predict behavior (Weinstein, 1993; Fishbein *et al.*, 2001). This is a much more complex question, which may be facilitated by answers to the first question. Finally, whether there are differences (or similarities) across different behaviors and situations has been addressed somewhat within

theories [e.g. (Madden *et al.*, 1992; Prochaska *et al.*, 1994)], but has rarely been addressed across theories. This is a question that may be better facilitated by answers to the more basic questions.

Finally, we should note that an issue that has been discussed in the literature is whether or not a single theory is appropriate across multiple behaviors, as compared to theories that are specific to certain behaviors. For instance, where illness avoidance and perceived threat are the most salient issues, a theory such as the HBM may be most appropriate (Biddle and Nigg, 2000; Murray-Johnson *et al.*, 2001). For behaviors that are more rational in nature and in which the intention-behavior link is strong, theories such as the TRA/TPB may be most appropriate (Biddle and Nigg, 2000; Murray-Johnson *et al.*, 2001). In addition, some suggest that stage models such as the TTM may be most applicable to deliberate behaviors (e.g. exercise) and less applicable to automatic behaviors that are simplistic [e.g. seatbelt use (Norman and Conner, 1996)]. Finally, a number of researchers point to the need for theoretical approaches to the *maintenance* of behavior change being distinct from initiation of behavior change [see (Wing *et al.*, 2000)]. Clearly, more empirical work on the issue of specific versus general theories is warranted.

What is the best way to move forward?

Is the proposition and study of numerous HBTs the best way for the field to move forward? We next present three possible directions that the field could move in.

The current direction: proliferation and testing of theories

Theory drives research. It serves as a guide for knowing what variables to measure, how to measure them, and how to combine them. It also serves as a framework for aiding researchers in developing and evaluating intervention approaches. If a researcher believes that existing theories are inappropriate or incomplete, he or she can extend upon an

existing theory or create a new theory altogether. When Ajzen and Madden (Ajzen and Madden, 1986) observed that the TRA could be improved, they proposed the TPB. Their research demonstrated that the addition of perceived behavioral control added variance in the prediction of health behaviors (Madden *et al.*, 1992) and subsequent research found success in using the TPB as a guide for developing interventions [see (Hardeman *et al.*, 2002)]. In addition, out of Prochaska and DiClemente's (Prochaska and DiClemente, 1983) work on smoking cessation came a new paradigm—a stage paradigm for understanding health behavior. This has resulted in successful intervention approaches across a variety of health behaviors [see (Burkholder and Evers, 2002)]. The current research model gives investigators complete autonomy to create their own theories and interventions based upon their own judgment.

Further, Glanz and Maddock (Glanz and Maddock, 2000) argue that out of what is sometimes a confusing literature, the best and brightest will emerge. That is, every health researcher could propose a theory, but only those best supported empirically and those that resonate with researchers would proliferate further inquiry. This is likely the current state of the literature, as numerous theories exist, but some theories are used much more than others. Researchers may use theory for other reasons than suggested by Glanz and Maddock (Glanz and Maddock, 2000), however. For instance, using a particular theory because it is easy to understand or it is the one learned during one's academic training. In addition, the fragmentation of the HBT literature because of multiple theories and its potential slowing of our understanding of health behavior have already been discussed as problems with this approach.

Another direction: the case for integration

If the case is made for theoretical integration, what would that mean? An integrated theory might take the constructs with the most support from varying theories and combine them into a single theory. This theory could then be subjected to rigorous

testing across behaviors and situations, and refined as necessary. An integrated theory would first require that theorists agree on common conceptualizations and names for similar constructs.

There have been various attempts to create integrated theories of health behavior, and a recent example is Fishbein's (Fishbein, 2000) integrated theory [see also (Institute of Medicine, 2002)]. The core constructs of the theory are essentially the TRA constructs with the addition of self-efficacy. The theory also includes constructs such as demographics and personality variables as well as skills and environmental constraints. Fishbein's (Fishbein, 2000) integrated model grew out of a theorists' workshop that took place in 1991, in which many prominent theorists came together to identify core determinants of health behavior. The theorists produced a chapter that listed and described eight variables believed to be most important to health behavior and specifically to safer sexual behavior [see (Fishbein *et al.*, 2001)]. They discussed similarities between constructs and theories, though a common set of terminology was not proposed. The way in which these constructs combine to effect behavior was *not* agreed upon and was discussed as an 'unresolved issue'. Although Fishbein (Fishbein, 2000) suggests one conceptualization, additional conceptualizations of the same variables could also be tested. However, it is likely that many researchers would disagree with these core determinants, which is a major difficulty with the integrated approach.

A new direction: the case for theory comparison and beyond

While both proliferation and integration of theories have many problems associated with them, comparison of theories, we believe, is the most compelling direction to move in. Why empirically compare theories of health behavior to one another? Several reasons could be given, not the least of which is that we may not truly know which theories are most accurate if we do not do so. Consumers faced with making decisions regarding buying products (e.g. a stereo system, an automobile) often rely on product comparisons conducted by magazines

such as Consumer Reports. How else can one know what product is best, if we do not compare one to the other? Theories are, in fact, academic products that are fashioned from well thought out conceptual work. Do they not justify comparison if they all claim to explain the same phenomena, yet have fundamental differences among them?

Theoretical comparisons are in many cases not difficult to conduct, they yield important results and we have the potential to learn much about HBT as a result of such comparisons. The fact that theories have so many similar constructs demands that we (1) try and discover what the best conceptualization of those constructs is, and (2) compare theories to discover how these constructs combine and result in the enactment of health behavior. Further, since many constructs in theories are similar or the same, when we measure one variable (e.g. self-efficacy) it may cut across many theories, allowing for a reasonable level of response burden on study participants.

Despite this, few data-based empirical comparisons exist in the literature. Weinstein (Weinstein, 1993) reported that out of 205 theoretically based articles published between 1974 and 1991, only 10 articles mentioned more than one theory and only four of these 10 were empirical comparisons. In order to examine an updated state of empirical comparisons in the literature, we conducted a comprehensive search of the PsycInfo database through June of 2003. We searched for articles that were classified in PsycInfo as *health-related* articles (using the keyword *health* which includes health behavior, health attitude, etc.), and included peer-reviewed articles, book chapters, books and dissertations. We searched for the major theories discussed in this article (HBM, TRA/TPB, TTM, SCT), using advanced search commands so that we would not count articles twice. In addition, the PsycInfo search examined the title, abstract and keywords for potential matches to these criteria.

As can be seen in Figure 1, the vast majority of articles utilize just one theory (though we should note that some of the TRA/TPB studies compared the TRA with the TPB and we did not distinguish between them because these theories are so similar to one another). Only 178 (6%) out of a total of 2901

unique citations contained two or more theories in the search record. Further, when we more closely examined these 178 citations, we found the following. First, nine articles were excluded for various reasons (e.g. they were improperly coded in PsycInfo and had no relevance here). This left 169 articles, which were broken down into various categories in Figure 2. As one can see, $N = 67$ were not




Fig. 1. Number of unique citations (total $N = 2901$) for individual versus multiple theories found in PsycInfo through June 2003.




Fig. 2. Categorization of $N = 169$ multiple theory citations found in PsycInfo through June 2003.

empirical studies, but rather were descriptive theoretical articles [e.g. (Sutton, 1987; Bandura, 1998)]. Next, $N = 18$ of the studies were intervention projects [e.g. (Pinto *et al.*, 2002)], while $N = 4$ were categorized as ‘other,’ which contained studies such as qualitative focus group investigations [e.g. (Levy and Bavendam, 1995)]. Finally, $N = 80$ were empirical articles that utilized two or more of the theories or concepts from the theories (these were correlational/behavioral prediction articles). Thus, it is clear that more investigators are using multiple theories to inform their empirical investigations. Of these 80 empirical articles, seven were published before 1990, 47 were published between 1990 and 1999, and 26 were published between 2000 and June of 2003.

However, the major caveat here is that most of these articles were not empirical comparisons of two or more theories. Rather, many were ‘predictors’ articles that looked at *selected* concepts derived from various theories and examined their ability to predict health behavior [e.g. (Manfredi *et al.*, 1998; Adih and Alexander, 1999)]. In fact, when we examined these 80 articles more closely, we identified only 13 published articles (in a journal or book chapter) that were true theoretical comparisons (0.4% of the 2901 total citations). Thus, though many more studies are utilizing multiple theories, empirical comparisons of these theories are still extremely rare in the research literature.

If more such comparisons were undertaken, we might start to have a clearer picture of the similarity or difference of constructs. The field could come to better agreement on what names to give to certain concepts, resulting in greater consensus and a shared conceptual language. And, although the end result might not be one unified theory of health behavior, we would certainly be moving in the direction of theories and models that truly integrate what we are learning from research. If we do not move in this direction, then we wish to pose a question. What is the point of the massive literature on theory testing? If it is to find support for numerous theories, then we are doing quite well. If it is to understand which concepts and theories are most related to health

behavior, then some changes to the way we conduct such research are necessary.

How should researchers empirically compare theories?

We next consider how one might empirically compare theories and concepts from theories. We believe that studies that empirically compare individual *concepts* from theories are important because they suggest commonalties (or refute the idea of commonalties) across theories. A specific example of this are the numerous types of attitudinal beliefs described in Table I. If one is able to say with confidence that some of these concepts are identical, then response burden on participants will be lessened. On the other hand, if these concepts are indeed shown to be conceptually distinct, then more concepts will need to be measured. Comparison of *concepts* can be examined using a variety of correlational techniques (e.g. regression) to assess whether or not unique variance is added when a second concept is considered above and beyond a first concept (Noar *et al.*, 2003). In addition, structural equation modeling (SEM) techniques can be used to examine measurement models of concepts in order to examine their similarity or difference (Jordan *et al.*, 2002).

A more complex issue is how one might carry out comparisons of entire theories as advocated in this article. In order to examine theory comparison articles in the literature, we compiled articles from the PsycInfo search conducted, and supplemented the search with additional articles identified from the reference sections of the 13 theory comparison articles. All studies had to meet the following criteria in order to be included. First, they had to be English language journal articles or book chapters (dissertations were excluded). Second, they had to be non-redundant studies. If results from one study were published multiple times, only one article was chosen. Finally, they had to truly test theories, rather than testing *selected components* from theories. Although our list of theoretical comparisons is not exhaustive, we believe that our

search captured a number of important comparisons in the literature.

Nineteen studies met the criteria and are described in Table II. All of the studies were correlational in nature and used survey data; none were lab-based or experimental studies. In terms of analytic techniques, by far the most common technique used was multiple regression, in which each theory was tested separately and the R^2 values as well as the standardized regression coefficients (β weights) were compared. This allows one to examine the overall variance accounted for in the dependent variable (DV) as well as examine which specific variables were responsible for prediction of that variance. A second analytic technique, used in far fewer of the studies, was SEM. The approach here is similar to regression in that the overall fit of models is examined using various SEM fit indices and R^2 , while the contribution of specific variables is examined by standardized path coefficients. While multiple regression is a strong technique for testing and comparing theories, many would consider SEM (and path analysis) to be superior in part because it allows for several layers of endogenous variables (e.g. attitudes predict intentions which in turn predict behavior).

A number of these studies have some strengths, as well as some having weaknesses. Rather than discuss one particular study as a model study, we focus on characteristics of strong comparisons (see Table III). Most of these characteristics are self-explanatory and thus we elaborate here only briefly. Some methodological strengths of studies included longitudinal designs, using SEM, having a strong sample size, and utilizing non-college samples and multiple samples. Some conceptual strengths included examining more than one behavior, more than one DV, more than two theories and examining an integrated model based on the results of the comparison. These methodological and conceptual strengths of studies are recommended in future theory comparison studies where possible. Further, theories vary in a number of ways such as which DV is most important and whether or not past behavior or demographics are important. Thus, we recommend that theory comparison studies in turn

test theories in a number of ways within a single study, including testing multiple DVs [e.g. (Bish *et al.*, 2001)] as well as testing theories with and without past behavior [e.g. (Quine *et al.* 2000)] and demographics controlled for [e.g. (Vanlandingham *et al.*, 1995)]. We also note that studies examining multiple health behaviors allow us to examine the important question of how generalizable HBTs are across behaviors. Finally, studies examining integrated models [e.g. (Wulfert *et al.*, 1996)] begin to move us in the direction of where this line of inquiry may ultimately take us.

A brief note on meta-analysis

One may wonder whether meta-analyses and systematic literature reviews have the potential to integrate and compare HBTs in ways we advocate in this article. First, we note that without a doubt, meta-analysis has helped synthesize the literature on correlates of health behavior in a way that few if any other techniques could do. However, meta-analyses and research reviews often examine effect sizes based on bivariate correlations [e.g. (Gerrard *et al.*, 1996; Sheeran *et al.*, 1999)] or significance ratios (Janz and Becker, 1984). Although these are excellent contributions to the literature, they contribute more to our understanding of the relationship of individual variables to behavior rather than theories. Some have recently used meta-analysis as more of a theory testing technique [e.g. (Albarracin *et al.*, 2001; Hagger *et al.*, 2002)] and we view this as an excellent step in the right direction.

Lipsey and Wilson (Lipsey and Wilson, 2001) note that it is still quite difficult to use meta-analysis to analyze findings from multivariate analyses such as multiple regression and SEM. Thus, while meta-analysis can certainly bring some consensus to this area, there are currently some difficulties in using it as a multivariate theory testing and comparison technique. In fact, we are not aware of any published studies that have used meta-analysis to compare HBTs (with the exception of TRA/TPB meta-analyses, which have just one variable to

Table II. Empirical comparisons of HBTs (N = 19)

Article	Behavioral domain	Study design	Sample(s)	Theories compared	DV	Comparative strategy
Bish <i>et al.</i> (2000)	cervical cancer screening	longitudinal (baseline, 3 months)	N = 133 females, mean age: 38.1 (UK)	HBM, TPB	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Boyd and Wandersman (1991)	safer sex	longitudinal (baseline, 3 months)	N = 190 undergraduates, mean age: 18.9 (US)	TRA, TABM	intention, behavior	multiple regression for complete theories; R^2 and <i>significance of</i> β weights compared
Conner and Graham (1993)	safer sex	cross-sectional	N = 218 undergraduates, age not reported (UK)	HBM, TPB	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Conner and Norman (1994)	health screening	longitudinal (baseline, 6 months)	N = 407 individuals, age not reported (UK)	HBM, TPB	intention, behavior	correlations for theoretical components. multiple regression for complete theories (path analysis); R^2 and β weights compared
Garcia and Mann (2003)	resisting dieting and breast self-exam	cross-sectional	study 1: N = 159 female undergraduates; study 2: N = 120 female undergraduates; ages not reported (US)	HBM, HBM plus self-efficacy, TRA, TPB, HAPA	intention	multiple regression for complete theories; R^2 and β weights compared
Hennig and Knowles (1990)	cervical cancer screening	cross-sectional	N = 144 females, mean age: 54 (US)	HBM, TRA	intention	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Hill <i>et al.</i> (1985)	breast self-examination, cervical cancer screening	cross-sectional	N = 123 females, median age: 34 (US)	HBM, TRA, SPM	intention	multiple regression for complete theories; R^2 and β weights compared
Kloeblen <i>et al.</i> (1999)	breast-feeding	cross-sectional	N = 1001 females, mean age: 23 (US)	TRA, TTM	intention, stage of change behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and partial R^2 compared
Mullen <i>et al.</i> , 1987	smoking, exercise, dietary habits	longitudinal (baseline, 8 months)	N = 326 individuals aged 17–65 (US)	HBM, TRA, PRECEDE		multiple regression for complete theories; R^2 and β weights compared
Murray-Johnson <i>et al.</i> (2001)	reproductive health	cross-sectional	N = 7540 individuals (Ghana); N = 3621 individuals (Nepal); N = 2000 individuals (Nicaragua); ages not reported	HBM, TRA, SCT, EPPM	exposure, knowledge, attitudes, fear and danger control, intention,	correlations for theoretical components. SEM for complete theories; SEM fit indices compared

Article	Behavioral domain	Study design	Sample(s)	Theories compared	DV	Comparative strategy
Oliver and Berger (1979)	inoculation behavior (flu shot)	cross-sectional	study 1: $N = 323$ undergraduates; study 2: $N = 469$ individuals; ages not reported (US)	HBM, TRA	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Quine <i>et al.</i> (2000)	bicycle helmet use	longitudinal (baseline, 1 month)	$N = 162$ males aged 11–18 (UK)	HBM, TPB	behavior	correlations for theoretical components. multiple regression for complete theories (path analysis); R^2 and β weights compared
Reid and Christensen (1988)	medication compliance for urinary tract infection	longitudinal (baseline, 10 days)	$N = 107$ undergraduate and other females aged 16–79 (US)	HBM, TRA	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Seibold and Roper (1979)	cervical cancer screening	cross-sectional	$N = 93$ undergraduate and other females aged 18–90 (US)	TRA, TABM	intention	multiple regression for theoretical components and complete theories; multiple correlation (R) and β weights compared
Seydel <i>et al.</i> (1990)	cancer prevention behaviors (e.g. breast self-exam, cancer screenings)	cross-sectional	study 1: $N = 358$ females, mean age: 48; study 2: $N = 256$ individuals, mean age: 38 (US)	HBM, PMT	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Vanlandingham <i>et al.</i> (1995)	safer sex	cross-sectional	$N = 1472$ males, mean age: 23 (Thailand)	HBM, TRA	behavior	correlations for theoretical components. logistic regression for complete theories; odds ratios and percent correctly classified compared
Warwick <i>et al.</i> (1993)	safer sex	longitudinal (baseline, 1 month)	$N = 138$ undergraduates, mean age: 18.6 (US)	HBM, TRA	intention, behavior	correlations for theoretical components. multiple regression for complete theories; R^2 and β weights compared
Wulfert and Wan (1995)	safer sex	two cross-sectional studies, one longitudinal study (baseline, 3 months)	study 1: $N = 496$ undergraduates, mean age: 20.3; study 2: $N = 421$ individuals, mean age: 46; study 3: $N = 105$ undergraduates, mean age: 20.6. (US)	HBM, TRA, SCT	intention, behavior	SEM for complete theories; SEM fit indices compared (overall indices and specific standardized path coefficients)
Wulfert <i>et al.</i> (1996)	safer sex	cross-sectional	$N = 153$ males, mean age: 37.4 (US)	HBM, TRA, SCT	behavior	correlations for theoretical components. SEM for complete theories; SEM fit indices compared (overall indices and specific standardized path coefficients)

DV = dependent variable; TABM = Triandis attitude behavior model; HAPA = health action process approach; SPM = subjective probability model; PRECEDE = Predisposing, reinforcing, enabling factors model; EPPM = extended parallel process model; PMT = Protection motivation theory; SEM = structural equation modeling.

manipulate to make it a theory comparison). We encourage researchers to examine novel ways to use meta-analysis as a theory testing and theory comparison technique.

Moving the dialogue forward

Since so much research on health behavior is theory based, it is crucial that the field ‘audit’ itself to be sure that we are moving in the right direction. Glanz *et al.*’s (Glanz *et al.*, 1997a) review of the health education literature published between mid-1992 and 1994 found that 526 articles (45%) of 1174 utilized some theory or model. This is no small amount of research and we as a field owe it to ourselves to continue this dialogue on the best way to move forward. Since researchers value and apply theory in the study of health behavior, we believe it remains an important task to refine theory and move toward consensus in the field where possible.

What theory or theories predict behavior most precisely, and what are the key principles of behavior change? We think it is fair to say that at this point in the literature, we are not entirely sure. Table I in this article presented some of the major concepts of HBTs and suggested some key principles of behavior change. It is now up to researchers in the

Table III. Summary of strong characteristics of theory comparison studies (total N = 19)

Study characteristic	No. of Studies
Longitudinal study	8/19
Used SEM	3/19
Included past behavior in some or all model tests	3/19
Included demographics in some or all model tests	4/19
Included non-college participants in some or all samples	15/19
Had strong sample size in one or more samples (>200)	10/19
Utilized multiple samples in model testing	5/19
Utilized samples from more than one country	1/19
Had >1 DV (e.g. intention and behavior)	11/19
Examined more than one behavior	4/19
Compared more than two theories	6/19
Empirically examined an integrated model	6/19

Table IV. Suggested important theory comparison questions for the field

Research questions	Examples of application
1. What is the extent of similarity or difference regarding constructs from differing theories that appear to be similar or the same in nature?	Is there any difference among behavioral beliefs (TRA), benefits and barriers (HMB), outcome expectancies (SCT), and decisional balance (TTM)? Are there substantive conceptual differences between perceived behavioral control (TPB) and self-efficacy (SCT) or are they essentially the same?
2. Are certain theories or elements of theories more useful in terms of predicting behavior or behavior change as compared to others?	Does the stage-based TTM or continuum-based TRA provide better prediction of behavior?
3. Are the combinatorial rules for one theory better supported empirically than for other theories?	Are health behaviors mediated by intention formation (as TRA/TPB suggest) or not (as the HBM suggests)?
4. Are certain theories or elements of theories better predictors of addictive behaviors (as opposed to non-addictive behaviors)?	Are SCT constructs better at predicting addictive behaviors, while TPB constructs are better at predicting non-addictive behaviors?
5. Are certain theories or elements of theories better predictors of one-time behaviors (e.g. vaccinations) as opposed to behaviors that must be maintained over time (e.g. exercise)?	Are HBM constructs better at predicting one-time behaviors, while constructs from the TTM better at predicting behaviors that must be maintained?
6. Are certain theories or elements of theories better predictors of cessation behaviors (e.g. smoking cessation) as opposed to behaviors that must be adopted (e.g. exercise)?	Do theories such as the TRA/TPB predict adoption behaviors better than cessation behaviors, or vice versa?
7. Are certain theories or elements of theories better predictors in different cultures?	Is self-efficacy a better predictor in cultures with more of a focus on individualism, and beliefs and norms better predictors in cultures with more of a focus on collectivism?
8. Is there one set of behavior change principles that can account for all health behaviors, or are they different according to different behaviors, cultures and contexts?	Questions 4–7 address this

field to uncover how these and other principles work together and result in the enactment of health behaviors, by asking important questions and putting them to empirical tests (see Table IV). We applaud attempts to move in this direction and integrate knowledge in this area, such as Fishbein's (Fishbein, 2000) integrative model and Prochaska *et al.*'s (Prochaska, 1992) TTM. However, the problem remains that these two theories, as one example, have many differences. Therefore, we must rely on empirical comparisons of such theories to understand which operates best. Rimer [(Rimer, 1997), p. 146] reminds us that 'Theory is not theology. Theory needs questioners more than loyal followers'. What the field needs are researchers who are willing to put these concepts and theories to the strongest possible tests, so we can progress further in understanding health behavior and health behavior change.

Acknowledgements

We gratefully acknowledge the thoughtful suggestions of the three anonymous reviewers.

References

- Adih, W.K. and Alexander, C.S. (1999) Determinants of condom use to prevent HIV infection among youth in Ghana. *Journal of Adolescent Health*, **24**, 63–72.
- Ajzen, I. and Fishbein, M. (1980) *Understanding Attitudes and Predicting Social Behavior*. Prentice-Hall, Englewood Cliffs, NJ.
- Ajzen, I. and Madden, T.J. (1986) Prediction of goal-directed behavior: attitudes, intentions and perceived behavioral control. *Journal of Experimental Social Psychology*, **22**, 453–474.
- Albaracin, D., Johnson, B.T., Fishbein, M. and Muellerleile, P.A. (2001) Theories of reasoned action and planned behavior as models of condom use: a meta-analysis. *Psychological Bulletin*, **127**, 142–161.
- Bandura, A. (1986) *Social Foundations of Thought and Action: A Social Cognitive Theory*. Prentice-Hall, Englewood Cliffs, NJ.
- Bandura, A. (1998) Health promotion from the perspective of social cognitive theory. *Psychology and Health*, **13**, 623–649.
- Becker, M.H. (ed.) (1974) The health belief model and personal health behavior. *Health Education Monographs*, **2**(4).
- Biddle, S.J.H. and Nigg, C.R. (2000) Theories of exercise behavior. *International Journal of Sport Psychology*, **31**, 290–304.
- Bish, A., Sutton, S. and Golombok, S. (2000) Predicting uptake of a routine cervical smear test: a comparison of the health belief model and the theory of planned behaviour. *Psychology and Health*, **15**, 35–50.
- Boyd, B. and Wandersman, A. (1991) Predicting undergraduate condom use with the Fishbein and Ajzen and the Triandis attitude-behavior models: implications for public health interventions. *Journal of Applied Social Psychology*, **21**, 1810–1830.
- Burkholder, G.J. and Evers, K.E. (2002) Application of the transtheoretical model to several problem behaviors. In Burbank, P.M. and Riebe, D. (eds), *Promoting Exercise and Behavior Change in Older Adults: Interventions with the Transtheoretical Model*. Springer, New York, pp. 85–145.
- Catania, J.A., Kegeles, S.M. and Coates, T.J. (1990) Towards an understanding of risk behavior: an AIDS risk reduction model (ARRM) *Health Education Quarterly*, **17**, 53–72.
- Connelly, J. (2002) The behavior change consortium studies: missed opportunities—individual focus with an inadequate engagement with personhood and socio-economic realities. *Health Education Research: Theory and Practice*, **17**(6), 691–695.
- Conner, M. and Graham, S. (1993) Situational and attitudinal influences upon students' intention to use condoms. In Schroeder, H., Reschke, K., Johnston, M. and Maes, S. (eds), *Health Psychology: Potential in Diversity*. S. Roderer, Regensburg, pp. 91–100.
- Conner, M. and Norman, P. (1994) Comparing the health belief model and the theory of planned behaviour in health screening. In Rutter, D.R. and Quine, L. (eds), *Social Psychology and Health: European Perspectives*. Avebury, Aldershot, pp. 1–24.
- Crosby, R.A., Kegler, M.C. and DiClemente, R.J. (2002) Understanding and applying theory in health promotion practice and research. In DiClemente, R.J., Crosby, R.A. and Kegler, M.C. (eds), *Emerging Theories in Health Promotion Practice and Research: Strategies for Improving the Public Health*. Jossey-Bass, San Francisco, CA, pp. 1–15.
- Cummings, K.M., Becker, M.H. and Maile, M.C. (1980) Bringing the models together: an empirical approach to combining variables used to explain health actions. *Journal of Behavioral Medicine*, **3**, 123–145.
- DiClemente, R.J., Crosby, R.A. and Kegler, M.C. (eds) (2002) *Emerging Theories in Health Promotion Practice and Research: Strategies for Improving the Public Health*. Jossey-Bass, San Francisco, CA.
- Fishbein, M. (2000) The role of theory in HIV prevention. *AIDS Care*, **12**, 273–278.
- Fishbein, M., Triandis, H.C., Kanfer, F.H., Becker, M., Middlestadt, S.E. and Eichler, A. (2001) Factors influencing behavior and behavior change. In Baum, A., Revenson, T.A. and Singer, J.E. (eds), *Handbook of Health Psychology*. Lawrence Erlbaum, Mahwah, NJ, pp. 3–17.
- Fisher, J.D. and Fisher, W.A. (1992) Changing AIDS-risk behavior. *Psychological Bulletin*, **111**, 455–474.
- Fisher, J.D. and Fisher, W.A. (2000) Theoretical approaches to individual-level change in HIV risk behavior. In Peterson, J.L. and DiClemente, R.J. (eds), *Handbook of HIV Prevention*. Kluwer Academic/Plenum, New York, pp. 3–55.
- Garcia, K. and Mann, T. (2003) From 'I wish' to 'I will': social-cognitive predictors of behavioral intentions. *Journal of Health Psychology*, **8**, 347–360.

- Gerrard, M., Gibbons, F.X. and Bushman, B.J. (1996) Relation between perceived vulnerability to HIV and precautionary sexual behavior. *Psychological Bulletin*, **119**, 390–409.
- Glanz, K. and Maddock, J. (2000) On judging models and theories: Research and practice, psychology and public health. *Journal of Health Psychology*, **5**, 151–154.
- Glanz, K., Lewis, F.M. and Rimer, B.K. (1997a) Linking theory, research and practice. In Glanz, K., Lewis, F.M. and Rimer, B.K. (eds), *Health Behavior and Health Education: Theory, Research and Practice*, 2nd edn. Jossey-Bass, San Francisco, CA, pp. 19–35.
- Glanz, K., Lewis, F.M. and Rimer, B.K. (eds) (1997b) *Health Behavior and Health Education: Theory, Research and Practice*, 2nd edn. Jossey-Bass, San Francisco, CA.
- Glanz, K., Rimer, B.K. and Lewis, F.M. (eds) (2002) *Health Behavior and Health Education: Theory, Research and Practice*, 3rd edn. Jossey-Bass, San Francisco, CA.
- Glasgow, R.E., Dziewaltowski, D.A., Estabrooks, P.A., Klesges, L.M. and Bull, S.S. (2002) Response from the behavior change consortium representativeness and translation work group: the issue is one of impact, not of world view or preferred approach. *Health Education Research*, **17**, 696–699.
- Godin, G. and Kok, G. (1996) The theory of planned behavior: a review of its applications to health-related behaviors. *American Journal of Health Promotion*, **11**, 87–98.
- Greenwald, A.G., Pratkanis, A.R., Leippe, M.R. and Baumgardner, M.H. (1986) Under what conditions does theory obstruct research progress? *Psychological Review*, **93**, 216–229.
- Hagger, M.S., Chatzisarantis, N.L.D. and Biddle, S.J.H. (2002) A meta-analytic review of the theories of reasoned action and planned behavior in physical activity: predictive validity and the contribution of additional variables. *Journal of Sport and Exercise Psychology*, **24**, 3–32.
- Hardeman, W., Johnston, M., Johnston, D.W., Bonetti, D., Wareham, N.J. and Kinmonth, A.L. (2002) Application of the theory of planned behaviour in behaviour change interventions: a systematic review. *Psychology and Health*, **17**, 123–158.
- Harrison, J.A., Mullen, P.D. and Green, L.W. (1992) A meta-analysis of studies of the health belief model with adults. *Health Education Research*, **7**, 107–116.
- Hausenblaus, H., Carron, A.V. and Mack, D.E. (1997) Application of the theories of reasoned action and planned behavior to exercise behavior: a meta-analysis. *Journal of Sport and Exercise Psychology*, **19**, 36–51.
- Hennig, P. and Knowles, A. (1990) Factors influencing women over 40 years to take precautions against cervical cancer. *Journal of Applied Social Psychology*, **20**, 1612–1621.
- Hill, D., Gardner, G. and Rassaby, J. (1985) Factors predisposing women to take precautions against breast and cervix cancer. *Journal of Applied Social Psychology*, **15**, 59–79.
- Institute of Medicine (2002) *Speaking of Health: Assessing Health Communication Strategies for Diverse Populations*. National Academies Press, Washington, DC.
- Janz, N.K. and Becker, M.H. (1984) The health belief model: a decade later. *Health Education Quarterly*, **11**, 1–47.
- Jones, B.T., Corbin, W. and Fromme, K. (2001) A review of expectancy theory and alcohol consumption. *Addiction*, **96**, 57–72.
- Jordan, P.J., Nigg, C.R., Norman, G.J., Rossi, J.S. and Benisovich, S.V. (2002) Does the transtheoretical model need an attitude adjustment? Integrating attitude with decisional balance as predictors of stage of change for exercise. *Psychology of Sport and Exercise*, **3**, 65–83.
- Kegler, M.C., Crosby, R.A. and DiClemente, R.J. (2002) Reflections on emerging theories in health promotion practice. In DiClemente, R.J., Crosby, R.A. and Kegler, M.C. (eds), *Emerging Theories in Health Promotion Practice and Research: Strategies for Improving the Public Health*. Jossey-Bass, San Francisco, CA, pp. 386–395.
- Kloeblen, A.S., Thompson, N.J. and Miner, K.R. (1999) Predicting breast-feeding intention among low-income pregnant women: a comparison of two theoretical models. *Health Education and Behavior*, **26**, 675–688.
- Levy, R.I. and Bavendam, T.G. (1995) Promoting women's urologic self-care: five single-case replications. *Research on Social Work Practice*, **5**, 430–441.
- Lipsey, M.W. and Wilson, D.B. (2001) *Practical Meta-analysis*. Sage, Newbury Park, CA.
- Madden, T.J., Ellen, P.S. and Ajzen, I. (1992) A comparison of the theory of planned behavior and the theory of reasoned action. *Personality and Social Psychology Bulletin*, **18**, 3–9.
- Manfredi, C., Lacey, L., P., Warnecke, R. and Petraitis, J. (1998) Sociopsychological correlates of motivation to quit smoking among low-SES African American women. *Health Education and Behavior*, **25**, 304–318.
- Mullen, P.D., Hersey, J.C. and Iverson, D.C. (1987) Health behavior models compared. *Social Science and Medicine*, **24**, 973–981.
- Murray-Johnson, L., Witte, K., Boulay, M., Figueroa, M.E., Storey, D. and Tweedie, I. (2001) Using health education theories to explain behavior change: a cross-country analysis. *International Quarterly of Community Health Education*, **20**, 323–345.
- Nigg, C.R., Allegrante, J.P. and Ory, M. (2002a) Theory-comparison and multiple-behavior research: common themes advancing health behavior research. *Health Education Research: Theory and Practice*, **17**, 670–679.
- Nigg, C. R., Allegrante, J. P. and Ory, M. (eds) (2002b) Special issue: Behavior Change Consortium (BCC). *Health Education Research*, **17**, 493–679.
- Noar, S.M., Laforge, R.G., Maddock, J.E. and Wood, M.D. (2003) Rethinking positive and negative aspects of alcohol use: suggestions from a comparison of alcohol expectancies and decisional balance. *Journal of Studies on Alcohol*, **64**, 60–69.
- Noar, S.M., Anderman, E.M., Zimmerman, R.S. and Cupp, P.K. (2004) Fostering achievement motivation in health education: Are we applying relevant theory to school-based HIV prevention programs? *Journal of Psychology and Human Sexuality*, in press
- Norman, P. and Conner, M. (1996) The role of social cognition models in predicting health behaviours: Future directions. In Conner, M. and Norman, P. (eds), *Predicting Health Behaviour*. Open University Press, Buckingham, pp. 179–225.
- Ogden, J. (2003) Some problems with social cognition models: a pragmatic and conceptual analysis. *Health Psychology*, **22**, 424–428.
- Oliver, R.L. and Berger, P.K. (1979) A path analysis of preventive health care decision models. *Journal of Consumer Research*, **6**, 113–122.
- Pinto, B., Friedman, R., Marcus, B.H., Kelley, H., Tennstedt, S. and Gillman, M.W. (2002) Effects of a computer-based,

- telephone-counseling system on physical activity. *American Journal of Preventive Medicine*, **23**, 113–120.
- Prochaska, J.O. and DiClemente, C.C. (1983) Stages and processes of self-change of smoking: Toward an integrative model of change. *Journal of Consulting and Clinical Psychology*, **51**, 390–395.
- Prochaska, J.O., DiClemente, C.C. and Norcross, J.C. (1992) In search of how people change: applications to addictive behaviors. *American Psychologist*, **47**, 1102–1114.
- Prochaska, J.O., Velicer, W.F., Rossi, J.S., Goldstein, M.G., Marcus, B.H., Rakowski, W., Fiore, C., Harlow, L.L., Redding, C.A., Rosenbloom, D. and Rossi, S. (1994) Stages of change and decisional balance for 12 problem behaviors. *Health Psychology*, **13**, 39–46.
- Quine, L., Rutter, D.R. and Arnold, L. (2000) Comparing the theory of planned behaviour and the health belief model: the example of safety helmet use among schoolboy cyclists. In Norman, P., Abraham, C. and Conner, M. (eds), *Understanding and Changing Health Behaviour: From Health Beliefs to Self-regulation*. Harwood, Amsterdam, pp. 73–98.
- Reed, L.D. and Christensen, D.B. (1988) A psychosocial perspective in the explanation of patients drug-taking behavior. *Social Science and Medicine*, **27**, 277–285.
- Rimer, B.K. (1997) Perspectives on intrapersonal theories of health behavior. In Glanz, K., Lewis, F.M. and Rimer, B.K. (eds), *Health Behavior and Health Education: Theory, Research and Practice*, 2nd edn. Jossey-Bass, San Francisco, CA, pp. 139–147.
- Rosen, C.S. (2000) Is the sequencing of change processes by stage consistent across health problems? A meta-analysis. *Health Psychology*, **19**, 593–604.
- Rosenstock, I.M., Strecher, V.J. and Becker, M.H. (1988) Social learning theory and the health belief model. *Health Education Quarterly*, **15**, 175–183.
- Seibold, D.R. and Roper, R.E. (1979) Psychosocial determinants of health care intentions: test of the Triandis and Fishbein models. *Communication Yearbook*, **3**, 625–643.
- Seydel, E., Taal, E. and Wiegman, O. (1990) Risk-appraisal, outcome and self-efficacy expectancies: cognitive factors in preventive behaviour related to cancer. *Psychology and Health*, **4**, 99–109.
- Sheeran, P., Abraham, C. and Orbell, S. (1999) Psychosocial correlates of heterosexual condom use: A meta-analysis. *Psychological Bulletin*, **125**, 90–132.
- Smedslund, G. (2000a) A pragmatic basis for judging models and theories in health psychology: The axiomatic method. *Journal of Health Psychology*, **5**, 133–149.
- Smedslund, G. (2000b) Psycho-Logic: A 'conceptual toolbox' and 'mind sharpener' for health psychologists: A response to commentaries. *Journal of Health Psychology*, **5**, 165–171.
- Spencer, L., Pagell, F., Hallion, M.E. and Adams, T.B. (2002) Applying the transtheoretical model to tobacco cessation and prevention: a review of the literature. *American Journal of Health Promotion*, **17**, 7–71.
- Strecher, V.J., DeVellis, B.M., Becker, M.H. and Rosenstock, I.M. (1986) The role of self-efficacy in achieving health behavior change. *Health Education Quarterly*, **13**, 73–92.
- Sutton, S. (1987) Social-psychological approaches to understanding addictive behaviours: attitude-behaviour and decision-making models. *British Journal of Addiction*, **82**, 355–370.
- Vanlandingham, M.J., Suprasert, S., Grandjean, N. and Sittitrai, W. (1995) Two views of risky sexual practices among northern Thai males: the health belief model and the theory of reasoned action. *Journal of Health and Social Behavior*, **36**, 195–212.
- Warwick, P., Terry, D. and Gallois, C. (1993) Extending the theory of reasoned action: the role of health beliefs. In Terry, D.J., Gallois, C. and McCamish, M. (eds), *The Theory of Reasoned Action: Its Application to AIDS-Preventive Behaviour*. International Series in Experimental Social Psychology. Pergamon Press, Elmsford, NY, vol. 28, pp. 117–133.
- Weinstein, N.D. (1993) Testing four competing theories of health-protective behavior. *Health Psychology*, **12**, 324–333.
- Werch, C.E., Pappas, D.M., Carlson, J.M., Edgemon, P., Sinder, J.A. and DiClemente, C.C. (2000) Evaluation of a brief alcohol prevention program for urban school youth. *American Journal of Health Behavior*, **24**, 120–131.
- Wing, R.R., Voorhees, C.C. and Hill, D.R. (eds) (2000) Maintenance of behavior change in cardiorespiratory risk reduction. *Health Psychology*, **19** (Suppl. 1).
- Wulfert, E. and Wan, C.K. (1995) Safer sex intentions and condom use viewed from a health belief, reasoned action and social cognitive perspective. *Journal of Sex Research*, **32**, 299–311.
- Wulfert, E., Wan, C.K. and Backus, C.A. (1996) Gay men's safer sex behavior: an integration of three models. *Journal of Behavioral Medicine*, **19**, 345–366.
- Zimmerman, R.S. and Vernberg, D. (1994) Models of preventive health behavior: comparison, critique and meta-analysis. In Albrecht, G. (ed.), *Advances in Medical Sociology, Health Behavior Models: A Reformulation*. JAI Press, Greenwich, CT, vol. 4, pp. 45–67.

Received on December 17, 2003; accepted on June 9, 2004