

OPEN

JAIDS Journal of Acquired Immune Deficiency Syndromes Publish Ahead of Print
DOI: 10.1097/QAI.0000000000002085

Do quality improvement initiatives improve outcomes for patients in antiretroviral programmes in low- and middle-income countries? a systematic review

S Hargreaves^{+,1}, K Rustage^{+,1}, LB Nellums^{+,1}, JE Bardfield¹, B Agins¹, P Barker¹, M R Massoud¹, N P Ford¹, M Doherty¹, G Dougherty¹, S Singh^{*,1}

⁺Joint First Authors

¹Department of HIV/AIDS (S Singh MD MSc, N Ford FRCPE, M Doherty MD PhD); International Health Unit, Section of Infectious Diseases and Immunity, Imperial College London, London, UK and The Institute for Infection and Immunity, St George's, University of London, London, UK (S Hargreaves FRCPE, K Rustage MSc, LB Nellums PhD); Healthqual, University of California San Francisco, Institute for Global Health Sciences (JE Bardfield MPH, BD Agins MD, MPH); URC, Chevy Chase, MD, USA and Institute for Healthcare Improvement, Boston USA (PM Barker MBCHB); University Research Co., LLC/ USAID ASSIST Project (M. R. Massoud MD MPH FACP); ICAP/Columbia University (Gillian Dougherty) BSN MPH

*Corresponding author: Dr Sally Hargreaves E) s.hargreaves@sgul.ac.uk

Conflicts of interest: Nothing to declare

This is an open access article distributed under the terms of the Creative Commons Attribution [License 4.0 \(CC BY\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: There have been a range of quality improvement (QI) and quality assurance (QA) initiatives in low and middle-income countries (LMICs) to improve antiretroviral therapy (ART) treatment outcomes for people living with HIV (PLHIV). To date, these initiatives have not been systematically assessed and little is known about how effective, cost-effective, or sustainable these strategies are in improving clinical outcomes.

Methods: We conducted a systematic review adhering to PRISMA guidelines (PROSPERO ID: CRD42017071848), searching Pubmed, MEDLINE, Embase, Web of Science, and the Cochrane database of controlled trials for papers reporting on the effectiveness of QI and QA initiatives in HIV programmes in LMICs in relation to ART uptake, care retention, adherence, viral load suppression, mortality, and other outcomes including cost-effectiveness and long-term sustainability.

Results: 1860 articles were found, of which 29 were included. QI approaches were categorised as: i) health systems approaches using QI methods; ii) QI learning networks including collaboratives; iii) standards-based methods that use QI tools to improve performance gaps; and iv) campaigns using QI methods. The greatest improvements were seen in ART uptake (median increase of 14.0% [IQR -9.0, 29.3]), adherence (median increase of 22.0% [IQR -7.0, 25.0]), and viral load suppression (median increase 26.0% [IQR -8.0, 26.0]).

Conclusions: QI interventions can be effective in improving clinical outcomes; however, there was significant variability, making it challenging to identify which aspects of interventions lead to clinical improvements. Standardising reporting and assessment of QI initiatives is needed, supported by national quality policies and directorates, and robust research.

Keywords: HIV, AIDS, quality improvement, quality assurance, ART, LMICs

Introduction

Considerable strides have been made in the scale-up of antiretroviral therapy (ART) in low- and middle-income countries (LMICs). By the end of 2017, 21.7 million people living with HIV (PLHIV) were receiving ART, with a reported fall of 48% in AIDS-related deaths since a peak in 2005.^{1,2} However, only 75% of the estimated 36.7 million PLHIV globally know their status, 59% were receiving ART, and 47% were virologically suppressed. In LMICs with the highest burden of HIV, coordinated action is urgently needed to achieve targets so 90% of all people living with HIV know their status, 90% of those diagnosed as HIV positive start ART, and 90% of all people receiving ART have durable viral suppression.³

There is increasing recognition of gaps along the cascade of care, and the need to strengthen the quality of service delivery.⁴⁵ Various quality improvement (QI) and quality assurance (QA) strategies have been implemented in ART programmes.⁶⁷ Many definitions of QI exist in the literature. The recently published WHO National Quality and Policy Strategy Manual (<http://apps.who.int/iris/bitstream/handle/10665/272357/9789241565561-eng.pdf?ua=1>) defines QI as “a change in process in a health-care system, service, or supplier for the purposes of increasing the likelihood of optimal clinical quality of care measured by positive health outcomes for individuals and populations.”⁸ For this review, we consulted with the Quality of Clinical HIV Care Technical Working Group (see Acknowledgments) to generate a narrower definition (Figure 1): “a method of improving programme quality using standard QI methodologies involving systems analysis, process investigation and analysis of results/indicators, developing solutions by teams, testing and measuring effects of changes, and implementing and following up improvement.” QA is defined as: “a process of external

measurement of performance against standards and expectation that action will be taken to improve performance” (Figure 1).

Across the global literature on QI initiatives globally, there is a high risk of bias, with studies predominantly from high-income countries, or not specifically focused on HIV treatment programmes.⁹ Furthermore, the evidence-base is limited by a lack of systematic or robust examination of evidence on the effectiveness of initiatives to improve quality in the delivery of ART programmes. As a result, there is a lack of consensus on effective, cost-effective, or sustainable approaches.

We undertook a systematic review to identify and synthesise evidence-base on the cost-effectiveness, acceptability, impact of QI and QA initiatives on key clinical outcomes in ART programmes in LMICs, and wider indicators.

Methods

This systematic review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines,¹⁰ and the protocol published on the PROSPERO database (CRD42017071848; https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=71848).

Inclusion and exclusion criteria

We included randomised controlled trials, observational studies, and grey literature reporting on the effectiveness of QI and QA initiatives implemented in ART programmes in low- and middle-income countries (LMICs) as defined by World Bank classification.¹¹

We considered evidence of effectiveness for pre-defined outcomes including:

- Key clinical outcomes: ART uptake, care retention, adherence, viral load suppression and mortality
- Other outcomes: screening, identification, and treatment of incident opportunistic infections, process indicators such as CD4/VL testing, acceptability to patients and/or service providers, prevention of mother to child HIV transmission
- Cost-effectiveness
- Wider impact (e.g. long-term sustainability)

There were no restrictions on language. Research was excluded if it was conducted in high-income countries, data for key outcomes were not present, or the main focus of the intervention was task-shifting, which is not considered a QI or QA intervention for the purposes of this review. In addition, papers only reporting on specific types of clinical care/interventions (e.g. prevention of mother to child HIV transmission) were not included. These studies represent a substantial and important literature, and merit separate reviews tailored to the specific methodologies and outcomes relevant to these distinct areas. Training, clinical mentoring, and supportive supervision, without the precise improvement actions defined as part of QI, were not included, as the focus was on quality improvement methods rather than the specific components of interventions; in addition, we note that supportive supervision and mentorship has its own literature, as part of broader quality assurance efforts. These decisions were made in consultation with the Quality of Clinical HIV Care Technical Working Group.

Search strategy

We searched the databases Pubmed, MEDLINE, Embase, Web of Science, and the Cochrane database of controlled trials from inception to 24/10/17. We used a Boolean search strategy with keywords relevant to quality improvement, quality assurance, and HIV, identified from

relevant research, previous related systematic reviews, and consultation with the World Health Organization and Quality of Clinical HIV Care Technical Working Group (see Supplementary information for full search strategy). Grey literature was obtained through a hand-search of web search engines, related websites, and submission by key experts in the Quality of Clinical HIV Care Technical Working Group. These experts were also formally invited to submit grey literature (conference abstracts, unpublished reports, presentations etc) relating to QI in ART programmes in LMICs. The bibliographies of included papers were cross-referenced, and key experts consulted, to identify additional research.

Data extraction and analysis

Title and abstract screening was carried out by three reviewers (KR, LBN, SH). Subsequent full-text screening was carried out by two reviewers (SH, KR). Discrepancies were resolved through discussion with a third reviewer. Screening was facilitated by the web-based application Rayyan.¹²

Data were extracted by two reviewers (KR, SH) using a piloted form on pre-defined outcomes determined by the study team through consultation with experts in the field. The outcomes included: ART uptake, retention in care, adherence, viral load suppression, mortality, and other wider outcomes (opportunistic infections, acceptability to patients and service providers, process indicators, cost-effectiveness, and long-term sustainability).

Quality and risk of bias were assessed by two reviewers (KB, LN) using a piloted critical appraisal tool which included indicators from the Joanna Briggs and Newcastle-Ottawa scales as relevant for the diverse study types.^{13,14} The average quality score was 69.1%. Quality scores were divided into tertiles, categorising papers in relation to whether they were scored in the low (22.7% – 59.1%), medium (63.6% – 72.7%), or high (77.3% - 100%) third of papers. Whilst study quality was assessed in order to indicate methodological rigour and

clarity and transparency in reporting, studies were not excluded on the basis of quality in order to strengthen the transparency of the review and comprehensively report on all available evidence, including both primary peer-reviewed research and the grey literature.

All included studies were categorised in relation to methodological approaches agreed in collaboration with the identified panel of experts (Figure 1): health systems approaches using quality improvement methods, standards-based methods that use QI tools to improve performance gaps, campaigns that use QI methods, and QI learning networks including collaboratives.

Summary analyses were carried out in Microsoft Excel and Stata 15 to show the distribution of percent increase reported for each outcome by study, and to calculate the mean and median increase in the percentage of patients with each clinical outcome to provide an indication of the reported impact of QI interventions across the available evidence base. We also compared the median increase in ART uptake by methodological approach and country setting, and in programmes focused on the prevention of mother to child transmission (PMTCT) compared to the general patient population. Where relevant data were not reported for the pre-defined outcomes, papers were not included in the syntheses.

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, writing of the report, or the decision to submit the paper for publication. All authors had full access to all data, and responsibility for the decision to submit for publication.

Results

Overview of included literature

1860 records were identified in the database searches, with 1073 publications being subject to title and abstract screening after removing duplicates. 101 publications were carried forward for full-text screening, in addition to 34 grey literature records. Of the 135 papers included in the full-text screening, 29 were included in the review (see Figure 2), including 14 were peer-reviewed^{15,16,25–28,17–24} and 15 grey literature articles (Supplementary Table 1).^{29,30,39–43,31–38}

There was significant variation in the quality of included studies (reported in Supplementary Table 1), with limited detail in relation to measures, outcomes or definitions used, observational designs across a majority of studies, and a lack of data to isolate the effects of interventions.

Many initiatives involved a multifaceted package of interventions. No studies were identified that reported solely on QA, therefore our results present no data on QA initiatives. There are a range of quality initiatives being evaluated in LMICs (Standards-based Management and Recognition, HIVQUAL/HEALTHQUAL [Note HIVQUAL changed its name to HEALTHQUAL], Breakthrough Series Collaborative, ASSIST), as well as multiple other approaches (quality collaboratives, Performance management-quality improvement, mentorship and training without specific QI component, amongst others).

We grouped approaches into four main categories with support from experts in the Quality of Clinical HIV Care Technical Working Group (Figure 1): health systems approaches using quality improvement methods (n = 20),^{21,22,34–43,24–26,29–33} Standards-based methods that use QI tools to improve performance gaps (n = 2),^{17,23} Campaigns that use QI methods (n = 1),²⁸ and QI Learning Networks including collaboratives (n = 6).^{15,16,18–20,27} Initiatives were carried out in 13 LMICs, including South Africa,^{15,19,20,28} Vietnam,^{16,35} Zambia,^{17,23} Nigeria,^{18,27}

Uganda,^{21,22,31,36,40} Mozambique,^{21,22} Namibia,^{21,22,29,34,42,43} Haiti,^{21,22,24,32,36,41} Thailand,^{25,26} Nicaragua,³³ Kenya,³⁸ Tanzania,³⁷ and Guyana,^{34,39} (Figure 3). Across the studies, improvements were reported for ART uptake (n=17), ART adherence (n=10), CD4 testing (n=6), retention (n=5), and cost (n=1) (Figure 4).

Overall impact of QI approaches

There was significant variation across the evidence regarding the influence of QI initiatives on clinical outcomes (Figure 5). The greatest improvement was seen in ART uptake (median increase of 14.0% (IQR -9.0, 29.3) of patients across sites), ART adherence (median increase of 22.0% (IQR -7.0, 25.0)), and viral load suppression (median increase 26.0% (IQR -8.0, 26.0)) (see Supplementary Table 1).

These outcomes should be considered in the context of other initiatives and national policy and programme changes that may have targeted ART uptake and adherence, and viral load suppression at the same time. Improvements may also be partly attributed to inclusion of programmes focused on the prevention of mother-to-child transmission (PMTCT; Supplementary Table 1). We carried out a sensitivity analysis to examine the increase in ART uptake in pregnant women compared to the general patient population across the evidence, identifying that the median percentage increase in patients initiating ART in programmes focused on PMTCT was 19.0% (IQR 13.5, 40.5) compared 13.0% (IQR 4.5, 29.3) for programmes directed at the general population (Table 1), though this was not statistically significant.

There was also significant variation in the effectiveness of QI interventions across initiatives implemented in different country settings. The highest median percentage improvement in ART uptake was seen in Vietnam (29.0%, IQR 29.0, 29.5), followed by Haiti (25.0%, IQR 13.0, 37.0), however there were limited data for other countries, and variability in the length

of follow up, making it challenging to make meaningful cross-country comparisons (Table 2). For example, three studies looking to improve ART uptake had follow-up periods of 8-,³⁵ 12-,³¹ and 24-months.³²

Effectiveness by QI methodological approach

(i) Health systems approaches using QI methods

20 studies (5 published; 15 grey literature) were categorised as health systems approaches using QI methods (e.g. including systems analysis, process investigation and analysis of results/indicators, developing solutions by teams, testing and measuring effects of changes, and implementing and following up improvement) (Figure 1). 14 studies reported outcomes related to ART uptake, generally showing a positive impact. In one study, the number of eligible children taking up treatment rose from 12 to 25%,³² whilst ART uptake rose from 61 to 90% in another health systems intervention.³⁵ Even in a previously well performing setting, ART uptake rose from 98 to 100% after the introduction of the intervention.²⁵ However, there were also two examples of a decline in ART uptake. In one such case, uptake fell from 82 to 76% over a 12-month period,³¹ whilst in another, CD4 testing for subsequent ART treatment in individuals with CD4 cell counts < 200/ μ l dropped from 100 to 90% of eligible individuals.²⁶ Overall, the median percentage increase in patients initiated on ART in programmes utilising this methodological approach was 13.0% (IQR 5.5, 34.3), with a range of -13.0% (decrease) to 59.0% (Table 1).

Among papers reporting health system approaches using QI methods, there were also nine studies reporting ART adherence (2 published; 7 grey literature), and five studies reporting on retention. Six of the nine studies reporting adherence outcomes showed improved adherence documentation and recording procedures, as opposed to being related directly to the level of adherence seen in individuals on treatment. One study reported that paediatric

adherence rose from 43 to 81%,³⁶ whilst a multi-centre study documented improvements in adherence from 90 to 97% in Namibia, 63 to 85% in Uganda, 66 to 97% in Mozambique, 29 to 83% in Haiti, and 56 to 80% in Guyana.³⁹ One study reported a decrease in adherence from 82 to 76%; however, this finding may be due to an associated improvement in documentation procedures and adherence assessment, which increased from 81 to 92%.³¹ Retention was also suggested to slightly increase across the studies (median increase 3.0%, IQR 3.0, 3.0; range 3.0 – 25.0).

Health systems approaches utilising QI methods were also associated with positive impacts on process indicators. CD4 testing rates over 6 months improved from 43 to 78% in one setting,³⁵ and from 10.8 to 20.5% when testing individuals upon enrolment and after 6 months of follow up.⁴¹ However, there were not always substantial gains or improvements.⁴⁴ Prophylaxis for opportunistic infections also increased. One study reported an increase from 12 to 95% of children receiving cotrimoxazole prophylaxis,⁴³ whilst another reported that prophylaxis peaked during the intervention at 84.8% of eligible individuals receiving treatment.⁴¹ Tuberculosis screening also increased across all reports, with one example citing an increase from 24 to 99%.²⁶

(ii) *QI learning networks across multiple sites (including collaboratives)*

Six peer-reviewed papers reported on QI learning networks (including collaboratives), which included coaching and mentoring of healthcare staff, alongside peer exchange to address performance gaps. One model adopted was the Breakthrough Series (BTS) collaborative,¹⁵ which brings facility teams together to learn QI methods, identify performance gaps, and plan implementation interventions, with follow up visits from quality mentors to coach teams on using QI methods and maintain momentum for improvement. Four studies reported on ART uptake outcomes, all of which showed improvements. Overall, the median increase in ART

uptake was 22.0% (IQR 12.8, 29.8) (Table 1). One study reported a district level increased uptake of 62 to 91% over 30 days.¹⁶ In another setting, monthly ART initiation rose 185.5% following initiation of the intervention, from 179 initiations per month to 511.¹⁹ ART adherence and retention were only reported in one study each. The study which included adherence outcomes reported improvements in documentation of adherence support and adherence assessment procedures, and it was not clear if there was an impact on patient adherence. In this study, documentation improved from 83 to 99% at the provincial level, and from 54 to 97% at the district level.¹⁶ In terms of ART retention, the one reporting study found no impact associated with the intervention, with no statistically significant difference in retention of post-partum women after 6-months in the intervention and control arms.¹⁸

The effects of QI interventions on process indicators, opportunistic infection and TB screening were sparsely reported. The studies reporting on process indicators indicated an improvement associated with the intervention. CD4 testing in the previous 6-months increased from 80 to 94% at the provincial level, and from 72 to 74% at the district level in one setting,¹⁶ and rates of early infant testing at 4-6 weeks increased in intervention sites from 25.3% to 48.8% in another study.¹⁸ Data quality was reported to have improved in another setting.²⁷ Cotrimoxazole prophylaxis prescription to combat opportunistic infections in eligible individuals increased in one study from 31 to 99% at the provincial level, and from 75 to 83% at the district level, whilst TB screening in this same study also increased from 15 to 100% at the provincial level and from 18 to 79% at the district level.¹⁶

One study further looked at the acceptability of the intervention to healthcare workers and patients. However, despite enthusiastic participation in the programme by healthcare workers, there was no measurable increase in patient satisfaction.²⁷

(iii) *Standards-based methods that use QI tools to improve performance gaps*

The two citations which utilised standard-based methods only reported on acceptability of the intervention (2 published, 0 grey literature).^{17,23} The studies initially sought to define performance criteria relevant to the context, before utilising outside support to measure pre-existing performance gaps (in relation to the new standards), which are subsequently addressed and re-assessed. In both studies, the authors indicated that healthcare worker perceptions of their work environment were positively impacted by the intervention, whilst they declined in comparison sites. Standards-based methods also increased ART readiness scores, and provider performance related to ART and PMTCT at the intervention sites versus comparison sites.¹⁷

(iv) *Campaigns that use QI methods*

Only one study reported a QI campaign among health workers (Figure 1) in two districts in KwaZulu-Natal, South Africa, between 2009 and 2010, and was included as a separate category upon consultation with the expert group.²⁸ This study reported on outcomes related to ART uptake among HIV-infected pregnant women and cost, finding that monthly referrals for ART rose from 798.7 (95% CI: 69-89) individuals to 188(95% CI: 167-209) in the intervention district, and monthly ART initiation concurrently rose from 21 (95% CI: 2-40) individuals to 124 (95% CI: 108-140), compared with much smaller increases in the control district (91 to 99 ART referrals and 39 to 55 initiations per month). In terms of cost, there was no formal analysis, however the authors state that they believed the interventions could be rapidly implemented with a low incremental cost, as no new personnel were added to the existing health system.

Discussion

This is the first systematic review to explore the evidence-base on the effectiveness of interventions to improve quality in ART programmes in LMICs, and to define and categorise methodological approaches being implemented in these settings. The review suggests that QI initiatives at site level, applied at multi-site learning collaboratives, and as part of a campaign, were associated with increased ART uptake and ART adherence across sites. However, there was variability in the effect across studies and approaches. Whilst the findings point to the potential effectiveness of QI interventions to improve quality, we should interpret the results with caution given the limitations of the study designs; i.e. lack of comparators or the contribution of other initiatives. This points to the need for more rigorous evaluation methods to assess the impact of quality improvement interventions, and innovative approaches to assessing quality in public health initiatives more broadly. It is also important to be aware of a bias towards publication of positive results.

The effect of other programme factors is noted in some cases, both in rapid improvement of clinical outcomes, and in an apparent worsening of outcomes in some cases, which could influence the interpretation of QI studies. For example, in Uganda, ART uptake decreased from 82 to 76% over 12 months,³¹ and in Thailand, CD4 testing coverage dropped from 100% to 90%.²⁶ As both of these studies were supported by authors of this manuscript, further information was obtained from the programme reports, and it is reported that in subsequent years, ART coverage increased, while the number of patients enrolled in care increased dramatically over that time period. As well, in Thailand, the drop in CD4 monitoring may have been related to a rapid expansion in the program from 12 hospitals in 2002 to 64 in 2006, thus highlighting the impact of other programme factors. The heterogeneity across QI initiatives makes it challenging to discern the relative benefit of specific QI approaches. Both increases and decreases in the research need to be

contextualised in relation to factors such as methodological variations, socio-economic conditions, changes in health expenditure, and other possible factors such as stockouts, and documentation challenges.

Our findings concur with a recent systematic review exploring one QI approach (collaboratives) mostly being evaluated in high-income countries, which reported that the evidence for effectiveness of initiatives that use QI collaboratives is positive, but that the effects cannot be predicted with great certainty with limited evidence of effectiveness, cost-effectiveness, and sustainability.⁷

Many QI initiatives involve a complex multifaceted package of interventions. For example, HEALTHQUAL is a public health model for capacity building and can involve multiple interventions as part of the effort to improve health systems. Studies may also report outcomes on initiatives involving QI and other capacity-building initiatives, and also adapt established QI models to local use.¹⁶ This means that it is often not straightforward to identify which particular aspect of a QI initiative, has led to improvements. A strength of QI initiatives is that they target multiple aspects of the health system which support clinical processes. However, these changes are often introduced simultaneously making it challenging to identify which factors contribute to improvements. However, it is hypothesised that this more 'holistic' approach is required for successful programmes, suggesting that QI initiatives and methods should be judged on the overall effectiveness of the programme, rather than attempting to elucidate which specific components are most integral to the benefits observed.⁴⁵

Strengths and limitations

The review utilised a rigorous approach, aligning with PRISMA guidelines, and was registered prospectively in PROSPERO. The results are driven by a comprehensive search strategy, which includes five databases and grey literature. The assessment of key clinical outcomes alongside other wider effects allowed a broad examination of evidence of effectiveness, enabling a robust synthesis to evaluate and inform QI initiatives. A formal meta-analysis was not deemed appropriate given the heterogeneity in QI approaches, study methods, and reporting. However, studies consistently reported percent increases in the pre-defined outcomes, allowing such data to be combined and summarised to provide an indication of the reported impact of QI interventions across the available evidence base.

Quality scores for included papers ranged from 22.7% to 100%, with a third of papers scoring below 60%, which were categorised as low quality. The significant variation in the quality of study designs as well as reported outcomes and their definitions highlights the need to improve methodological rigour in evaluation and research in this area. The evidence base would be strengthened by efforts to adopt emerging QI evaluation designs.⁴⁶ The decision to retain papers and not exclude on the basis of quality was essential in order to demonstrate the range of quality across the available evidence base, and the need to strengthen study quality, benefitting the transparency and impact of this review.

There was limited reporting of QI initiatives in both the published and grey literature, with few peer-reviewed studies of the effectiveness of QI/QA initiatives, or publicly available evaluations. For many organisations, QI is part of standard technical assistance for countries, and reports of QI work may be disseminated only internally within the support organisations or funding bodies.

Another key limitation is that improvements may not be solely attributed to the QI interventions, but also wider initiatives targeting HIV programmes or healthcare systems, or improvements in documentation and data recording procedures. For example, changes in national policies to shift CD4 eligibility criteria or HIV testing campaign or expansion of sites/task shifting/decentralisation effort would improve certain outcomes unrelated to the specific QI efforts. Future research should aim to address these concerns, and where interventions are implemented, these wider contextual factors should be acknowledged through the adoption of implementation science frameworks.

The WHO recently published recommendations for Standard Programme Reporting standards,⁴⁷ which could be enhanced with practitioners agreeing on a common set of clinical and health system outcomes. It has also been proposed that the design and evaluation of such programmes should be mutually informing for both improvers and evaluators. The Framework for Learning about Improvement and the Evaluation Continuum, for example, provides a structure to improve the design, implementation, and evaluation of improvement programmes. This could enable both the generation of more robust and generalisable data, and improve understandings of the mechanisms contributing to improvement, and how to scale-up improvement efforts.⁴⁸

Evaluation of programmatic quality could be further enhanced through improved reporting mechanisms to assess implementation, which could be based on commonly agreed core components for QI programmes.⁴⁹ Standardised quality scores based on national guidelines for care recommendations have been developed in similar contexts assessing adherence to such guidelines, and have been shown to be associated with decreased mortality in these settings.⁵⁰ The implementation of such procedures across all settings would enable a fuller comparison of QI programmes, alongside their respective standards of quality both pre- and post- intervention.⁴⁹

The ability to assess the impact of QI interventions on clinical outcomes, or determine which specific approaches result in the greatest improvements is also limited by observational study designs. However, a key strength of the approaches across the included studies is the development and implementation of interventions in the community context at which they are aimed, which is likely to increase the success of QI initiatives.⁴⁵

Most of the included studies had externally funded implementing partners providing staff to support data collection, implementation, and evaluation. This reliance on external technical support raises a question about the sustainability and scalability of these interventions. Many interventions included participation from management and frontline teams of the local health system, but did not demonstrate the ability to undertake QI efforts using local data systems, or sustainability or scalability beyond the life of the project.

The use of external or parallel data systems also circumvented the poor quality of routinely collected data – a major obstacle to the implementation, scale up, sustainability, and evaluation of QI interventions. Many of the studies reported in the review used data that was collected independently of the local routine data reporting systems, and the future performance of those initiatives may be dependent on continued support. There are some examples of large scale QI programming that used routine data systems, showing that is possible to improve local data systems to the point where they can be used for QI approaches. While one of the aims of this review was to assess sustainability, the identified literature did not specifically address this question so no data are presented.

Conclusions and further research

Our findings support the effectiveness of QI in ART programmes, with the greatest improvement in clinical outcomes seen in ART uptake, adherence, and viral load suppression, and in programmes focused on PMTCT. Though the evidence suggests QI initiatives are associated with improved clinical outcomes, there was significant heterogeneity across approaches, settings, and reporting, making it challenging to identify best practices and to understand what specific aspects of these interventions lead to significant and sustainable clinical improvements in an LMIC context. Furthermore, there was very limited evidence on the cost-effectiveness of these interventions, and a need for formal economic analyses to determine the cost implications of QI initiatives.

The findings point to the need to better utilise standard evaluation designs and reporting methods.⁵¹⁴⁷ Whilst programmatic quality reporting focusing on the implementation of interventions against established guidelines would enhance our ability to compare the effectiveness of interventions across diverse settings, it could also be tailored to the specific context in which programmes are being delivered, for example primary healthcare sites as opposed to specialty care.^{5,49,50,52}

A further key consideration for practice is the need to embed QI initiatives within national efforts to improve health systems. Most of the studies in this review were supported or conducted by implementing partners with external funding, using external data collection and analysis systems. The question remains what approaches will be needed to support the capacity of health systems in LMICs to undertake these approaches through their existing quality management infrastructure without a need for external partners and funding. In a related field, the WHO-led Network for Improving Quality of Care for Maternal Newborn and Child Health Care is driving the approach of moving care away from NGO-led,

fragmented and often unsustainable individual quality improvement projects toward country-led initiatives.⁵³ WHO is also providing guidance to governments on how to design and organise national QI programming through development of their National Quality Policy and Strategies, an approach which supports the use of QI in ART programmes to reach 90-90-90 targets within the context of Universal Health Coverage^{54,55} across HIV programmes globally.⁵³

Research is needed to evaluate the effectiveness, cost-effectiveness, and long-term sustainability of quality improvement interventions, and identify which elements contribute to improved clinical outcomes in LMICs. This will support efforts to achieve the 90-90-90 goals,⁵⁶ and aligns with the recent *Lancet Global Health* Commission on high-quality health in the SDG Era.⁵⁷ Ultimately, evidence-based QI methods must be integrated into broader efforts to ensure the delivery of high-quality care in low resource settings, within the context of the *WHO Framework on integrated people-centred health services*.^{58,59}

Acknowledgements

We acknowledge the support of all members of the Quality of Clinical HIV Care Technical Working Group. SH, LBN, and KR acknowledge funding from WHO, the Wellcome Trust, the UK National Institute for Health Research Imperial Biomedical Research Centre, the European Society for Clinical Microbiology and Infectious Diseases (ESCMID) through an ESCMID Study Group (ESGITM) research grant/

References

- 1 UNAIDS. Global AIDS Update 2016. 2016.
http://www.unaids.org/sites/default/files/media_asset/global-AIDS-update-2016_en.pdf (accessed Feb 21, 2018).
- 2 UNAIDS. Fact sheet - Latest global and regional statistics on the status of the AIDS epidemic. 2018.
http://www.unaids.org/sites/default/files/media_asset/UNAIDS_FactSheet_en.pdf (accessed Feb 21, 2018).
- 3 UNAIDS. 90-90-90: An ambitious treatment target to help end the AIDS epidemic. Geneva, Switzerland, 2014 http://www.unaids.org/sites/default/files/media_asset/90-90-90_en.pdf (accessed Feb 21, 2018).
- 4 UNAIDS. Ending AIDS: Progress Towards the 90-90-90 Targets. Geneva, Switzerland, 2017
http://www.unaids.org/sites/default/files/media_asset/Global_AIDS_update_2017_en.pdf (accessed March 26, 2018).
- 5 The Lancet Global Health. Are we ready for a quality revolution? *Lancet Glob Heal* 2018; **6**: e121.
- 6 Joint United Nations Programme on HIV/AIDS (UNAIDS). On the Fast-Track to end AIDS: UNAIDS 2016–2021 Strategy. Geneva, Switzerland, 2016
http://www.unaids.org/sites/default/files/media_asset/20151027_UNAIDS_PCB37_15_18_EN_rev1.pdf (accessed Jan 17, 2018).
- 7 Schouten LMT, Hulscher MEJL, van Everdingen JJE, Huijsman R, Grol RPTM. Evidence for the impact of quality improvement collaboratives: systematic review.

- BMJ* 2008; **336**: 1491–4.
- 8 World Health Organization. A Handbook for National Quality Policy and Strategy: A practical approach for developing policy and strategy to improve quality of care. 2018.
- 9 Buckley GJ, Pittluck RE. Improving Quality of Care in Low- and Middle-Income Countries: Workshop Summary. 2015 DOI:10.17226/21736.
- 10 Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group TP. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- 11 The World Bank. World Bank Country and Lending Groups – World Bank Data Help Desk. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> (accessed Feb 21, 2018).
- 12 Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 2016; **5**: 210.
- 13 Wells G, Shea B, O’Connell D. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2009.
- 14 Joanna Briggs Institute. The Joanna Briggs Institute Critical Appraisal tools for use in JBI Systematic Reviews - Checklist for Case Series. 2016. <http://joannabriggs.org/research/critical---appraisal---tools.html> (accessed Aug 8, 2017).
- 15 Barker P, Barron P, Bhardwaj S, Pillay Y. The role of quality improvement in achieving effective large-scale Prevention of Mother-To-Child Transmission of HIV in South Africa. *Aids* 2015; **29**: S137–43.
- 16 Cosimi LA, Dam H V., Nguyen TQ, *et al.* Integrated clinical and quality improvement

- coaching in Son La Province, Vietnam: a model of building public sector capacity for sustainable HIV care delivery. *BMC Health Serv Res* 2015; **15**: 269.
- 17 Kols A, Kim YM, Bazant E, Necochea E, Banda J, Stender S. A standards-based approach to quality improvement for HIV services at Zambia Defence Force facilities: Results and lessons learned. *Aids* 2015; **29**: S145–53.
- 18 Oyeleledun B, Oronsaye F, Oyelade T, *et al.* Increasing retention in care of HIV-positive women in PMTCT services through continuous quality improvement-breakthrough (CQI-BTS) series in primary and secondary health care facilities in Nigeria: A cluster randomized controlled trial. The Lafiyan Jikin. *J Acquir Immune Defic Syndr* 2014; **67**: S125–31.
- 19 Webster P, Sibanyoni M, Malekutu D, Mate K, Venter W, Barker P. Using quality improvement to accelerate highly active antiretroviral treatment coverage in South Africa. *BMJ Qual Saf* 2012; **21**: 315–24.
- 20 Youngleson MS, Nkurunziza P, Jennings K, Arendse J, Mate KS, Barker P. Improving a mother to child HIV transmission programme through health system redesign: Quality improvement, protocol adjustment and resource addition. *PLoS One* 2010; **5**. DOI:10.1371/journal.pone.0013891.
- 21 Bardfield K, Agins B, Palumbo M, Wei A, Morris J. Improving rates of cotrimoxazole prophylaxis in resource-limited settings: Implementation of a quality improvement approach. *Int J Qual Heal Care* 2014; **26**: 613–22.
- 22 Massoud MR, Shakir F, Livesley N, *et al.* Improving care for patients on antiretroviral therapy through a gap analysis framework. *Aids* 2015; **29**: S187–94.
- 23 Bazant E, Sarkar S, Banda J, *et al.* Effects of a performance and quality improvement

- intervention on the work environment in HIV-related care: A quasi-experimental evaluation in Zambia. *Hum Resour Health* 2014; **12**: 1–11.
- 24 Joseph JP, Jerome G, Lambert W, Almazor P, Cupidon CE, Hirschhorn LR. Going beyond the vertical: Leveraging a national HIV quality improvement programme to address other health priorities in Haiti. *Aids* 2015; **29**: S165–73.
- 25 Lolekha R, Chunwimaleung S, Hansudewechakul R, *et al.* Pediatric HIVQUAL-T: Measuring and improving the quality of pediatric HIV care in Thailand, 2005-2007. *Jt Comm J Qual Patient Saf* 2010; **36**: 541–51.
- 26 Thanprasertsuk S, Supawitkul S, Lolekha R, *et al.* HIVQUAL-T: Monitoring and improving HIV clinical care in Thailand, 2002-08. *Int J Qual Heal Care* 2012; **24**: 338–47.
- 27 Osibo B, Oronsaye F, Alo O, *et al.* Using small tests of change to improve PMTCT services in Northern Nigeria: Experiences from implementation of a continuous quality improvement and breakthrough series program. *J Acquir Immune Defic Syndr* 2017; **75**: S165–72.
- 28 Ngidi W, Reddy J, Luvuno Z, Rollins N, Barker P, Mate KS. Using a campaign approach among health workers to increase access to antiretroviral therapy for pregnant HIV-infected women in South Africa. *J Acquir Immune Defic Syndr* 2013; **63**: 133–9.
- 29 Bardfield J, Agins B, Akiyama M, *et al.* A quality improvement approach to capacity building in low- and middle-income countries. *Aids*. 2015; **29**: S179–86.
- 30 Bardfield J, Agins B, Palumbo M, Wei A, Celestin N, Lamothe N. A Quality Improvement Approach To Scale-up of ART in Resource Limited Settings. ; : 1339.

- 31 Behumbiize P, Ssendiwala J, Kayita G, *et al.* Organizational Structure and Capacity for Evaluating the Quality of Care among HIV Clinics in Northern Uganda. 2006.
- 32 Bijou S, Labbe C, Nancy R, *et al.* Use of an electronic medical record to implement, monitor, and improve HEALTHQUAL clinical care indicator performance rates. 2005.
- 33 Broughton E, Nunez D, Moreno I. Cost-Effectiveness of Improving Health Care to People with HIV in Nicaragua. *Nurs Res Pract* 2014; **2014**: 1–6.
- 34 HealthQual International. Performance Measurement Data Report 2015. San Francisco, CA, US, 2015.
- 35 HealthQual International. HIVQUAL Vietnam: National and local leadership support for improvement in HIV care & treatment. 2012.
- 36 Kayita G, Ssendiwala J, Lamothe N, Palumbo M, Birchard R, Agins B. Building a sustainable national model to improve pediatric and maternal health care in Haiti and Uganda. .
- 37 Kimaro J, Kihwele D, Stover K, Shrestha R, Rumisha D, Fatta K. Empowering community groups to support access and retention in HIV care in Muheza , Tanzania. 2015.
- 38 Mohamed I, Wanyungu J, Abass M, Amagove M, Ngugi E, Oggot C. Improving quality of care for people living with HIV and AIDS in Kenya: A case of Coast province. 2010; : 54.
- 39 Palumbo M, Birchard R, Geis M, *et al.* A public health approach to building sustainable national quality management programs in low- and middle-income countries. ; : 8599.
- 40 Ssendiwala J, Kayita G, Palumbo M, Musin- J. National Scale up of Quality

- Improvement Activities : A Case Study of Uganda. .
- 41 Thimothe G, Duval N, Lauture D, *et al.* Application of a National Electronic Health Records System for Measuring Performance for Improvement In Systems of Care for Persons Living with Hiv Measurement of Effect Changes. .
- 42 USAID Applying Science to Strengthen and Improve Systems Project. USAID ASSIST Project Experience Improving HIV Services. Bethesda, MD, 2014
https://www.usaidassist.org/sites/assist/files/usaid_assist_experience_improving_hiv_services_ada_june2014_0.pdf.
- 43 USAID. Lessons Learned from Applying Collaborative Improvement Methodologies to Strengthen the Performance and Productivity of HIV Human Resources - Technical Report. 2016.
- 44 Lolekhha R, Chunwimaleung S, Hansudewechakul R, *et al.* Pediatric HIVQUAL-T: measuring and improving the quality of pediatric HIV care in Thailand, 2005-2007. *Jt Comm J Qual Patient Saf* 2010; **36**: 541–51.
- 45 Batalden PB, Davidoff F. What is quality improvement; and how can it transform healthcare? *Qual Saf Health Care* 2007; **16**: 2–3.
- 46 Massoud MR, Kimble LE, Goldmann D, Ovretveit J, Dixon N, Salzburg Global Seminar Session 565—‘Better Health Care: how do we learn about improvement?’ *Int J Qual Heal Care* 2018; **30**: 1–4.
- 47 Kågesten AE, Tunçalp Ö, Portela A, Ali M, Tran N, Gülmezoglu AM. Programme Reporting Standards (PRS) for improving the reporting of sexual, reproductive, maternal, newborn, child and adolescent health programmes. *BMC Med Res Methodol* 2017; **17**: 117.

- 48 Barry, D, Kimble, L, Nambiar, B, Parry, G, Jha, A, Chattu, VK, Massoud, MR, Goldmann D. A framework for learning about improvement: embedded implementation and evaluation design to optimize learning. *Int J Qual Heal Care* 2018; **30**: 10–4.
- 49 Bardfield J, Palumbo M, Geis M, Jasmin M, Agins BD, NOA Working Group T. A National Organizational Assessment (NOA) to Build Sustainable Quality Management Programs in Low- and Middle-Income Countries. *Jt Comm J Qual Patient Saf* 2016; **42**: 325–30.
- 50 Opondo C, Allen E, Todd J, English M. Association of the Paediatric Admission Quality of Care score with mortality in Kenyan hospitals: a validation study. *Lancet Glob Heal* 2018; **6**: e203–10.
- 51 SQUIRE. SQUIRE | SQUIRE 2.0 Guidelines. 2017. <http://www.squire-statement.org/index.cfm?fuseaction=page.viewpage&pageid=471> (accessed March 27, 2018).
- 52 Johnston S, Kendall C, Hogel M, McLaren M, Liddy C. Measures of Quality of Care for People with HIV: A Scoping Review of Performance Indicators for Primary Care. *PLoS One* 2015; **10**: e0136757.
- 53 Adeniran A, Likaka A, Costello A, *et al.* Leadership, action, learning and accountability to deliver quality care for women, newborns and children. *Bull World Heal Organ* 2018; **96**: 222–4.
- 54 World Health Organization. Chapter 11 - Quality Improvement (QI): Introduction to Quality Improvement. http://www.who.int/hiv/pub/imai/om_11_quality_improvement.pdf (accessed Feb 21, 2018).

- 55 World Health Organization. Quality of care - A process for making strategic choices in health systems. 2006.
http://www.who.int/management/quality/assurance/QualityCare_B.Def.pdf (accessed Feb 21, 2018).
- 56 Barker PM, Reid A, Schall MW. A framework for scaling up health interventions: lessons from large-scale improvement initiatives in Africa. *Implement Sci* 2015; **11**: 12.
- 57 Kruk ME, Pate M, Mullan Z. Introducing The Lancet Global Health Commission on High-Quality Health Systems in the SDG Era. *Lancet Glob Heal* 2017; **5**: e480–1.
- 58 Nambiar B, Hargreaves DS, Morroni C, *et al.* Improving health-care quality in resource-poor settings. *Bull World Health Organ* 2017; **95**: 76–8.
- 59 World Health Organization. WHO | WHO Framework on integrated people-centred health services. WHO. 2018. <http://www.who.int/servicedeliverysafety/areas/people-centred-care/en/> (accessed March 7, 2018).

Figure captions

Figure 1. Definitions and categorisation of QI/QA interventions used for this systematic review.

Figure 2: PRISMA flow diagram

Figure 3: Proportion of QI initiatives reported in the included literature.

Figure 4. Number of included studies reporting improvements in key outcomes.

Figure 5. Median percentage increase in outcomes across QI/QA initiatives

Table 1. Changes in ART uptake by population and QI/QA approach

	Population		QI approach	
	Maternity	General	Health system approach using QI methods	QI learning network
Mean % (SD)	28.0 (17.6)	19.0 (20.0)	20.0 (21.3)	23.0 (12.0)
Median % (range)	19.0 (12.0, 56.0)	13.0 (-13.0, 59.0)	13.0 (-13.0, 59.0)	22.0 (12.0, 41.0)
Interquartile range (IQR) 25%, 75%	13.5, 40.5	4.5, 29.3	5.5, 34.3	12.8, 29.8

Table 2. Changes in ART uptake by country of study

	South Africa	Vietnam	Haiti	Thailand	Namibia	Uganda	Mozambique	Guyana
Mean % (SD)	23.0 (14.1)	29.0 (0.6)	26.0 (14.5)	2.0	22.0 (23.4)	18.0 (28.4)	-13.0	5.0
Median % (range)	14.0 (12.0, 41.0)	29.0 (29.0 - 30.0)	25.0 (13.0, 40.0)	2.0	15.0 (3.0, 56.0)	9.5 (-6.0, 59.0)	-13.0	5.0
Interquartile range (IQR) 25%, 75%	12.0, 21.5	29.0, 29.5	13.0, 37.0	--	9.0, 28.3	3.75, 23.8	--	--

Figure 1. Definitions and categorisation of QI/QA interventions used for this systematic review.

Definitions:

A **quality improvement intervention**: a method of improving programme quality using standard QI methodologies involving systems analysis, process investigation and analysis of results/indicators, developing solutions by teams, testing and measuring effects of changes, and implementing and following up improvement

A **quality assurance intervention**: a process of external measurement of performance against standards and expectation that action will be taken to improve performance

Methodological categorisations of reported QI initiatives:

1. Health systems approaches that use QI methods: A method of improving programme quality involving process investigation and analysis of results/indicators, developing solutions by teams, testing and measuring effects of changes, and implementing and following up improvement.

2. Standards-based methods that use QA and QI tools to improve performance gaps: A model which combines assessment of performance through external measurement of a core set of standards (QA), together with a QI process to implement of best practices, follow up measurement of progress and rewarding achievement through social/peer recognition mechanisms.

3. QI learning networks: A structured approach that uses QI methods to promote improvement activities and successful improvement strategies across multiple networked facilities in a health system. Network structure offers face to face opportunities and a virtual platform to build improvement skills, share improvement and technical knowledge, monitor performance, and spread improvement principles.

4. QI campaigns: An intensive improvement approach to improve care in a particular clinical area and designed to influence change in a large geopolitical area (region or an entire country). Campaigns are designed to mobilise large groups of providers – hundreds or thousands – to focus on the selected area of care that has been identified as an urgent priority. Senior officials and opinion leaders visibly champion the cause of the campaign and facilities and communities are engaged through public programmes and the media. QI methods are used to adapt well developed implementation strategies to local context.

Figure 2. PRISMA flow diagram





Figure 3. Proportion of QI initiatives reported in the included literature.



ACCEPTED

Figure 4. Proportion of included studies reporting improvements in key outcomes




Figure 5. Percentage increase in outcomes across QI/QA initiatives

