

Probing Hairpin Structures of Small DNAs by Nondenaturing Polyacrylamide Gel Electrophoresis

Ulrich Baumann,^{*,1} Ronald Frank,[†] and Helmut Blöcker[†]

^{*}Max-Planck-Institut für Biophysikalische Chemie, D-3400 Göttingen, Federal Republic of Germany, and [†]GBF, DNA-Synthesegruppe, Braunschweig, Federal Republic of Germany

Received March 7, 1989

The influence of temperature on the electrophoretic mobility of small DNAs, capable of forming hairpin structures, is investigated under nondenaturing conditions. Three series of hairpin-forming DNAs containing different numbers of thymidine, deoxyadenosine, and deoxyguanosine residues in their loop, and an identical sequence in the helical region, are analyzed. All show enhanced electrophoretic mobility if they adopt the hairpin conformation. The same quantitative relationship between hairpin formation and increase in electrophoretic mobility is observed for all of the three series. The constancy of this increase suggests a dependence of electrophoretic acceleration on the length of the helical region. A possible application of nondenaturing electrophoresis is monitoring the hairpin/coil transition. Another possible application is the detection of dimers formed by partially self-complementary sequences. This dimer formation is detected for completely complementary DNAs, whereas sequences which might form imperfect double helices, especially those with three bulged-out nucleotides, prefer hairpin formation. The possible applications are experimentally approached and discussed. © 1989 Academic Press, Inc.

The secondary and tertiary structures of nucleic acids influence their electrophoretic mobility in polyacrylamide and agarose gels. In the high molecular weight range the most significant effect is the enhanced mobility of double-stranded, compared to single-stranded, DNA revealed by application of a gradient of a denaturing agent (1). Supercoiled plasmids move faster than the relaxed form; the number of supercoils is directly related to the electrophoretic behavior of a plasmid (2,3). Bended double-stranded DNA shows reduced mobility (4), as well as DNA containing cruciforms (5) in compar-

ison to linear structures. Single-stranded DNAs, generated from blunt end duplexes, migrate at different rates due to variations in secondary structure of both strands (6,7). Furthermore, the two-dimensional separation of tRNAs is based on their different tertiary structures (8).

Hairpins formed by concatemers in the range of 8–80 nucleotides exhibit enhanced mobility, although denatured by urea (9). In this study rather small oligodeoxynucleotides, capable of forming single hairpin structures, are investigated by nondenaturing polyacrylamide gel electrophoresis. The temperature dependence of electrophoretic mobility reveals enhanced mobility of the hairpin form. As the temperature range of transition from hairpin to random coil of the DNAs investigated is known from independent studies (10,11), the electrophoretic mobility can be unambiguously correlated to their secondary structure. Thus, the method promises to be useful for monitoring hairpin/random coil transition and might supplement NMR and temperature dependence of ultraviolet absorption. Its advantages include the application of more physiological concentrations and reduced consumption of materials.

MATERIALS AND METHODS

Materials

The hairpin-forming DNAs 5'-d(CTATCCTAA₆TAGGATAG) as well as hp(dG)₃₋₅ and hp(dA)_{3,4,6}² were synthesized by phosphotriester chemistry on cellulose disks acting as segmental support (12). The hairpins hp(dT)₃₋₆ and hp(dA)₅ were synthesized according to Ref. (13). Oligo(dA)₁₂₋₁₈, oligo(dT)₁₂₋₁₈ (both in a 5' phosphorylated and dephosphorylated form), oligo(dC)₁₂₋₁₈, oligo(dA)₁₉₋₃₀, and oligo(dAG)₅, all bearing a 5' phosphate group, were from Pharmacia (Freiburg, FRG).

² Abbreviations used: hp(dG)₃₋₅, hp(dA)₃₋₆, hp(dT)₃₋₆ = 5'-d(ATCCTAN₃₋₆TAGGAT), N being either dG, dA, or dT; hp = hairpin.

¹ To whom correspondence should be addressed.

a b c d e f g h i b






TABLE 1

Hairpin-forming DNA	Mobility x (cm)		Acceleration (%) $x_{\text{exp.}}/x_{\text{calc.}} \cdot 100 - 100$	Average acceleration (%)
	Calc.	Exp.		
Hairpin (dT) ₃	7.00	8.10	15.7	17.4
Hairpin (dT) ₄	6.77	7.95	17.4	
Hairpin (dT) ₅	6.51	7.70	18.8	
Hairpin (dT) ₆	6.37	7.45	16.9	
Hairpin (dA) ₃	8.76	10.10	15.0	15.3
Hairpin (dA) ₄	8.47	9.70	14.5	
Hairpin (dA) ₅	8.25	9.65	17.0	
Hairpin (dA) ₆	8.06	9.25	14.8	
Hairpin (dG) ₃	7.38	8.40	13.8	15.2
Hairpin (dG) ₄	7.01	8.05	14.8	
Hairpin (dG) ₅	6.67	7.80	16.9	

Note. The increased electrophoretic mobility of the hairpin form is expressed in relation to the mobility of random coil which is set to 100%. The calculation of the theoretical mobility (calc.) assuming a single-stranded structure was performed as described in the text, these values are compared with the experimental values (exp.) revealed from electrophoresis at 10°C.

The basis of the calculation are the experimental values of homooligonucleotides $x(T_n)$, $x(A_n)$, etc. These values are derived directly from gels run at different temperatures. As oligo(dG)_n forms clusters consisting of several molecules (17), the mobility value of oligo(dG)_n was indirectly determined from that of p(dGA)₅. For this purpose, first the value of (dA)₅ was determined from that of oligo(dA)₁₂₋₁₈ by averaging the ratio $r_A = x(A_{n-1})/x(A_n)$ for each gel. This ratio is rather constant. By calculating $r_G = \sqrt[5]{x(A_5)/x(AG_5)}$ the ratio $x(G_{n-1})/x(G_n)$ was determined. By means of this ratio, first the mobility of (dG)₁ and then that of (dG)_n were determined. This determination is less accurate than the directly determined values for other homooligonucleotides. But one must consider that all of the hairpin series, except hp(dG)₃₋₅, contain only two deoxyguanosine residues. An experimental error of 10% for the mobility of oligo(dG)_n would result only in an error of 1.5% for mobility of a DNA hairpin. As all hairpins bear no 5' phosphate group whereas oligo(dC)_n and p(dAG)₅ contain it, the effect of one additional phosphate group was evalu-

FIG. 2. Electrophoretic behavior of hp(dT)₃₋₆ below and above the hairpin/random coil transition. In the respective lane 0.05 A₂₆₀ unit of the hairpin-forming DNA is applied and the electrophoresis is performed together with single-stranded marker oligonucleotides. Three different conditions, (A) nondenaturing at 60°C, (B) nondenaturing at 10°C, and (C) denaturing at ca. 50°C. Lane a (dT)₁₂₋₁₈, lane b (dA)₁₂₋₁₈, lane c p(dA)₁₂₋₁₈, lane d p(dAG)₅, lane e hp(dT)₃, lane f hp(dT)₄, lane g hp(dT)₅, lane h hp(dT)₆, and lane i p(dC)₁₂₋₁₈. The two outer lanes in (C) are changed, compared to A and B, as indicated.




FIG. 3. Comparison of $hp(dA)_{3-6}$ and $hp(dG)_{3-5}$ with the electrophoretic mobility of $hp(dT)_{3-6}$. (A) Electrophoresis at 60°C and (B) at 10°C , both under nondenaturing conditions. Lane a $(dT)_{12-18}$, lane b $(dA)_{12-18}$, lane c $p(dGA)_5$, lane d $p(dC)_{12-18}$, and different hairpin-forming DNAs as indicated; d for deoxy is omitted to simplify the figure.

ated by comparing the 5' phosphorylated and dephosphorylated form of $(dA)_{12-18}$ and $(dT)_{12-18}$. The effect was small (see Fig. 2), and the 5' phosphorylated molecules show the same mobility as a dephosphorylated oligonucleotide shortened by half a nucleotide, this was considered in the determination of $x(C_n)$ and $x(G_n)$.

Theoretical values of mobility calculated from a gel run at 60°C are in good agreement with experimental values indicating that all DNAs are present as random coil at this temperature. The mobilities of each of the hairpin forms are given in Table 1. The values are expressed as excess percentage assuming 100% for the random coil. These values lie between 13.8 and 17.4%. They are significantly greater than the deviation of experimental and theoretical values at 60°C which are in the range $\pm 2\%$. The apparent size of a DNA hairpin is three to four nucleotides smaller than the true size. The average values of acceleration are 17.2% for dT-containing series, 15.2% for dG series, and 15.3% for dA series. The last two series are shown in Fig. 3 in comparison to thymidine containing DNA hairpins. The constancy of relative enhancement indicates, at least for the dG- and dA-containing series, that the effect caused by hairpin

formation is not dependent on the sequence of the loop but is only dependent on the length of the helical region. If this is true a means to determine the size of the double-helical region of a hairpin structure would be established.

Application of the Enhanced Mobility of DNA Hairpins

First, one can determine if a small DNA might adopt a hairpin conformation. If an indication for hairpin formation is given, one can determine the thermal dependence of the transition and the stability of a hairpin structure. In Fig. 4 the electrophoretic pattern at 40°C for $hp(dT)_{3-6}$ in comparison with $hp(dA)_{3-6}$ is shown. This temperature is within phase transition of all hairpin-forming DNAs. The oligonucleotide $hp(dT)_3$ exhibits an acceleration of 7.4%, $hp(dT)_4$ of 7.8%, the five-membered loop 3.8%, and the six membered loop 0%. The values for full hairpin form listed in Table 1 are around 17%. A hairpin electrophoresed at its T_m would hence give a mobility increase of about 8%. It is concluded that bigger loops $hp(dT)_5$ and $hp(dT)_6$ have a lower T_m as 40°C , whereas T_m of small loops is very near




FIG. 4. Electrophoretic pattern of the DNA hairpins within helix/coil transition at $T = 40^\circ\text{C}$. (A) The series $\text{hp}(\text{dT})_{3-6}$ and (B) the series $\text{hp}(\text{dA})_{3-6}$ compared with $\text{hp}(\text{dT})_{3-6}$. Lane a $(\text{dA})_{12-18}$, lane b $\text{hp}(\text{dT})_3$, lane c $\text{hp}(\text{dT})_4$, lane d $\text{hp}(\text{dT})_5$, lane e $\text{hp}(\text{dT})_6$, lane g $\text{hp}(\text{dA})_6$, lane h $\text{hp}(\text{dA})_5$, lane i $\text{hp}(\text{dA})_4$, and lane k $\text{hp}(\text{dA})_3$. The label A_n indicates the mixture of the four hairpins $\text{hp}(\text{dA})_{3-6}$. (C) An enlarged section from a separation of $\text{hp}(\text{dT})_{3-6}$ (indicated by T_n) is shown in comparison with $\text{hp}(\text{dA})_{3-6}$ and single-stranded $(\text{dA})_{12-18}$. The lanes for $(\text{dA})_{12-18}$ and T_n are from the same gel; lane A_n is from a different gel. Note the different band distances revealed by the two hairpin series and the enhanced distances between hairpins from the same series differing in one monomer.

to this temperature. This observation supports results from NMR (10) that the most stable hairpin DNA from the series containing four to seven thymidines in the loop is that with a four-membered loop and that stability decreases with increase in loop size to seven nucleotides. As the behavior of $\text{hp}(\text{dT})_3$ is close to the full hairpin

form, it is concluded that this hairpin structure is at least as stable as $\text{hp}(\text{dT})_4$. The reported T_m values are 44°C for $\text{hp}(\text{dT})_3$ (11) and $\text{hp}(\text{dT})_4$, 40°C for $\text{hp}(\text{dT})_5$, and 36°C for $\text{hp}(\text{dT})_6$ (10). The values are determined at 15 mM Na^+ and are thus slightly higher than our estimation, because no salt is present during electrophoresis. The series containing deoxyadenosine residues in the loop shows that the hairpin structure of $\text{hp}(\text{dA})_3$ is considerably more stable than $\text{hp}(\text{dA})_4$, because both DNAs are separated by a greater distance as expected for one additional nucleotide (see Fig. 4C). Thus, the method can also be used to improve separation of hairpin-forming DNAs with slightly different stability if electrophoresis is performed near the average T_m . The separation of single-stranded and hairpin-forming DNAs may be improved by variation of the electrophoresis temperature.

This simple method clearly shows that oligonucleotides with a three-membered loop do not exist as a dimer with three bulged-out nucleotides in the concentration range applied. To prove this fact, $\text{hp}(\text{dT})_3$ and $\text{hp}(\text{dA})_3$, which are fully complementary, were annealed at 0°C and run in parallel with $\text{hp}(\text{dA})_3$ in the presence of 10




FIG. 5. Dimer formation of completely complementary $\text{hp}(\text{dT})_3$ and $\text{hp}(\text{dA})_3$ at 0°C . The hairpin-forming DNAs $\text{hp}(\text{dT})_3$ and $\text{hp}(\text{dA})_3$, $0.1 A_{260}$ unit of each, were heated for 5 min to 65°C , then annealed in a buffer consisting of 10 mM Tris/HCl , pH 7.2, 10 mM NaCl , and 10 mM MgCl_2 for 5 min at 0°C , total volume $4\ \mu\text{l}$ (lane d), and electrophoresed at 0°C (A) and 60°C (B) in a gel containing 10 mM magnesium acetate. The hairpin-forming DNA $\text{hp}(\text{dA})_3$ was used as a control demonstrating no dimer formation under these conditions. Lane a $(\text{dT})_{12-18}$, lane b $(\text{dA})_{12-18}$, lane c $\text{p}(\text{dAG})_5$, lane d mixture of $\text{hp}(\text{dA})_3$ and $\text{hp}(\text{dT})_3$, and lane e $\text{hp}(\text{dA})_3$.

mM Mg^{2+} (see Fig. 5). Under these conditions a dimer of $hp(dA)_3$ and $hp(dT)_3$ is formed and is stable during electrophoresis, whereas $hp(dA)_3$ shows the molecular weight of the monomer. This indicates that in the concentration range considered, 0.12–0.3 mM, hairpin-forming DNAs with a three-membered loop do not exist as an imperfect duplex even in the presence of 10 mM Mg^{2+} . The partial dimer formation detected by NMR (11) might be possible in the NMR concentration range being 10- to 30-fold higher.

All the hairpin series tested have an identical helix length and their relative increase of mobility is quantitatively similar. Therefore, it was presumed that the degree of acceleration is only dependent on the length of the helical region. To test this assumption, the electrophoretic mobility of a hairpin-forming structure with six adenosine residues in the loop, the same loop as in $hp(dA)_6$, and a prolonged helix consisting of 8 instead of 6 bp was analyzed. The sequence was 5'-d(CTATCCTAA₆TAGGATAG). The electrophoretic pattern, shown in Fig. 6, exhibits as expected an acceleration effect of the hairpin form. Interestingly, it shows two bands at this temperature and only one main band under denaturing conditions. Quantitative evaluation gives an increase in mobility of 12.9% for the intense, upper band and 16% for the less intense, faster moving band.

DISCUSSION

With regard to the unusual electrophoretic mobility of DNA structures revealed by previous studies two groups are distinguishable. Single-stranded DNA is assumed to behave normally. Structures causing electrophoretic acceleration are closed circular (18) and supercoiled DNA, present in plasmids (2), as well as double strands in the high molecular weight range (1). Slower moving structures are bended DNA, those DNAs and RNAs bearing single- and double-stranded parts in the same molecule as well as stable cruciforms (4,5). It is concluded that all structural elements leading to closer packing, but not rendering the overall rod-like shape of molecules, increase the electrophoretic mobility. Elements leading to complex branching, in contrast, result in decrease of DNA mobility. Therefore, it is not surprising that the formation of a single hairpin of the investigated oligonucleotides cause electrophoretic acceleration. The change of shape is similar to that of plasmids during supercoiling. The degree of acceleration is mainly dependent on the size of the helical region, because all hairpins reveal similar values independent of their loop size and nucleotide content. This dependence on the helix length, however, seems not to be very sensitive. The extension of the helix from 6 to 8 bp does not significantly increase the accelerating effect. The effect of helix extension is only examined for one hairpin, further investigations are necessary. The emergence of two bands for this oligonucleo-




FIG. 6. Increase of the mobility of a hairpin-forming DNA with prolonged helix. The mobility of the hairpin 5'-d(CTATCCTAA₆TAGGATGA) (lane f) was determined at 60°C (A) and at 5°C (B) under standard conditions. The emergence of two bands at 5°C might be due to the formation of two different hairpins as shown below the gel patterns; indication d for deoxy and of the 5' end are omitted to clarify the picture. Lane a (dT)₁₂₋₁₈, lane b (dA)₁₂₋₁₈, lane c p(dAG)₅, lane d $hp(dA)_3$, lane e $hp(dA)_5$, and lane g p(dA)₁₉₋₃₀.

tide, on the other hand, indicates that this bigger DNA might form two different hairpin structures with different lengths of the helical region which are possibly formed and also shown in Fig. 6.

The T_m values for dG and dA series were not determined independently and might significantly differ from those of dT series. We assumed a similar stability for the three series. Two facts support the assumption: First, the agreement between theoretical and experimental mobilities at high temperature and second, the constancy of the accelerating effect at low temperature. Thus, the method presented may supplement established techniques as NMR and temperature dependence of ultraviolet absorption. This method requires much less material and also requires very simple equipment. It is certainly limited to small oligonucleotides, because additional structural elements are easily introduced with increase of size; these would prevent unambiguous interpretation. In addition, the amount of information regarding distinct base pairs and stacking of loop nucleotides which have been evaluated by NMR (19) or accessibility to a single-strand-specific nuclease (20,21) is more limited when this method is applied. The ques-

tion if oligonucleotides can possibly form very small loops consisting only of two or three nucleotides, on the other hand, can be answered more easily. The fully complementary oligonucleotides hp(dA)₃ and hp(dT)₃ form a dimer, easily detected by its molecular weight, whereas partially self-complementary hp(dA)₃ prefers hairpin formation. A similar observation, although regarding a two-membered loop of a completely self-complementary sequence, has been reported by Xodo *et al.* (22). NMR measurements, on the other hand, cannot distinguish both possible structures in such a simple manner.

ACKNOWLEDGMENTS

We thank Drs. D. Möbius and L. De Maeyer (Göttingen, FRG), for support of the work. For critical comments on the manuscript we thank D. Mellerick (Göttingen, FRG). Part of the work was funded by Bundesministerium für Forschung und Technologie.

REFERENCES

1. Fischer, S. G., and Lerman, L. S. (1980) *Proc. Natl Acad. Sci. USA* **77**, 4420-4424.
2. Keller, W. (1975) *Proc. Natl Acad. Sci. USA* **72**, 4876-4880.
3. Gellert, M., Mizuuchi, K., O'Dea, M. H., and Nash, H. A. (1976) *Proc. Natl Acad. Sci. USA* **73**, 3872-3876.
4. Diekmann, S., and Wang, J. C. (1985) *J. Mol. Biol.* **186**, 1-11.
5. Gough, G. W., and Lilley, D. M. J. (1985) *Nature (London)* **313**, 154-156.
6. Hayward, G. S. (1972) *Virology* **49**, 342-344.
7. Maxam, A. M., and Gilbert, W. (1977) *Proc. Natl Acad. Sci. USA* **74**, 560-564.
8. Fradin, A., Gruhl, H., and Feldmann, H. (1975) *FEBS Lett.* **50**, 185-189.
9. Frank, R., Müller, D., and Wolff, C. (1981) *Nucleic Acids Res.* **9**, 4967-4979.
10. Haasnot, C. A. G., de Bruin, S. H., Berendsen, R. G., Janssen, H. G. J. M., Binnendijk, T. J. J., Hilbers, C. W., van der Marel, G. A., and van Boom, J. H. (1983) *J. Biomol. Struct. Dyn.* **1**, 115-129.
11. Haasnot, C. A. G., de Bruin, S. H., Hilbers, C. W., van der Marel, G. A., and van Boom, J. H. (1985) *J. Biosci.* **8** (Suppl.), 767-780.
12. Frank, R., Heikens, W., Heisterberg-Moutsis, G., and Blöcker, H. (1983) *Nucleic Acids Res.* **11**, 4365-4377.
13. van Boom, J. H., van der Marel, G. A., Westerink, H., van Boeckel, C. A. A., Mellema, J.-R., Altona, C., Hilbers, C. W., Haasnot, C. A. G., de Bruin, S. H., and Berendsen, R. G. (1982) *Cold Spring Harbor Symp. Quant. Biol.* **47**, 403-409.
14. Donis-Keller, H., Maxam, A. M., and Gilbert, W. (1977) *Nucleic Acids Res.* **4**, 2527-2538.
15. Nadeau, J. G., and Gilham, P. T. (1985) *Nucleic Acids Res.* **13**, 8259-8274.
16. Frank, R., and Köster, H. (1979) *Nucleic Acids Res.* **6**, 2069-2086.
17. Lipsett, M. N. (1964) *J. Biol. Chem.* **239**, 1250-1255.
18. Johnson, P. H., and Grossman, L. I. (1977) *Biochemistry* **16**, 4217-4225.
19. van de Ven, F. J. M., and Hilbers, C. W. (1988) *Eur. J. Biochem.* **178**, 1-38.
20. Baumann, U., Frank, R., and Blöcker, H. (1986) *Eur. J. Biochem.* **161**, 409-413.
21. Baumann, U., Lehmann, U., Schweltnus, K., van Boom, J. H., and Kuhn, H. (1987) *Eur. J. Biochem.* **170**, 267-272.
22. Xodo, L. E., Manzini, G., Quadrifoglio, F., van der Marel, G. A., and van Boom, J. H. (1988) *Biochemistry* **27**, 6321-6326.