



UNIVERSITY OF LEEDS

This is a repository copy of *Changes in implicit wanting and explicit liking and wanting for food after gastric bypass surgery*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/147102/>

Version: Published Version

Proceedings Paper:

Redpath, T, Price, RK, Finlayson, G orcid.org/0000-0002-5620-2256 et al. (5 more authors) (2018) Changes in implicit wanting and explicit liking and wanting for food after gastric bypass surgery. In: Proceedings of the Nutrition Society. Irish Section Meeting, 20-22 Jun 2018 Cambridge University Press , E96.

<https://doi.org/10.1017/S0029665118001003>

© The Authors 2018. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Irish Section Meeting, 20–22 June 2018, Targeted approaches to tackling current nutritional issues

Changes in implicit wanting and explicit liking and wanting for food after gastric bypass surgery

T. Redpath¹, R.K. Price¹, G. Finlayson², A. Boyd¹, F. Naseer¹, D. Liddy¹, C.W. Le Roux³ and M.B.E Livingstone¹

¹Nutrition Innovation Centre for Food and Health (NICHE), Ulster University, Coleraine, BT52 1SA, ²Institute of Psychological Sciences, University of Leeds, West Yorkshire, LS2 9JT and ³Diabetes Complications Research Centre, University College Dublin, Ireland.

Gastric bypass surgery (GB) is a safe, effective treatment for morbid obesity⁽¹⁾. Aside from reduced energy consumption, patients report changes in food preferences and appetite that may contribute to weight loss post-surgery^(2,3). Food preferences can be separated into two constructs; 'liking' (pleasure) and 'wanting' (motivation/desire to eat)⁽⁴⁾ which may be influenced by physiological and psychological changes post-surgery. The aim of this study was to investigate changes in food preferences ('liking' and 'wanting') in GB patients from pre- to post-surgery, compared to weight-stable controls.

Nineteen patients (45.2 ± 12.5 years, Body Mass Index (BMI): 45.3 ± 6.4 kg/m²) and sixteen time-matched controls (44.7 ± 15.6 years, BMI: 25 ± 4.4 kg/m²) completed the computer-based Leeds Food Preference Questionnaire, which assesses preferences for sweet/savoury foods and low-fat/high-fat foods⁽⁵⁾, at baseline (1-month pre-surgery) and 3-months post-surgery. Explicit liking and wanting (conscious hedonic feelings/desire to consume) were measured using Visual Analogue Scales, whilst implicit wanting (unconscious motivational expression) was measured using food selection and reaction times in a forced-choice task.

There were no significant differences in food preferences between patients and controls at baseline. BMI for controls remained stable (+0.1 ± 1.3 kg/m², p = 0.73), whilst BMI in patients significantly decreased (−8.3 ± 2.7 kg/m², p = <0.01). GB patients expressed a significant decrease in the explicit liking (F(1,33) = 10.42, p = 0.03), explicit wanting (F(1,33) = 5.76, p = 0.02) and implicit wanting (F(1,33) = 4.92, p = 0.03) for sweet foods after surgery. However, there were no significant changes in preferences for high-fat foods in patients from pre to post-surgery. There were no significant changes in preferences for controls for sweet or high-fat foods.

In conclusion, patients express a significant decrease in preferences for sweet foods post-surgery. As changes in implicit unconscious motivational expression of food preferences may be more predictive of actual food consumption⁽⁶⁾ these findings highlight a possible mechanism responsible for the success of GB as an obesity treatment.

1. Colquitt JL, Pickett K *et al.* (2014) *Cochrane Database Syst Rev* **8**, CD003641.
2. Mathes CM & Spector AC (2012) *Physiol Behav* **107**, 476–483.
3. Miras AD & Le Roux CW (2013) *Nat Rev Gastroenterol Hepatol* **10**, 575–584.
4. Berridge KC & Robinson TE (2003) *Trends Neurosci* **26**, 507–513.
5. Finlayson G, King N & Blundell J (2008) *Appetite* **50**, 120–127.
6. Pool E, Sennwald V *et al.* (2016) *Neurosci Biobehav Rev* **63**, 124–142.