



Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Phytochemistry 66 (2005) 1536–1539

PHYTOCHEMISTRY

www.elsevier.com/locate/phytochem

Chickpea wilt: identification and toxicity of 8-*O*-methyl-fusarubin from *Fusarium acutatum*

Subramaniam Gopalakrishnan ^a, Michael H. Beale ^b,
Jane L. Ward ^b, Richard N. Strange ^{a,*}

^a Department of Biology, Darwin Building, University College London, Gower Street, London WC1E 6BT, UK

^b National Centre for Plant and Microbial Metabolomics, Rothamsted Research, West Common, Harpenden, Herts AL5 2JQ, UK

Received 13 April 2005; accepted 25 April 2005

Available online 15 June 2005

Abstract

Fusarium acutatum was isolated from wilting chickpea plants in Pakistan. Filtrates from cultures grown on a defined liquid medium caused permanent wilting of chickpea cuttings and killed cells, isolated enzymically from healthy plants, in a bioassay. Toxic activity was retained by a cyano solid phase extraction cartridge and the toxin was isolated by elution from the cartridge in acetonitrile and Si-gel thin layer chromatography of the eluate. Analytical HPLC of the compound on a cyano column with diode array detection gave a single peak with a homogeneous spectrum and λ_{\max} at 224 and 281 nm. NMR and mass spectral studies showed that the toxin was 8-*O*-methyl-fusarubin. The pure compound caused permanent wilting of chickpea cuttings and the LD50 value in the cell bioassay was 327 ng/ml.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: *Cicer arietinum*; Leguminosae; Chickpea; *Fusarium acutatum*; Wilt; Bioassay; NMR; Mass spectrum; Naphthoquinone; Phytotoxin

1. Introduction

Wilt is regarded as the most important disease of chickpea on the Indian subcontinent and the causal organism of the disease is usually ascribed to *Fusarium oxysporum* f. sp. *ciceri* but other species and formae speciales of *Fusarium* also cause wilting of a variety of plants (Jiménez-Díaz et al., 1993; Srivastava et al., 1984; Di Pietro et al., 2003). The fungus that is the subject of the present investigation and which was isolated from wilted chickpea plants grown in the Thal area of Pakistan, did not accord morphologically with *F. oxy-*

sporum f. sp. *ciceri* but was identified as *Fusarium acutatum*. This organism was originally reported as a pathogen of pigeon pea by O'Donnell and Cigelnik (1997) and described as a new species by Nirenberg and O'Donnell (1998). More recently, Kurmut et al. (2000) showed that it was the cause of root rot of *Vicia faba* L. in the Sudan.

Toxins of importance in plant disease syndromes are usually isolated from axenic cultures of pathogens. For example, the solanapyrone toxins which are produced by another pathogen of chickpea, *Ascochyta rabiei*, were isolated from cultures of the plant pathogen grown on a medium containing chickpea seed extract (Alam et al., 1989). Later a defined medium was formulated which allowed production of the toxins (Chen and Strange, 1991). In this investigation, essentially the same medium was used to grow cultures of *F. acutatum*. Filtrates were

* Corresponding author. Tel.: +44 020 8660 3770; fax: +44 020 7679 7096.

E-mail addresses: r.strange@ucl.ac.uk, rs003f4775@blueyonder.co.uk (R.N. Strange).

toxic to cuttings of chickpea and cells isolated from leaflets of the plant and the compound responsible was isolated and identified.

2. Results and discussion

When *F. acutatum* was grown on a defined medium consisting of Czapek-Dox nutrients and additional cations (Hamid and Strange, 2000), two types of cultures were obtained, those that were red and those that were colourless. Filtrates of the cultures were tested for their toxicity to cells isolated from chickpea leaflets using fluorescein diacetate to differentiate live and dead cells. This compound readily enters live cells with intact plasmamembranes and, once inside the cell, is metabolised by esterases to give free fluorescein. As plasmamembranes are impermeable to fluorescein, the compound accumulates and imparts a yellow-green fluorescence to such cells, which may be viewed under a fluorescence microscope. Using this assay to assess the toxicity of the two types of filtrates, those that were red were far more toxic than those that were colourless. Maximum activity of red cultures was attained after incubation of the fungus for 12 days at 20°C and exceeded 70 units of activity/ml culture filtrate where one unit of activity was defined as the amount of toxin required to kill 50% of the cells isolated from chickpea leaflets. In contrast, filtrates from colourless cultures did not exceed 10 units/ml.

The majority of the toxic activity of filtrates from red cultures of *F. acutatum* partitioned into ethyl acetate or could be extracted by a cyano-SPE cartridge. Thin-layer chromatography (TLC) of the ethyl acetate fraction, or of the acetonitrile eluate from the cyano cartridge, with CHCl₃–MeOH (8:1) gave a red spot at *R_f* 0.57. When the silica from the TLC plate was eluted with acetonitrile only the area occupied by the red spot was active in the assay. RP-HPLC of the eluate on a cyano column with 10% acetonitrile as mobile phase revealed a single major peak with absorption maxima at 224 and 281 nm.

Cuttings of chickpea plants placed in culture filtrates of the pathogen or the pure toxin wilted permanently within 4 h. Such cuttings took up the equivalent of 1.5 and 25.5 µg of the toxin, respectively. In the cell bioassay, one unit of activity was the equivalent of 327 ng/ml of purified toxin.


¹H and ¹³C NMR data were collected at 400 MHz in a CDCl₃ solution. In the ¹H-spectrum the compound displayed one aromatic singlet at δ6.73 and a highly deshielded singlet at δ13.16 which corresponded to an aromatic hydroxyl group. These signals and the presence of two methoxy singlets at δ4.03 and 4.00 suggested a 1,2,4-trisubstituted naphthoquinone arrangement. Additional to the naphthoquinone signals the ¹H-spectrum also contained a 2H double-doublet at δ4.75

which is characteristic of a CH₂ adjacent to oxygen. Two further geminally coupled (*J* = 18 Hz) 1H-doublets at δ2.53 and 2.84 indicated the presence of a further isolated methylene moiety. The spectrum also contained a 3H singlet at δ1.55 indicating a slightly deshielded aliphatic methyl group. Comparison of the observed signals and their coupling arrangements with the literature on known naphthoquinone natural products, showed that the isolated toxin corresponded to those of the previously reported 8-*O*-methyl-fusarubin (**1**) (Steyn et al., 1979; Tatum et al., 1985). Interestingly, two numbering systems have been utilised for this compound, which is also known as 9-*O*-methyl-fusarubin (Tatum et al., 1985). We have chosen to use that of Steyn et al. (1979). ¹³C NMR data was collected to confirm structure assignment. All chemical shifts were in agreement with those previously reported (Steyn et al., 1979) as can be seen in Table 1.

Electrospray mass spectroscopy (ESI-MS) was carried out by direct infusion of a methanol solution of the isolated toxin. Atmospheric Pressure Chemical Ionisation (APCI) methods, deployed in the negative ionisation mode, indicated a molecular weight of 334. This was higher than the expected molecular weight of 320 for 8-*O*-methyl-fusarubin, although an ion corresponding to 320 was also observed in the spectrum. However, Tatum et al. (1985) showed that the addition of acidic methanol to 8-*O*-methyl-fusarubin causes the formation of 3-*O*-methyl-8-*O*-methyl-fusarubin (**2**) resulting from the methylation of the hydroxy group at C-3. The molecular weight of this compound is 334 and thus the observation of this ion in the mass spectrum, as an artefact of solubilization (**1**) in acidic methanol, served as further confirmation of the structural assignment.

Table 1
¹³C NMR chemical shifts in CDCl₃ solution using TMS as internal standard

Carbon atom	Observed chemical shift	Literature chemical shift (Steyn et al., 1979)
1	58.7	57.7
3	94.4	93.1
4	31.2	31.7
5	148.5	147.6
6	155.7	155.0
7	102.5	103.0
8	156.0	155.3
9	180.0	179.0
10	189.2	188.7
11	114.3	113.7
12	109.9	108.9
13	144.9	144.6
14	136.9	136.8
15	29.3	28.3
6-OMe	56.8	56.1
8-OMe	56.4	56.4



- (1) R = H. 8-O-Methyl-fusarubin
 (2) R = Me. 3-O-Methyl-8-O-methyl-fusarubin

Naphthazarin toxins, produced by species of *Fusarium*, of which 8-*O*-methyl-fusarubin is one, have been implicated in disease symptoms in citrus (van Rensburg et al., 2001), cotton (Bell et al., 2003) and pea (Holenstein and Défago, 1983). Their activity has been attributed to the production of the superoxide radical, (O_2^-) H_2O_2 and semiquinone radicals of the compounds (Medentsev and Akimenko, 1992). Such compounds attack membranes and could be responsible for the loss of the semi-permeability of the plasmamembrane as found in the cell assay reported in this paper. An attack on plasmamembranes could also explain wilting as chickpea is likely to depend in part on the turgor of parenchyma cells surrounding the stele for support. As a parallel, it is pertinent to point out that stem breakage, a prominent symptom of Ascochyta blight of chickpea, can be reproduced by solanapyrone A, one of the toxins produced by the causal agent of the disease, *A. rabiei*. Here, too, the loss of turgor of parenchyma cells surrounding the stele was invoked as an explanation of the phenomenon (Hamid and Strange, 2000).

3. Experimental

3.1. Isolation, spore production and storage of *F. acutatum*

Fungi were isolated from wilted chickpea seedlings collected in the Thal area of the Punjab, Pakistan. Collar regions from diseased seedlings were excised, surface sterilised with 2% sodium hypochlorite for 2 min, rinsed in distilled water and placed on petri plates containing Komada's medium (KM), ostensibly specific for *F. oxysporum* (Komada, 1975). Plates were incubated at $25 \pm 2^\circ\text{C}$ in an illuminated incubator with a photoperiod of 16 h for 5–7 d. Fungi developing from the chickpea material were sub-cultured and single spored on PDA. Spores were produced in shake culture on Czapek-Dox Cation Liquid Medium (CDCLM; 30 ml in 250 ml flasks) consisting of Czapek Dox liquid medium (Oxoid) supplemented with $ZnSO_4 \cdot 7H_2O$, 0.05 g l^{-1} ; $CuCl_2 \cdot 6H_2O$, 0.02 g l^{-1} ; $MnCl_2 \cdot 4H_2O$, 0.02 g l^{-1} ; $CaCl_2 \cdot 2H_2O$, 0.1 g l^{-1} ; $CoCl_2 \cdot 6H_2O$, 0.02 g l^{-1} (Hamid and Strange, 2000). After 5 days, mycelium was removed by filtration through four layers of muslin cloth and

the spores pelleted by centrifuging at 3000g for 15 min. They were resuspended in sterile distilled water and centrifuged twice more before finally resuspending them at 10^7 spores/ml and storing them in liquid nitrogen.

3.2. Toxin production

Roux bottles (1 l) containing 100 ml CDCLM were inoculated with spore suspension of *F. acutatum* ($100 \mu\text{l}$; 10^7 spores/ml) and incubated at 20°C for 12 d. Cultures were filtered through four thicknesses of muslin and the filtrates were centrifuged at 10,000g for 30 min. MeOH was added to the supernatants to 20% (v/v) and after aging for 4–16 h the preparation was filtered again through four thicknesses of muslin. The MeOH was removed by film evaporation at $<35^\circ\text{C}$ and the aqueous remainder filtered through a 0.5 mm filter (Nalgene, Rochester, USA).

3.3. Solid phase extraction

Filtrates (100–150 ml) were introduced into a cyano SPE cartridge (5 g; Jones Chromatography, UK) after solvation and equilibration of the cartridge with 30 ml each of acetonitrile and ultra purified water. After washing with water (5 ml) and drying in a current of air the toxin was eluted in acetonitrile (10 ml).

3.4. Bioassay

Essentially the procedure of Hamid and Strange (2000) was followed.

3.4.1. Thin layer chromatography

The acetonitrile eluates were spotted on Silica gel 60 F₂₅₄ TLC sheets with a concentrating zone and developed in $CHCl_3$ –MeOH (8:1). Silica corresponding to zones containing a red compound at R_f 0.57 was scraped from the plate and eluted in acetonitrile.

3.4.2. HPLC

The acetonitrile eluates (20 μl) from TLC were injected into a cyano column (150 mm \times 4.6 mm, protected with a guard column [20 \times 4.6 mm] of the same material; Jones Chromatography, UK) and the column developed with 10% acetonitrile as mobile phase. Data were acquired on a Philips HPLC system consisting of a PU4100 quaternary pump, PU4021 diode array detector and a personal computer equipped with PU6003 diode array software for data handling.

3.4.3. NMR

NMR was carried out on a Bruker Avance 400 MHz instrument. 8-*O*-Methyl-fusarubin (1) ^1H NMR (400 MHz, $CDCl_3$, TMS as internal standard): δ 1.55 (3H,

s, 15-H₃), 2.53 (1H, *d*, *J* = 18 Hz, 4-H_{ax}), 2.84 (1H, *d*, *J* = 18 Hz, 4-H_{eq}), 4.00 (3H, *s*, 8-OMe), 4.03 (3H, *s*, 6-OMe), 4.75 (2H, *dd*, *J* = 18, 2.8 Hz, 1-H₂), 6.73 (1H, *s*, 7-H), 13.16 (1H, *s*, 5-OH); ¹³C NMR (100 MHz, CDCl₃, TMS as internal standard): δ29.33 (15-C), 31.20 (4-C), 56.40 (8-OMe), 56.82 (6-OMe), 58.66 (1-C), 94.36 (3-C), 102.49 (7-C), 109.89 (12-C), 114.28 (11-C), 136.91 (14-C), 144.94 (13-C), 148.47 (5-C), 155.70 (6-C), 155.97 (8-C), 180.01 (9-C), 189.23 (10-C).

3.4.4. ESI-MS

Direct infusion ESI-MS was carried out on a ThermoFinnigan LCQ by Atmospheric Pressure Chemical Ionisation (APCI) in negative ionisation mode.

Acknowledgements

We thank Mr. Muhammad Hanif for making the isolations of fungi and the European Union for support with the grant “Ascorab” (Contract No. ICA4-CT-2000-30003). We thank Jason Ejimadu (RRES) for mass spectrometry.

References

- Alam, S.S., Bilton, J.N., Slawin, A.M.Z., Williams, D.J., Sheppard, R.N., Strange, R.N., 1989. Chickpea blight – production of the phytotoxins solanapyrone A and C by *Ascochyta rabiei*. *Phytochemistry* 28, 2627–2630.
- Bell, A.A., Wheeler, M.H., Liu, J.G., Stipanovic, R.D., 2003. United States Department of Agriculture – Agricultural Research Service studies on polyketide toxins of *Fusarium oxysporum* f. sp. *vasinfectum*: potential targets for disease control. *Pest Management Science* 59, 736–747.
- Chen, Y.M., Strange, R.N., 1991. Synthesis of the solanapyrone phytotoxins by *Ascochyta rabiei* in response to metal cations and development of a defined medium for toxin production. *Plant Pathology* 40, 401–407.
- Di Pietro, A., Madrid, M.P., Caracuel, Z., Delgado-Jarana, J., Roncero, M.I.G., 2003. *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus. *Molecular Plant Pathology* 4, 315–325.
- Hamid, K., Strange, R.N., 2000. Phytotoxicity of solanapyrones A and B produced by the chickpea pathogen *Ascochyta rabiei* (Pass.) Labr. and the apparent metabolism of solanapyrone A by chickpea tissues. *Physiological and Molecular Plant Pathology* 56, 235–244.
- Holenstein, J., Défago, G., 1983. Inheritance of naphthazarin production and pathogenicity to pea in *Nectria haematococca*. *Journal of Experimental Botany* 34, 927–935.
- Jiménez-Díaz, R.M., Crino, P., Halila, M.H., Mosconi, C., Trapero-Casas, A.T., 1993. Screening for resistance to *Fusarium* wilt and *Ascochyta* blight in chickpea. In: Singh, S.N., Saxena, M.C. (Eds.), *Breeding for Stress Tolerance in Cool Season Food Legumes*. John Wiley and Sons, New York, pp. 77–95.
- Komada, H., 1975. Development of a selective medium for the quantitative isolation of *Fusarium oxysporum* from natural soils. *Review of Plant Protection Research* 8, 114–124.
- Kurmut, A., Nirenberg, H.I., Bochow, H., 2000. *Fusarium acutatum* Nirenberg and O'Donnell a causal agent of root rot of *Vicia faba* L. in Sudan. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft* 377, 89–90.
- Medentsev, A.G., Akimenko, V.K., 1992. Mechanism of phytotoxic action of naphthoquinone pigments of the fungus *Fusarium decemcellulare*. *Phytochemistry* 31, 77–79.
- Nirenberg, H.I., O'Donnell, K., 1998. New *Fusarium* species and combinations within the *Gibberella fujikuroi* species complex. *Mycologia* 90, 434–458.
- O'Donnell, K., Cigelnik, E., 1997. Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* 7, 103–116.
- Srivastava, S.K., Singh, S.N., Khare, M.N., 1984. Assessment of yield losses in some promising grain cultivars due to *Fusarium* wilt. *Indian Journal of Plant Protection* 12, 125–126.
- Steyn, P.S., Wessels, P.L., Marasas, W.F.O., 1979. Pigments from *Fusarium moniforme* Sheldon. *Tetrahedron* 35, 1551–1555.
- Tatum, J.H., Baker, R.A., Berry, R.E., 1985. Naphthoquinones produced by *Fusarium oxysporum* isolated from citrus. *Phytochemistry* 24, 457–459.
- van Rensburg, J.C.J., Labuschagne, N., Nemeč, S., 2001. Occurrence of *Fusarium*-produced naphthazarins in citrus trees and sensitivity of rootstocks to isomarticin in relation to citrus blight. *Plant Pathology* 50, 258–265.