

Indian J. Genet., 72(3): 376-379 (2012)

Short Communication

## Validation of foreground and background SSR markers for introgression of QTL governing leaf glossiness into a sorghum variety - NTJ 2

G. Lakshmidevi, P. Janila<sup>1\*</sup>, G. Anuradha and A. Shivashankar

Institute of Biotechnology, Acharya N G Ranga Agricultural University, Rajendranagar, Hyderabad 500 030  
<sup>1</sup>ICRISAT, Patancheru 502 324

(Received : November 2011; Revised : June 2012; Accepted: July 2012)

### Abstract

Leaf glossiness and trichomes on leaf surface, two important component traits governing resistance to shoot fly in sorghum is controlled by Quantitative trait loci (QTLs) located on linkage group SBI-05. Foreground and background markers were validated for transfer of the QTLs to NTJ2, a popular variety of sorghum. For foreground selection, five SSR markers, Xtxp268, Xisep1111, Xtxp065, Xtxp303, and Xiabt440 in the region of the QTL were identified from 17 tested markers. From a total of 167 SSR markers spanning the entire genome, 35 polymorphic SSR markers, distributed on all the 10 chromosomes were identified for background selection. The F<sub>1</sub>, BC<sub>1</sub>F<sub>1</sub> and F<sub>2</sub> generation plants were screened using foreground markers. Based on the results obtained, the hybridity of the F<sub>1</sub>'s was confirmed and marker-homozygote individuals in F<sub>2</sub> generation were identified for subsequent backcrossing.

**Key words:** Sorghum, shoot fly, QTL, leaf glossiness, foreground selection, background selection, introgression

Sorghum [*Sorghum bicolor* (L.) Moench] is the fifth most important cereal crop in the world, cultivated in 40.5 m ha with a production of 55.6 m t during 2010 [1]. In India it is cultivated in an area of 7.7 m ha. Shoot fly (*Atherigona soccata*) is one of the most important pests in India causing grain and stover losses. The average grain losses in the country are estimated to be 5% [2].

Host plant resistance can play a major role in minimizing the extent of losses and is compatible with other tactics of pest management, including the use of natural enemies and chemical control [3]. Resistance to shoot fly in sorghum is known and various component traits contributing to resistance have been identified, of which, leaf glossiness, leaf trichomes are found to be very important [4]. Several sources of resistance have been identified and used in breeding programs but the complex nature of resistance has limited the success through conventional breeding approaches [3].

The use of molecular markers in breeding programs can be promising to improve quantitative traits. Once major QTLs are detected they can be transferred into desirable parents through marker assisted backcrossing (MABC) where foreground and background markers are used to select target loci and recipient genome, respectively. We report validation of foreground and background markers for introgression of QTL associated with leaf glossiness and thus confer shoot fly resistance in sorghum into an elite parent, NTJ 2 through MABC. The target QTL is on linkage group SBI-05. The foreground validation is important as all the primer pairs in the region of QTL are not expected to detect polymorphism between two parents, thus polymorphism of markers in the target QTL region is essential to begin MABC. The background markers

\*Corresponding author's e-mail: [p.janila@cgiar.org](mailto:p.janila@cgiar.org)

Published by Indian Society of Genetics & Plant Breeding, F2, First Floor, NASC Complex, PB#11312, IARI, New Delhi 110 012  
Online management by [indianjournals.com](http://indianjournals.com)

hasten the introgression of QTL for leaf glossiness into NTJ 2; evident from the simulation studies, where it was shown that equivalent levels of the recurrent parent genome may be recovered by BC<sub>4</sub> or BC<sub>3</sub> using a marker assisted backcrossing approach [5], thus saving two or three backcross generations. Leaf glossiness is an important component trait contributing to resistance to shoot fly in sorghum and the QTLs governing the trait were reported [6].

NTJ 2 (NandyalTellaJonna 2), used as recurrent parent is a high yielding postrainy season adapted, sweet stalk variety developed at Acharya N G Ranga Agricultural University and released in 1990. It is a pureline selection from E-1966, a *zerazera* land race obtained from ICRISAT, Patancheru. It is valued for grain and fodder yield and quality and is also promoted as sweet-sorghum variety. The donor parent, J 2779-P4 was obtained from ICRISAT, Patancheru, India. This parent, bred at ICRISAT is a backcross derivative in the background of BTx623 with QTL on SBI 05 governing leaf glossiness introgressed from a shoot fly resistant germplasm accession, IS 18551. Crosses were made in 2010 rainy season to raise F<sub>1</sub> plants in 2010-11 post rainy season. The elite parent, NTJ 2 was used as female parent to tap its maternal effects of the variety, if any. The F<sub>1</sub>'s were backcrossed and selfed to obtain BC<sub>1</sub>F<sub>1</sub> and F<sub>2</sub> seeds, respectively. The BC<sub>1</sub>F<sub>1</sub> and F<sub>2</sub> plants were raised in glass house during 2011 summer season.


The DNA from parents (NTJ 2 and J 2779-P4), F<sub>1</sub>, F<sub>2</sub> and BC<sub>1</sub>F<sub>1</sub> generations plants was extracted and checked for quality. Polymerase chain reactions (PCR) were performed in 5 µl reaction volumes and PCR reactions were carried out in a GeneAmp<sup>®</sup> PCR System 9700 thermal cycler (Applied Biosystems, USA) with a Touchdown (61-51) program. Capillary electrophoresis of denatured pooled products was performed using ABI3130x/DNA genetic analyzer. Allele calling was done by GENOTYPER 3.7 v software (Applied Biosystems). The segregation of SSR markers in F<sub>2</sub> generation was tested by Chi-square test.

A set of 17 SSR markers (Xisep0338, Xisep1029, Xisep1041, Xisep1111, Xisep1133, Xisep1107, Xisep1109, Xisep1208, Xtxp14, Xtxp15, Xtxp065, Xtxp112, Xtxp268, Xtxp303, Xiabt67, Xiabt215, Xiabt440t) were identified for foreground selection based on their location close to desired QTL on SBI 05 governing leaf glossiness from the published data [6]. Five primer pairs were polymorphic between the parents,

NTJ 2 and J 2779-P4 that can be used for foreground selection of the target QTL. The differences in allele size among the parents varied between 6 and 11 bp. The five SSR markers, Xtxp268, Xisep1111, Xtxp065, Xtxp303, and Xiabt440 can be used for foreground selection of the target QTL. The F<sub>1</sub> plants were confirmed by the presence of marker-heterozygotes that represent the alleles from both the parents (Fig 1). The F<sub>1</sub> plants confirming to hybridity were backcrossed to NTJ 2 and also selfed.

For background screening, a total of 167 markers spanning the entire genome (24 on SBI-01, 23 on SBI-02, 23 on SBI-03, 15 on SBI-04, 25 on SBI-05, 12 on SBI-06, 11 on SBI-07, 9 on SBI-08, 12 on SBI-09 and 13 on SBI-10) were identified for the study from published data [6]. Of which, 35 primers (Xisep0114, Xisep0132, Xisep0209, Xisep0327, Xisep0422, Xisep0504, Xisep0506, Xisep0522, Xisep0608, Xisep0612, Xisep0716, Xisep0809, Xisep0949, Xisep1008, Xisep1031, Xisep1032, Xisep1042, Xisep1150, Xisep1231, Xisep1241, Xtxp001, Xtxp009, Xtxp025, Xtxp027, Xtxp149, Xisep1046, Xtxp278, Xtxp088, Xtxp141, Xmsbcir300, Xmsbcir238, Xmsbcir 248, Xisep 938, Xmsbcir223, and Xgap342) were found to be polymorphic between parents, which can distinguish them. The 35 primers are distributed on all the 10 linkage groups and can be used for background selection to recover the maximum genomic content of recurrent parent in early backcross generations. The linkage group, SBI-01 has highest number of polymorphic markers.

Fifteen BC<sub>1</sub>F<sub>1</sub> seed were harvested, from which none confirmed for the target QTL as the markers amplified only A type (recurrent parent type) allele. The 88 F<sub>2</sub> seedlings that were genotyped with 35 validated background markers and scoring is given as A/B/H/-/O, where A represents the homozygote for the recurrent parent allele; B represents the homozygote for the donor parent allele; H represents the heterozygote for both of the parent alleles; '-' represents the missing data point (due to PCR failure or DNA isolation failure); and 'O' represents the presence of an unexpected, off-type allele. Since SSRs are co-dominant markers, they are expected to segregate in 1:2:1 ratio in F<sub>2</sub> generation. The segregation of A type (homozygote for allele from P<sub>1</sub> parent, NTJ 2), H type (Heterozygote) and B type (homozygote for allele from P<sub>2</sub> parent, J 2779-P4) was tested for goodness of fit for 1:2:1 ratio. Three makers



**Fig. 1. Graphical representation (chromotogram) of the SSR markers, Xtxp065, Xtxp0303, Xtxp0268 and Xiabt0440 in  $F_1$  and parents analyzed through ABI Prism 3700**

showed p levels > 0.15 indicating high probability of goodness of fit to the ratio, 1:2:1, while one primer showed a p level < 0.005. Thus all the four markers showed goodness to fit to the expected ratio of 1:2:1 ratio. Among them, the markers, Xtxp268, Xtxp303, Xiabt 440 have high probability for goodness to fit (Table 1).

Of the 88  $F_2$  plants, 18 plants were positive *i.e.*, B/H type all the markers. Among them, five are 'B type' at all the four marker loci, 3 are 'H type' at all marker loci and 9 are a combination of B and H type and five plants were identified for morphological similarity with NTJ 2. Among the five individual plants identified for NTJ 2 phenotype, one plant has 'B type' at all four marker loci and that was backcrossed to NTJ 2 to obtain  $BC_1F_1$  seed. Background selection with validated markers will be done after deriving  $BC_1$ ,  $BC_2$  and advanced generations, *i.e.*, after backcrossing the selected  $F_2$  plants to NTJ 2. The selected NTJ 2 type plants in  $BC_2F_2$  are selfed to obtain  $BC_2F_3$  progenies which will be phenotyped for yield performance and shoot fly resistance, and stable lines will be advanced to multilocation testing.

**Table 1. Segregation of SSR markers in  $F_2$  generation and goodness of fit to 1:2:1 ratio**

Marker/allele	No. of $F_2$ plants			
	Observed	Expected	Chi-square value	Probability
Xtxp0268				
A	17	15		
H	25	28		
B	16	15		
Total	58	58	0.65	0.1-0.9
Xtxp065				
A	33	21		
H	28	41		
B	22	21		
Total	83	83	11.02	<0.005
Xtxp303				
A	24	16		
H	29	33		
B	12	16		
Total	65	65	5.48	0.05-0.1
Xiabt440				
A	12	13		
H	26	27		
B	15	13		
Total	53	53	0.42	0.1-0.9

**References**

1. FAOSTAT, 2011 [www.faostat.org](http://www.faostat.org) (verified on 02.08.2012).
2. **Sharma H. C.** 1993. Host plant resistance to insects in sorghum and its role in integrated pest management. *Crop Protection*, **12**: 11-34.
3. **Kumar A. A., Reddy B. V. S., Sharma H. C. and Ramaiah B.** 2008. Shoot fly (*Atherigonasoccata*) resistance in improved grain sorghum hybrids. *Journal of SAT Agricultural Research* **6**.
4. **Dhillon M. K., Sharma H. C., Singh R. and Naresh J. S.** 2005. Mechanisms of resistance to shoot fly, *Atherigonasoccata* in sorghum. *Euphytica*, **144**: 301-312.
5. **Ramu P., Deshpande S. P., Senthilvel S., Jayashree B., Billot C., Deu L., Ananda Reddy M. and Hash C. T.** 2010. In silico mapping of important genes and markers available in the public domain for efficient sorghum breeding. *Molecular Breed.*, **26**: 409-418.
6. **Folkertsma R. T., Sajjanar G. M., Reddy B. V. S., Sharma H. C. and Hash C. T.** 2003. Genetic mapping of QTL associated with sorghum shoot fly (*Atherigonasoccata*) resistance in sorghum (*Sorghum bicolor*). *In: Final Abstracts Guide. Plant and Animal Genome*, **9**: 11-15.