

Research article

Alteration in emergency theatre prioritisation does not alter outcome for acute appendicitis: comparative cohort study

Stefano Partelli, Sabina Beg, Juliette Brown, Soumil Vyas and Hemant M Kocher*

Address: Barts and the London HPB Centre, The Royal London Hospital, Whitechapel, London E1 1BB, UK

Email: Stefano Partelli - stefano.partelli@bartsandthelondon.nhs.uk; Sabina Beg - sabina.beg@bartsandthelondon.nhs.uk;

Juliette Brown - juliette.brown@bartsandthelondon.nhs.uk; Soumil Vyas - soumil.vyas@bartsandthelondon.nhs.uk;

Hemant M Kocher* - hemant.kocher@bartsandthelondon.nhs.uk

* Corresponding author

Published: 8 June 2009

Received: 4 February 2009

World Journal of Emergency Surgery 2009, **4**:22 doi:10.1186/1749-7922-4-22

Accepted: 8 June 2009

This article is available from: <http://www.wjes.org/content/4/1/22>

© 2009 Partelli et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Despite dedicated emergency theatre, emergency surgery can be often delayed due to competing urgencies, suggesting a need for innovative theatre time management.

Aim: To investigate if a change in the emergency theatre prioritisation affects outcomes for a common urgent operation such as appendicectomy.

Methods: We prospectively recorded data from 67 patients undergoing appendicectomy, for two cohorts of patients: before and after change in theatre prioritisation: Group 1 (Jan-Mar) and 2 (Aug-Oct) respectively. Demographic and peri-operative data, time from admission to surgery, postoperative length of stay and total length of stay and complications were compared.

Results: The two groups were comparable with regards to gender, age, time of admission and histological confirmation of appendicitis. No differences between the two groups were found regarding time from admission to surgery (24.4 (95% CI 11.2;27.6) hours versus 16.1 (95% CI 10.4;21.7) hours, Mann-Whitney U test, $p = 0.35$), postoperative length of stay (90.8 (95% CI 61.4;120.1) hours versus 70 (95% CI 48.3;91.6) hours, Mann-Whitney U test, $p = 0.25$) and total length of stay (115.2 (95% CI 84.6;145.7) hours versus 86 (95% CI 61.6;110.4) hours, Mann-Whitney U test, $p = 0.07$) as well as complication or re-admission rates.

Conclusion: A change in the emergency theatre prioritisation does not affect outcome for appendicectomy. Provision of a second emergency theatre could be a solution to reduce the delays in acute surgical operations.

Background

Appendicectomy is amongst the commonest acute surgical operation of intermediate nature, which if not treated in a timely manner could be life-threatening. During the working week (Monday-Friday) urgent and emergency surgery is often delayed until the elective operating list has

finished [1], particularly for non-life or limb threatening situations. The American College of Surgeons recommend the provision of a dedicated trauma operating theatre [2]; this intervention could reduce the incidence of complications [3]. In the UK, the National Confidential Enquiry into Patient Outcome and Death (NCEPOD)

annually recommends changes in management policies affecting patient outcomes based on national audits. In 1992 NCEPOD recommended the provision of dedicated emergency theatres in the UK[4]. Several authors have reported improvement in the quality of emergency services by providing easy access to theatres during daytime and effectively minimising out-of-hours operating [5-9]. Apart from these two instances, we could not uncover any other national audit or guidelines.

Nevertheless NCEPOD report in 2003 suggested that only 58% of all NHS hospitals (in the UK), had a designated theatre for emergency surgery during daytime [10]. Furthermore, even the presence of a single dedicated emergency operating theatre may not be sufficient for a tertiary referral centre, catering to a diverse, socio-economically deprived population and offering specialist trauma surgical services (which takes precedence over most other urgent surgical procedures) [11]. We have previously shown that precisely for this particular reason, common operations such as abscess drainage and appendicectomy stay longer in hospital [11].

We, therefore, convinced the hospital management for a change in emergency theatre utilisation. In the absence of additional space for another parallel day-time emergency theatre, the hospital management implemented a change in emergency theatre prioritisation. Hence we audited whether such a change affected outcomes for appendicectomy.

Methods

For the purpose of this study, in order to obtain two comparable homogenous groups we prospectively collected anonymous data over two time periods: January–March 2008 (Group 1) and August–October 2008 (Group 2). The intervening period (April 2008 – July 2008), was the transition period whilst the below mentioned changes were implemented but were inconsistent with allocation; therefore this period was not analysed. All patients admitted at the Royal London Hospital (RLH) with suspected acute appendicitis were included. Demographic, operative and post-operative details were obtained; time of admission, time of operation, and time of discharge were prospectively recorded.

Before April 2008, the dedicated emergency operating theatres at the RLH worked on "first come first serve" policy, with the flexibility of allowing for immediate surgery, at the clinical discretion of the surgeons and anaesthetists concerned. After April 2008, the dedicated emergency theatre was divided in 3 sessions of 3.5 hours each (divided between 0800 hours to 1830 hours for the five working days), with sessions being systematically allocated to each surgical sub-speciality (General Surgery, Orthopaedics

and Trauma, Vascular/Trauma Surgery, Neurosurgery, Renal Surgery, Gynaecologic Surgery, Plastic Surgery and Oral and Maxillofacial Surgery) on different days of each working week. Additionally, an "open session" allowed for any unscheduled emergency operating.


Statistical analysis

Distribution of continuous variables are reported as median and interquartile range (IQR) (25th; 75th centiles). Categorical variables are presented as numbers and percentages. The comparison between subgroups was carried out using Student's t test, or Mann-Whitney U test, (for continuous variables). Qualitative data were compared by the Chi square test or Fisher's exact test when necessary. Statistical analyses were performed in SPSS 16.0 for Windows software (SPSS Inc, Chicago, Illinois, USA). For all comparisons, a two-sided $p < 0.05$ was considered statistically significant.


Results

Demographic and clinical details are summarized in table 1 with no differences between groups. For the entire cohort of 67 patients the distribution of time of admission (figure 1a), the distribution of time of surgery (figure 1b), showed no difference, allowing us to compare two groups for any delays to theatre. Figure 1c demonstrates time required from decision to operate to time for surgery, again demonstrating no difference (Mann-Whitney U test, $p = 0.349$). A comparison using mean and 95% confidence interval suggested absence of type II error, though, of course, this cannot be entirely ruled out. Thus no differences between the two groups were found regarding time from admission to surgery (24.4 (95% CI 11.2;27.6) hours versus 16.1 (95% CI 10.4;21.7) hours, Mann-Whitney U test, $p = 0.35$), postoperative length of stay (90.8 (95% CI 61.4;120.1) hours versus 70 (95% CI 48.3;91.6) hours, Mann-Whitney U test, $p = 0.25$) and total length of stay (115.2 (95% CI 84.6;145.7) hours versus 86 (95% CI 61.6;110.4) hours, Mann-Whitney U test, $p = 0.07$).


Four patients had post-operative complications: 3 of these were operated within 5–10 hours from admission while the remaining one was operated 18 hours after the admission. In all the 4 patients requiring readmission within a week of discharge, the appendicectomy was performed with a delay of more than 10 hours. Table 1 summarises the final histological examination with a trend to more complicated appendicitis in group 1 (Fisher's exact test, $p = 0.07$). Figure 2c demonstrates that there was no difference in the overall length of stay (Mann-Whitney U test, $p = 0.072$), duration of delay to surgery (Mann-Whitney U test, $p = 0.35$) and length of postoperative stay in hospital (Mann-Whitney U test, $p = 0.25$).



a



b



c

Figure 1
Distribution of patients admitted, with a suspected diagnosis of appendicitis, during the day clustered by time of admission (a), time of operation (b) and delay from making to diagnosis to operation (c) across both groups and overall.

Table 1: Demographic and clinical details

Period	Group 1	Group 2	p	Test
	January–March	August–October		
Number of patients (n)	36	31	-	
Males (n)	27	17	0.08	Fisher's exact
Age (mean;95% CI)	20.7 (16.6;24.7)	25 (19;31)	0.36	Mann-Whitney U
Perioperative antibiotics (n)	15	15	0.63	Fisher's exact
Complications (n)	4	0	0.12	Fisher's exact
Confirmed appendicitis	33	28	1	Fisher's exact
Appendix histology*				
Normal	3	4	0.07	Fisher's exact
Inflamed	19	20		
Necrosed	11	2		
Perforated	3	5		

Discussion

Our audit in a comparable cohort of patients over two different time periods, after a change in theatre prioritisation policy, did not demonstrate any significant differences in the outcome after appendicectomy. The intention of implementing this change was to effectively reduce waiting times to emergency surgery and hence length of hospital stay – but clearly the present study has failed to demonstrate this effect. There could be numerous reasons for this finding.

Foremost, this could be due to the small sample size, which will require a multi-centre study. Such a study could be hampered by non-homogeneity of the profile of emergency workload. Our hospital is one of the premier trauma units in the UK and the only site of the only Helicopter emergency medical service (HEMS) in London. Despite this, numerically at least emergency general surgery accounts for 64.2% of all the emergency surgical workload with abscesses and acute appendicitis being the two most frequent reasons for requiring theatre [11]. Of course, trauma as well as vascular operations, because of the complexity of pre-operative and operative work and multiple team involvement, take longer duration and therefore occupy a prominent part of the emergency theatre schedule.

Some authors have suggested an increase in post-appendicectomy complications and longer hospital stay associated to the delay to surgery [12,13], whilst others have failed to demonstrate this trend [14-17]; although, of course most patients would prefer immediate surgical

procedure [18]. In our cohort only four patients had a complication; of those, three were operated within 10 hours from admission and only one after 18 hours. Our data doesn't demonstrate significant changes in outcome after the appendicectomy, despite changes in theatre prioritisation. The median length of hospital stay was 76 hours, comparable to other publications [13,14].

Delay to surgery is associated with an increased incidence of complications and length of hospital stay after appendicectomy [12,13,19]. Analyzing a large series of 1081 patients, Ditillo et al [12] from the Yale University, USA demonstrated that in adult patients with acute appendicitis, the risk of developing advanced pathology and post-operative complications increases with time; particularly, those risks rise proportional to delay. These results were corroborated by another study from the Wellington Hospital, New Zealand [13] reporting a significant increase in the rate of complicated appendicitis and morbidity when time to surgery exceeded 24 hours. Furthermore, a study from the Massachusetts General Hospital Von Tittle et al [19] reported a incidence of perforation of nearly 90% among 40 patients who had a delay of 72 hours or more after the onset of symptoms. On the other hand others have failed to demonstrate this trend [14-17]. Stahlfeld et al. [15] found no difference in operative time, length of stay, wound infections and antibiotic use in patients operated less than 10 hours from the admission. Similar results were shown by Abou-Nukta et al [14] in a cohort of 309 patients when the delays was 12 to 24 hours. Therefore it seems that a short delay (12–24 hours) to surgery does not significantly alter the outcomes after appen-




Figure 2
Comparison of time from admission to surgery (a), postoperative length of stay (b) and total length of stay (c) between the two groups. Box and whisker graphs represent median \pm inter-quartile range.

dicectomies. However, a greater delay (more than 24 hours) can increase the rate of complications. Delay in carrying out appendicectomy may be due to failure to diagnose the condition accurately, thus resulting in higher incidence of complicated appendicitis (necrosis or perforation) [20]. Over a 25 year period, with increasing use of CT scan and laparoscopy, however there has not been any associated decrease in rate of perforated appendicitis[21]. In our first cohort (group 1), there was a trend towards a delay of mean of 24 hours which may explain a trend towards more complicated appendicitis (table 1).

The median time from admission to operation, the median postoperative and total length of hospital stay were minimally reduced after the changing the theatre prioritisation scheme but these results failed to reach a statistical significance. Utilization of the operating theatre (OT) should not only to guarantee that the greatest number of cases are done, but also consider the costs involved [22]. When additional OT capacity is available, it should be planned with multiple variables in mind such as sub-specialities with the greatest contribution margin per OT hour, as well as those that have minimal need for limited resources such as intensive care unit beds[23]. Mainly due to financial circumstances it is difficult to provide one or more dedicated emergency OTs even if it is strongly desired based on clinical needs [24]. Day case surgery can be severely affected by the increase of emergency admissions. *Nasr et al* reported that 40% of all planned elective surgical operations were cancelled, mainly due to bed unavailability because of the overflow of emergency admissions [25]. Robb et al confirmed the increasing role of the bed unavailability in the cancellation of elective surgical cases and additionally demonstrated cost implications[26]. Vinukondaya et al reported that emergency surgery during the operating list is the reason for cancellation of elective surgery in the 13.9% of the cases [27].

In other countries the main cause for emergency surgery delays is not due to the absence of a dedicated emergency OT. Data from 498 patients from the University College Hospital of Ibadan, Nigeria, over a three-month period showed that only in 38% of cases booked for an emergency operation, surgery was carried out. The main reason for cancellation was surgeon's unavailability [28]. Changing the operating theatre policy, as demonstrated in this article, allows surgeons to designate and inform the patient more accurately the time of his/her operation. However, it did not necessarily reduce the waiting times to surgery. We feel that provision of a second emergency theatre at all times would be an effective solution to this problem. Patients would be operated upon promptly. This would reduce waiting times to surgery and facilitate quicker discharges from hospital, thereby increasing turnover. This would also be satisfactory for the patients; bed

management for the elective patients, thereby increasing volumes of elective work load and shortening waiting list times. The increased costs involved in running the second additional theatres should be balanced against the cost of reduced length of hospital stay. Taking an example from emergency laparoscopic cholecystectomy versus elective cholecystectomy after conservative management, the increased immediate operative cost is neutralized by the reduced length of stay and quicker return to work [29]. More detailed cost – benefit analysis involving multiple hospitals and larger number of patients would be required to lend credible evidence to support this belief.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SP, SB, JB, and SV collected data under supervision of HMK. HMK initiated the project; did the analysis and wrote the paper with SP. HMK will act as a guarantor for the manuscript.

Acknowledgements

We thank all the medical and nursing staff of the wards and theatres of the surgical services for taking care of patients and helping in data collection. We thank Mr Ajit Abraham & Mr Mike Walsh, Consultant Surgeons for spearheading the theatre change programme and Ms Ceri Cranston, Theatre Manager for implementing the changes with rigor.

References

- Wyatt MG, Houghton PW, Brodribb AJ: **Theatre delay for emergency general surgical patients: a cause for concern?** *Ann R Coll Surg Engl* 1990, **72(4)**:236-8.
- American College of Surgeons Trauma Program** [<http://www.facs.org/trauma>]
- Bhattacharyya T, et al.: **The value of the dedicated orthopaedic trauma operating room.** *J Trauma* 2006, **60(6)**:1336-40. discussion 1340-1
- The Report of the National Confidential Enquiry into Perioperative Deaths 1990.** NCEPOD, London; 1992.
- Sweetnam DI, Williams JR, Britton DC: **An audit of the effect of a 24-hour emergency operating theatre in a district general hospital.** *Ann R Coll Surg Engl* 1994, **76(2 Suppl)**:56-8.
- Lovett BE, Katchburian MV: **Emergency surgery: half a day does make a difference.** *Ann R Coll Surg Engl* 1999, **81(1)**:62-4.
- Calder FR, Jadhav V, Hale JE: **The effect of a dedicated emergency theatre facility on emergency operating patterns.** *J R Coll Surg Edinb* 1998, **43(1)**:17-9.
- Barlow AP, et al.: **An emergency daytime theatre list: utilisation and impact on clinical practice.** *Ann R Coll Surg Engl* 1993, **75(6)**:441-4.
- Scriven MW, et al.: **The use and impact of a daily general surgical emergency operating list in a district general hospital: a prospective study.** *Ann R Coll Surg Engl* 1995, **77(3 Suppl)**:117-20.
- The 2003 Report of the National Confidential Enquiry into Perioperative Deaths.** NCEPOD, London; 2003.
- Mai-Phan TA, et al.: **Emergency room surgical workload in an inner city UK teaching hospital.** *World J Emerg Surg* 2008, **3**:19.
- Ditillo MF, Dziura JD, Rabinovici R: **Is it safe to delay appendectomy in adults with acute appendicitis?** *Ann Surg* 2006, **244(5)**:656-60.
- Omundsen M, Dennett E: **Delay to appendicectomy and associated morbidity: a retrospective review.** *ANZ J Surg* 2006, **76(3)**:153-5.

14. Abou-Nukta F, et al.: **Effects of delaying appendectomy for acute appendicitis for 12 to 24 hours.** *Arch Surg* 2006, **141(5)**:504-6.
15. Stahlfeld K, et al.: **Is acute appendicitis a surgical emergency?** *Am Surg* 2007, **73(6)**:626-9.
16. Clyde C, et al.: **Timing of intervention does not affect outcome in acute appendicitis in a large community practice.** *Am J Surg* 2008, **195(5)**:590-2.
17. Eldar S, et al.: **Delay of surgery in acute appendicitis.** *Am J Surg* 1997, **173(3)**:194-8.
18. Sideso E, Richards T, Galland RB: **Appendicectomy deferred to a CEPOD list: the patients' opinion.** *Surgeon* 2008, **6(4)**:198-200.
19. Von Tittle SN, McCabe CJ, Ottinger LW: **Delayed appendectomy for appendicitis: causes and consequences.** *Am J Emerg Med* 1996, **14(7)**:620-2.
20. Chung CH, Ng CP, Lai KK: **Delays by patients, emergency physicians, and surgeons in the management of acute appendicitis: retrospective study.** *Hong Kong Med J* 2000, **6(3)**:254-9.
21. Livingston EH, et al.: **Disconnect between incidence of nonperforated and perforated appendicitis: implications for pathophysiology and management.** *Ann Surg* 2007, **245(6)**:886-92.
22. Viapiano J, Ward DS: **Operating room utilization: the need for data.** *Int Anesthesiol Clin* 2000, **38(4)**:127-40.
23. Wachtel RE, Dexter F: **Tactical increases in operating room block time for capacity planning should not be based on utilization.** *Anesth Analg* 2008, **106(1)**:215-26.
24. Collins C: **The standards for emergency surgical services.** *J R Soc Med* 2001, **94(Suppl 39)**:13-5.
25. Nasr A, et al.: **Impact of emergency admissions on elective surgical workload.** *Ir J Med Sci* 2004, **173(3)**:133-5.
26. Robb WB, et al.: **Are elective surgical operations cancelled due to increasing medical admissions?** *Ir J Med Sci* 2004, **173(3)**:129-32.
27. Vinukondaiah K, Ananthkrishnan N, Ravishankar M: **Audit of operation theatre utilization in general surgery.** *Natl Med J India* 2000, **13(3)**:18-21.
28. Windokun A, Obideyi A: **Audit of emergency theatre utilisation.** *Afr J Med Med Sci* 2002, **31(1)**:59-62.
29. Germanos S, Gourgiotis S, Kocher HM: **Clinical update: early surgery for acute cholecystitis.** *Lancet* 2007, **369(9575)**:1774-6.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

