

Accepted Manuscript

Coffee, Tea, and Cardiovascular Morbidity and Mortality

Carl J. Lavie M.D. , James J. DiNicolantonio Pharm D ,
James H. O’Keefe M.D.

PII: S1050-1738(18)30239-1
DOI: <https://doi.org/10.1016/j.tcm.2018.11.016>
Reference: TCM 6616

To appear in: *Trends in Cardiovascular Medicine*



Please cite this article as: Carl J. Lavie M.D. , James J. DiNicolantonio Pharm D ,
James H. O’Keefe M.D. , Coffee, Tea, and Cardiovascular Morbidity and Mortality, *Trends in
Cardiovascular Medicine* (2018), doi: <https://doi.org/10.1016/j.tcm.2018.11.016>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Coffee, Tea, and Cardiovascular Morbidity and Mortality

Carl J. Lavie M.D. (1), James J. DiNicolantonio, Pharm D (2), James H. O'Keefe, M.D. (2)

Department of Cardiovascular Diseases

John Ochsner Heart and Vascular Institute

Ochsner Clinical School - The University of Queensland School of Medicine

New Orleans, Louisiana (1)

Saint Luke's Mid America Heart Institute

Kansas City, Missouri (2)

Direct correspondence to Dr. Lavie:

John Ochsner Heart and Vascular Institute

Ochsner Clinical School - The University of Queensland School of Medicine

1514 Jefferson Highway

New Orleans, Louisiana 70121

Phone: 504-842-1281

E-mail: clavie@ochsner.org

Caffeine is probably the most widely consumed stimulant worldwide, as the majority of adults worldwide consume caffeine on a daily basis, most commonly in the forms of caffeinated coffee and tea. Coffee, in particular, is one of the most commonly consumed pharmacologically active beverages across the world, with over 50% of Americans above the age of 18 drinking coffee on a daily basis, and another 30% of the population consuming it occasionally. More than 4 million metric tons of tea are consumed each year, and tea is consumed by two-thirds of all adults worldwide. Because of the remarkably high prevalence of the frequent consumption of these beverages, even small health effects of coffee and tea could translate into large effects on public health, positively or negatively, as we and others have discussed many times during the past five years. (1 - 12) In the past decades, there was concern that the stimulatory effects of these beverages could raise blood pressure (BP), trigger atrial and/or ventricular arrhythmias, and generally cause adverse effects on health and well-being.

In the current review in *Trends in Cardiovascular Medicine*, Voskoboinik and colleagues (13) reviewed the cardiovascular (CV) effects of these caffeinated beverages, and reviewed their impact on chronic diseases, especially CV diseases (CVD). Although acutely caffeine in large doses can increase BP, in general, these agents have been shown to have neutral to beneficial effects on the risk of developing hypertension (HTN). Regular coffee and caffeine, in a dose-dependent fashion, confer beneficial effects on glucose metabolism, including significant reductions in the risk for type 2 diabetes mellitus (T2D); maximal protection against T2D is noted with six or more cups per day. Some of the benefits may be due to reduction of obesity, either directly by increasing metabolism or indirectly or via displacing high caloric beverages. Although short-term use of unfiltered coffee may worsen lipids, this does not occur with tea or filtered coffee. (1 -13)

More importantly, coffee and tea consumption have not been associated with increases in CVD. Although the impact on secondary prevention of coronary heart disease (CHD) may be neutral, there seems to be modest impact to protect against incident CVD. Regular coffee consumption and tea consumption appear to reduce the risk of stroke, despite its potential vasoconstrictive effects on the cerebrovascular bed. Since coffee may protect against the development of CHD and HTN, two strong risk factors for heart failure (HF), it may protect against the development of HF, as well as benefit patients with established HF, possibly by having a diuretic effect and without worsening dysrhythmias. In fact, in population studies, caffeine and coffee may be neutral or even actually reduce the risk of atrial fibrillation, without worsening ventricular arrhythmias. (2,13-15)

More importantly, large scale studies suggest that coffee and tea consumption may reduce the risk of both all-cause and CVD mortality, with consumptions as high as four to five cups per day being associated with the greatest benefits, (1, 2, 13) although our data years ago suggested potential toxicity at doses of five cups per day or higher for younger individuals. (4)

Although caffeine has potential beneficial effects, we believe it is probably prudent to keep upper doses at 200 to 300 mg/day (two to three cups of coffee), moreover, these benefits and safety may not extend to certain energy drinks, which can contain excessively high caffeine doses in addition to high levels of added sugar. Not surprisingly, energy drinks have been associated with serious potential toxicity. (13)

In addition to CVD benefits, coffee has been associated with lower risks for breast, endometrial, prostate, and colorectal malignancies. Moreover, coffee (particularly caffeinated coffee) bestows other potential benefits, including slowing cognitive decline, reducing risks for depression, suicide, obesity, asthma, and liver disease. (1-3, 6, 8) Realizing that observational data do not prove cause and effect, it is

difficult to recommend coffee and tea as a prevention strategy. Additionally, the potential benefits need to be weighed against potential risks, generally due to high caffeine content, including anxiety, insomnia, headaches, tremulousness, palpitations, low birth weight, and increased risk of fractures in women. However, the overall evidence is quite reassuring, and probably enough for us to encourage lovers of these beverages to continue to enjoy them and possibly reap the benefits of chronic disease and CVD prevention. We applaud Voskoboinik and colleagues (13) for their outstanding review in this issue of *TCM*.

References

1. O'Keefe JH, Bhatti SK, Patil HR, DiNicolantonio JJ, Lucan SC, Lavie CJ. Effects of habitual coffee consumption on cardiometabolic disease, cardiovascular health, and all-cause mortality. *J Am Coll Cardiol* 2013;62:1043-1051.
2. O'Keefe JH, DiNicolantonio JJ, Lavie CJ. Coffee for cardioprotection and longevity. *Prog Cardiovasc Dis* 2018;61:38-42.
3. Bhatti SK, O'Keefe JH, Lavie CJ. Coffee and tea: perks for health and longevity? *Curr Opin Clin Nutr Metab Care* 2013;16:688-697.
4. Liu J, Sui X, Lavie CJ, Hebert JR, Earnest CP, Zhang J, Blair SN. Association of coffee consumption with all-cause and cardiovascular disease mortality. *Mayo Clin Proc* 2013;88:1066-1074.
5. Yan Y, Sui X, Yao B, Lavie CJ, Blair SN. Is there a dose-response relationship between tea consumption and all-cause, CVD, and cancer mortality? *J Am Coll Nutr* 2017;36:281-286.
6. Patil H, Lavie CJ, O'Keefe JH. Cuppa Joe: friend or foe? Effects of chronic coffee consumption on cardiovascular and brain health. *Mo Med* 2011;108:431-438.
7. DiNicolantonio JJ, O'Keefe JH, Lavie CJ. Reply: Effects of habitual coffee consumption on vascular function. *J Am Coll Cardiol* 2014;63:607.
8. Tsujimoto T, Kajio H, Sugiyama T. Association between caffeine intake and all-cause and cause-specific mortality: a population-based prospective cohort study. *Mayo Clin Proc* 2017;92:1190-1202.
9. Joyner MJ. Caffeine: a not so stiff drink. *Mayo Clin Proc* 2018;93:558-559.
10. Liu J, Sui X, Blair SN, Lavie CJ. In reply: Association of coffee consumption with all-cause and cardiovascular disease mortality. *Mayo Clin Proc* 2013;88:1493-1494.

11. DiNicolantonio JJ, Lucan SC, O'Keefe JH. Is coffee harmful? If looking for longevity, say yes to the coffee, no sugar. *Mayo Clin Proc* 2014;89:577.
12. Liu J, Sui X, Blair SN. In reply: is coffee harmful? If looking for longevity, say yes to the coffee, no sugar. *Mayo Clin Proc* 2014;89:576-577.
13. Voskoboinik A, Koh Y, Kistler PM. Cardiovascular effects of caffeinated beverages. *Trends Cardiovasc Med* 2018; in press.
14. Conen D, Chiuve SE, Everett BM, Zhang SM, Buring JE, Albert CM. Caffeine consumption and incident atrial fibrillation in women. *Am J Clin Nutr* 2010;92:509-514.
15. Shen J, Johnson VM, Sullivan LM, Jacques PF, Magnani JW, Lubitz SA, et al. Dietary factors and incident atrial fibrillation: the Framingham Heart Study. *Am J Clin Nutr* 2011;93:261-266.