

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This author's accepted manuscript may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

The full details of the published version of the article are as follows:

TITLE: Retrospective evaluation of factors influencing transfusion requirements and outcome in cats with pelvic injury (2009–2014): 122 cases

AUTHORS: Poppy Gant, Imola Asztalos, Elvin Kulendra, Karla Lee, Karen Humm

JOURNAL: Journal of Veterinary Emergency and Critical Care

PUBLISHER: Wiley

PUBLICATION DATE: 20 June 2019

DOI: <https://doi.org/10.1111/vec.12852>

1 **Abstract:**

2 **Objective** – To characterize a population of cats with pelvic trauma and evaluate factors influencing transfusion
3 requirement and outcome.

4 **Design** – Retrospective case series (2009-2014)

5 **Setting** – University teaching hospital

6 **Animals** - One hundred and twelve client-owned cats with pelvic trauma

7 **Interventions** – None

8 **Measurements and main results** - Twenty-one (18.8%) cats received a transfusion. Most cats required only one fresh
9 whole blood transfusion (85.8%). Packed cell volume at admission was significantly lower in cats that required
10 transfusion but was not associated with hospitalization time or survival to discharge. Increasing Animal Trauma
11 Triage (ATT) score at admission was significantly associated with transfusion requirement ($p=0.0001$) and non-
12 survival to discharge ($P=0.03$). Number of pelvic fractures was not associated with transfusion requirement but cats
13 with sacroiliac luxations and pubic fractures were more likely to require a transfusion ($p=0.0015$ and $p=0.0026$
14 respectively). However, fracture type was not associated with survival to discharge. Most cats (86%) required a
15 surgical procedure and half of transfusions were administered pre-operatively. No surgical comorbidities were
16 associated with transfusion requirement or survival. Transfusion requirement was associated with longer length of
17 hospitalization but not survival to discharge.

18 **Conclusions** – Transfusion requirement in this population of cats with pelvic fractures was fairly high. Transfusion
19 requirement was associated with lower packed cell volume, higher ATT score at admission, longer length of
20 hospitalization and certain types of pelvic fractures. Transfusion requirement was not associated with surgical co-
21 morbidities, surgical intervention or survival to discharge. Lower ATT score at admission was associated with survival
22 to discharge.

23

24 ***Abbreviations:***

25 ATT Animal Trauma Triage Score

26 ISS Injury Severity Score

27 FWB Fresh Whole Blood

28 PCV Packed Cell Volume

29 AFAST Abdominal Focused Assessment with Sonography for Trauma

30

31 **Introduction:**

32 Pelvic fractures in cats account for approximately 25% of all reported fractures and are usually a result of
33 high impact blunt trauma.¹ These forces, when applied to the rigid box-like structure of the pelvis, often result in
34 multiple bilateral fractures and pelvic instability. Consequently, pelvic hemorrhage can occur, originating from bone
35 or from disruption of local vasculature.²

36 High impact trauma is also commonly associated with injury to other body systems. These sites may act as
37 additional sources of hemorrhage. Injury to the thorax and abdomen in particular are thought to contribute
38 significantly to morbidity and mortality. Additional soft tissue, urinary tract, and neurologic injuries are also
39 encountered.³

40 Traumatic pelvic fractures in people are associated with high mortality rates in the first 24 hours. This is
41 most commonly associated with acute, severe hemorrhage from the pelvis or abdomen.⁴⁻⁶ Predicting transfusion
42 requirement and outcome in this patient group has therefore received much attention. Age⁷, presence of shock on
43 arrival⁷⁻¹⁰ and admission hematocrit,^{8,10} have all been associated with requirement for blood products. However,
44 multiple large studies have been unable to consistently predict transfusion requirement based on the type of pelvic
45 fracture, with various classification schemes being trialed based on direction of force, degree of displacement and
46 rotational stability.^{6-8,11,12} This is likely because isolated major hemorrhage from the pelvis is thought to be rare.^{4,13}
47 Overall assessment of concurrent injuries has instead been shown to be more useful in predicting transfusion
48 requirement and mortality in human pelvic fracture patients.^{5,6,13,14} This is often presented as an Injury Severity
49 Score (ISS): an anatomical scoring system of 6 body regions to numerically describe the overall severity of injury.¹⁵
50 Mortality in this patient group has been associated with age^{6,14,16}, admission hematocrit/hemoglobin⁵, systolic
51 arterial blood pressure^{5,14,17} and transfusion requirement.^{6,16,18} Pre and post operative hemoglobin levels have been
52 associated with length of hospitalization.¹⁹

53 Although there is little information regarding the use of transfusions in feline trauma patients, it is the

54 perception of the authors that acute, severe hemorrhage is an uncommon presentation in cats with pelvic fractures.
55 However, whilst rarely needed during initial resuscitation, blood transfusions are considered to be a fairly frequent
56 requirement to stabilize patients for further interventions.²

57 Up to 58.6% of cats with pelvic fractures have been reported to have concurrent injuries,²⁰ and the
58 presence of abdominal injury or soft tissue trauma has been associated with non-survival to discharge in general
59 feline trauma cases.³ However, to the authors' knowledge, the incidence of significant hemorrhage, and consequent
60 requirement for transfusion and outcome prediction, has not been specifically investigated in feline patients. The
61 aim of the current study was to perform a retrospective review of cats that sustained traumatic pelvic fractures, to
62 document their need for blood transfusion and to identify predictors of transfusion requirement and outcome. Our
63 principal null hypothesis was that receiving a blood transfusion would have no impact on survival to discharge in cats
64 with pelvic trauma.

65

66

67 ***Materials and Methods:***

68 The medical records database at a university teaching hospital was searched for cats presenting with pelvic
69 trauma between January 2009 and January 2014. Cats were eligible for study inclusion if they had sustained any
70 bony pelvic injury (fractures and/or sacroiliac luxation), had complete medical records (detailing initial physical
71 examination findings, packed cell volume, treatment given and patient outcome), and had radiographs or computed
72 tomography images of the pelvis reviewed at the time by a board certified diagnostic imager. Age, sex, breed and
73 body weight were recorded. Both primary and referral cases were included. An Animal Trauma Triage Score (ATT)
74 ¹¹, an anatomical numerical scoring system for assessing polytrauma, similar to the ISS in people, was determined
75 retrospectively based on the physical examination findings at the time of presentation.

76 A blood transfusion was defined as administration of any hemoglobin-containing blood product including
77 Oxyglobin™^[a]. The type, and number, of transfusions provided, whether they were administered pre, intra or post-
78 operatively and the packed cell volume (PCV) prior to transfusion were recorded for all cases.

79 The types of pelvic injury documented on the imaging reports were recorded as sacroiliac luxation, sacral
80 fracture, ilial fracture, acetabular fracture, ischial fracture, and pubic fracture. Management of pelvic fractures
81 (surgical or conservative treatment) was also recorded. Additional co-morbidities requiring surgical intervention
82 were determined from the medical records, then classified by body system affected (cardiovascular, neurological,
83 urinary, integumentary, gastrointestinal, ocular). Integumentary systems injuries were included if abrasions, partial
84 thickness lacerations, or full thickness lacerations involving deep tissues but without penetration into a body cavity
85 were noted in the medical record.

86 Length of hospitalization and survival to discharge were recorded. If a patient died, then it was noted
87 whether this was secondary to cardiopulmonary arrest or euthanasia.

88

89

90 ***Statistical Methods:***

91 For statistical analysis of study objectives, cats were classified as either transfusion status positive or
92 negative, whether they received any surgical intervention and whether they survived to discharge. Population
93 characteristics (age, ATT score, presence of surgical co-morbidities, type of pelvic fracture and PCV at admission) and
94 outcome criteria (requirement for surgery, length of hospitalization and mortality) were then compared between
95 transfusion status, requirement for surgical procedure and survival to discharge groups.

96 Continuous data sets were analyzed for normality using the Shapiro Wilk test. A Student's t-test was used
97 to compare the means of continuous data. Categorical data were presented as median and range. A Fisher's exact
98 test was used to compare categorical data. Ordinal data was presented as median and range. Chi-square for trend
99 analysis was used to examine the relationship between ATT and requirement for blood transfusion or survival. The
100 remaining data were presented descriptively. Statistical analysis was performed using Graphpad software^b. Results
101 were considered significant with a P value <0.05.

102

103

104 ***Results:***

105 One hundred and twelve cats were identified with pelvic fractures over the study period. All data could be
106 obtained from the medical records and therefore no cats were excluded from the study. The cats had a median age
107 of 40 months (range 4 months to 192 months). There was no significant difference between the ages of the cats
108 based on requirement for transfusion, surgical intervention or survival to discharge. There were more males than
109 females (63 and 49 respectively) and 92.8% were neutered. Median weight was 4.18kg (range 2.2 – 6.18kg). Fourteen
110 breeds were represented with Domestic Shorthairs being the most common (63.4%). The majority of cases were un-
111 witnessed trauma but thought to be vehicular related. Seventy-nine cats (70.5%) were presented to a primary care
112 practice before referral.

113 Twenty-one (18.6%) cats received a blood transfusion. Average time from admit to transfusion was 3.6 days
114 (range 1 to 6 days). All cats that received a transfusion underwent surgery, with 12 transfusions (50%) being
115 administered pre-operatively, 3 intra-operatively (12.5%) and 9 post-operatively (37.5%). Eighteen cats received one
116 transfusion (85.8%) and 3 cats received 2 transfusions (14.2%). Eighteen cats received fresh whole blood (FWB) alone
117 and 3 cats received both FWB and Oxyglobin™. Cats receiving Oxyglobin™ all went on to receive a FWB transfusion.

118 The median ATT score for all cats was 4. Median ATT score for cats receiving a transfusion was 6 (range 3-
119 11) compared to 4 (range 0-9) for those not receiving a transfusion. A chi-square test for trend showed a significant
120 linear association between increasing ATT at admission and proportion of cats receiving a blood transfusion ($P =$
121 0.0001). Median ATT score for cats surviving to discharge was 4 (range 0-10) compared to 6 (range 2-11) in those
122 that did not survive. A chi-square test for trend showed a significant linear association between a lower ATT score
123 at admission and survival ($P < 0.03$).

124 In total, 96 of 112 cats (86%) had a surgical procedure. The majority of cats (75%) underwent fracture
125 stabilization. Some of these required an additional procedure with twenty-five cats (22.3%) undergoing surgery for
126 injuries sustained to another body system. Fifteen cats (13.4%) sustained skin wounds and 9 cats (8.0%) had urinary
127 tract trauma requiring surgical management. One cat required an enucleation (0.9%). There was no significant
128 difference between the frequency of surgical co-morbidities in the transfusion or survival groups. There was also no
129 significant difference between the transfusion requirement or the ATT scores for cats that did or did not undergo a
130 surgical procedure.

131 Diagnostic imaging revealed 76 out of 112 cats had sacroiliac luxation(s) (67.9%), 58 had pubic fractures
132 (51.8%), 54 had ischial fractures (48.2%), 45 had ilial fractures (40.2%), 19 had acetabular fractures (17.0%), 11 had
133 coxofemoral luxations (9.8%), 6 had lumbosacral luxations (5.3%) and 6 had sacral wing fractures (5.3%). There was
134 no significant difference in the number of pelvic fractures in those cats that required a transfusion and those that
135 didn't. However, significantly more cats that required a transfusion had sacroiliac luxation and pubic fractures
136 compared to those cats that weren't transfused ($p=0.0015$ and $p=0.0026$ respectively). Type of pelvic fracture was
137 not associated with survival to discharge.

138 The mean PCV at admission was significantly lower in the cats that required transfusion (23% [SD 6.6%])
139 compared to the group that weren't transfused (29.2% [SD 7.0%]) ($p=0.0005$). There was no correlation between
140 PCV at admission and hospitalization time, irrespective of transfusion status. PCV was not associated with survival
141 to discharge. When comparing the PCV of first opinion and referral cases, for the cats that were transfused, the
142 mean PCV was not significantly different (first opinion = 25.6% [SD=8.5%]; referred = 20% [SD=6.3%]). For the cats
143 that weren't transfused, the PCV was significantly different (first opinion = 35.5% [SD=6.2%]; referred = 27.4 [5.8%];
144 $p<0.0001$).

145 All cats were hospitalized for at least 24 hours and 102 cats survived to discharge (91.1%). Mean length of
146 hospitalization for all cats was 7.0 days (SD=4.0). After excluding those that were euthanized, length of
147 hospitalization was significantly longer in cats that received a transfusion (mean=9.4days [SD=5.2days]) compared
148 to those that didn't (mean=6.4days [SD3.5days]) ($p0.002$). Of the 10 cats that did not survive, 2 animals underwent
149 cardiorespiratory arrest and the remainder were euthanized. Only 4 out of the 10 underwent a surgical procedure
150 (two underwent fracture stabilization; two other cats had cystotomy tubes placed). Requirement for transfusion or
151 surgery was not associated with survival to discharge.

152

153 **Discussion:**

154 The results of this study indicate that, in this referral population of cats with pelvic fractures, transfusion
155 requirement was fairly high, with 18.6% of cats requiring a FWB transfusion during hospitalization. Transfusion
156 requirement was associated with lower PCV and increased ATT score at admission, as well as sacroiliac luxation and
157 pubic fractures. Transfusion requirement was not associated with surgical co-morbidities, surgical intervention or

158 survival to discharge but was associated with longer length of hospitalization. Overall survival to discharge was high
159 and a higher ATT score at admission was the only finding associated with non-survival to discharge.

160 The signalment in this population was similar to that reported in other studies, with younger, male cats
161 being most commonly represented.^{21,22} Age was not associated with transfusion requirement or survival to
162 discharge as has been reported in human studies. This is likely because the geriatric population in this study was
163 small, with only 5.4% of cats greater than 10 years old. With a small geriatric component, pre-existing age-related
164 co-morbidities that could impact the ability to tolerate anemia, or impact survival, were difficult to evaluate.

165 Nearly 1 in 5 cats required a transfusion, although no transfusions were administered in the first 24 hours
166 and no cat received more than one FWB transfusion. Transfusion is therefore unlikely to be required for immediate
167 resuscitation in this population as all cats survived at least 24 hours from presentation. This is useful information for
168 the clinician discussing financial implications with owners and also when planning blood product supplies. It may
169 also explain why transfusion requirement had no association with mortality in contrast to human medicine. People
170 with pelvic fractures can require multiple transfusions during initial stabilization and this may ultimately result in a
171 much greater proportion of the circulating blood volume being replaced. This can predispose to a coagulopathic
172 state and increased mortality.²³ Cats that received a transfusion were hospitalized for significantly longer than those
173 that did not. This may be associated with the time required time required to procure FWB or overall injury severity,
174 as cats requiring a transfusion also had a higher ATT score.

175 The finding that the ATT score was associated with transfusion requirement and also survival to discharge
176 is similar to the relationship between ISS and transfusion requirement and mortality in people with pelvic fractures.
177 A previous study also demonstrated this relationship in dogs.²⁶ This same canine study also suggested that
178 hemoperitoneum, as detected by abdominal focused assessment with sonography for trauma (AFAST) scans, is a
179 common finding in canine pelvic fracture patients with high ATT scores.²⁶ Therefore, although the cats in this study
180 were not routinely screened for hemoabdomen, given the data available in dogs, inclusion of serial AFAST evaluation
181 may have further helped identify cats at risk of requiring a transfusion.

182 In this population, only 50% of all transfusions were administered preoperatively. It is unclear given the
183 retrospective nature of this study whether this was a result of difficulty obtaining feline blood products, or whether
184 the patients truly did not require blood until intra- or post-operatively. One concern, if related to difficulty obtaining

185 the blood for transfusion, is that this could mean administration was not always at the optimum time for the patient.
186 Unlike in people and dogs, there are no widely available storage facilities for feline blood products in the United
187 Kingdom and much of Europe. Transfusion availability therefore often relies on the location of a suitable donor and
188 collection of blood on demand, both of which can significantly delay transfusion administration. Given these
189 impediments to transfusion, being able to better predict transfusion requirements in feline trauma patients would
190 aid in their timely management, especially if transfusions are required prior to further interventions.

191 The finding that cats requiring transfusion had significantly lower PCVs on presentation may help to prompt
192 clinicians to consider that significant hemorrhage has occurred and blood transfusion may be required. However,
193 PCV alone is not the sole indication for transfusion administration, and it should also be remembered that this value
194 will be influenced by volume and hydration status. Studies in dogs and humans have shown a better correlation
195 between lactate or base excess and transfusion requirement and mortality compared to initial PCV.^{24 25}

196 The major cause of anemia in this population of cats was thought to be blood loss due to the initial trauma
197 that caused the pelvic fracture. However, critically ill cats presenting secondary to trauma may develop anemia via
198 a variety of pathways including frequent phlebotomy, oxidative damage and a poor regenerative response. In a
199 retrospective study, 33% of critically ill cats developed anemia whilst hospitalized. Given over one third of
200 transfusions were given post-operatively, it's likely anemia was multifactorial in these patients. Intra-operative
201 factors such as surgical time, blood loss or anesthetic factors, which could not be determined retrospectively, may
202 also have had an impact.

203 Concurrent surgical injuries were seen in 22.3% of this population. Although the frequency of surgical co-
204 morbidities was not associated with transfusion status or survival to discharge, they may have implications for early
205 stabilization of patients. For example, in this study, all trauma to the urinary tract was managed surgically rather
206 than medically and therefore cats required stabilization prior to/in order to undergo general anesthesia. Only
207 surgical co-morbidities were included in this study as they were more consistently recorded and described in the
208 patient records. However, in omitting medical comorbidities from the study, which are more variable in severity and
209 therefore difficult to consistently recognize in a retrospective study, we could have failed to identify other factors
210 that may have influenced outcome.

211 The finding that sacroiliac luxations and pubic fractures were both seen more frequently in the transfusion
212 group could theoretically be useful. However, as previously discussed, significant hemorrhage from the pelvis alone
213 is thought to be rare in people and concurrent injuries are more useful in predicting transfusion requirement. Further
214 studies are required to determine if this is also the case in cats, although the finding that higher ATTs were associated
215 with transfusion requirement suggests this may be possible. Given the overall frequency of sacroiliac luxations and
216 pubic fractures in this population, it is also not feasible to use these injuries alone to predict transfusion requirement
217 clinically. However, they should perhaps prompt more thorough investigation of pelvic and extra-pelvic hemorrhage
218 and closer monitoring of PCV.

219 Survival to discharge in this population was high at 91% and our null hypothesis was rejected as no
220 association was found between survival and transfusion requirement. The low mortality rates seen in this study are
221 consistent with other veterinary studies, including a report of 12-14% mortality in dogs presenting with blunt trauma.
222 ²⁷ However, cats with the most severe trauma may die very soon after the injury, therefore never presenting to a
223 veterinary facility for treatment. As such, this study may be a poor representation of the overall population of cats
224 that suffer pelvic trauma and survival rates may not be as high. Furthermore, the ATT scores of referral cases may
225 also not reflect the initial primary care assessment and may have been significantly higher in some cats, depending
226 on the varying levels of care prior to arrival at our facility.

227 This study has a few other limitations to note. It was not possible to access digital radiographs between
228 2009 and 2011. Therefore, although each radiograph was reported by a board certified diagnostic imager, none of
229 the radiographs were reviewed specifically for this study. It was therefore not possible to investigate whether
230 specific characteristics of sacroiliac luxation and pubic fractures, for example, degree of cranio-caudal or medio-
231 lateral displacement provide better prediction of transfusion requirement, hospitalization time or survival. However,
232 in dogs, increased lateral pelvic compression was not associated with increased intra-abdominal injury (including
233 hemoperitoneum) and in humans the literature is inconsistent. Although not necessarily required for surgical
234 planning, CT would also have offered better analysis of pelvic fractures compared to radiography.²⁵

235 A further limitation of this study is that more cats may have benefitted from, rather than required, blood
236 products but may have not received them due to limited resources. This could be from a lack of synthetic product
237 availability, variable presence of donor colonies and reliance on client owned donors. Given there are no

238 standardized transfusion 'trigger' protocols in veterinary medicine, clinician decision making about when to
239 transfuse patients also introduces some variability.

240

241 Conclusions:

242 This study highlights that cats presenting with pelvic fractures require transfusion fairly frequently, although in
243 this population, no cat required more than one FWB transfusion. Transfusion requirement was associated with lower
244 PCV and higher ATT score on admission; presence of sacroiliac luxations and pubic fractures and longer
245 hospitalisation times. Transfusion requirement was not associated with surgical co-morbidities or surgical
246 intervention. This data supports our null hypothesis as only ATT score was associated with survival to discharge and
247 not transfusion requirement.

248

249

250

251

252 **Footnotes:**

253 a. Oxyglobin™ Biopure Corporation, Cambridge, MA

254 b. GraphPad Prism 7, GraphPad Software, La Jolla California USA

255

256

257

258

259

260 **References:**

261 1. Phillips IR. A survey of bone fractures in the dog and cat. J Small Anim Pract 1979;20(11):661–74.

262 2. Meeson R, Corr S. Management of Pelvic Trauma Neurological Damage, Urinary Tract Disruption and Pelvic
263 Fractures. J Feline Med Surg 2011;13(5):347–61.

264 3. Zulauf D, Kaser-Hotz B, Hässig M, Voss K, Montavon P. Radiographic examination and outcome in consecutive

- 265 feline trauma patients. *Vet Comp Orthop Traumatol*. 2008;21(1):36–40.
- 266 4. Vaidya R, Scott AN, Tonnos F, Hudson I, Martin AJ, Sethi A. Patients with pelvic fractures from blunt trauma.
267 What is the cause of mortality and when? *The American Journal of Surgery*. 2016;211(3):495–500.
- 268 5. Holstein JH, Culemann U, Pohlemann T. Working Group Mortality in Pelvic Fracture Patients. What are
269 predictors of mortality in patients with pelvic fractures? *Clin Orthop Relat Res*. 2012 Aug;470(8):2090–7.
- 270 6. Smith W, Williams A, Agudelo J, Shannon M, Morgan S, Stahel P, et al. Early predictors of mortality in
271 hemodynamically unstable pelvis fractures. *J Orthop Trauma*. 2007;21(1):31-7.
- 272 7. Starr AJ, Griffin DR, Reinert CM, Frawley WH, Walker J, Whitlock SN, et al. Pelvic Ring Disruptions: Prediction
273 of Associated Injuries, Transfusion Requirement, Pelvic Arteriography, Complications, and Mortality. *J*
274 *Orthop Trauma*. 2002;16(8):553–61.
- 275 8. Blackmore CC, Cummings P, Jurkovich GJ, Linnau KF, Hoffer EK, Rivara FP. Predicting Major Hemorrhage in
276 Patients with Pelvic Fracture. *J Trauma Acute Care Surg*. 2006;61(2):346–52.
- 277 9. Magnussen RA, Tressler MA, Obremesky WT, Kregor PJ. Predicting blood loss in isolated pelvic and
278 acetabular high-energy trauma. *J Orthop Trauma*. 2007;21(9):603–7.
- 279 10. Bramos A. Predictors of Bleeding From Stable Pelvic Fractures. *Arch Surg*. 2011;146(4):407.
- 280 11. White CE, Hsu JR, Holcomb JB. Hemodynamically unstable pelvic fractures. *Injury*. 2009;40(10):1023–30.
- 281 12. Manson T, O'Toole RV, Whitney A, Duggan B, Sciadini M, Nascone J. Young-Burgess classification of pelvic
282 ring fractures: does it predict mortality, transfusion requirements, and non-orthopaedic injuries? *J Orthop*
283 *Trauma*. 2010;24(10):603–9.
- 284 13. Rittmeister M, Lindsey RW, Kohl HW. Pelvic fracture among polytrauma decedents. Trauma-based mortality
285 with pelvic fracture--a case series of 74 patients. *Arch Orthop Trauma Surg*. 2001;121(1-2):43–9.
- 286 14. O'Sullivan REM, White TO, Keating JF. Major pelvic fractures: identification of patients at high risk.
287 *Bone Joint J*. 2005;87(4):530–3.
- 288 15. Baker SP, O'Neill B, Haddon W, Long WB. The injury severity score: a method for describing patients with
289 multiple injuries and evaluating emergency care. *J Trauma*. 1974;14(3):187–96.
- 290 16. Wang H, Phillips JL, Robinson RD, Duane TM, Buca S. Predictors of mortality among initially stable adult pelvic

- 291 trauma patients in the US: Data analysis from the National Trauma Data Bank. *Injury*. 2015;46(11):2113-7.
- 292 17. Costantini TW, Coimbra R, Holcomb JB. Current management of hemorrhage from severe pelvic fractures:
293 Results of an American Association for the Surgery of Trauma multi-institutional trial. *J Trauma Acute Care*
294 *Surg*. 2016;80(5):717-23.
- 295 18. Palmcrantz J, Hardcastle TC, Naidoo SR, Muckart DJJ, Ahlm K, Eriksson A. Pelvic fractures at a new level 1
296 trauma centre: who dies from pelvic trauma? *Orthop Surg*. 2012;29(4):216–21.
- 297 19. Halm EA, Wang JJ, Boockvar K, Penrod J, Silberzweig SB, Magaziner J, et al. The effect of perioperative anemia
298 on clinical and functional outcomes in patients with hip fracture. *J Orthop Trauma*. 2004;18(6):369–74.
- 299 20. Lanz OI. Lumbosacral and pelvic injuries. *Vet Clin North Am Small Anim Pract*. 2002;32(4):949–62.
- 300 21. Bookbinder PF, Flanders JA. Characteristics of pelvic fracture in the cat. A 10-year retrospective study. *VCOT*
301 *Archive*. 1992;3: 37-42.
- 302 22. Messmer M, Montavon PM. Pelvic fractures in the dog and cat: a classification system and review of 556
303 cases. *Vet Comp Orthop Traumatol*. 2004;17(4):167.
- 304 23. Maegele M, Paffrath T, Bouillon B. Acute traumatic coagulopathy in severe injury: incidence, risk
305 stratification, and treatment options. *Dtsch Arztebl Int*. 2011;108(49):827-35.
- 306 24. Stillion JR, Fletcher DJ. Admission base excess as a predictor of transfusion requirement and mortality in dogs
307 with blunt trauma: 52 cases (2007–2009). *J Vet Emerg Crit Care*. 2012;22(5):588–94.
- 308 25. Husain FA, Martin MJ, Mullenix PS, Steele SR, Elliott DC. Serum lactate and base deficit as predictors of
309 mortality and morbidity. *Am J Surg*. 2003;185(5):485-91.
- 310 26. Hoffberg JE, Koenigshof AM, Guiot LP. Retrospective evaluation of concurrent intra-abdominal injuries in
311 dogs with traumatic pelvic fractures: 83 cases (2008-2013). *J Vet Emerg Crit Care*. 2016;26(2):288–94.
- 312 27. Simpson SA, Syring R, Otto CM. Severe blunt trauma in dogs: 235 cases (1997-2003). *J Vet Emerg Crit Care*
313 2009 20;19(6):588–602.

314