



Міністерство освіти і науки України  
Харківський національний університет імені В. Н. Каразіна  
Медичний факультет

**XVI МІЖНАРОДНА НАУКОВА КОНФЕРЕНЦІЯ  
СТУДЕНТІВ, МОЛОДИХ ВЧЕНИХ ТА ФАХІВЦІВ**

# **АКТУАЛЬНІ ПИТАННЯ СУЧАСНОЇ МЕДИЦИНИ**

**Тези  
доповідей**

**28-29 березня  
2019 року**

до 25-річчя від дня відродження кафедр  
внутрішньої медицини, хірургічних хвороб,  
загальної та клінічної імунології  
та алергології медичного факультету  
Харківського національного університету  
імені В.Н. Каразіна



МІНІСТЕРСТВО ОСВІТИ І НАУКИ УКРАЇНИ  
MINISTRY OF EDUCATION AND SCIENCE OF UKRAINE

ХАРКІВСЬКИЙ НАЦІОНАЛЬНИЙ УНІВЕРСИТЕТ ІМЕНІ В.Н. КАРАЗИНА  
V.N. KARAZIN KHARKIV NATIONAL UNIVERSITY

XVI Міжнародна наукова конференція студентів,  
молодих вчених та фахівців

## **АКТУАЛЬНІ ПИТАННЯ СУЧАСНОЇ МЕДИЦИНИ**

ТЕЗИ ДОПОВІДЕЙ  
(28-29 березня 2019 року, м. Харків, Україна)

16<sup>th</sup> International Scientific Conference of Students, Young  
Scientists and Specialists

## **TOPICAL ISSUES OF MODERN MEDICINE**

ABSTRACTS  
(March 28-29 2018, Kharkiv, Ukraine)

ХАРКІВ – 2019

COMMUNITY-ACQUIRED PNEUMONIA ON THE BACKGROUND OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE: CLINICAL CASE	
Boakye – Yiadom Valentina.....	320
INFECTIVE ENDOCARDITIS IN INTRAVENOUS DRUG USER	
Bogun N. Yu., Brynza M. S., Khmara V. T., Babaeva E.Sh.....	321
DIAGNOSTICS ISSUES OF SYSTEMIC SCLEROSIS ON THE EXAMPLE OF CLINICAL CASE	
Brahmbatt Harshkumar Kirankumar.....	322
EFFECT OF THE SESSION ON THE PSYCHOEMOTION STATE OF KNMU STUDENTS	
Bulynina O., Senna D. ....	323
COMPARATIVE EVALUATION OF RESULTS OF SURGICAL TREATMENT OF ACUTE ADHESIVE INTESTINAL OBSTRUCTION	
Chubuk I. V., Voronova D. I. ....	324
POST-STROKE APHASIA: CURRENT INTERVENTION APPROACHES FOCUSING ON INDIVIDUAL'S PARTICIPATION	
Daher Q. ....	325
RELATIONSHIP BETWEEN REPEATED EXPOSURE TO TRAUMA IN WAR AND MENTAL HEALTH IN EASTERN MEDITERRANEAN REGION AND UKRAINE	
Daher Q. ....	326
CHANGES IN LIVER FUNCTION IN PATIENTS WITH JUVENILE IDIOPATHIC ARTHRITIS	
Dankwah Philip Sedem, Panko N. O. ....	327
THE EFFECT OF POOR SLEEP ON STUDENT HEALTH PREVENTION OF INSOMNIA	
Dankwah Philip Sedem.....	328
ASSESSMENT OF THE HEPATOPROTECTIVE AND ANTI-OXIDANT POTENTIALS OF THE TWIGS OF ANDROGRAPHIS PANICULATA IN CCL4- INDUCED LIVER DAMAGE IN WISTAR ALBINO RATS.	
David O. L. ....	330
THE EFFECT OF CHIKUNGUNYA VIRUS ON THE NERVOUS SYSTEM	
David O. L. ....	331
COMPARATIVE STUDY OF THE TREATMENT OF EXTERNAL GENITAL WARTS WITH MONOTHERAPY AND COMBINATION THERAPY	
David Prasad Catherine Preethi, Gradil O. G. ....	332
INFLUENCE OF IL-6 ON FORMATION PATHOLOGY OF MYOCARDIUM IN PATIENTS FOR TYPE 2 DIABETES MELLITUS	
Dehtiar K. O., Onikova A. O. ....	332
MODERN MEDICAL APPROACH TO MANAGEMENT OF VAGINISMUS	
Dlamini T. T. ....	333
EXERCISE TOLERANCE AND PHYSICAL ACTIVITY IN CHILDREN WITH CHRONIC HEALTH CONDITIONS	
Dsouza Allwin Jenifer ....	334
A CASE OF UNEXPLORED DYSPEPSIA	
Dwayat Yazan ....	335
THE DEVELOPMENT OF ATHEROSCLEROSIS IN A PATIENT WITH DIABETES MELLITUS	
Edema T. F. ....	336
THE RELATIONSHIP OF THE FUNCTIONAL ACTIVITY OF THE PITUITARY-THYROIDAL SYSTEM AND SEROTONIN PRODUCTION IN EARLY AGE CHILDREN	
Edema T. F., Itepu A. E. ....	337
THE COMBINED EFFECT OF SUCCINIC AND ASCORBIC ACIDS	

use the term “unexplored dyspepsia”. This primary syndromic diagnosis requires either empirical symptomatic or anti-*Helicobacter pylori* treatment, or further examination of the patient, especially esophagogastroduodenoscopy (EGDS) in order to establish organic or functional causes of dyspepsia.

**Clinical case.** The man, 49 years old, with complaints of constipation up to 5 days, pain in the upper abdomen, persistent, decreasing after eating, feeling of bloating in the epigastric region after eating, 12 kg weight loss since April 2018. Was repeatedly examined, took treatment without a pronounced effect. Anamnesis vitae is unremarkable. Physical examination: BMI = 24 kg / m<sup>2</sup>. The tongue is wet, at the root is coated with white bloom. On palpation, the abdomen is soft, painless. On auscultation, the abdominal intestinal peristalsis saved. Stool - a tendency to constipation, last stool 2 days ago with a laxative. The rest of physical examination is unremarkable. EGDS (02.01.19) shows: Chronic superficial antrum gastritis. Papular antral gastropathy 1 degree. Colonoscopy (28.08.18): dolichosigma, hypomotor syndrome. CT scan of the abdominal organs (30.10.18): pathological lesions in the projection of the stomach and intestinal loops are not defined. Clinical diagnosis: Unexplored dyspepsia dyspepsia: postprandial distress syndrome, syndrome of epigastric pain. Irritable bowel syndrome with constipation with colon dyskinesia of hypomotor type.

Pharmacological treatment: Omeprazole 20 mg twice a day, domperidone 10 mg thrice a day.

Taking into account the presence of “red flag” such as significant mass loss in association with non-conclusive results of EGDS we highly recommend patient to undergo EGDS with biopsy and rapid urease test to confirm the diagnosis of chronic gastritis and clarify its type in order to prescribe optimal therapy.

**Conclusion.** Dyspepsia is a common, long-recognized condition with a number of upper abdominal symptoms. But diagnosing this condition requires exclusion of organic diseases of digestive tract.

UDC 616.132.2-004.6:616.379-008.64

### **THE DEVELOPMENT OF ATHEROSCLEROSIS IN A PATIENT WITH DIABETES MELLITUS**

*Edema Temituoyo Folusho*

*V. N. Karazin Kharkiv National University, School of Medicine, Kharkiv, Ukraine*

*Department of Internal Medicine*

Scientific supervisors: Zhuravka N. V., assistant; Shop I. V., PhD, associate professor

**Introduction.** Numerous studies have shown that patients with diabetes mellitus have accelerated atherosclerotic vascular disease, and major advances in understanding its pathogenesis have made. Violation of glycemic homeostasis has a direct stimulating effect on the formation, increase in the number and size of atherosclerotic plaques. Factors that may play an important role in the more rapid progression of atherosclerosis in patients with diabetes mellitus: inflammation, neovascularization and hemorrhage inside plaques. As a result, patients with diabetes mellitus and coronary heart disease are subject to higher morbidity and mortality have compared with patients without diabetes.

**Clinical case.** The man, 55 years old, with complaints of attacks of dyspnea at night, stopped in a sitting position on their own, sometimes dyspnea attacks are accompanied by pain in the chest, transient rises in blood pressure, headaches. Anamnesis is remarkable significant for arterial hypertension (AH) (max 240/180 mm Hg, adapted to 130-140/80 mm Hg), he takes antihypertensive therapy, chest pains since 2012, In May 2015, during ECG holter, a pathological number of ventricular extrasystoles and paroxysms of unstable ventricular tachycardia were detected. In June 2015 - coronary angiography followed by stenting of PKA. In December 2018 due to the deterioration of the condition - re-coronary angiography followed by stenting of the PNA. In January 2019 the deterioration. Diabetes mellitus since 1998. Anamnesis vitae is unremarkable. Physical examination: BMI = 40 kg / m<sup>2</sup>. BP 140/80 mm Hg. Oedema of lower third of both shins. HR =80 bpm, heart sounds are muted. Pulsation on a. dorsalis pedis and a. tibialis posterior are reduced. The rest of physical examination is unremarkable. Complete blood count (17.01.19) shows erythrocytosis. Fasting blood glucose (17.01.19) - 17, 95 mmol/l. Echocardiography (17.01.19): atherosclerotic cardiosclerosis, aorta atherosclerosis, LV hypertrophy. Electrocardiography (ECG) (16.01.19): Atrial fibrillation with heart rate 75-112/ min. Hypertrophy of the myocardium of the left ventricle. Subendocardial ischemia.

**Clinical diagnosis:** Ischemic heart disease. Stable angina, II functional class. Arterial hypertension, II stage, 3 degree. CVD risk very high. Atherosclerosis of the coronary arteries. PKA stenting (22.05.15) PNA stenting (13.12.18) Ventricular extrasystolic arrhythmia with episodes of unstable ventricular tachycardia. Heart failure III FC, stage C. Diabetes mellitus II type, severe degree, decompensation. Diabetic polyneuropathy.

**Conclusion.** Atherosclerotic changes develop not only faster, but also earlier in patients with diabetes, while reducing their life expectancy. In patients with diabetes compared with those without this disease, even though a normal in cholesterol level of low-density lipoproteins in the blood achieved, poor glycemic control in patients with diabetes mellitus is associated with the progression of atherosclerotic changes.

UDC 616.8-008.6:[616.32+616.441-008.64]-053.4

## THE RELATIONSHIP OF THE FUNCTIONAL ACTIVITY OF THE PITUITARY-THYROIDAL SYSTEM AND SEROTONIN PRODUCTION IN EARLY AGE CHILDREN

*Edema T. F., Itepu A. E.*

*V. N. Karazin Kharkiv National University, School of Medicine, Kharkiv, Ukraine*

*Department of Pediatrics No. 2*

Scientific Supervisor: Shlieienkova H. O., PhD, assistant

**Introduction.** Serotonin plays a significant role in the regulation of the formation and functioning of the Central Nervous System. It is formed by conversion of the amino acid L - tryptophan. In the embryonic period, like thyroid hormones, it affects glia proliferation, differentiation of neurons, myelination of axons and accelerates the maturation of the nervous system. In the early neonatal period it affects branching of neurons. At any age serotonin acts as a neurotransmitter of synaptic transmission of nerve impulses.

**The aim of the study.** To study the functional activity of serotonin and pituitary-thyroid system and to evaluate their impact on mental development and neurological status of young children.