


LETTER

Open Access



Effect of high-flow nasal therapy on dyspnea, comfort, and respiratory rate

Andrea Cortegiani^{1*} , Claudia Crimi², Alberto Noto³, Yigal Helviz⁴, Antonino Giarratano¹, Cesare Gregoretti¹ and Sharon Einav⁴

Letter to the Editor

Systematic reviews comparing the effect of high-flow nasal treatment (HFNT) to conventional oxygen therapy (COT) or noninvasive ventilation (NIV) have focused on major clinical outcomes (i.e., endotracheal intubation, mortality) [1–3]. None have explored weaker outcomes that may nonetheless be important from the patient's perspective, yet physiopathological mechanisms suggest that the HFNT may provide some advantage in this regard [4, 5]. We therefore systematically reviewed all randomized (RCTs) and crossover trials enrolling patients either post-extubation or during acute respiratory failure (ARF), comparing HFNT to COT or NIV and reporting data about dyspnea, comfort, and respiratory rate (RR) (PROSPERO CRD42019119536).

Full search strategy, detailed study methods, reference lists, and risk of bias assessments are reported in Additional file 1.

Twenty-four relevant studies were identified and included: for patients post-extubation, ten RCTs and one crossover trial and, for patients in ARF, eight RCTs and five crossover trials.

The summary of our findings is presented in the Table 1. More studies compared the effects of HFNT vs COT rather than vs NIV. Overall, there seems to

be a trend showing that HFNT is probably not inferior to COT in most studies and perhaps better than NIV in terms of dyspnea, comfort, and decreasing of RR in some studies.

Heterogeneity in case-mix, the tools used for outcome assessment and measurement time-points precluded performance of meta-analysis. Neither patients nor treating clinicians were blinded to the intervention in any of the trials, introducing a high risk of detection bias. Differences in HFNT settings (i.e., flow and temperature) and a lack of full description for weaning criteria or protocol may have also contributed to the diversity in findings with regard to comfort and dyspnea.

In this analysis of the literature, the use of HFNT during ARF or post-extubation seems to be not clearly associated with improvements in comfort, dyspnea, and RR since findings from the most recent available evidence were inconsistent. However, in this regard, HFNT does not seem inferior to either COT or NIV. Future research should be focused in assessing patient-reported outcomes using appropriate standardized and validated measures in order to investigate the comparative effectiveness of the different respiratory support strategies.

* Correspondence: cortegiania@gmail.com

¹Department of Surgical, Oncological and Oral Science (Di.Chir.On.S.), Section of Anesthesia, Analgesia, Intensive Care and Emergency, Policlinico Paolo Giaccone, University of Palermo, Via del vespro 129, 90127 Palermo, Italy
Full list of author information is available at the end of the article



Table 1 Summary of findings in studies of the HFNT with regard to dyspnea, comfort, and respiratory rate

Study	Type	Design	Intervention (N)	Control (N)	Treatment methods	Measurement method	Dyspnea	Comfort	Respiratory rate
Bell N. [6] <i>Emerg Med Australas</i> 2015	AHRF	RCT	HFNT (48)	COT (52)	HFNT: flow 50 L/m, FIO2 30% titrated to SpO2 95% COT: discretion of the treating physician	Dyspnea: Borg Scale Comfort: Likert Scale	HFNT ^s	HFNT ^s (1 h)	HFNT ^s (2 h)
Frat J.P. [7] <i>N Engl J Med</i> 2015	AHRF	RCT	HFNT (106)	COT (94) NIV (110)	HFNT: flow 50 L/m, FIO2 100% then titrated to SpO2 92% COT: O2 titrated to SpO2 92% NIV: PSV PEEP from 2 up to 10 fIO2 adjusted to SpO2 92%	Dyspnea: Likert Scale Comfort: VAS	HFNT ^s	HFNT ^s	HFNT ^s (1 h)
Lemiale V. [8] <i>Crit Care</i> 2015	AHRF (Immunocompromised)	RCT	HFNT (52)	COT (52)	HFNT: flow from 40 up to 50 L/m, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: VAS Comfort: VAS	NS	NS	NS
Jones P.G. [9] <i>Respir Care</i> 2016	AHRF	RCT	HFNT (172)	COT (150)	HFNT: flow 40 L/m, 37 °C, FIO2 28% COT: FIO2 titrated to clinical needs	Dyspnea: Survey questions Comfort: Survey questions	NS	Overall comfort: NS "Dry my nose": HFNT ^s "In future I prefer": COT ^s "This method is worst": HFNT ^s	NS
Doshi P. [10] <i>Ann Emergency Med</i> 2017	AHRF	RCT	HFNT (104)	NIV (112)	HFNT: flow from 35 L/m up to 40 L/m, T ^m between 35 and 37 °C NIV: IPAP from 10 up to 20 cmH2O, EPAP from 5 up to 10 cmH2O, FIO2 100%	Dyspnea: Borg Scale Comfort: NA	NA	NA	NA
Makdee O. [11] <i>Ann Emergency Med</i> 2017	AHRF (CPE)	RCT	HFNT (63)	COT (65)	HFNT: flow from 35 up to 60 L/m, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: VAS Comfort: NA	NS	NA	HFNT ^s (15, 30, 60 min)
Azoulay E. [12] <i>JAMA</i> 2018	AHRF (Immunocompromised)	RCT	HFNT (388)	COT (388)	HFNT: flow 50 L/min, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: Dyspnea Score Comfort: VAS	NS	NS	HFNC ^s (6 h)
Spoletini G. [13] <i>J Crit Care</i> 2018	AHRF (On NIV)	RCT	HFNT (23)	COT (24)	HFNT: flow 35 L/m, FIO2 titrated to SpO2 92% (hypoxic) or to 88–92% (hypercapnic) COT: flow adjusted to maintain the same SpO2	Dyspnea: Borg Scale Comfort: VAS	NS	HFNT ^s	NS
Cuquemelle E. [14] <i>Respir Care</i> 2012	AHRF	Crossover	HFNT (37)	COT (37)	HFNT: flow 40 L/m, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: NA Comfort: Dyness	NA	HFNT ^s	NA
Schwabbauber N. [15] <i>BMC Anesthesiol</i> 2014	AHRF	Crossover	HFNT (14)	COT (14) NIV (14)	HFNT: flow 55 L/m, FIO2 60% COT: Venturi mask FIO2 60% NIV: PSV FIO2 60% PEEP 5 cmH2O PS 6–8 ml/kg PBW	Dyspnea: Borg Scale Comfort: NRS	HFNT vs. COT HFNT ^s vs. NIV vs. NIV	HFNT vs. COT HFNT ^s vs. NIV	HFNT vs. COT HFNT vs. NIV COT vs. NIV ^s
Vargas F. [16] <i>Respir Care</i> 2015	AHRF	Crossover	HFNT (n = 12)	COT (12) CPAP	HFNT: flow 60 L/m, T 37 °C, FIO2 same as COT COT: O2 titrated to SpO2 90%	Dyspnea: Dyspnea Score Comfort: NRS	NS	NS	HFNT ^s vs. COT HFNT vs. CPAP

Table 1 Summary of findings in studies of the HFNT with regard to dyspnea, comfort, and respiratory rate (Continued)

Study	Type	Design	Intervention (N)	Control (N)	Treatment methods	Measurement method	Dyspnea	Comfort	Respiratory rate
Mauri T. [17] <i>Am J Respir Crit Care Med</i> 2017	AHRF	Crossover	HFNT (15)	COT (15)	CPAP: 5 cmH2O FIO2 same as COT HFNT: flow 40 L/m, FIO2 titrated to SpO2 90–95% COT: Airvo2 face mask 12 L/min same FIO2	Dyspnea: DeltaPes Comfort: NA	HFNT ^s	NA	HFNT ^s
Sklar M.C. [18] <i>Ann Intensive Care</i> 2018	ARF (Exacerbation of cystic fibrosis)	Crossover	HFNT (15)	NIV (15)	HFNT: flow 55 L/m, T° 34 or 37 °C FIO2 titrated to SpO2 92% NIV: FIO2 titrated to SpO2 92%, setting as previously adjusted	Dyspnea: VAS Comfort: VAS	NS	NS	NS
Parke R. [19] <i>Br J Anaesth</i> 2013	Post-extubation (Cardiac surgery)	RCT	HFNT (169)	COT (171)	HFNT: flow 45 L/m, FIO2 titrated to SpO2 93% COT: O2 titrated to SpO2 93%	Dyspnea: NA Comfort: NRS	NA	HFNT ^s	NA
Maggiore S.M. [20] <i>Am J Respir Crit Care Med</i> 2014	Post-extubation	RCT	HFNT (53)	COT (52)	HFNT: flow 50 L/m, FIO2 titrated to SpO2 92–98% (hypoxic) or to 88–95% (hypercapnic) COT: O2 titrated to SpO2 92–98% (hypoxic) or 88–95% (hypercapnic)	Dyspnea: NA Comfort: NRS	NA	Interface: HFNT ^s (from 12 h) Dyyness: HFNT ^s (from 24 h)	HFNT ^s (from 1 h)
Conley A. [21] <i>Intensive Care Med</i> 2015	Post-extubation (Cardiac)	RCT	HFNT (81)	COT (74)	HFNT: flow 35 up to 50 L/min, T 37 °C, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: Borg Scale Comfort: NA	COT ^s (8 h)	NA	NS
Stephan F. [22] <i>JAMA</i> 2015	Post-extubation (Cardiac)	RCT	HFNT (414)	NIV (416)	HFNT: flow 50 L/m, FIO2 titrated to SpO2 92–98% NIV: PEEP and PS adjusted to RR < 25/min and TV 8 ml/kg, FIO2 SpO2 92–98%	Dyspnea: Borg Scale Comfort: NRS	NS	NS	HFNT ^s (1 h, 1 day, 2 days, 3 days)
Futier E. [23] <i>Intensive Care Med</i> 2016	Post-extubation (Abdominal or thoracic)	RCT	HFNT (108)	COT (112)	HFNT: flow 50–60 L/m, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: NA Comfort: NRS	NA	NS	NA
Hernandez G. (a) [24] <i>JAMA</i> 2016	Post-extubation (Low-risk extubation failure)	RCT	HFNT (264)	COT (263)	HFNT: flow 10 L/m titrated in 5 L step until discomfort, FIO2 to SpO2 92%, T 37 °C COT: O2 titrated to SpO2 92%	Dyspnea: NA Comfort: NA	NA	NA	NA
Hernandez G. (b) [25] <i>JAMA</i> 2016	Post-extubation (High-risk extubation failure)	RCT	HFNT (290)	NIV (314)	HFNT: flow 10 L/m titrated in 5 L step until discomfort, FIO2 to SpO2 92%, T 37 °C NIV: PEEP and PS adjusted to RR 25/min, SpO2 92%, pH 7.35	Dyspnea: NA Comfort: NA	NA	NA	NA
Fernandez R. [26] <i>Ann Intensive Care</i> 2017	Post-extubation (High-risk extubation failure)	RCT	HFNT (78)	COT (77)	HFNT: flow 40 L/min (adjusted on tolerance), T 37 or 34 °C, FIO2 titrated to SpO2 92–95% COT: O2 titrated to SpO2 92–95%	Dyspnea: NA Comfort: NA	NA	NA	NA
Yu Y. [27] <i>Can Respir J</i> 2017	Post-extubation (Thoracic)	RCT	HFNT (56)	COT (54)	HFNT: flow from 35 to 60 L/m, FIO2 titrated to SpO2 95% COT: O2 titrated to SpO2 95%	Dyspnea: NA Comfort: Rates of throat/nasal pain	NA	HFNT ^s	HFNT ^s (1 h, 2 h, 6 h, 24 h, 48 h, 72 h)
Song H.Z. [28] <i>Clinics (Sao</i>	post-extubation	RCT	HFNT (30)	COT (30)	HFNT: flow 60 L/m, FIO2 titrated to SpO2 94–98% (hypoxic) or to 88–92% (hypercapnic)	Dyspnea: NA Comfort: VAS	NA	HFNT ^s (interface) HFNT ^s (dryness)	HFNT ^s

Table 1 Summary of findings in studies of the HFNT with regard to dyspnea, comfort, and respiratory rate (Continued)

Study	Type	Design	Intervention (N)	Control (N)	Treatment methods	Measurement method	Dyspnea	Comfort	Respiratory rate
Paulo) 2017									
Rittayamai N. [29] <i>Respir Care</i> 2014	post-extubation	Crossover	HFNT (17)	COT (17)	COT: O2 titrated to SpO2 94–98% (hypoxic) or 88–92%(hypercapnic) HFNT: flow 35 L/m, FIO2 titrated to SpO2 94% COT: O2 titrated to SpO2 94%	Dyspnea: VAS Comfort: VAS	HFNT [§] (10, 15, 30 min)	NS	HFNT [§] (5, 10, 15, 30 min)

AHFR acute hypoxemic respiratory failure, ARF acute respiratory failure, CPAP continuous positive airway pressure, COT conventional oxygen therapy, HFNT high-flow nasal treatment, h hours, IPAP inspiratory positive airway pressure, N number of patients, NA not available, NIV noninvasive ventilation, NS not statistically significant, PES esophageal pressure, PSV pressure support ventilation, RCT randomized controlled trial, VAS visual analog scale

[§]Comparison between intervention and control with a statistically significant *p* value in favor of (the specified intervention)

Additional file

Additional file 1: List of included studies, search strategy, and risk of bias assessment. Detailed study methods, reference list of included studies, search strategy, risk of bias assessment. (DOCX 520 kb)

Abbreviations

ARF: Acute respiratory failure; COT: Conventional oxygen therapy; HFNT: High-flow nasal therapy; NIV: Noninvasive ventilation; RCT: Randomized controlled trial; RR: Respiratory rate

Acknowledgements

We would like to thank Dr. Filippo Sanfilippo for his help in registering the manuscript to PROSPERO.

Funding

None.

Availability of data and materials

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

Authors' contributions

AC, CC, AN, YE, AG, CG, and ES contributed substantially to the conception and design of the study, the acquisition of data, or the analysis and interpretation of the data. AC, CC, AN, YE, AG, CG, and ES drafted or provided critical revision of the article and approved the final version of the manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

Cesare Gregoretti received fees for consultancies or lectures from Orion Pharma, ResMed, Medtronic, Philips, Air Liquide and EOVE, and travel cost reimbursement from Fisher & Paykel. All other authors declared that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Surgical, Oncological and Oral Science (Di.Chir.On.S.), Section of Anesthesia, Analgesia, Intensive Care and Emergency, Policlinico Paolo Giaccone, University of Palermo, Via del vespro 129, 90127 Palermo, Italy. ²Respiratory Medicine Unit, A.O.U. "Policlinico-Vittorio Emanuele", Department of Clinical and Experimental Medicine, University of Catania, Via Santa Sofia 78, 95123 Catania, Italy. ³Department of Anesthesia and Critical Care, A.O.U. Policlinico "G. Martino", University of Messina, Via Consolare Valeria 1, 98100 Messina, Italy. ⁴Intensive Care Unit of the Shaare Zedek Medical Medical Centre and Hebrew University Faculty of Medicine, Jerusalem, Israel.

Received: 23 April 2019 Accepted: 13 May 2019

Published online: 05 June 2019

References

- Helviz Y, Einav S. A systematic review of the high-flow nasal cannula for adult patients. *Crit Care*. 2018;22:71.
- Cortegiani A, Crimi C, Sanfilippo F, Noto A, Di Falco D, Grasselli G, Gregoretti C, Giarratano A. High flow nasal therapy in immunocompromised patients with acute respiratory failure: a systematic review and meta-analysis. *J Crit Care*. 2019;50:250–6.
- Rochweg B, Granton D, Wang DX, Helviz Y, Einav S, Frat JP, Mekontso-Dessap A, Schreiber A, Azoulay E, Mercat A, Demoule A, Lemiale V, Pesenti A, Riviello ED, Mauri T, Mancebo J, Brochard L, Burns K. High flow nasal cannula compared with conventional oxygen therapy for acute hypoxemic respiratory failure: a systematic review and meta-analysis. *Intensive Care Med*. 2019. <https://doi.org/10.1007/s00134-019-05590-5>.
- Spoletini G, Cortegiani A, Gregoretti C. Physiopathological rationale of using high-flow nasal therapy in the acute and chronic setting: a narrative review. *Trends Anaesth Crit Care*. 2019. <https://doi.org/10.1016/j.tacc.2019.02.001>.
- Cortegiani A, Accurso G, Mercadante S, et al. High flow nasal therapy in perioperative medicine: from operating room to general ward. *BMC Anesthesiol*. 2018;18:166.
- Bell N, Hutchinson CL, Green TC, Rogan E, Bein KJ, Dinh MM. Randomised control trial of humidified high flow nasal cannulae versus standard oxygen in the emergency department. *Emerg Med Australas*. 2015;27:537–41.
- Frat JP, Thille AW, Mercat A, Girault C, Ragot S, Perbet S, Prat G, Boulain T, Morawiec E, Cottreau A, Devaquet J, Nseir S, Razazi K, Mira JP, Argaud L, Chakarian JC, Ricard JD, Wittebole X, Chevalier S, Herbland A, Fartoukh M, Constantin JM, Tonnelier JM, Pierrot M, Mathonnet A, Beduneau G, Deletage-Metreau C, Richard JC, Brochard L, Robert R, Group FS, Network R. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med*. 2015;372:2185–96.
- Lemiale V, Mokart D, Mayaux J, Lambert J, Rabbat A, Demoule A, Azoulay E. The effects of a 2-h trial of high-flow oxygen by nasal cannula versus Venturi mask in immunocompromised patients with hypoxemic acute respiratory failure: a multicenter randomized trial. *Crit Care*. 2015;19:380.
- Jones PG, Kamona S, Doran O, Sawtell F, Wilsher M. Randomized Controlled Trial of Humidified High-Flow Nasal Oxygen for Acute Respiratory Distress in the Emergency Department: The HOT-ER Study. *Respir Care*. 2016;61:291–9.
- Doshi P, Whittle JS, Bublewicz M, Kearney J, Ashe T, Graham R, Salazar S, Ellis TW Jr, Maynard D, Dennis R, Tillotson A, Hill M, Granado M, Gordon N, Dunlap C, Spivey S, Miller TL. High-Velocity Nasal Insufflation in the Treatment of Respiratory Failure: A Randomized Clinical Trial. *Ann Emerg Med*. 2018;72(73-83):e75.
- Makdee O, Monsomboon A, Surabenjawong U, Praphruetkit N, Chaisirin W, Chakorn T, Permpikul C, Thiravit P, Nakornchai T. High-Flow Nasal Cannula Versus Conventional Oxygen Therapy in Emergency Department Patients With Cardiogenic Pulmonary Edema: A Randomized Controlled Trial. *Ann Emerg Med*. 2017;70:465–472 e462.
- Azoulay E, Lemiale V, Mokart D, Nseir S, Argaud L, Pene F, Kontar L, Bruneel F, Klouche K, Barbier F, Reignier J, Berrahil-Meksen L, Louis G, Constantin JM, Mayaux J, Wallet F, Kouatchet A, Peigne V, Theodose I, Perez P, Girault C, Jaber S, Oziel J, Nyunga M, Terzi N, Bouadma L, Lebert C, Lautrette A, Bige N, Raphaelen JH, Papazian L, Darmon M, Chevret S, Demoule A. Effect of High-Flow Nasal Oxygen vs Standard Oxygen on 28-Day Mortality in Immunocompromised Patients With Acute Respiratory Failure: The HIGH Randomized Clinical Trial. *JAMA*. 2018;320:2099–107.
- Spoletini G, Mega C, Pisani L, Alotaibi M, Khoja A, Price LL, Blasi F, Nava S, Hill NS. High-flow nasal therapy vs standard oxygen during breaks off noninvasive ventilation for acute respiratory failure: A pilot randomized controlled trial. *J Crit Care*. 2018;48:418–25.
- Cuquemelle E, Pham T, Papon JF, Louis B, Danin PE, Brochard L. Heated and humidified high-flow oxygen therapy reduces discomfort during hypoxemic respiratory failure. *Respir Care*. 2012;57:1571–7.
- Schwabbauer N, Berg B, Blumenstock G, Haap M, Hetzel J, Riessen R. Nasal high-flow oxygen therapy in patients with hypoxic respiratory failure: effect on functional and subjective respiratory parameters compared to conventional oxygen therapy and non-invasive ventilation (NIV). *BMC Anesthesiol*. 2014;14:66.
- Vargas F, Saint-Leger M, Boyer A, Bui NH, Hilbert G. Physiologic Effects of High-Flow Nasal Cannula Oxygen in Critical Care Subjects. *Respir Care*. 2015; 60:1369–76.
- Mauri T, Turrini C, Eronia N, Grasselli G, Volta CA, Bellani G, Pesenti A. Physiologic Effects of High-Flow Nasal Cannula in Acute Hypoxemic Respiratory Failure. *Am J Respir Crit Care Med*. 2017;195:1207–15.
- Sklar MC, Dres M, Rittayamai N, West B, Grieco DL, Telias I, Junhasavasdikul D, Rauseo M, Pham T, Madotto F, Campbell C, Tullis E, Brochard L. High-flow nasal oxygen versus noninvasive ventilation in adult patients with cystic fibrosis: a randomized crossover physiological study. *Ann Intensive Care*. 2018;8:85.
- Parke R, McGuinness S, Dixon R, Jull A. Open-label, phase II study of routine high-flow nasal oxygen therapy in cardiac surgical patients. *Br J Anaesth*. 2013;111:925–31.
- Maggiore SM, Idone FA, Vaschetto R, Festa R, Cataldo A, Antonicelli F, Montini L, De Gaetano A, Navalesi P, Antonelli M. Nasal high-flow versus

- Venturi mask oxygen therapy after extubation. Effects on oxygenation, comfort, and clinical outcome. *Am J Respir Crit Care Med.* 2014;190:282–8.
21. Corley A, Bull T, Spooner AJ, Barnett AG, Fraser JF. Direct extubation onto high-flow nasal cannulae post-cardiac surgery versus standard treatment in patients with a BMI ≥ 30 : a randomised controlled trial. *Intensive Care Med.* 2015;41:887–94.
 22. Stephan F, Barrucand B, Petit P, Rezaiguia-Delclaux S, Medard A, Delannoy B, Cosserant B, Flicoteaux G, Imbert A, Pilorge C, Berard L, Bi POPSG. High-Flow Nasal Oxygen vs Noninvasive Positive Airway Pressure in Hypoxemic Patients After Cardiothoracic Surgery: A Randomized Clinical Trial. *JAMA.* 2015;313:2331–9.
 23. Futier E, Paugam-Burtz C, Godet T, Khoy-Ear L, Rozencwajg S, Delay JM, Verzilli D, Dupuis J, Chanques G, Bazin JE, Constantin JM, Pereira B, Jaber S, Os i. Effect of early postextubation high-flow nasal cannula vs conventional oxygen therapy on hypoxaemia in patients after major abdominal surgery: a French multicentre randomised controlled trial (OPERA). *Intensive Care Med.* 2016;42:1888–98.
 24. Hernandez G, Vaquero C, Colinas L, Cuenca R, Gonzalez P, Canabal A, Sanchez S, Rodriguez ML, Villasclaras A, Fernandez R. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Testo Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients: A Randomized Clinical Trial. *JAMA.* 2016a;316:1565–74.
 25. Hernandez G, Vaquero C, Gonzalez P, Subira C, Frutos-Vivar F, Rialp G, Laborda C, Colinas L, Cuenca R, Fernandez R. Effect of Postextubation High-Flow Nasal Cannula vs Conventional Oxygen Therapy on Reintubation in Low-Risk Patients: A Randomized Clinical Trial. *JAMA.* 2016b;315:1354–61.
 26. Fernandez R, Subira C, Frutos-Vivar F, Rialp G, Laborda C, Masclans JR, Lesmes A, Panadero L, Hernandez G. High-flow nasal cannula to prevent postextubation respiratory failure in high-risk non- hypercapnic patients: a randomized multicenter trial. *Ann Intensive Care.* 2017;7:47.
 27. Yu Y, Qian X, Liu C, Zhu C. Effect of High-Flow Nasal Cannula versus Conventional Oxygen Therapy for Patients with Thoracoscopic Lobectomy after Extubation. *Can Respir J.* 2017;2017:7894631.
 28. Song HZ, Gu JX, Xiu HQ, Cui W, Zhang GS. The value of high-flow nasal cannula oxygen therapy after extubation in patients with acute respiratory failure. *Clinics (Sao Paulo).* 2017;72:562–7.
 29. Rittayamai N, Tscheikuna J, Rujjwit P. High-flow nasal cannula versus conventional oxygen therapy after endotracheal extubation: a randomized crossover physiologic study. *Respir Care.* 2014;59:485–90.