


**KAZAN FEDERAL UNIVERSITY  
A. E. ARBUZOV INSTITUTE OF ORGANIC AND PHYSICAL  
CHEMISTRY  
RUSSIAN SCIENTIFIC FOUNDATION  
RUSSIAN FOUNDATION OF BASIC RESEARCH  
RUSSIAN ACADEMY OF SCIENCES  
FEDERAL AGENCY OF SCIENTIFIC ORGANIZATIONS  
D.I. MENDELEEV RUSSIAN CHEMICAL SOCIETY**



**VIII<sup>th</sup> INTERNATIONAL SYMPOSIUM  
«DESIGN AND SYNTHESIS OF SUPRAMOLECULAR  
ARCHITECTURES»**

**II<sup>nd</sup> YOUTH SCHOOL ON SUPRAMOLECULAR AND  
COORDINATION CHEMISTRY**

**April 25-29, 2016  
Kazan, Russia**

## **INTERNATIONAL ADVISORY COMMITTEE**

Prof. Alexander Konovalov - Chairman  
Prof. Vladimir Arslanov (Russia)  
Prof. Antony Coleman (France)  
Prof. Vladimir Fedin (Russia)  
Prof. Olga Fedorova (Russia)  
Prof. Yulia Gorbunova (Russia)  
Prof. Sergei Gromov (Russia)  
Prof. Sylvie Ferlay (France)  
Prof. Mir Wais Hosseini (France)  
Prof. Vitaly Kalchenko (Ukraine)  
Prof. Burkhard Koenig (Germany)  
Prof. Janusz Lipkowski (Poland)  
Prof. David Reinhoudt (Netherlands)  
Prof. Kinga Suwinska (Poland)

## **ORGANIZING COMMITTEE**

Prof. Igor Antipin - Chairman  
Prof. Svetlana Solovieva - Secretary  
Dr. Vladimir Burilov  
Dr. Yulia Elistratova  
Prof. Valery Gorbachuk  
Nataly Ivanova  
Dr. Alla Mirgorodskaya  
Dr. Diana Mironova  
Dr. Anton Muravev  
Prof. Asiya Mustafina  
Dr. Alexander Ovsyannikov  
Dr. Farida Valeeva  
Prof. Lucia Zakharova  
Dr. Albina Ziganshina  
Dr. Marat Ziganshin

## ELUCIDATION OF HYBRID MULTI-PORPHYRIN SUPRAMOLECULAR ASSEMBLIES BY 2D NMR SPECTROSCOPY

I. A. Khodov, N. Zh. Mamardashvili

*Institute of Solution Chemistry of Russian Academy of Science, Ivanovo, Russia*

*iakh@isc-ras.ru*

Two-dimensional nuclear magnetic resonance (NMR) spectroscopy studies of supramolecular assemblies were carried out. The chemical structure of supramolecular assemblies were confirmed by the two-dimensional NMR. It was found that two-dimensional rotating-frame Overhauser spectroscopy and double-quantum filtered correlation spectroscopy for this system are not sufficiently informative. Moreover, the explanation of experimental data is complicated by the fact that spectral signals of the supramolecular assemblies molecules at the same time depend on two factors: a variation of the electron density distribution and a change in the shielding upon complex formation. Thus questions about the chemical structure of the complexes formed and their size remains open. Therefore, to obtain true evidence of the fact of information about their size and complex formation, other way should be applied. We have used an approach to elucidate the chemical structures for complex that is based on the analysis of two-dimensional high-resolution diffusion-ordered NMR spectroscopy.

1. I.A. Khodov, G.A. Alper, G.M. Mamardashvili, N.Z. Mamardashvili . *J. Mols. Str.*, 2015, **1090**, 174-180.

**Acknowledgements** This work was supported by Russian Scientific Foundation, project № 14-13-00232.