

Kent Academic Repository

Full text document (pdf)

Citation for published version

Frain, Kelly May and van Dijl, Jan Maarten and Robinson, Colin (2019) The Twin-Arginine Pathway for Protein Secretion. *EcoSal Plus*, 8 (2). ISSN 2324-6200.

DOI

<https://doi.org/10.1128/ecosalplus.ESP-0040-2018>

Link to record in KAR

<https://kar.kent.ac.uk/74552/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

The Twin-arginine pathway for protein secretion

Kelly M. Frain¹, Jan Maarten van Dijl^{2*} and Colin Robinson¹

¹The School of Biosciences, University of Kent, Canterbury, United Kingdom

²University of Groningen, University Medical Center Groningen, Department of Medical Microbiology, Groningen, The Netherlands

*Correspondence to: J.M. van Dijl, University of Groningen (UMCG), Department of Medical Microbiology, Hanzeplein 1, P.O. Box 30001, 9700 RB Groningen, the Netherlands. E-mail: j.m.van.dijl01@umcg.nl

Running title: The bacterial Tat pathway

Summary

The Tat pathway for protein translocation across bacterial membranes stands out for its selective handling of fully folded cargo proteins. In this review, we provide a comprehensive summary of our current understanding of the different known Tat components, their assembly into different complexes and their specific roles in the protein translocation process. In particular, this overview focuses on the Gram-negative bacterium *Escherichia coli* and the Gram-positive bacterium *Bacillus subtilis*. Using these organisms, we discuss structural features of Tat complexes alongside mechanistic models that allow for the Tat pathway's unique protein proofreading and transport capabilities. Finally, we highlight recent advances in exploiting the Tat pathway for biotechnological benefit, the production of high value pharmaceutical proteins.

INTRODUCTION

About 20-30% of proteins synthesised in the bacterial cytoplasm are destined for extra-cytoplasmic locations (1). They pass the cytoplasmic membrane using specialised transport systems, involving gated pores, energy, and signal peptides to direct protein export. Two major protein export systems are known, namely the general Secretory (Sec) pathway and the Twin-Arginine Translocation (Tat) pathway (Figure 1). Most proteins use the Sec pathway common to all domains of life. The Tat pathway, the focus of this review, is more exclusive. For example, it has only ~30 native substrates in the Gram-negative bacterium *Escherichia coli*, and is not universally conserved (2).

Twin-arginine signal peptides

Specific N-terminal signal peptides direct proteins to the Sec or Tat pathways. Upon membrane translocation, the signal peptide is removed by signal peptidase to release the mature protein (3,4,5,6,7). Bacterial Sec and Tat signal peptides have a core tripartite structure: a positively charged N-terminal domain, a hydrophobic H-domain and a C-terminal domain (3,8). The signal peptidase cleavage site, usually Ala-x-Ala, resides in the C-domain (Figure 2). Tat signal peptides are defined by the twin-arginine motif S-R-R-x-F-L-K (x is any polar amino acid), joining the N- and H- domains (9,10). The RR-residues are essential for protein export (11,12) although an RR>KR substitution may be tolerated (12,13,14,15,16). Tat signal peptides are around 30 residues long, while Sec signal peptides range between 17-24 residues (17). To avoid Sec, Tat signal peptides are less hydrophobic than Sec signal peptides (18), and often contain basic residues as 'Sec avoidance motif' (19,20).

THE TAT PATHWAY

The Tat pathway was discovered in the 1990's (21,22). Three thylakoidal membrane proteins were identified as essential for protein translocation via Tat, namely Tha4, Hcf106 and cpTatC

(23,24,25,26). Homologues were found in bacteria, archaea and mitochondria (26,27). In *E. coli*, these were, respectively, named TatA, TatB and TatC (9,28,29). Their importance for protein export was highlighted by protein mislocalisation in mutant strains (27,28).

Studies on thylakoidal and bacterial Tat pathways showed that they transport complex fully-folded proteins requiring cofactor insertion or oligomerisation (30). Accordingly, Tat facilitates many processes, including cell division and cell envelope biogenesis in *E. coli* (31,32). Tat mutants of other organisms, e.g. *Agrobacterium* and *Salmonella*, are impaired in quorum sensing, motility and pathogenesis (33,34). Unlike Sec, Tat even exports complex proteins from eukaryotes, as first evidenced for tightly folded dihydrofolate reductase or GFP provided with Tat signal peptides (35,36,37,38,39).

Tat Genes

The *E. coli* Tat system is constitutively expressed from the *tatABCD* operon, where *tatA* is expressed ~25- and ~50-fold more than *tatB* and *tatC*, respectively (40). Elsewhere in the genome, *tatE* is expressed. Presumably, *tatB* and *tatE* originate from gene duplications of *tatA* (26,41). Δ *tatABCDE* strains are viable, but display pleiotropic phenotypes (42).

Tat systems minimally function with TatA-like and TatC-like proteins, although this varies between organisms (43). The Gram-positive bacterium *Bacillus subtilis* expresses two minimal Tat pathways from distinct *tatAdCd* and *tatAyCy* operons (5). *TatAdCd* is co-expressed with its only known substrate, PhoD, during phosphate-starvation (44,45). *TatAyCy* is expressed constitutively as is the case for its substrates EfeB (YwbN), QcrA and YkuE (5,46,47,48,49). The *tatAc* gene is constitutively expressed elsewhere in the genome (50).

***Escherichia coli* Tat Components**

TatA channels proteins across the inner membrane. Additionally, *E. coli* has two TatA-like proteins, TatB and TatE, involved in protein translocation (51). In contrast, *B. subtilis* has three TatA proteins, TatAd, TatAy and TatAc, but no TatB (5,45,52). TatA-like proteins can often be interchanged, functioning both intra- and inter-bacteria. In *E. coli*, TatE can substitute TatA (41), *B. subtilis* TatAd can substitute TatA and TatB (53), and TatAc forms active translocases with TatCd and TatCy (54) that functionally replace *E. coli* TatA and TatE (55). This functional overlap suggests a universal Tat translocation mechanism (53,56) (Table 1).

TatA-like proteins share structural features, including a short N-terminal domain exposed to periplasm or cell wall (57), a single transmembrane (TM) helix linked to an amphipathic helix (APH) positioned against the cytoplasmic membrane side (58), and an unstructured cytoplasmic tail (59,60,61,62)(Figure 3). Only few substitutions in TatA block its export function. In *E. coli* TatA, Gly33 in the “hinge region” connecting the TM helix and the APH is essential for export (63) as is Phe39 that anchors the APH to the membrane (64). Other APH residues and Gln8 in the TM helix are also imperative (65).

TatE shares 57% similarity with TatA (9). Despite its low abundance (40), TatE can substitute TatA (41), being a regular constituent of the Tat translocase (66). TatE interacts with Tat signal peptides, partially preventing premature signal peptide cleavage (67). Homologues function in many other bacteria as exemplified by *B. subtilis* TatAc, showing 45% amino acid sequence similarity to TatE (50).

TatB is functionally different from TatA. It binds Tat signal peptides and subsequent mature proteins. TatB’s APH and C-terminal region are longer than those of TatA (64,68,69)(Figure 3), and only mutations in “the hinge” and APH cause translocation defects (70). Single N-terminal amino acid substitutions in TatA allow complementation of Δ *tatB* strains (71,72). This bifunctionality of TatA mirrors the minimal TatAC systems in Gram-positive bacteria (5,73).

TatC is the largest Tat protein aiding substrate binding (74,75). It has 6 TM helices and an N-in C-in topology (Figure 3)(76). Crystallisation of *Aquifex aeolicus* TatC revealed that this protein resembles

a baseball glove or a cupped hand (77). TatC shows restricted structural flexibility, and is unlikely to undergo major conformational changes during translocation (78). A notable surface feature of TatC is the conserved Glu165 in *A. aeolicus* or Glu170 in *E. coli* (12,78). This residue is close to the binding pocket for Tat signal peptides, and its position in the membrane may perturb bilayer structure (79). Other important residues reside in TatC's cytoplasmic N-region and first cytoplasmic loop (75,80).

***Bacillus subtilis* Tat Components**

B. subtilis is the model organism for Gram-positive bacteria. Like all Gram-positive bacteria except *Actinomycetes* (81), *B. subtilis* has only TatA and TatC subunits (5,45). Accordingly, *B. subtilis* TatA is bifunctional, and able to functionally replace both both *E. coli* TatA and TatB.

TatAdCd exists as a complex of ~230 kDa, while TatAd alone assembles into complexes of ~270 kDa (53). TatAd has essentially the same structure as *E. coli* TatA (58,82). Mutations in the N-terminus of heterologously expressed TatAd block translocation in Δ *tatB* *E. coli* cells, implying their importance for TatAd's bi-functionality. Further, electron microscopy (EM) showed that TatAd complexes are too small and homogeneous to serve as pores for passage of differently-sized Tat substrates (44). TatCd is very similar to *E. coli* TatC with 6 TM helices, and it both binds and stabilises TatAd (83,84).

TatAyCy and TatAy form complexes of ~200 kDa (52). The conserved hinge region of TatAy is equally important as in TatAd and *E. coli* TatA (82,85). The Pro2 residue in the N-terminal extracytoplasmic region of TatAy is key for bifunctionality and Pro2 mutations interfere with TatAy's protein export function (50,82,85). The P2A mutation causes formation of fibril-like TatAy-TatCy assemblies of varying lengths, as shown by EM and atomic force microscopy (86). Compared to other TatC proteins, TatCy's C-terminus is extended by five amino acids that are important for EfeB export (87). Further, depending on the substrate, the N-terminus and the first cytoplasmic loop of TatC, TatCd and TatCy are important for translocation (88).

TatAc is dispensable for Tat function in *B. subtilis* (5). However, when expressed in *E. coli*, TatAc interacts with TatCd and TatCy facilitating export of Tat substrates, like AmiA, AmiC and TorA (54). While TatAc can substitute TatA and TatB in *E. coli* (55), it cannot replace TatAd or TatAy in *B. subtilis*. Nonetheless, TatAc enhances TatAyCy function in *Bacillus* (50).

Tat Complexes

The largest native Tat substrates are heterodimeric formate dehydrogenases of ~150 kDa (89). However, Tat components themselves are relatively small, so the translocase must coalesce with multiple copies of each individual subunit. Live cell imaging fluorescence microscopy showed the TatABC system assembling on demand to newly synthesised Tat substrates (90). Two major complexes exist: TatBC and TatA. TatBC forms a receptor that binds Tat substrates at the membrane. This triggers proton-motive force (PMF-)dependent TatA recruitment and oligomerisation to form an active translocase (91,92). Starting from substrate-bound TatBC, the translocation event is slow, taking 1-3 minutes (93). Disassembly of TatA from TatBC requires substrate export (94).

TatB and TatC interact in a 1:1 stoichiometry (95), and several TatBC complexes bind substrate(s) (96). Low resolution EM structures of TatBC revealed a hemispherical morphology, with an internal cavity that could accommodate the signal peptide (74). Seven copies of TatBC fit into the 11-17nm reconstruction, and apparently more than one substrate binds at once (97). Cross-linking showed TatC is the primary interaction site of the signal peptide (92,98,99,100), which inserts deeply into TatC by adopting a hairpin-like conformation (77,98). Subsequently, the signal peptide exposes its C-region at the extracytoplasmic membrane side to signal peptidase (101,102). The signal peptide's H-domain interacts with TatB's TM helix (92) and Tat precursors initially bind to TatB at multiple sites (103). NMR data indicate that TatB's extended C-terminus is flexible to facilitate substrate binding (68). Each TatC monomer has two TatB contact sites (75,102,103,104,105). Intriguingly, the TM helix of TatB seems close to where TatA initiates translocase oligomer assembly, suggesting TatB is a

regulatory surrogate of TatA (106).

Coevolution analysis predicts TatA and TatB binding along TM5/6 of TatC, indicating a cluster of polar amino acids on TatC forms contact with polar side chains of TatA or TatB (107,108). Cross-linking studies indicate that, upon substrate docking, TatA binding to the TM6 of TatC is reduced, while TatA and TatB binding at the polar site of TatC increased. As the polar cluster site is adjacent to the docking site of the signal peptide (66), this docking event could cause the conformational rearrangements. Signal peptide binding also alters TatC's resting state arrangement of head-to-tail to tail-to-tail. This opens the complex, allowing TatA access to the vacated polar cluster site and placing TatA adjacent to the concave face of TatC where TatA nucleation has been suggested (104).

TatA facilitates protein translocation, either by forming a pore (109) or by weakening the membrane (110). While TatA is not obligatory for TatBC formation or substrate docking (111), trace amounts are associated with TatBC, possibly representing nucleation points for TatA oligomers (112,113). TatA complexes in *E. coli* are in the 100-500 kDa size range (114), which can be resolved at 34 kDa intervals suggesting modular formation of three or four complexes at a time. Self-oligomerisation involves TatA's TMH (59,68,115).

MECHANISM OF TAT TRANSLOCATION

The mechanism of Tat translocation is still not fully resolved. The 'translocation pore model' theorises that TatA forms a channel/pore for protein passage (109). More recent data favor the 'membrane-destabilisation model' (110,116).

Low-resolution images of TatA revealed a pore-like complex of varying diameters (8.5–13 nm) that would accommodate varying substrate sizes (109,117). A lid-like feature was identified which presumably resides on the cis-side of the inner membrane, as if the transporter were a trap door (118). Thus, TatA's APH could form a hairpin to align with its TM helix, forming a pore for hydrophilic

cargo passage (119). Upon substrate docking onto TatBC, TatA protomers would assemble to suit the substrate size in an oligomeric ring format (120,121). Each TatA protein could be 'zipped together' by 7 salt bridges. Indeed, salt-dependence for the translocase was reported (85). However, this model is less attractive as APH insertion into the bilayer was not observed in other studies (57,122), while solution NMR indicated the APH fans outwards (59).

The second model proposes TatA complexes weaken/destabilise the membrane due to TatA's short TM domain (59,110), restricting membrane thickness to its own length. Destabilisation only happens when Tat substrate is bound. When no substrate is bound, TatA's APH immerses itself into the membrane, elongating the membrane. Upon substrate association, TatA re-orientates its APH outside of the membrane (122). The 'switch' from one state to another relates also to interactions between substrate and TatA's APH. TatA does not recognise the 'RR' motif, but its APH interacts with the C-domain of signal peptides (123,124), suggesting the mature protein also interacts with TatA (92,103,125).

Energy Requirements

The Tat system is powered by the PMF, consisting of a pH gradient (ΔpH) and the electric potential ($\Delta\psi$) across the membrane and Tat can use both (126,127,128,129,130). Accordingly, a proton- H^+ antiporter mechanism was suggested (131). It was estimated that a counter flow of 7.9×10^4 protons is needed per protein (132). The PMF is not required for protein targeting or binding to TatBC, but is necessary for TatA oligomerisation and the actual export process (92,112,133,134).

Tat Proofreading

Generally, Tat substrates only exit the cytosol once they are fully folded and contain their cofactor (138,139,140,141,142). The respective proofreading is extremely tight in *B. subtilis* (166,167), but

less stringent in the thylakoidal Tat system as unfolded proteins are also transported (35). Requirements for conformational stability were studied using non-native Tat substrates, like PhoA and antibody fragments. These are only exported in oxidising conditions where disulphide bonds form (38,148,149,152). Nevertheless, human growth hormone (hGH) and interferon α 2b are also exported via Tat without disulphide bonds formed (143), suggesting that they assume near-native states as shown for *E. coli* CueO, which is exported without its copper cofactor bound (144). Further, unstructured and hydrophilic polypeptides of 100-120 residues are tolerated by Tat, but a short hydrophobic stretch stops export (145,146,147). It thus seems that proofreading senses conformational flexibility. Mutations with less stringent proofreading were identified in *E. coli* TatABC (153), indicating that proofreading is undertaken by the Tat translocon. Proofreading may also involve chaperones known as Redox Enzyme Maturation Proteins (REMPs) (135). This is exemplified by TorD, facilitating Tat-dependent export of the trimethylamine-*N*-oxide (TMAO) reductase TorA in *E. coli* (136,137). Lastly, Tat may recruit quality control proteases to clear the translocase, as evidenced in *B. subtilis* for TatAyCy and the wall protease WprA (150,151).

FUTURE PERSPECTIVES FOR BIOTECHNOLOGY

Export of fully folded proteins is advantageous for biotechnological applications. Owing to their rapid growth, high yield, ease of scale-up and cost-effectiveness, ~30% of biotherapeutics are synthesised in bacteria (154). Preferably, these proteins are exported to the oxidising periplasm with an N-terminal signal peptide. This is the usual approach in *B. subtilis*, where most translocated proteins are secreted into the medium (155,156). In *E. coli*, export to the periplasm decreases proteolysis and eases downstream purification as the periplasm only contains 4% of the proteome (157,158). Here, proteins are released by osmotic shock (159). Recently it was shown that biotherapeutics can be exported via Tat (143,160). Moreover, Tat overexpression in so-called TatExpress cells resulted in 5-fold increased export of hGH (161). The 'gold standard' of bacterial

protein production, protein 'secretion' to the medium, was achieved when *B. subtilis* TatAdCd was expressed in $\Delta tatABCDE$ strains, as this led to protein release from the periplasm (162). Current efforts to apply Tat in biotechnology focus on different biotherapeutics and increased protein yields. It is anticipated that this area will expand rapidly once the mechanisms underlying Tat-dependent protein export are fully understood.

REFERENCES

1. **Holland IB.** 2004. Translocation of bacterial proteins - An overview. *Biochimica et Biophysica Acta - Molecular Cell Research* **1694**:5–16.
2. **Palmer T, Sargent F, Berks BC.** 2011. The Tat protein export pathway. In *EcoSal- Escherichia coli and Salmonella: Cellular and Molecular Biology*. Eds Böck, A Curtiss III, R Kaper, JB Karp, PD Neidhardt, FC Nyström T, Slauch JM, Squires CL, Ussery D, editor. ASM Press.
3. **von Heijne G.** 1990. The Signal peptide. *J Membr Biol* **115**:195–201.
4. **Lüke I, Handford JI, Palmer T, Sargent F.** 2009. Proteolytic processing of *Escherichia coli* twin-arginine signal peptides by LepB. *Arch Microbiol* **191**:919–25.
5. **Jongbloed JDH, Grieger U, Antelmann H, Hecker M, Nijland R, Bron S, Van Dijl JM.** 2004. Two minimal Tat translocases in *Bacillus*. *Mol Microbiol* **54**:1319–25.
6. **Dalbey RE, Wang P, van Dijl JM.** 2012. Membrane proteases in the bacterial secretion and quality control pathway. *Microbiol Mol Biol Rev* **76**:311-30.
7. **Sakaguchi M, Tomiyoshi R, Kuroiwa T, Mihara K, Omura T.** 1992. Functions of signal and signal-anchor sequences are determined by the balance between the hydrophobic segment and the N-terminal charge. *Proc Natl Acad Sci* **89**:16–9.
8. **Berks BC.** 1996. A common export pathway for proteins binding complex redox cofactors? *Mol Microbiol* **22**: 393–404.

9. **Sargent F, Bogsch EG, Stanley NR, Wexler M, Robinson C, Berks BC, Palmer T.** 1998. Overlapping functions of components of a bacterial Sec-independent protein export pathway. *EMBO J* **17**:3640–50.
10. **Berks BC, Palmer T, Sargent F.** 2003. The Tat protein translocation pathway and its role in microbial physiology. *Advances in Microbial Physiology* **47**: 187–254.
11. **Stanley NR, Palmer T, Berks BC.** 2000. The twin arginine consensus motif of Tat signal peptides is involved in Sec-independent protein targeting in *Escherichia coli*. *J Biol Chem* **275**: 11591–6.
12. **Buchanan G, Sargent F, Berks BC, Palmer T.** 2002. A genetic screen for suppressors of *Escherichia coli* Tat signal peptide mutations establishes a critical role for the second arginine within the twin-arginine motif. *Arch Microbiol* **177**:107–12.
13. **Chaddock AM, Mant a, Karnauchov I, Brink S, Herrmann RG, Klösgen RB, Robinson C.** 1995. A new type of signal peptide: central role of a twin-arginine motif in transfer signals for the delta pH-dependent thylakoidal protein translocase. *EMBO J* **14**:2715–22.
14. **Halbig D, Hou B, Freudl R, Sprenger GA, Klösgen RB.** 1999. Bacterial proteins carrying twin-R signal peptides are specifically targeted by the Δ pH-dependent transport machinery of the thylakoid membrane system. *FEBS Lett* **447**:95–8.
15. **Hinsley AP, Stanley NR, Palmer T, Berks BC.** 2001. A naturally occurring bacterial Tat signal peptide lacking one of the “invariant” arginine residues of the consensus targeting motif. *FEBS Lett* **497**:45–9.
16. **DeLisa MP, Samuelson P, Palmer T, Georgiou G.** 2002. Genetic analysis of the twin arginine translocator secretion pathway in bacteria. *J Biol Chem* **277**:29825–31.
17. **Tullman-Ercek D, Delisa MP, Kawarasaki Y, Iranpour P, Ribnicky B, Palmer T, Georgiou G.** 2007. Export pathway selectivity of *Escherichia coli* twin arginine translocation signal peptides. *J Biol Chem* **282**:8309-8316.
18. **Cristóbal S, De Gier JW, Nielsen H, Von Heijne G.** 1999. Competition between Sec- and TAT-dependent protein translocation in *Escherichia coli*. *EMBO J* **18**:2982–90.

19. **Bogsch E, Brink S, Robinson C.** 1997. Pathway specificity for a Δ pH-dependent precursor thylakoid lumen protein is governed by a "Sec-avoidance" motif in the transfer peptide and a "Sec-incompatible" mature protein. *EMBO J* **16**:3851–9.
20. **Mould RM, Robinson C.** 1991. A proton gradient is required for the transport of two luminal oxygen-evolving proteins across the thylakoid membrane. *J Biol Chem* **266**:12189–93.
21. **Klöggen RB, Brock IW, Herrmann RG, Robinson C.** 1992. Proton gradient-driven import of the 16 kDa oxygen-evolving complex protein as the full precursor protein by isolated thylakoids. *Plant Mol Biol* **18**:1031–4.
22. **Clark SA, Theg SM.** 1997. A folded protein can be transported across the chloroplast envelope and thylakoid membranes. *Mol Biol Cell* **8**:923–34.
23. **Mori H, Summer EJ, Ma X, Cline K.** 1999. Component specificity for the thylakoidal Sec and Δ pH-dependent protein transport pathways. *J Cell Biol* **146**:45–55.
24. **Settles AM, Yonetani A, Baron A, Bush DR, Cline K, Martienssen R.** 1997. Sec-independent protein translocation by the maize Hcf106 protein. *Science* **278**:1467–70.
25. **Cline K, Mori H.** 2001. Thylakoid Δ pH-dependent precursor proteins bind to a cpTatC-Hcf106 complex before Tha4-dependent transport. *J Cell Biol* **154**:719–29.
26. **Wu LF, Ize B, Chanal A, Quentin Y, Fichant G.** 2000. Bacterial twin-arginine signal peptide-dependent protein translocation pathway: evolution and mechanism. *J Mol Microbiol Biotechnol* **2**:179–89.
27. **Bogsch EG, Sargent F, Stanley NR, Berks BC, Robinson C, Palmer T.** 1998. An essential component of a novel bacterial protein export system with homologues in plastids and mitochondria. *J Biol Chem* **273**:18003–6.
28. **Sargent F, Stanley NR, Berks BC, Palmer T.** 1999. Sec-independent protein translocation in *Escherichia coli*. A distinct and pivotal role for the TatB protein. *J Biol Chem* **274**:36073–82.
29. **Palmer T, Sargent F, Berks BC.** 2005. Export of complex cofactor-containing proteins by the bacterial Tat pathway. *Trends Microbiol* **13**:175–80.

30. **Rodrigue A, Chanal A, Beck, B, Muller M, Wu LF.** 1999. Co-translocation of a periplasmic enzyme complex by a hitchhiker mechanism through the bacterial tat pathway. *J Biol Chem* **274**: 13228-8.
31. **Bernhardt TG, De Boer PAJ.** 2003. The *Escherichia coli* amidase AmiC is a periplasmic septal ring component exported via the twin-arginine transport pathway. *Mol Microbiol* **48**:1171–82.
32. **Ize B, Stanley NR, Buchanan G, Palmer T.**2003. Role of the *Escherichia coli* Tat pathway in outer membrane integrity. *Mol Microbiol* **48**:1183–93.
33. **Ding Z, Christie PJ.** 2003. *Agrobacterium tumefaciens* twin-arginine-dependent translocation is important for virulence, flagellation, and chemotaxis but not type IV secretion. *J Bacteriol* **185**:760–771.
34. **Craig M, Sadik AY, Golubeva YA, Tidhar A, Slauch JM.** 2013. Twin-arginine Translocation System (*tat*) Mutants of *Salmonella* are Attenuated Due to Envelope Defects, not Respiratory Defects. *Mol Microbiol* **89**:887-902.
35. **Hynds PJ, Robinson D, Robinson C.** 1998. The Sec-independent twin-arginine translocation system can transport both tightly folded and malfolded proteins across the thylakoid membrane. *J Biol Chem* **273**:34868–74.
36. **Santini CL, Bernadac A, Zhang M, Chanal A, Ize B, Blanco C, Wu, LF.** 2001. Translocation of Jellyfish Green Fluorescent Protein via the Tat System of *Escherichia coli* and Change of Its Periplasmic Localization in Response to Osmotic Up-shock. *J Biol Chem* **276**:8159–64.
37. **Alanen HI, Walker KL, Lourdes Velez Suberbie, M, Matos, CFRO, Bönisch S, Freedman RB, Keshavarz-Moore E, Ruddock L, Robinson, C.** 2014. Efficient export of human growth hormone, interferon α 2b and antibody fragments to the periplasm by the *Escherichia coli* Tat pathway in the absence of prior disulfide bond formation. *Biochimica et Biophysica Acta - Molecular Cell Research* **1853**:756–763.
38. **Delisa MP, Tullman D, Georgiou G.** 2003. Folding quality control in the export of proteins by the bacterial twin-arginine translocation pathway. *PNAS* **100**:6115–6120.

39. **Richter S, Brüser T.** 2005. Targeting of unfolded PhoA to the TAT translocon of *Escherichia coli*. *J Biol Chem* **280**:42723–30.
40. **Jack RL, Sargent F, Berks BC, Sawers G, Palmer T.** 2001. Constitutive expression of *Escherichia coli* tat genes indicates an important role for the twin-arginine translocase during aerobic and anaerobic growth. *J Bacteriol* **183**:1801–4.
41. **Baglieri J, Beck D, Vasisht N, Smith CJ, Robinson C.** 2012. Structure of TatA paralog, TatE, suggests a structurally homogeneous form of Tat protein translocase that transports folded proteins of differing diameter. *J Biol Chem* **287**:7335–7344.
42. **Stanley NR, Findlay K, Berks BC, Palmer T.** 2001. *Escherichia coli* strains blocked in Tat-dependent protein export exhibit pleiotropic defects in the cell envelope. *J Bacteriol* **183**:139–44.
43. **Goosens VJ, Monteferrante CG, Van Dijl JM.** 2014. The Tat system of Gram-positive bacteria. *Biochimica et Biophysica Acta - Molecular Cell Research* **1843**:1698–706.
44. **Beck D, Vasisht N, Baglieri J, Monteferrante CCG, van Dijl JM, Robinson C, Smith C.** 2009. Subcellular localization of TatAd of *Bacillus subtilis* depends on the presence of TatCd or TatCy. *J Biol Chem* **191**:4410–8.
45. **Jongbloed JDH, Martin U, Antelmann H, Hecker M, Tjalsma H, Venema G, Bron S, van Dijl JM, Muller J.** 2000. TatC is a specificity determinant for protein secretion via the twin-arginine translocation pathway. *J Biol Chem* **275**:41350–7.
46. **van der Ploeg R, Mäder U, Homuth G, Schaffer M, Denham EL, Monteferrante CG, Miethke MA, Harwood CR, Winter T, Hecker M, Antelmann H, van Dijl JM.** 2011. Environmental salinity determines the specificity and need for tat-dependent secretion of the YwbN protein in *Bacillus subtilis*. *PLoS One* **6**: 18140.
47. **Monteferrante CG, Miethke M, Van Der Ploeg R, Glasner C, Van Dijl JM.** 2012. Specific targeting of the metallophosphoesterase YkuE to the *Bacillus* cell wall requires the twin-arginine translocation system. *J Biol Chem* **287**:29789–800.

48. **Miethke M, Monteferrante CG, Marahiel MA, van Dijk JM.** 2013. The *Bacillus subtilis* EfeUOB transporter is essential for high-affinity acquisition of ferrous and ferric iron. *Biochim Biophys Acta - Mol Cell Res* **1833**: 2267-78.
49. **Goosens VJ, Otto A, Glasner C, Monteferrante CC, Van Der Ploeg R, Hecker M, Becher D, van Dijk JM.** 2013. Novel twin-arginine translocation pathway-dependent phenotypes of *Bacillus subtilis* unveiled by quantitative proteomics. *J Proteome Res* **12**:796-807.
50. **Goosens VJ, De-San-Eustaquio-Campillo A, Carballido-López R, van Dijk JM.** 2015. A Tat ménage à trois - The role of *Bacillus subtilis* TatAc in twin-arginine protein translocation. *Biochim Biophys Acta - Mol Cell Res* **1853**:2745-53.
51. **Sargent F, Gohlke U, De Leeuw E, Stanley NR, Palmer T, Saibil HR, Berks BC.** 2001. Purified components of the *Escherichia coli* Tat protein transport system form a double-layered ring structure. *Eur J Biochem* **268**:3361-7.
52. **Goosens VJ, Monteferrante CG, Van Dijk JM.** 2014. The Tat system of Gram-positive bacteria. *Biochimica et Biophysica Acta - Molecular Cell Research* **1843**:1698-706.
53. **Barnett JP, Eijlander RT, Kuipers OP, Robinson C.** 2008. A minimal tat system from a gram-positive organism: A bifunctional TatA subunit participates in discrete TatAC and TatA complexes. *J Biol Chem* **283**:2534-42.
54. **Monteferrante CG, Baglieri J, Robinson C, van Dijk JM.** 2012. TatAc, the third TatA subunit of *Bacillus subtilis*, can form active twin-arginine translocases with the TatCd and TatCy subunits. *Appl Environ Microbiol* **78**:4999-5001.
55. **Beck D, Vasisht N, Baglieri J, Monteferrante CG, van Dijk JM, Robinson C, Smith CJ.** 2013. Ultrastructural characterisation of *Bacillus subtilis* TatA complexes suggests they are too small to form homooligomeric translocation pores. *Biochim Biophys Acta - Mol Cell Res* **1833**:1811-9.
56. **Porcelli I, De Leeuw E, Wallis R, Van den Brink-van der Laan E, De Kruijff B, Wallace BA, Palmer T, Berks, BC.** 2002. Characterization and membrane assembly of the TatA component of the *Escherichia coli* twin-arginine protein transport system. *Biochemistry* **41**:13690-7.

57. Koch S, Fritsch MJ, Buchanan G, Palmer T. 2012. Escherichia coli TatA and TatB proteins have N-out, C-in topology in intact cells. *J Biol Chem* **287**:4420–31.
58. Hu Y, Zhao E, Li H, Xia B, Jin C. 2010. Solution NMR structure of the TatA component of the twin-arginine protein transport system from gram-positive bacterium *Bacillus subtilis*. *J Am Chem Soc* **132**:15942–4.
59. Rodriguez F, Rouse SL, Tait CE, Harmer J, De Riso A, Timmel CR, Sansom M, Berks BC, Schnell JR. 2013. Structural model for the protein-translocating element of the twin-arginine transport system. *Proc Natl Acad Sci U S A* **110**:1092-101.
60. Müller SD, De Angelis AA, Walther TH, Grage SL, Lange C, Opella SJ, Ulrich AS. 2007. Structural characterization of the pore forming protein TatAd of the twin-arginine translocase in membranes by solid-state ¹⁵N-NMR. *Biochim Biophys Acta - Biomembr* **1768**:3071–9.
61. Walther TH, Grage SL, Roth N, Ulrich AS. 2010. Membrane alignment of the pore-forming component TatAd of the twin-arginine translocase from *Bacillus subtilis* resolved by solid-state NMR spectroscopy. *J Am Chem Soc* **132**:15945–56.
62. Lange C, Müller SD, Walther TH, Bürck J, Ulrich AS. 2007. Structure analysis of the protein translocating channel TatA in membranes using a multi-construct approach. *Biochim Biophys Acta - Biomembr* **1768**:2627–34.
63. Barrett CML, Ray N, Thomas JD, Robinson C, Bolhuis A. 2003. Quantitative export of a reporter protein, GFP, by the twin-arginine translocation pathway in *Escherichia coli*. *Biochem Biophys Res Commun* **304**:279–84.
64. Hicks MG, De Leeuw E, Porcelli I, Buchanan G, Berks BC, Palmer T. 2003. The *Escherichia coli* twin-arginine translocase: Conserved residues of TatA and TatB family components involved in protein transport. *FEBS Lett* **539**:61–7.
65. Greene NP, Porcelli I, Buchanan G, Hicks MG, Schermann SM, Palmer T, Berks BC. 2007. Cysteine scanning mutagenesis and disulfide mapping studies of the TatA component of the bacterial twin arginine translocase. *J Biol Chem* **282**:23937–45.

66. Eimer E, Fröbel J, Blümmel AS, Müller M. 2015. TatE as a regular constituent of bacterial twin-arginine protein translocases. *J Biol Chem* **290**:29281–9.
67. Eimer E, Kao WC, Fröbel J, Blümmel AS, Hunte C, Müller M. 2018. Unanticipated functional diversity among the TatA-type components of the Tat protein translocase. *Sci Rep* **8**:1–12.
68. Zhang Y, Wang L, Hu Y, Jin C. 2014. Solution structure of the TatB component of the twin-arginine translocation system. *Biochim Biophys Acta - Biomembr* **1838**:1881–8.
69. Lee PA, Buchanan G, Stanley NR, Berks BC, Palmer T. 2002. Truncation analysis of TatA and TatB defines the minimal functional units required for protein translocation. *J Bacteriol* **184**:5871–9.
70. Barrett CML, Mathers JE, Robinson C. 2003. Identification of key regions within the *Escherichia coli* TatAB subunits. *FEBS Lett* **537**:42–6.
71. Blaudeck N, Kreutzenbeck P, Müller M, Sprenger GA, Freudl R. 2005. Isolation and characterization of bifunctional *Escherichia coli* TatA mutant proteins that allow efficient Tat-dependent protein translocation in the absence of TatB. *J Biol Chem* **280**:3426–32.
72. Barrett CML, Freudl R, Robinson C. 2007. Twin arginine translocation (Tat)-dependent export in the apparent absence of TatABC or TatA complexes using modified *Escherichia coli* TatA subunits that substitute for TatB. *J Biol Chem* **282**: 36206–13.
73. Jongbloed JDH, Van Der Ploeg R, Van Dijk JM. 2006. Bifunctional TatA subunits in minimal Tat protein translocases. *Trends in Microbiol* **14**:2-4.
74. Tarry MJ, Schäfer E, Chen S, Buchanan G, Greene NP, Lea SM, Berks, BC. 2009. Structural analysis of substrate binding by the TatBC component of the twin-arginine protein transport system. *Proc Natl Acad Sci U S A* **106**:13284–9.
75. Kneuper H, Maldonado B, Jäger F, Krehenbrink M, Buchanan G, Keller R, Muller M, Berks BC, Palmer, T. 2012. Molecular dissection of TatC defines critical regions essential for protein transport and a TatB-TatC contact site. *Mol Microbiol* **85**:945–61.
76. Behrendt J, Standar K, Lindenstrauß U, Brüser T. 2004. Topological studies on the twin-arginine translocase component TatC. *FEMS Microbiol Lett* **234**: 303–8.

77. **Ramasamy S, Abrol R, Suloway CJM, Clemons WM.** 2013. The glove-like structure of the conserved membrane protein tatC provides insight into signal sequence recognition in twin-arginine translocation. *Structure* **21(5):**777–88.
78. **Rollauer SE, Tarry MJ, Graham JE, Jääskeläinen M, Marcoux J, Mcdowell MA, Rodriguez F, Roversi P, Stansfeld PJ, Robinson, CV, Sansom, M., Palmer T, Högbom M, Berks BC, Lea SM.** 2013. Structure of the TatC core of the twin-arginine protein transport system. *Nature* **492:**210–4.
79. **Blümmel AS, Drepper F, Knapp B, Eimer E, Warscheid B, Müller M, Frobel J.** 2017. Structural features of the TatC membrane protein that determine docking and insertion of a twin-arginine signal peptide. *J Biol Chem* **292:**21320–9.
80. **Holzappel E, Eisner G, Alami M, Barrett CML, Buchanan G, Lüke I, Betton JM, Robinson C, Palmer T, Monser M, Muller M.** 2007. The entire N-terminal half of TatC is involved in twin-arginine precursor binding. *Biochemistry* **46:**2892–8.
81. **Schaerlaekens K, Schierová M, Lammertyn E, Geukens N, Anné J, Van Mellaert L.** 2001. Twin-arginine translocation pathway in *Streptomyces lividans*. *J Bacteriol* **183:**6727–32.
82. **Barnett JP, Lawrence J, Mendel S, Robinson C.** 2011. Expression of the bifunctional *Bacillus subtilis* TatAd protein in *Escherichia coli* reveals distinct TatA/B-family and TatB-specific domains. *Arch Microbiol* **193:**583–94.
83. **Schreiber S, Stengel R, Westermann M, Volkmer-Engert R, Pop OI, Müller JP.** 2006. Affinity of TatCd for TatAdelucidates its receptor function in the *Bacillus subtilis* twin arginine translocation (Tat) translocase system. *J Biol Chem* **281:** 19977–84.
84. **Nolandt O V., Walther TH, Roth S, Bürck J, Ulrich AS.** 2009. Structure analysis of the membrane protein TatCd from the Tat system of *B. subtilis* by circular dichroism. *Biochim Biophys Acta - Biomembr* **1788:**2238–44.
85. **Van Der Ploeg R, Barnett JP, Vasisht N, Goosens VJ, Pöther DC, Robinson C, van Dijk JM.** 2011. Salt sensitivity of minimal twin arginine translocases. *J Biol Chem* **286:**43759–70.

86. **Patel R, Vasilev C, Beck D, Monteferrante CG, Van Diji JM, Hunter CN, Smith C, Robinson C.** 2014. A mutation leading to super-assembly of twin-arginine translocase (Tat) protein complexes. *Biochim Biophys Acta - Mol Cell Res* **1843**:1978–86.
87. **Eijlander RT, Jongbloed JDH, Kuipers OP.** 2009. Relaxed specificity of the *Bacillus subtilis* TatAdCd translocase in Tat-dependent protein secretion. *J Bacteriol* **91**:196–202.
88. **Eijlander RT, Kolbusz MA, Berendsen EM, Kuipers OP.** 2009. Effects of altered TatC proteins on protein secretion efficiency via the twin-arginine translocation pathway of *Bacillus subtilis*. *Microbiology* **155**:1776–85.
89. **Berks BC, Sargent F, Palmer T.** 2000. The Tat protein export pathway. *Mol Microbiol* **35**:260–274.
90. **Rose P, Fröbel J, Graumann PL, Müller M.** 2013. Substrate-Dependent Assembly of the Tat Translocase as Observed in Live *Escherichia coli* Cells. *PLoS One* **8**:1–17.
91. **Berks BC, Lea SM, Stansfeld PJ.** 2014. Structural biology of tat protein transport. *Curr Opin Struct Biol* **27**:32–7.
92. **Alami M, Lüke I, Deitermann S, Eisner G, Koch HG, Brunner J, Müller, M.** 2003. Differential interactions between a twin-arginine signal peptide and its translocase in *Escherichia coli*. *Molecular Cell* **12**:937–946.
93. **Celedon JM, Cline K.** 2012. Stoichiometry for binding and transport by the twin Arginine translocation system. *J Cell Biol* **197**:523–34.
94. **Alcock F, Baker MAB, Greene NP, Palmer T, Wallace MI, Berks BC.** 2013 Live cell imaging shows reversible assembly of the TatA component of the twin-arginine protein transport system. *Proc Natl Acad Sci* **110**:E3650–9.
95. **Bolhuis A, Mathers JE, Thomas JD, Claire M, Barrett L, Robinson C.** 2001. TatB and TatC Form a Functional and Structural Unit of the Twin-arginine Translocase from *Escherichia coli*. *J Biol Chem* **276**:20213–9.
96. **Behrendt J, Brüser T.** 2014. The TatBC complex of the tat protein translocase in *Escherichia coli* and its transition to the substrate-bound TatABC complex. *Biochemistry* **53**:2344–54.

97. **Ma X, Cline K.** 2010. Multiple precursor proteins bind individual Tat receptor complexes and are collectively transported. *EMBO J* **29**:1477–88.
98. **Gérard F, Cline K.** 2006. Efficient twin arginine translocation (Tat) pathway transport of a precursor protein covalently anchored to its initial cpTatC binding site. *J Biol Chem* **281**:6130–5.
99. **Kreutzenbeck P, Kröger C, Lausberg F, Blaudeck N, Sprenger GA, Freudl R.** 2007. *Escherichia coli* twin arginine (Tat) mutant translocases possessing relaxed signal peptide recognition specificities. *J Biol Chem* **282**:7903–11.
100. **Zoufaly S, Fröbel J, Rose P, Flecken T, Maurer C, Moser M, Müller M.** 2012. Mapping precursor-binding site on TatC subunit of twin arginine-specific protein translocase by site-specific photo cross-linking. *J Biol Chem* **287**:13430–41.
101. **Fröbel J, Rose P, Lausberg F, Blümmel A-S, Freudl R, Müller M.** 2012. Transmembrane insertion of twin-arginine signal peptides is driven by TatC and regulated by TatB. *Nat Commun* **3**:1311.
102. **Blümmel AS, Haag LA, Eimer E, Müller M, Fröbel J.** 2015. Initial assembly steps of a translocase for folded proteins. *Nat Commun* **6**:7234.
103. **Maurer C, Panahandeh S, Jungkamp A-C, Moser M, Müller M.** 2010. TatB functions as an oligomeric binding site for folded Tat precursor proteins. *Mol Biol Cell* **21**:4151–61.
104. **Rollauer SE, Tarry MJ, Graham JE, Jääskeläinen M, Marcoux J, McDowell MA, Rodriguez F, Roversi P, Stansfeld PJ, Robinson, CV, Sansom, M., Palmer T, Högbom M, Berks BC, Lea SM.** 2013. Structure of the TatC core of the twin-arginine protein transport system. *Nature* **492**:210–4.
105. **Lee PA, Orriss GL, Buchanan G, Greene NP, Bond PJ, Punginelli C, Jack RL, Sansom, MSP, Berks BC, Palmer T.** 2006. Cysteine-scanning mutagenesis and disulfide mapping studies of the conserved domain of the twin-arginine translocase TatB component. *J Biol Chem* **281**:34072–85.
106. **Cline K.** 2015. Mechanistic aspects of folded protein transport by the twin arginine translocase (Tat). *J Biol Chem* **290**:16530–8.
107. **Alcock F, Stansfeld PJ, Basit H, Habersetzer J, Baker MA, Palmer T, Wallace MI, Berks, BC.** 2016. Assembling the tat protein translocase. *Elife* **5**.

108. **Habersetzer J, Moore K, Cherry J, Buchanan G, Stansfeld PJ, Palmer T.** 2017. Substrate-triggered position switching of TatA and TatB during Tat transport in *Escherichia coli*. *Open Biol* **7**:8.
109. **Gohlke U, Pullan L, McDevitt CA, Porcelli I, de Leeuw E, Palmer T, Berks, BC.** 2005. The TatA component of the twin-arginine protein transport system forms channel complexes of variable diameter. *Proc Natl Acad Sci U S A* **202**:10482–6.
110. **Brüser T, Sanders C.** 2003. An alternative model of the twin arginine translocation system. *Microbiol Res* **158**:7–17.
111. **Orriss GL, Tarry MJ, Ize B, Sargent F, Lea SM, Palmer T, Berks, BC.** 2007. TatBC, TatB, and TatC form structurally autonomous units within the twin arginine protein transport system of *Escherichia coli*. *FEBS Lett* **581**:4091–7.
112. **Aldridge C, Ma X, Gerard F, Cline K.** 2014. Substrate-gated docking of pore subunit Tha4 in the TatC cavity initiates Tat translocase assembly. *J Cell Biol* **205**:51–65.
113. **Hauer RS, Schlesier R, Heilmann K, Dittmar J, Jakob M, Klösigen RB.** 2013. Enough is enough: TatA demand during Tat-dependent protein transport. *Biochim Biophys Acta - Mol Cell Res* **1833**:957–65.
114. **Oates J, Barrett CML, Barnett JP, Byrne KG, Bolhuis A, Robinson C.** 2005. The *Escherichia coli* twin-arginine translocation apparatus incorporates a distinct form of TatABC complex, spectrum of modular TatA complexes and minor TatAB complex. *J Mol Biol* **346**:295–305.
115. **White GF, Schermann SM, Bradley J, Roberts A, Greene NP, Berks BC, Thomas, AJ.** 2010. Subunit organization in the TatA complex of the twin arginine protein translocase: A site-directed EPR spin labeling study. *J Biol Chem* **285**:2294–301.
116. **Hou B, Heidrich ES, Mehner-Breitfeld D, Brüser T.** 2018. The TatA component of the twin-arginine translocation system locally weakens the cytoplasmic membrane of *Escherichia coli* upon protein substrate binding. *J Biol Chem* **293**:7592-7605.
117. **Sargent F, Berks BC, Palmer T.** 2006. Pathfinders and trailblazers: A prokaryotic targeting system for transport of folded proteins. *FEMS Microbiology Letters* **254**:198–207.

118. **Gouffi K, Gérard F, Santini CL, Wu LF.** 2004. Dual Topology of the *Escherichia coli* TatA Protein. *J Biol Chem* **279**:11608–15.
119. **Walther TH, Gottselig C, Grage SL, Wolf M, Vargiu A V., Klein MJ, Ulrich AS.** 2013. Folding and self-assembly of the TatA translocation pore based on a charge zipper mechanism *Cell* **152**:316–26
120. **Dabney-Smith C, Mori H, Cline K.** 2006. Oligomers of Tha4 organize at the thylakoid Tat translocase during protein transport. *J Biol Chem* **281**:5476–83.
121. **Chan CS, Zlomislic MR, Tieleman DP, Turner RJ.** 2007. The TatA subunit of *Escherichia coli* twin-arginine translocase has an N-in topology. *Biochemistry* **46**:7396–404.
122. **Aldridge C, Storm A, Cline K, Dabney-Smith C.** 2012. The chloroplast twin arginine transport (Tat) component, Tha4, undergoes conformational changes leading to Tat protein transport. *J Biol Chem* **287**:34752–63.
123. **Taubert J, Hou B, Risselada HJ, Mehner D, Lünsdorf H, Grubmüller H, Brüser, T.** 2015. TatBC-Independent TatA/Tat substrate interactions contribute to transport efficiency. *PLoS One* **10**:1–24.
124. **Pal D, Fite K, Dabney-Smith C.** 2013. Direct Interaction between a Precursor Mature Domain and Transport Component Tha4 during Twin Arginine Transport of Chloroplasts. *PLANT Physiol* **161**:990–1001.
125. **Taubert J, Brüser T.** 2014. Twin-Arginine translocation-Arresting protein regions contact Tata and TatB. *Biol Chem* **395**:827–36.
126. **Mould RM, Shackleton JB, Robinson C.** 1991. Transport of proteins into chloroplasts. Requirements for the efficient import of two luminal oxygen-evolving complex proteins into isolated thylakoids. *J Biol Chem* **266**:17286–9.
127. **Mori H, Cline K. A.** 2002. twin arginine signal peptide and the pH gradient trigger reversible assembly of the thylakoid Δ pH/Tat translocase. *J Cell Biol* **157**:205–10.
128. **DeLisa MP, Lee P, Palmer T, Georgiou G.** 2004. Phage Shock Protein PspA of *Escherichia coli* Relieves Saturation of Protein Export via the Tat Pathway. *J Bacteriol* **186**:366–73.
129. **Finazzi G, Chasen C, Wollman FA, De Vitry C.** 2003. Thylakoid targeting of Tat passenger

- proteins shows no Δ pH dependence in vivo. *EMBO J* **22**:807–15.
130. **Braun NA, Davis AW, Theg SM.** 2007. The chloroplast tat pathway utilizes the transmembrane electric potential as an energy source. *Biophys J* **93**:1993–8.
131. **Musser SM, Theg SM.** 2000;. Proton transfer limits protein translocation rate by the thylakoid Δ pH/Tat machinery. *Biochemistry.* **39**:8228–33.
132. **Alder NN, Theg SM.** 2003. Energetics of protein transport across biological membranes: A study of the thylakoid Δ pH-dependent/cpTat pathway. *Cell* **112**:231–42.
133. **Gérard F, Cline K.** 2007. The thylakoid proton gradient promotes an advanced stage of signal peptide binding deep within the Tat pathway receptor complex. *J Biol Chem* **282**:5263–72.
134. **Bageshwar UK, Musser SM.** 2007. Two electrical potential-dependent steps are required for transport by the *Escherichia coli* Tat machinery. *J Cell Biol* **179**:87–99.
135. **Turner RJ, Papish AL, Sargent F.** 2004. Sequence analysis of bacterial redox enzyme maturation proteins (REMPs). *J Microbiol* **50**:225–38.
136. **Tranier S, Mortier-Barrière I, Ilbert M, Birck C, Iobbi-Nivol C, Méjean V, Samama, JP.** 2002..Characterization and multiple molecular forms of TorD from *Shewanella massilia*, the putative chaperone of the molybdoenzyme TorA. *Protein Sci* **11**:2148–57.
137. **Chan CS, Chang L, Rommens KL, Turner RJ.** 2009. Differential interactions between tat-specific redox enzyme peptides and their chaperones. *J Bacteriol* **191**:2091–101.
138. **Halbig D, Wiegert T, Blaudeck N, Freudl R, Sprenger GA.** 1999. The efficient export of NADP-containing glucose-fructose oxidoreductase to the periplasm of *Zymomonas mobilis* depends both on an intact twin- arginine motif in the signal peptide and on the generation of a structural export signal induced by cofactor bind. *Eur J Biochem* **263**:543–51.
139. **Matos CFRO, Robinson C, Di Cola A.** 2008. The Tat system proofreads FeS protein substrates and directly initiates the disposal of rejected molecules. *EMBO J* **27**:2055–63.
140. **Goosens VJ, Monteferrante CG, Van Diji JM.** 2014. Co-factor insertion and disulfide bond

- requirements for twin-Arginine translocase-dependent export of the *Bacillus subtilis* Rieske protein QcrA. *J Biol Chem* **289**:13124–31.
141. **Sanders C, Wethkamp N, Lill H.** 2001. Transport of cytochrome c derivatives by the bacterial Tat protein translocation system. *Mol Microbiol* **41**:241–6.
142. **Sutherland GA, Grayson KJ, Adams NBP, Mermans DMJ, Jones AS, Robertson AJ, Auman, DB, Brindley AA, Sterpone F, Tuffery P, Derreumaux P, Dutton PL, Robinson C, Hitchcock A, Hunter CN.** 2018. Probing the quality control mechanism of the *Escherichia coli* twin-arginine translocase with folding variants of a *de novo* -designed heme protein. *J Biol Chem* **293**:6672-6681
143. **Alanen HI, Walker KL, Lourdes Velez Suberbie, M, Matos, CFRO, Bönisch S, Freedman RB, Keshavarz-Moore E, Ruddock L, Robinson, C.** 2014. Efficient export of human growth hormone, interferon α 2b and antibody fragments to the periplasm by the *Escherichia coli* Tat pathway in the absence of prior disulfide bond formation. *Biochimica et Biophysica Acta - Molecular Cell Research* **1853**:756–763.
144. **Stolle P, Hou B, Brüser T.** 2016. The Tat substrate CueO is transported in an incomplete folding state. *J Biol Chem* **291**:13520–8.
145. **Cline K, McCaffery M.** 2007. Evidence for a dynamic and transient pathway through the TAT protein transport machinery. *EMBO J* **26**:3039–49.
146. **Richter S, Lindenstrauß U, Lücke C, Bayliss R, Brüser T.** 2007. Functional tat transport of unstructured, small, hydrophilic proteins. *J Biol Chem* **282**: 33257–64.
147. **Lindenstrauß U, Brüser T.** 2009. Tat transport of linker-containing proteins in *Escherichia coli*. *FEMS Microbiol Lett* **295**:135–40.
148. **Austerberry JI, Dajani R, Panova S, Roberts D, Golovanov AP, Pluen, A, van der Walle CF, Uddin S, Warwicker J, Derrick JP, Curtis, R.** 2017. The effect of charge mutations on the stability and aggregation of a human single chain Fv fragment. *European Journal of Pharmaceutics and Biopharmaceutics* **115**: 18–30.
149. **Jones AS, Austerberry JI, Dajani R, Warwicker J, Curtis R, Derrick JP, Robinson, C.** 2016.

Proofreading of substrate structure by the Twin-Arginine Translocase is highly dependent on substrate conformational flexibility but surprisingly tolerant of surface charge and hydrophobicity changes. *Biochim Biophys Acta - Mol Cell Res* **1863**:3116–24.

150. **Monteferrante CG, Mackichan C, Marchadier E, Prejean MV, Carballido-López R, Van Dijl JM.** 2013. Mapping the twin-arginine protein translocation network of *Bacillus subtilis*. *Proteomics* **13**:800-11.

151. **Krishnappa L, Monteferrante CG, van Dijl JM.** 2012. Degradation of the twin-arginine translocation substrate YwBn by extracytoplasmic proteases of *Bacillus subtilis*. *Appl Environ Microbiol* **78**:7801-4.

152. **Panahandeh S, Maurer C, Moser M, DeLisa MP, Müller M.** 2008. Following the path of a twin-arginine precursor along the TatABC translocase of *Escherichia coli*. *J Biol Chem* **283**:33267–75.

153. **Rocco MA., Waraho-Zhmeyev D, DeLisa MP.** 2012. Twin-arginine translocase mutations that suppress folding quality control and permit export of misfolded substrate proteins. *Proc Natl Acad Sci* **109**:13392–7.

154. **Overton TW.** 2014. Recombinant protein production in bacterial hosts. *Drug Discov Today* **19**:590–601.

155. **Pooley HM, Merchante R, Karamata D.** 1996. Overall Protein Content and Induced Enzyme Components of the Periplasm of *Bacillus subtilis*. *Microb Drug Resist* **2**:9–15.156.

156. **Van Dijl JM, Braun PG, Robinson C, Quax WJ, Antelmann H, Hecker M, Muller J, Tjalsma H, Bron S, Jongbloed JD.** 2002. Functional genomic analysis of the *Bacillus subtilis* Tat pathway for protein secretion. *J Biotechnol* **98**:243–54.

157. **Dröge MJ, Boersma YL, Braun PG, Buining RJ, Julsing MK, Selles KGA van Dijl JM, Quax WJ.** 2006. Phage display of an intracellular carboxylesterase of *Bacillus subtilis*: Comparison of Sec and Tat pathway export capabilities. *Appl Environ Microbiol* **72**:4589-95.

158. **Goosens VJ, van Dijl JM.** 2017. Twin-arginine protein translocation. In: Current Topics in Microbiology and Immunology. *Curr Top Microbiol Immunol* **404**:69-94.

159. **French C, Keshavarz-Moore E, Ward JM.** 1996. Development of a simple method for the recovery of recombinant proteins from the *Escherichia coli* periplasm. *Enzyme Microb Technol* **19**:332–8.
160. **Matos CFRO, Robinson C, Alanen HI, Prus P, Uchida Y, Ruddock LW, Freedman R, Keshavarz-Moore E.** 2014. Efficient export of prefolded, disulfide-bonded recombinant proteins to the periplasm by the Tat pathway in *Escherichia coli* CyDisCo strains. *Biotechnol Prog* **30**:281–90.
161. **Browning DF, Richards KL, Peswani AR, Roobol J, Busby SJW, Robinson C.** 2017. *Escherichia coli* “TatExpress” strains super-secrete human growth hormone into the bacterial periplasm by the Tat pathway. *Biotechnol Bioeng* **114**:2828–36.
162. **Albiniak AM, Matos CFRO, Branston SD, Freedman RB, Keshavarz-Moore E, Robinson C.** 2013. High-level secretion of a recombinant protein to the culture medium with a *Bacillus subtilis* twin-arginine translocation system in *Escherichia coli*. *The FEBS Journal* **280**:3810–21.
163. **Kim JY, Kim Y-GG, Lee GM.** 2012. CHO cells in biotechnology for production of recombinant proteins: current state and further potential. *Appl Microbiol Biotechnol* **93**:917–30.
164. **Mattanovich D, Branduardi P, Dato L, Gasser B, Sauer M, Porro D.** 2012. Recombinant Protein Production in Yeasts. *Recombinant Gene Expression* **824**:329–58.
165. **Behrendt J, Lindenstrauss U, Bruser T.** 2007. The TatBC complex formation suppresses a modular TatB-multimerization in *Escherichia coli*. *FEBS Lett* **581**:4085–90.
166. **Barnett JP, Van Der Ploeg R, Eijlander RT, Nenninger A, Mendel S, Rozeboom R, Kuipers OP, Van Dijk JM, Robinson C.** 2009. The twin-arginine translocation (Tat) systems from *Bacillus subtilis* display a conserved mode of complex organization and similar substrate recognition requirements. *FEBS journal* **276**:232–243.
167. **Kolkman MA, van der Ploeg R, Bertels M, van Dijk M, van der Laan J, van Dijk JM, Ferrari E.** 2008. The twin-arginine signal peptide of *Bacillus subtilis* YwbN can direct either Tat- or Sec-dependent secretion of different cargo proteins: secretion of active subtilisin via the *B. subtilis* Tat pathway. *Appl Environ Microbiol.* **74**:7507–7513.

168. van der Ploeg R, Monteferrante CG, Piersma S, Barnett JP, Kouwen TR, Robinson C, van Dijk JM. 2012. High-salinity growth conditions promote Tat-independent secretion of Tat substrates in *Bacillus subtilis*. *Appl Environ Microbiol.* **78**:7733-7744.

<i>E. coli</i>	Molecular mass of gene product (kDa)	Molecular mass of complex (kDa)	Reference
TatA	11	100-500	(114)
TatB	18	100	(111)
TatC	30	220	(111)
TatE	7	50-110	(41)
TatABC		440-580	(165)
<i>B. subtilis</i>	Molecular mass of gene product (kDa)	Molecular mass of complex (kDa)	Reference
TatAd	7	270*	(53)
TatCd	27	100	(54)
TatAy	6	200*	(166)
TatCy	28	66	(54)
TatAc	6	100	(54)
TatAdCd		230	(54)
TatAyCy		200	(54)

Table 1. **Molecular mass of Tat proteins and complexes in *E. coli* and *B. subtilis***

The predicted molecular mass of *E. coli* and *B. subtilis* Tat proteins is according to gene sizes. The molecular mass of complexes was approximated mostly by blue-native gel electrophoresis, except those determined by gel filtration chromatography, as indicated with a *

Figure legends:

Figure 1. **Two alternative protein translocation pathways, Sec and Tat.**

Proteins always originate from translating ribosomes (R). Their N-terminal signal peptide (OmpA or TorA in this overview) will direct the nascent polypeptide chain to the correct translocase (Sec or Tat, respectively), which may be aided by chaperones. The unfolded Sec protein is transferred to SecA where it is threaded through the SecYEG channel in the plasma membrane, powered by repeated cycles of ATP binding and hydrolysis. In the oxidising periplasm, the unfolded protein assumes its tertiary fully folded state. The Tat-dependently translocated protein is fully folded within the cytoplasm where it may also acquire its cofactor. Once directed to TatBC, TatA protomers are recruited to translocate the protein across the cytoplasmic membrane. Energy required for this process is created by the proton-motive force. Please note that mRNA molecules are schematically represented by an interrupted line, synthesized proteins by uninterrupted lines, and translocase subunits by cylinders.

Figure 2. **Comparison of Tat (TorA) and Sec (OmpA) signal peptides**

The structure of Tat and Sec signal peptides includes three regions, namely a basic N-domain (blue), a hydrophobic H-domain (red) and a polar C-domain (green). A signal peptidase cleavage site (AxA) is positioned prior to the mature protein (pink). The amino acid sequences of the TorA and OmpA signal peptides are specified. Tat signal peptides (above) have a consensus motif containing 'twin-arginines', while Sec signal peptides do not contain this motif. Sec signal peptides tend to be shorter, with fewer residues in their N and H-domains, than Tat signal peptides.

Figure 3. **Schematic representation of *E. coli* TatABC components in the plasma membrane**

Three essential components form the Gram-negative bacterial Tat complex, namely TatA, TatB and TatC. TatA/E and TatB have similar topologies in that they have one TM helix domain with a short periplasmic N-terminal region, a tilted APH and an unstructured C-terminus on the cytoplasmic side

of the plasma membrane. Notably, TatB is larger than TatA with a longer C-terminal tail. TatC is significantly bigger as it containing 6 membrane-embedded helices with both the C- and N-terminal ends residing in the cytoplasm. The helices 5 and 6 do not fully span the membrane, which may contribute to TatC's function.




Figure 1




Figure 2




Figure 3