



APR 3 1974



THE PUMMELO IN HAWAII

Warren Yee
Specialist in Horticulture



ex libris



UNIVERSITY OF HAWAII
LIBRARY

Acknowledgments

The Honolulu Lions Club and the College of Tropical Agriculture gratefully acknowledge the help of the many home gardeners who participated in the 1971 and 1972 pummelo contests. Only with the assistance of these community gardeners was the locating and cataloging of local outstanding pummelo seedlings possible, and these trees are now being observed and evaluated. Both the Lions Club and the College hope that home gardeners will continue to notify the Extension Horticulturist of outstanding pummelos observed anywhere in the State.

Perhaps Hawaii's people will one day be able to serve to their friends another interesting and delightful Hawaii-grown fruit.

Members of the Agriculture Committee, Honolulu Lions Club, include:

Warren Y. J. Yee, Chairman
Thomas Vance, Vice-Chairman
Henry C. Bush
Jerry A. Y. Ching
Walter N. Fujikami
Glenn A. Lee
Wui Lung
Clarence Williams

Members of the College of Tropical Agriculture, University of Hawaii, for this project include:

Richard A. Hamilton
Bunki Kumabe
Warren Y. J. Yee

What Is a Pummelo?

Pummelos and grapefruits are closely related, and indeed the grapefruit was derived from the pummelo, probably by natural hybridization. But their differences are numerous. The pummelo looks like a grapefruit, but it is larger, up to five pounds in weight, and more varied in shape sometimes oval or pear-shaped rather than round. It is sweeter, although pummelo can also be highly acid and bitter. The rinds are very often thicker, and the flesh is more firm; segments can be separated and membranes removed, making it easy to eat with the fingers. The pummelo is one more idea for Hawaii citrus growers.

THE PUMMELO IN HAWAII

Warren Y. J. Yee
Specialist in Horticulture

The pummelo (*Citrus maxima*) is indigenous to the Malayan peninsula and East Indian archipelago, and has spread from there to South China, India, and finally to other parts of the world. Other names by which it is known are *shaddock* in Barbados (named after a Captain Shaddock), *pompelmoes* in Indonesia, *pamplemousse* in French Oceania, *buntan* or *zabon* in Japan, *Som-O* in Thailand, and pummelo or *poo look* in Hawaii. In the Hawaiian Islands, pummelos are grown from sea level up to 1500 feet elevation, and usually the trees are located in home gardens below 200 feet elevation. The temperature range in areas where they are grown varies from a low of 60–65 F in winter to a high of 85–90 in late summer and early fall.

Uses

Pummelos are principally eaten as fresh fruit or used as an ingredient in fruit salads. Sometimes the rind is sliced and made into candied peel or cooked and eaten as a vegetable. The fruit is popular with the Chinese and is prominently displayed by them in religious ceremonies, during New Year's festivities, and as a decorative piece in the home; it is ideal for these purposes because of its long shelf life and attractiveness. Pummelo trees are often used to accent the entrances to Chinese gardens and homes.

When pummelos are eaten as fresh fruit, the covering or capillary membrane around each segment is peeled off, and in good quality pummelos this membrane is easily removed. At the same time, the walls of exposed juice sacs should be strong enough to keep the juice from leaking out.

Varieties

Hawaii has a great many seedling pummelos in home gardens, but most of them are of poor eating quality. They may be bitter, seedy, thick skinned, poorly segmented, or very acid (Figure 1). Poor quality seedling pummelos are generally unpalatable as fresh fruit. Among the many seedlings and imported varieties grown in Hawaii, however, some are of good to excellent quality, including 'Au,' 'Chandler,' 'Diamond Head,' 'Ho,' 'Kao Pan,' 'Kau,' 'Pauthel,' 'Sakata,' and others.




Figure 1. Undesirable characteristics of pummelos include thick skin and a disarray of segments of varying sizes.




Figure 2.
'Au'—a pink-fleshed, slightly acid variety from Kauai.

Figure 3.
'Ho'—a light-yellow-fleshed variety which has produced fruits prolifically at Puna Street below Alewa Heights in Honolulu.



Au (Figure 2)

Seedling planted at Kapaa Homestead, Kauai, more than 40 years ago by the late Tai Hon Au. Shape oval to collared.* Rind dark yellow, ½ inch thick, and surface slightly rough. Flesh dark pink around segment membrane, and lemon colored toward the center. Segments well formed, small to medium, membrane fairly easy to remove. Juice sacs fine and compact. Flavor slightly acid to acid with slight bitterness. Best eaten when full ripe. Weight: 1½–2½ pounds. Season: October to January.

Chandler

Pink variety resulting from a 'Siamese Pink' pollen parent and a 'Siamese Sweet' seed parent, bred and released by Dr. James Cameron, University of California at Riverside. Shape oblate to globose. Rind yellow, about ½ inch thick, and smooth. Flesh moderately juicy and pink. Juice sacs loosely held together. Segments well formed and large. Flavor slightly acid but sweet. Weight: 3–3½ pounds. Season: October to November.

Diamond Head

Large, attractive, light-pink variety. Shape subglobose to spherical. Rind attractive yellow and about ¾ inch thick. Flesh moist and slightly pink near rind. Segments medium and uniform with juice sacs easily broken. Flavor slightly acid and sweet. Weight: about 3 pounds. Season: October to November.

Ho (Figure 3)

Seedling from Waialua, Oahu. Shape broadly pyriform. Rind yellow, 3/8 inch thick. Flesh moist and pale yellow. Juice sacs compact but easily broken. Segments small to medium and well formed with membrane slightly difficult to remove. Flavor slightly acid but sweet with a trace of bitterness. Weight: 1½–2 pounds. Season: September to November.

Kao Pan (Figure 4)

Introduced from Thailand by the U.S.A. and obtained from the University of California Citrus Research Center by the Hawaii Agricultural Experiment Station. Shape subglobose to spherical. Rind bright yellow, ½ inch thick, and smooth. Flesh lime colored and juicy. Juice sacs large and easily separated from segment membrane. Segments well formed. Flavor sweet and slightly acid. Weight: about 3 pounds. Season: October to November.

*In some varieties other shapes can appear on the same tree, but normally one form predominates.




Figure 4. 'Kao Pan'—a variety popular in Thailand. Trees grown at Waimanalo and Poamoho on Oahu have been prolific bearers.

Kao Phuang

Imported from Thailand by the University of California at Riverside. Shape broadly pyriform with short neck. Rind yellow and about $\frac{3}{8}$ inch thick. Flesh juicy and greenish in color. Segments large and uniform with juice sacs held firmly to base of segment membrane. Flavor moderately acid but pleasant. Weight: 2–3½ pounds. Season: October to December.

Kau (Figure 5)

Seedling from seed brought from China and planted by Nancy Kau, 650 Ninth Avenue, Kaimuki, Oahu. Shape broadly obovoid. Rind yellow, about $\frac{1}{2}$ inch thick, and easy to peel. Flesh moderately moist and tinged with pink next to rind. Segments large and well formed with membrane easily separated. Juice sacs firm and fairly compact. Flavor slightly acid, but sweet. Bitterness slight and infrequent. Weight: 3–4 pounds. Season: September to October.

Pauthel (Figure 6)

Seedling grown by Paul Wong, Aliikoa Street, Kaimuki, Oahu. Shape pyriform to obovate. Rind light yellow and about $\frac{3}{8}$ inch thick. Flesh moderately juicy and pale yellow. Juice sacs firm and compact. Segments well formed and large with membrane easily removed. Flavor sweet and slightly acid. Weight: 2–3 pounds. Season: August to September.




Figure 5. 'Kau'—a medium- to large-sized pummelo with flesh slightly tinged with pink. Excellent quality.



Figure 6. 'Pauthel'—a pale-yellow-fleshed variety of excellent quality and flavor. Considered an early variety.




Figure 7. 'Sakata'—a lime-colored pyriform fruit of excellent flavor.

Sakata (Figure 7)

Discovered and grown by Mrs. Hatsu Sakata, 45-623 Kealahala Road, Kaneohe, Oahu. Shape pyriform to obovate. Rind light yellow, 3/4 inch thick, and smooth. Flesh lime colored and juicy. Juice sacs medium and easily broken. Segments medium in size and well formed. Flavor slightly acid but sweet. Weight: 1½-2½ pounds. Season: August to September.

Siamese

Deep pink variety. Shape oblate. Rind yellow and about ¼ inch thick. Flesh juicy. Segments large but lacking in uniformity. Juice sacs tender and easily separated from segment membrane. Flavor moderately acid. Weight: 2½-3½ pounds. Season: September to October.

Climate

Pummelos grow well from sea level up to 2000 feet elevation, and in most of the areas where other citrus are cultivated in Hawaii, they also flourish. Rainfall is not as important for growth if supplementary irrigation is provided. As is true for many other trees, wind is detrimental to pummelos, making it often necessary to provide windbreaks, especially on the windward side of the Islands. Trees should be planted in full sunlight for optimum growth and yields. In Hawaii, temperature seems to be a critical factor affecting fruit quality and bearing season, and the best quality fruits so far have been produced in warm locations below 500 feet elevation.

Soil

Pummelos will grow on a wide range of soils. Well-drained soils with good textures and depths are considered ideal, examples of which include the red or reddish-brown soils of Lahaina, Maui, and Waipahu, Oahu. On sandy, infertile soil, such as that found on the coastal areas of Oahu, the introduction of compost, red topsoil, and an adequate fertilizer program will help to promote normal growth.

Season

The season of maturity for pummelos in Hawaii extends from August to January; however, September to November is the period of greatest production. Off-season fruits are sometimes available in limited quantities during other times of the year.

Harvesting

Some varieties can be harvested from the tree with good quality fruits when the skin is still greenish yellow; however, most varieties must be fully yellow before harvesting. Fruits harvested prematurely are more acid and drier than they should be. Those harvested just before they fall naturally tend to be less acid, sweeter, and juicier.

Culture

Propagation

The three most common methods of propagating pummelos are by seeds, air layering (marcottage), and grafting. At present, most pummelo trees are grown from seeds in home gardens; the trend, however, is toward air-layered and grafted trees.

Fruits from trees *grown from seed* are usually different from and of poorer quality than improved selections and varieties which are propagated asexually. Asexually propagated trees with known fruit quality, therefore, are usually recommended for home garden planting, but the grower or hobbyist who is interested in finding variations can experiment by planting seeds of one of the better pummelos. Trees grown from seed generally take 5 years or more to bear fruit.

Air layering is presently the most common method of reproducing trees of one variety, popular primarily because it is a simple method of propagation. Trees produced in this way frequently set fruit in 5-gallon containers in 2 to 3 years and are attractive displays in garden shops.

Pummelos may be *grafted* on a number of different rootstocks, but, since little research information is available, it is satisfactory in most cases to graft pummelos on pummelo rootstock. It usually takes approximately a year to grow a rootstock large enough for grafting. Among rootstocks upon which pummelos have been successfully grafted are rough lemon and *heennaran*. Trees grafted on rough lemon more than 15 years ago have shown no signs of incompatibility and are still producing well at Kihei, Maui.

Transplanting

Seedlings, grafted trees, or air-layered trees should be transplanted into the garden, normally when they reach a height of 1½ feet or larger and have well-matured leaves. In transplanting, the hole should be approximately 2 feet wide and 1½ feet deep to give the newly transplanted tree ample room to begin growth. The hole should be dug in an area far enough from other trees to avoid roots competing for space. One pound of treble superphosphate or 2 pounds of superphosphate should be placed at the bottom of the hole and covered with 1 inch of soil. The plant is then placed in the hole, and the root system is covered with soil so that the soil level is approximately 1½ to 2 inches higher than it was in the container. This is to insure that the soil level at the base of the tree after the soil settles into the hole will be about the same as before. Then a handful of complete fertilizer, such as 10-10-10, or any other fertilizer with a similar ratio of nitrogen (N), phosphorus (P₂O₅), and potash (K₂O), is applied by scattering it into the basin over the surface of the hole. Finally the tree is watered.

Fertilization

After planting, about a handful of fertilizer should be applied every second month during the first year of growth. The amount is gradually increased each following year, and one suggestion for the schedule is as follows:

2nd year—every 3 months—¼ to ½ pound

3rd year—every 4 months—1 to 2 pounds

4th year—every 4 months—2 pounds

5th year and thereafter—a minimum of 1 to 1½ pounds per inch trunk diameter per year which may be divided into 2 or 3 applications per year.

The time and amount of fertilization for bearing pummelo trees have not been fully determined, but recommendations made for other citrus trees may be applicable.

Irrigation

Young transplanted trees require plenty of water and should be irrigated at least twice a week during the first year. A weekly irrigation should be sufficient thereafter for the drier sections of the State. As the trees increase in size, the area irrigated should also be enlarged to cover at least 2 feet beyond the drip line of the branches.

References

- Reuther, Walter, H. J. Webber, and L. D. Batchelor, eds. 1967. The citrus industry, Vol. 1, Rev. Ed. Univ. Calif. Div. Agr. Sci. Pp. 534-538.
- Reitz, H. J., C. D. Leonard, et al. 1964. Recommended fertilizers and nutritional sprays for citrus. Florida Agr. Exp. Sta. Bull. 536B. 23pp.

The Hawaii Cooperative Extension Service provides equal opportunities in its programs and employment.

Hawaii residents may order single copies of publications free of charge from county offices. Quantities of any one publication are sold at cost. Quotations will be furnished on written request to Agricultural Publications and Information Office, College of Tropical Agriculture, 2500 Dole Street, Krauss Hall Room 107, Honolulu, Hawaii 96822. Price per copy to bulk users, 15¢ plus postage.

COOPERATIVE EXTENSION SERVICE OFFICES

Oahu County

Wahiawa Office
Wahiawa Civic Center Bldg.
910 California Avenue
P.O. Box 610
Wahiawa, Hawaii 96786
Telephone: 622-4185
Honolulu Office
429-B Waiakamilo Road
Honolulu, Hawaii 96817
Telephone: 847-0871
Kaneohe Office
State Office Bldg.
45-260 Waikalua Road
P.O. Box 607
Kaneohe, Hawaii 96744
Telephone: 247-0421

Mauī-Molokai County

Mauī-Molokai County Office
70 High Street
Post Office Bldg.
P.O. Box 870
Wailuku, Hawaii 96793
Telephone: 244-3242
244-3254
Kula Office
Kealahou Community Hall, Waiakoa
P.O. Box 185
Kula, Hawaii 96790
Telephone: 878-1275
Molokai Office
State Office Bldg., Kaunakakai
P.O. Box 158
Kaunakakai, Hawaii 96748
Telephone: 553-5445
Kauai County
Kauai County Office
State Office Bldg., Lihue
P.O. Box 1588
Lihue, Hawaii 96766
Telephone: 245-4471

Hawaii County

Hilo Office
Federal Bldg., Hilo
P.O. Box 1882
Hilo, Hawaii 96720
Telephone: 935-2804
Kamuela Office
P.O. Box 237
Kamuela, Hawaii 96743
Telephone: 885-4695
Kau Office
State Office Bldg., Naalehu
P.O. Box 96
Naalehu, Hawaii 96772
Telephone: 929-7012
Kona Office
Kainaliu
P.O. Box 208
Kealahou, Hawaii 96750
Telephone: 322-3761

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. C. Pears Wilson, Director, Cooperative Extension Service, College of Tropical Agriculture, University of Hawaii, Honolulu, Hawaii 96822.

CIRCULAR 483—MARCH 1974—5M