



Cite this: *Dalton Trans.*, 2014, **43**, 16227

Syntheses of sterically shielded stable carbenes of the 1,2,4-triazole series and their corresponding palladium complexes: efficient catalysts for chloroarene hydrodechlorination†

Nataliya V. Glinyanaya,^a Vagiz Sh. Saberov,^a Nikolai I. Korotkikh,^{*a} Alan H. Cowley,^b Rachel R. Butorac,^b Daniel A. Evans,^b Tatyana M. Pekhtereva,^a Anatolii F. Popov^a and Oles P. Shvaika^a

The new sterically shielded 1,3,4-trisubstituted 1,2,4-triazol-5-ylidenes **8b–d** were synthesized by a three step method starting from 2-phenyl-1,3,4-oxadiazole. The syntheses of palladium complexes **9a–d** and **10a–d** (including the sterically shielded derivatives **9c,d** and **10a–d**) were carried out *via* the reactions of the stable carbenes **8a–d** with palladium halogenide salts in THF or toluene solution. Complexes **9c,d** and **10a–d** were found to be excellent catalysts for the reductive dechlorination (hydrodechlorination) of *p*-dichlorobenzene. The structures of **8c**, **9a,b**, and **10a** were determined by single-crystal X-ray diffraction.

Received 7th May 2014,
Accepted 24th June 2014

DOI: 10.1039/c4dt01353k

www.rsc.org/dalton

Introduction

Recently, stable carbenes have proven to be versatile reagents and catalysts for many important organic reactions.^{1–3} Heteroaromatic imidazol-2-ylidenes and 1,2,4-triazol-5-ylidenes are particularly well known examples of stable carbenes.⁴ The former type has been studied extensively; however, as individual compounds, 1,2,4-triazol-5-ylidenes are relatively unexplored. The first stable carbene of this type, namely a 1,3,4-triphenyl substituted derivative, was prepared and investigated by Enders and co-workers.^{5,6} Subsequently, new types of 1-adamantyl- and 1-*tert*-butyl substituted 1,2,4-triazol-5-ylidenes have been described^{7,8} along with their transformations upon heating⁹ and reaction with a malonic ester.¹⁰ Biscarbenes with enhanced thermal and chemical stabilities featuring conjugated 3,3'- and 4,4'-bis-1,2,4-triazol-5-ylidenes have been synthesized and used as ligand supports for copper(i) complexes.^{11,12} In a series of studies 1,2,4-triazol-5-ylidene complexes were prepared *in situ* from the corresponding salts (for the latest see,

for example, ref. 13–15). Furthermore, 1,3,4-triphenyl-1,2,4-triazol-5-ylidene has proven to be not only an effective catalyst for organic reactions (*e.g.* benzoin and Stetter condensations),² but also a ligand for metal catalyzed reactions.³ To the best of our knowledge, sterically shielded carbenes of this general type have not been synthesized thus far. However, such compounds could prove to be useful for catalytic applications and for the development of new methods for haloarene hydrodehalogenation.

Halogenaromatic compounds (haloarenes) are widely used as *e.g.* pesticides, pharmaceuticals, and high-temperature fluids. Unfortunately, most of the aforementioned materials are highly toxic. Moreover, these toxic haloarenes have accumulated in large amounts throughout the world. Several of these haloarene compounds require particular attention, specifically DDT, hexachlorobenzene, dioxins, polychlorodibenzofurans (furans), and polychlorobiphenyls.¹⁶ All of the foregoing compounds were included in the Stockholm Convention of May 22nd 2001 in a list of the 12 most hazardous “persistent organic pollutants” (POPs). The safe management of these troublesome toxic POPs represents a persistent environmental concern.

The most common pathways for haloarene detoxification are as follows: (1) oxidative methods (combustion at high temperatures or oxidation in critical water, *etc.*), (2) reductive methods such as catalytic reduction with hydrogen, reduction with metals and nanometals, and (3) biodegradation methods.¹⁷ The first group of methods for haloarene detoxification has some disadvantages such as the use of high

^aThe L.M. Litvinenko Institute of Physical Organic and Coal Chemistry, Ukrainian Academy of Sciences, 70, R. Luxemburg, Donetsk, 83114, Ukraine.
E-mail: nkorotkikh@ua.fm; Fax: +(380 62)311-68-30; Tel: +(380 62)311-68-35

^bDepartment of Chemistry, The University of Texas at Austin, 1 University Station A5300, Austin, Texas 78712-0165, U.S.A.. E-mail: acowley@cm.utexas.edu;
Tel: +1(512)471-74-84

† Electronic supplementary information (ESI) available: X-ray crystallographic data for **8c**, **9a,b**, **10c**; NMR spectra; CIF files for **8c**, **9a,b**, **10c**. CCDC 1000720 (**8c**), 1000718 (**9a**), 1000719 (**9b**) and 1000721 (**10a**). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4dt01353k

temperature and/or high pressure conditions, the difficulty in recovering the vaporized polychloroarenes, and the high risk of producing dioxins. The second group of detoxification methods also presents some unfavorable aspects, particularly those that concern the formation of corrosive hydrohalogenides and the high pressures that are necessary for hydrogenation and acidic reductions,¹⁸ the use of dangerous or expensive reducing reagents such as alkali metals and metalorganic compounds,¹⁹ microwave techniques,²⁰ all of which result in low decomposition rates or efficiencies. For example, the promising use of metallic calcium in ethanol for hydrodehalogenation requires large quantities of the metal (10–40 fold excess per 1 Cl) and involves the formation of several by-products of the ring reduction.^{21–23} Noble metal nanoparticles allow recycling of the catalyst. However, they require approximately 1 mol% Pd to achieve the necessary activity for detoxification.^{24,25} Furthermore, biodegradation methods with aerobic microbes are only suitable for the reduction of low halogenated arenes and hence they are not useful as a general detoxification method (such organisms are contained in small amounts in the first few mm of soil).^{16,17} Moreover, anaerobic microorganisms are suitable for reductive dechlorination (hydrodechlorination) which results in the conversion of more chlorinated hydrocarbons. However, such species only react with *m*- and *p*-substituted compounds and do not activate *o*-substituted haloarenes.

Thus, among the known methods, catalytic reductive dehalogenation (hydrodehalogenation) promises the greatest potential for success. In the interest of achieving high catalytic activities and cost effectiveness, catalytic reductive dehalogenation could represent the best solution for the neutralization of toxic POPs. Moreover, palladium carbene complexes are well known to be the suitable catalysts for the foregoing reaction. For example, it is known that an *in situ* generated catalyst based on Pd(dba)₂ and 1,3-disubstituted imidazol-2-ylidenes affords high yields of hydrodehalogenation products when carried out in dioxane at 100 °C (TON up to 28–50, TOF up to 28–50 h⁻¹).²⁶ Compound **1** (Fig. 1) provides moderate catalytic

efficiencies for the reactions of monochlorobenzenes at room temperature²⁷ (TON 100, TOF 200 h⁻¹) and complex **2** has been shown to be active for the hydrodehalogenation of *p*-dichlorobenzene at 60 °C (TON 194, TOF 111 h⁻¹).²⁸ Furthermore, the use of microwave irradiation at 120 °C resulted in the highest rate for this reaction. In this case, a virtually quantitative yield of the product was obtained in only 2 min (TON 4000, TOF 120 000 h⁻¹ for *p*-chlorotoluene; TON 3800, TOF 114 000 h⁻¹ for *p*-dichlorobenzene).²⁸

The catalytic efficiency of the nickel complex **3** that was prepared *via* the reaction of Ni(PPh₃)₂Br₂ with the corresponding carbene²⁹ is not high for the hydrodebromination transformation of *p*-bromofluorobenzene or for the hydrodechlorination of *p*-chlorofluorobenzene (activated haloarenes) (TON 15–22 and TOF 4–44 h⁻¹ at 25 °C).

The nickel(i) complex **4** was used for the hydrodefluorination of fluorobenzenes.³⁰ It was found that in the presence of Et₂CHONa at 100 °C catalysts of the type L₂Ni are less efficient for hydrodehalogenation (TON 12 and TOF 4 h⁻¹ for L = IMes) than LNi (TON 18–31 and TOF 6–10 h⁻¹, respectively). Taken collectively, complexes **1–4** exhibit insufficient catalytic activities. However, the highest efficiencies for polyhalogenated arenes were obtained with the dimeric (IPrPdCl₂)₂ catalyst **5** (R = Dipp)³¹ (for the hydrodehalogenation of 1,2,4,5-tetrachlorobenzene in isopropanol at 80 °C TON 19 650 and TOF 819 h⁻¹ and for the reaction of 2,2',3,3'-tetrachlorobiphenyl with a TON of 10 000 and with a TOF 417 h⁻¹, respectively).

Given the foregoing information, sterically shielded 1,2,4-triazol-5-ylidenes were investigated as ligands for haloarene hydrodehalogenation catalysis. In the present contribution, we describe (i) the syntheses of new sterically shielded carbenes 1-alkyl-3,4-diaryl-1,2,4-triazol-5-ylidenes; (ii) the syntheses of mono- and biscarbene palladium halogenide complexes of the 1,2,4-triazol-5-ylidenes; (iii) the X-ray crystal structures of the sterically shielded carbene 1-*tert*-butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene; three PdI₂ and PdCl₂ carbene complexes bis-[1-*tert*-butyl-3-phenyl-4-(4-bromophenyl)-1,2,4-triazol-5-yliden]palladium chloride, bis-[1-*tert*-butyl-3-phenyl-4-(4-bromophenyl)-1,2,4-triazol-5-yliden]palladium iodide and the sterically shielded [1-*tert*-butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-yliden]palladium chloride; (iv) catalytic efficiencies of the foregoing complexes with respect to the hydrodechlorination of *p*-dichlorobenzene.





Fig. 1 Palladium and nickel catalysts for the hydrodehalogenation reactions (**1**, **2**, **5** R = Dipp (2,6-diisopropylphenyl); **3**, **4** R = Mes (2,4,6-trimethylphenyl)).

Results and discussion

Synthesis of carbenes and their palladium complexes

In order to obtain the desired carbene complexes, along with the known carbene **8a**,⁹ a series of new stable carbenes **8b–d** was synthesized in the following three stages: (1) syntheses of sterically hindered 4-substituted 3-phenyl-4-aryl-1,2,4-triazoles **6b,c** *via* the ring transformation of 2-phenyl-1,3,4-oxadiazole with the appropriate anilines, (2) quaternization of 1,2,4-triazoles **6b,c** using sterically hindered alkylhalogenides (*tert*-butyl iodide and 1-bromoadamantane), (3) deprotonation of



Scheme 1 Syntheses of carbenes **8a-d**. Reaction conditions: (a) ArNH_2 , TFA, *o*-DCB, 190 °C, 12 h; (b) (1) RX, AcOH, 120 °C, 10–18 h; (2) NaClO_4 ; (c) *t*-BuOK, PhMe, *i*-PrOH, r.t. Compounds: **6a** Ar = 4- BrC_6H_4 ; **6b** Ar = Mes; **6c** Ar = Dipp; **7a**, **8a** R = *t*-Bu; Ar = 4- BrC_6H_4 ; **7b**, **8b** R = 1-Ad; Ar = Mes; **7c**, **8c** R = *t*-Bu; Ar = Dipp; **7d**, **8d** R = 1-Ad; Ar = Dipp; Dipp = 2,6-*i*-Pr₂C₆H₃.




Fig. 2 Mono and biscarbene palladium complexes (**9a** X = Cl, R = *t*-Bu, Ar = *p*- BrC_6H_4 ; **9b** X = I, R = *t*-Bu, Ar = *p*- BrC_6H_4 ; **9c** X = I, R = Ad, Ar = Mes; **9d** X = I, R = *t*-Bu, Ar = Dipp; **10a** X = Cl, R = *t*-Bu, Ar = Dipp; **10b** X = I, R = *t*-Bu, Ar = Dipp; **10c** X = Cl, R = Ad, Ar = Dipp; **10d** X = I, R = Ad, Ar = Dipp).

the aforementioned salts **7b-d** using potassium *tert*-butoxide in a mixture of toluene and isopropanol akin to previously published work from our laboratory⁷ (Scheme 1), most of which (**8b-d**) include sterically hindered groups (*t*-Bu or 1-Ad in position 1 with Mes or Dipp in position 4).

The syntheses of the new biscarbene complexes **9a-d** (Fig. 2) were carried out *via* the direct reaction of the free carbenes **8a-d** with palladium halogenides (chloride and iodide) in a molar ratio of 2 : 1 in THF solution according to the pertinent literature method.³² An analogous method was used for the syntheses of the monocarbene complexes **10a-d** using the appropriate molar ratio of 1 : 1. For purposes of comparison, the known complex **5** was prepared *via* the reaction of the appropriate 1,3-diarylimidazol-2-ylidene with $\text{Pd}(\text{PhCN})_2\text{Cl}_2$ according to the literature procedure.³¹

Compounds **6a-c**, **7a-d**, **8a-d**, **9a-e**, **10a-d** were characterized by ¹H and ¹³C NMR spectroscopy, and their elemental compositions were established on the basis of elemental analysis. The ¹³C NMR spectra of the sparingly soluble complexes **9c,d** and **10b,d** were recorded in the solid state. In the case of the ¹H NMR spectra of triazoles **6b,c** with the exception of the specific signals for the mesityl or dipp groups, the resonances of the *meso*-protons CHN were evident (δ 8.73, 8.88 ppm in DMSO-*d*₆). In the case of the spectra of the triazolium salts **7a-d** the NMR signals were shifted significantly downfield (δ 10.61–10.91 ppm). It is noteworthy that two forms of the triazole **6c** were detected in CDCl₃ solution. The NMR spectrum of one of these forms was characterized on the basis of different signals for the methyl groups (δ 0.90, 1.10 ppm) and the presence of an upfield signal for the methine CH-groups (δ 2.37 ppm) of the corresponding isopropyl fragments. The

other form was evident from the presence of only one signal for the methyl groups (δ 1.19 ppm) and a downfield signal for the methine CH-groups (δ 3.02 ppm).

The *meso*-proton CHN resonances for these forms were found at 8.10 and 8.30 ppm, respectively. Presumably, the second form relates to a symmetrical structure with a close to 90° angle for the N(4)-aryl group. On the other hand, the alternate form could be a rotational form with a smaller bond angle. In DMSO-*d*₆ solution the second form converted to the first one (the pertinent signals were located at δ 0.88, 1.09, and 2.24 ppm, respectively). Presumably for the same reason, in DMSO-*d*₆ solution, salt **7c** shows only the first form of the isomers (δ 0.88, 1.09, and 2.24 ppm). However, the NMR spectrum of salt **7d** in CDCl₃ corresponds to only the first form (δ 0.91, 1.21, and 2.24 ppm, respectively). Interestingly, compounds **6b,c** were also obtained recently³³ by a similar method to the one described above. Following purification, only one form of this compound was isolated.

The most useful information, for carbenes **8a-d**, are the ¹³C NMR signals for the carbene carbon atom C(5) (δ 211.1–212.7 ppm). It was found that the NMR signals for complexes **9a-d** and **10a-d** are shifted upfield (δ 153.3–171.9 ppm). Interestingly, two closely spaced resonances were detected for compound **9b** in CDCl₃ solution. These resonances can be assigned to the *trans-trans* and *cis-trans* isomers of **9b** (the first designation relates to the disposition of the rotating triazole rings with respect to each other and the second one relates to the disposition of the halogen atoms at the metal center). The first isomer was isolated in a pure state and subsequently characterized on the basis of a single crystal X-ray diffraction study. The ¹H and ¹³C NMR spectra for complex **10a** in CDCl₃ solution exhibited two forms, presumably due to rotation of the N(4)-aromatic ring. The presence of two different signals of the *tert*-butyl protons confirmed this interpretation (δ 2.18 and 2.29 ppm). However, only one methyl signal was evident in the ¹³C NMR spectrum (δ 31.4 ppm), along with one signal for the *ipso*-C atom (δ 64.3 ppm). Furthermore, the presence of two groups of methyl and CH-resonances is evident in the ¹H NMR spectra (δ 0.54–0.70 ppm and 1.16–1.57 ppm, and 2.50, 2.61 ppm, respectively). These resonances indicate the existence of non-equivalent isopropyl and methyl groups in both isomers. However, only one isomer was detected in the single crystal X-ray structure (see below).

The solid state ¹³C NMR spectra of complexes **9c,d** and **10c,d** exhibited only one isomeric form for each compound (single resonances are apparent for the *i*-Pr CH, Ad *ipso*-C, C(3) and C(5)Pd atoms at δ 47.4–49.1, 64.1–65.6, 148.2–156.4, and 154.9–171.9 ppm, respectively). The C(5)Pd signals for the chloride complexes (**10a,c**) were shifted upfield (δ 153.3–154.9 ppm) relative to those for similar resonances (δ 167.0–171.9 ppm) that were observed for the iodide complexes (**9c,d** and **10b,d**). Four resonances were evident for the carbenoid carbon of complex **10b** and can be explained by the presence of four fixed rotational forms in the solid state. The ¹H NMR spectrum for **10b** in DMSO-*d*₆ solution indicates that only

two forms of the complex were present in solution (for the *i*-Pr group CH₃C δ 0.62, 1.36; CHC 2.94 ppm and CH₃C δ 0.83, 1.25; CHC 4.11 ppm and one signal of the *t*-Bu CH₃C group).

X-ray diffraction studies

The single crystals of compound **8c** were grown from toluene solution and studied subsequently by X-ray diffraction. The structure of **8c** is presented in Fig. 3 and the bond lengths (Å), bond orders, bond angles and dihedral angles (degrees) for structure **8c** are presented in Table 1. As can be seen from these data, the planar structure of **8c** is reminiscent of those for the 1,2,4-triazol-5-ylidenes.^{9,10,34} The bond angle at the carbene carbon atom is approximately 100.5° and the high bond order of 1.718 for the C(1)–N(3) bond implies a substantial ylide contribution to the carbene structure. Moreover, the C(2)–aromatic ring is twisted relative to the plane of the triazole ring by 36.3° (the bond order is 1.284), and furthermore the N(1)–aromatic ring is twisted almost orthogonally with respect to the triazole ring (87.4°) (the bond order is 1.139).

The single crystals of the palladium complexes **9a,b** and **10a** that were used for the X-ray diffraction studies were grown from acetonitrile. The crystal structures of **9a,b** and **10a** are displayed in Fig. 4–6 and the bond lengths, bond orders, bond angles and dihedral angles are presented in Tables 2–4.

Complexes **9a,b** adopt predominant *trans-trans* isomeric orientations. The bond orders for N(2)–C(1) in these molecules (*p* 1.770–1.782) closely resemble that of the initial carbene




Fig. 3 POV-Ray diagrams for 1-*tert*-butyl-3-phenyl-4-(2,6-diisopropyl-phenyl)-1,2,4-triazol-5-ylidene **8c** with thermal ellipsoids displayed at 50% probability. All hydrogen atoms have been removed for clarity.

Table 1 Selected bond lengths and angles for carbene **8c**

Bond	Length (Å)	Bond order	Angle	°
C(1)–N(3)	1.3491(14)	1.718	N(3)–C(1)–N(1)	100.53(9)
C(1)–N(1)	1.3843(15)	1.516	N(1)–C(2)–N(2)–N(3)	–0.48(11)
C(2)–N(1)	1.3860(14)	1.506	N(1)–C(1)–N(3)–C(21)	177.70(10)
C(2)–N(3)	1.3134(14)	1.923	N(1)–C(2)–C(9)–C(10)	36.3(6)
C(2)–C(15)	1.4781(15)	1.284	C(1)–N(1)–C(3)–C(4)	87.40(13)
C(3)–N(1)	1.4498(13)	1.139	—	—
N(2)–N(3)	1.3834(13)	1.336	—	—
N(3)–C(21)	1.4917(14)	0.898	—	—




Fig. 4 POV-Ray diagrams for bis-(1-*tert*-butyl-3-phenyl-4-bromophenyl)-1,2,4-triazol-5-yliden)palladium chloride **9a** with thermal ellipsoids displayed at 50% probability. All hydrogen atoms have been removed for clarity.




Fig. 5 POV-Ray diagrams for bis-(1-*tert*-butyl-3-phenyl-4-bromophenyl)-1,2,4-triazol-5-yliden)palladium iodide **9b** with thermal ellipsoids displayed at 50% probability. All hydrogen atoms have been removed for clarity.




Fig. 6 POV-Ray diagrams for 1-*tert*-butyl-3-phenyl-4-(2,6-diisopropyl-phenyl)-1,2,4-triazol-5-yliden)palladium chloride **10a** with thermal ellipsoids displayed at 50% probability. All hydrogen atoms and an acetonitrile molecule have been removed for clarity.

Table 2 Selected bond lengths and angles for complex **9a**

Bond	Length (Å)	Bond order	Angle	°
C(1)–N(2)	1.338(4)	1.782	N(1)–C(1)–N(2)	103.7(3)
C(1)–N(1)	1.368(5)	1.609	C(1)–Pd(1)–Cl(1)	89.70(10)
C(2)–N(3)	1.308(5)	1.954	Cl(1)–Pd(1)–Cl(1a)	180.00(5)
C(2)–N(1)	1.377(4)	1.557	N(3)–C(2)–N(1)–C(1)	–0.2(4)
C(2)–C(9)	1.469(5)	1.330	N(1)–C(1)–N(2)–N(3)	–1.7(4)
C(3)–N(1)	1.443(4)	1.178	N(1)–C(2)–C(9)–C(10)	36.3(6)
N(2)–N(3)	1.386(4)	1.323	C(4)–C(3)–N(1)–C(1)	69.3(5)
N(2)–C(15)	1.500(5)	0.851	Pd(1)–C(1)–N(1)–C(2)	–174.1(2)
C(6)–Br(1)	1.895(4)	—	Pd(1)–C(1)–N(2)–C(15)	–4.4(6)
C(1)–Pd(1)	2.025(4)	—	N(1)–C(1)–Pd(1)–Cl(1)	86.0(3)
Pd(1)–Cl(1)	2.3301(10)	—	N(2)–C(1)–Pd(1)–Cl(1)	92.4(4)

Table 3 Selected bond lengths and angles for complex **9b**

Bond	Length (Å)	Bond order	Angle	°
C(1)–N(2)	1.340(7)	1.770	N(1)–C(1)–N(2)	103.4(4)
C(1)–N(1)	1.368(7)	1.609	C(1)–Pd(1)–I(1)	90.02(15)
C(2)–N(3)	1.308(7)	1.954	I(2)–Pd(1)–I(1)	176.59(2)
C(2)–N(1)	1.386(7)	1.506	N(3)–C(2)–N(1)–C(1)	–0.2(6)
C(2)–C(9)	1.474(7)	1.305	N(1)–C(1)–N(2)–N(3)	0.8(6)
C(3)–N(1)	1.435(7)	1.224	N(1)–C(2)–C(9)–C(10)	12.3(10)
N(2)–N(3)	1.376(6)	1.374	C(4)–C(3)–N(1)–C(1)	69.3(5)
N(2)–C(15)	1.491(7)	0.902	Pd(1)–C(1)–N(1)–C(2)	–179.7(4)
C(6)–Br(1)	1.907(5)	—	Pd(1)–C(1)–N(2)–C(15)	–6.3(9)
C(1)–Pd(1)	2.034(5)	—	N(1)–C(1)–Pd(1)–I(1)	85.8(5)
Pd(1)–I(1)	2.6014(7)	—	N(2)–C(1)–Pd(1)–I(1)	–97.6(5)

Table 4 Selected bond lengths and angles for complex **10a**

Bond	Length (Å)	Bond order	Angle	°
C(1)–N(2)	1.329(4)	1.833	N(1)–C(1)–N(3)	104.6(2)
C(1)–N(1)	1.366(4)	1.621	C(1)–Pd(1)–Cl(1)	92.36(8)
C(2)–N(3)	1.384(4)	1.517	Cl(1)–Pd(1)–Cl(2)	173.88(3)
C(2)–N(1)	1.309(4)	2.142	C(1)–N(1)–N(2)–C(2)	0.4(3)
N(1)–N(2)	1.375(3)	1.379	N(3)–C(1)–N(1)–N(2)	0.1(3)
C(2)–C(7)	1.482(4)	1.264	C(19)–C(14)–N(3)–C(1)	102.1(3)
N(3)–C(14)	1.457(4)	1.098	N(3)–C(2)–C(7)–C(8)	151.0(3)
C(3)–N(1)	1.514(4)	0.770	Pd(1)–C(1)–N(3)–C(2)	175.50(19)
C(1)–Pd(1)	1.963(3)	—	Pd(1)–C(1)–N(2)–C(15)	–6.3(9)
Pd(1)–Cl(1)	2.3078(8)	—	N(1)–C(1)–Pd(1)–Cl(1)	–84.3(3)
Pd(1)–N(4)	2.075(3)	—	N(1)–C(1)–Pd(1)–Cl(2)	89.6(3)

(1.799).¹⁰ For comparison, similar features were observed for the linear bistriazolylidene complexes of copper(i) chloride^{9,16} and the related bistriazolylidene chelate complex of copper(i) iodide.¹¹

The dihedral angles of the aromatic rings in molecules **9a,b** are also close to that of the initial carbene. The C(2)-nucleus is twisted relative to the triazole ring by 36.3° (**9a**) and 12.3° (**9b**), respectively *versus* 29.9° for the initial carbene.

Furthermore, the N(1) nucleus is twisted with respect to the same plane by 69.3° (**9a,b**) *versus* 51.1° for the initial carbene. The bond orders for C(2)–C(9) and N1–C(3) are 1.31–1.33 and 1.18–1.22, respectively, since the C(2) nucleus is naturally more conjugated with the triazole ring than with the N(1)-nucleus.

The bond angles at the carbenoid carbon atom (103°) are significantly larger than those for the carbene (100.3°) and are noticeably smaller than that observed for the salt complex (110°).^{7,34} Furthermore, the Pd–C bond distances (2.025, 2.034 Å) are notably larger than the sum of the covalent radii for these atoms (1.95 Å) and significantly larger than the sum of covalent radius of the carbon atom and the ionic radius of palladium(II) (1.75 Å) that is indicative of a C–Pd coordination bond in these complexes.

The Pd–Cl bond distances in complex **9a** (2.330 Å) are intermediate in length between those of covalent and ionic bonds (the sum of the covalent radii is 2.19 Å, while the sum of the ionic radii is 2.67 Å). Overall, however it is closer to a covalent bond (the extent of Pd(1)–Cl(1) ionization is approximately 29%). In the case of complex **9b**, the Pd–I bond (2.60 Å) is also intermediate in character with respect to covalency *vs.* ionicity (the sum of the covalent radii is 2.53 Å and the sum of the ionic radii is 3.06 Å). Overall, however the Pd–I bond more closely resembles a covalent bond (the extent of Pd(1)–I(1) ionization is 13%). Thus, complexes **9a,b** possess typical structures with C–Pd coordination bonds and shortened Pd–halogen bonds.

Compound **10a** is a monocarbene–acetonitrile palladium complex. The bond order of the N(1)–C(1) bond in this molecule (*p* 1.833) is larger than that of the initial carbene (1.718). The Pd–C bond distance (1.963 Å) is close to the sum of the covalent radii of the constituent atoms (1.95 Å) and significantly larger than the sum of the covalent radius of the carbon atom and the ionic radius of palladium(II) (1.75 Å). The latter observation, along with the bond orders, indicates that there is a stronger interaction between the carbene carbon and the palladium atoms than in the case of the biscarbene complexes **9a,b**. Moreover, the Pd–Cl bond in the structure of **10a** (2.308 Å) is intermediate in length between covalent and ionic bonds (the sum of the covalent radii is 2.19 Å and the sum of the ionic radii is 2.67 Å), but closer to that of a covalent bond (the extent of Pd(1)–Cl(1) ionization is approximately 25%).

The bond angle of the carbenoid carbon atom (103.4°) is significantly larger than that of the initial carbene (100.5°) however the salt carbenoid angle is larger by comparison (110°).^{7,34} Furthermore, the C(2) nucleus is twisted relative to that of the triazole ring by 151.0° as opposed to 29.9° for the initial carbene. Furthermore, the N(3)-nucleus is twisted with respect to the triazole ring by 102.1° *versus* 51.1° for the same carbene. The bond orders for C(2)–C(7) and N(3)–C(14) are 1.264 and 1.098, respectively. Thus, complex **10a** adopts a typical structure with a strong C–Pd coordination bond and a shortened bond Pd–Cl.

Catalytic tests

Test reactions for the hydrodechlorination catalysis were carried out by heating a reaction mixture of the haloarene with potassium or sodium isopropoxides, which had been generated from potassium or sodium *tert*-butoxide or hydroxide, in the presence of a catalyst in isopropanol at 80 °C. Control of the conversion (on 1 chlorine atom) was calculated according to the masses of the sodium or potassium chlorides, both of

Table 5 Catalytic efficiencies of carbene complexes **9a–d**, **10a–d**, **5** for the reaction of *p*-dichlorobenzene with potassium *tert*-butoxide in isopropanol at 80 °C

Catalyst	Catalyst amount (mol%)	Reaction time (h)	Conv. (%)	TON per 1 Cl	TOF (h ⁻¹)
9a	0.1	8	25	500	62
9b	0.1	8	11	220	28
9c	0.1	8	70	1400	175
9d	0.1	8	38	760	95
10a	0.1	8	66	1320	165
10b	0.1	8	58	1160	145
10c	0.1	8	72	1140	180
10c^a	0.01	24	40	8000	334
10d	0.01	24	100	20 000	840
5	0.01	8	63	12 600	1560
5	0.01	24	88	17 600	734

^aThe reagent is potassium *tert*-butoxide to which 1 mol% lithium *tert*-butoxide has been added.

Table 6 Catalytic efficiencies of carbene complexes **9d**, **10b,c**, **5** for the reaction of *p*-dichlorobenzene with sodium *tert*-butoxide in isopropanol at 80 °C

Catalyst	Catalyst amount (mol%)	Reaction time (h)	Conv. (%)	TON per 1 Cl	TOF (h ⁻¹)
9d	0.01	24	53	10 600	440
10b	0.01	24	36	7200	300
10c	0.01	6	77	15 400	2230
5	0.01	24	32	6400	266

Table 7 Catalytic efficiencies of carbene complexes **9d**, **10a–d** for the reaction of *p*-dichlorobenzene with sodium hydroxide in isopropanol at 80 °C

Catalyst	Catalyst amount (mol%)	Reaction time (h)	Conv. (%)	TON per 1 Cl	TOF (h ⁻¹)
10c	0.1	8	100	2000	250
10c	0.01	24	34	6800	284
10d	0.01	24	21	4200	176
5	0.01	24	96	19 200	800

which are insoluble in isopropanol. The most efficient catalysis trials were examined on the basis of gas chromatography. The catalytic efficiencies were estimated using the values of TON and TOF.

A summary of the results pertaining to the studies of the catalytic efficiencies for the carbene complexes **5**, **9a–d**, and **10a–d** with respect to the model compound *p*-dichlorobenzene and a variety of bases is presented in Tables 5–7.

Examination of Table 5 reveals that the hydrodehalogenation of *p*-dichlorobenzene with potassium *tert*-butoxide proceeds efficiently with the majority of the carbene catalyst complexes that were tested. The most rapid reaction was observed with the triazolylidene monocarbene complex **10d** (after 24 h of the reaction the maximum TON and TOF values reached 20 000 and 840 h⁻¹ respectively). The efficiency of

complex **10c** is relatively low (TON 1140 cycles after 8 h), however, in the presence of 1 mol% lithium *tert*-butoxide the TON is enhanced up to 8000 cycles after 24 h.

When sodium *tert*-butoxide was used for these hydrodehalogenation reactions (Table 6), a significant increase in the efficiency indices by almost an order of magnitude was observed for the triazolylidene complexes **9d**, **10b,c** (TON 7200–15 400, TOF 300–2230 h⁻¹). In fact, these values exceeded those for the etalon complex **5** (TON 6400, TOF 266 h⁻¹).

The hydrodechlorination reaction was examined in the presence of sodium hydroxide and isopropanol using complexes **5** and **10c,d**. The use of sodium hydroxide is particularly appropriate in view of the possible development of a technology for haloarene (production wastes) neutralization (detoxification) since it is an inexpensive reagent. It was found (Table 7) that the triazolylidene complexes **10c,d** exhibited high efficiencies for the reaction of *p*-dichlorobenzene with sodium hydroxide (TON 2000–6800, TOF 176–284 h⁻¹). However, for **10d** this represents threefold decrease with respect to the use of potassium *tert*-butoxide.

Interestingly, the known complex **5** appeared to be somewhat more effective for the reaction with sodium hydroxide (TON 19 200, TOF 800 h⁻¹), than those with potassium *tert*-butoxide (TON 17 600, TOF 734 h⁻¹) and sodium *tert*-butoxide (TON 6400, TOF 266 h⁻¹).


On the basis of gas chromatographic data, noticeable quantities of haloarenes were not observed in the experiments that resulted in quantitative conversion of the initial haloarene.

In conclusion, the highest efficiency is apparent for the catalysts that feature the sterically bulky 1-(1-adamantyl)-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene ligand.

The high TON of 20 000 cycles (**10d**) with potassium *tert*-butoxide and TON 15 400 with sodium *tert*-butoxide exceed the efficiency level of complex **5** (TON 17 600 and 6400, respectively). Another factor that contributes to the catalytic efficiency is the presence of iodide as the second ligand which raises the efficiency of the catalyst (TON 20 000 cycles for **10d**, TON 8000 cycles for **10c**).

As far as we know, the mechanism of haloarene (*i.e.* *p*-dichlorobenzene) hydrodehalogenation^{2,3} involves the formation of palladium(0) complex (**D**) as the first step *via* the interaction of the palladium(II) complex with metal isopropoxide, followed by reaction with the haloarene **E** thereby affording the palladium(II) species (**F**) (Scheme 2).

The activated complex **F** reacts subsequently with the metal isopropoxide *via* the nucleophilic substitution of the halogen, thus forming the adduct of the H-arene with palladium species **H** due to decomposition of the previously formed isopropoxide complex **G**. Finally, decomposition of the latter produces H-arene **I** and the palladium(0) complex **D**. The enthalpy changes that take place during chlorobenzene conversion catalyzed by complex **10a** were evaluated on the basis of DFT, B3LYP/5, 3-21G calculations (Scheme 2, in kcal mol⁻¹). These calculations indicate that a strong exothermic reaction takes place between palladium(0) compound **D** and chlorobenzene



Scheme 2 Mechanism of haloarene hydrodehalogenation.

($\Delta H -48.0 \text{ kcal mol}^{-1}$) and that the conversion of compound **G** to **H** is endothermic ($\Delta H +30.7 \text{ kcal mol}^{-1}$). An important stage in the foregoing process may be the interaction of the initial precatalyst **A** with the metal isopropoxide ($\Delta H -16.0 \text{ kcal mol}^{-1}$), which is the initial stage in the formation of these molecular complexes. As might be expected, the stabilities of these complexes depend on steric protection of the palladium complex.

A possible reason for the increase in catalytic efficiency for sterically shielded carbene complexes, particularly **10c,d**, is attributed to the steric protection which results in an increased stability of the intermediate Pd(0) species in the reaction medium.

Conclusions

Along with the syntheses of the new biscarbene palladium complexes **9a,b** from the known carbene **8a**, a series of sterically shielded 1,3,4-trisubstituted 1,2,4-triazol-5-ylidenes **8b-d** and their palladium complexes **9c,d**, **10a-d** have been synthesized in the present work. It was found that sterically shielded compounds (**9c,d**, **10a-d**) are effective catalysts for *p*-dichlorobenzene hydrodechlorination. The efficiencies for the monocarbene complexes **10c,d** were shown to be somewhat higher than those for the biscarbene complexes **9c,d**. However, the latter results are strongly dependent on the nature of the reagent.

Experimental section

General

^1H NMR and ^{13}C NMR spectra were recorded using a Bruker Avance II 400 spectrometer (400 MHz for ^1H NMR spectra and 100 MHz for ^{13}C NMR spectra) in DMSO- d_6 , C_6D_6 or CDCl_3 solution or in the solid state. The ^1H NMR and ^{13}C NMR chemical shifts are reported relative to tetramethylsilane (TMS) (solution) and sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS) (solid state). The X-ray diffraction data were collected

using a Rigaku AFC12 Saturn 724+ CCD diffractometer equipped with a Rigaku XStream low temperature device that is operated at 100 K, a Rigaku SCX-Mini Diffractometer equipped with a Rigaku XStream low temperature device that is operated at 100 K, or a Nonius-Kappa CCD diffractometer with an Oxford cryostream 600 that is operated at 153 K. All instruments used a graphite-monochromated Mo $K\alpha$ radiation source ($\lambda = 0.71075 \text{ \AA}$). Corrections were applied for Lorentz and polarization effects. The structures were solved by direct methods and refined by full-matrix least-squares cycles on F^2 using the Siemens SHELXTL PLUS 5.0 (PC) software package³⁵ and PLATON.³⁶ All non-hydrogen atoms were refined anisotropically. The hydrogen atoms were placed in fixed, calculated positions using a riding model.

Thin-layer chromatography was performed on silica gel with chloroform or a 10 : 1 mixture of chloroform and methanol as the eluent, followed by development with iodine. Elemental analyses were carried out at the Analytical Laboratory of the Litvinenko Institute of Physical Organic and Coal Chemistry. The quantum chemical calculations were performed with the help of Firefly 8.0 program.

3,4-Diaryl-1,2,4-triazoles (**6b,c**) (general procedure)

3,4-Diaryl-1,2,4-triazoles were obtained according to the modified procedure of the work³⁷ by the ring transformation of 2-phenyl-1,3,4-oxadiazole with the respective anilines in the presence of trifluoroacetic acid at 190 °C. The mixture of aromatic amine (20 mmol) and trifluoroacetic acid (20 mmol) in *o*-dichlorobenzene (1–2 mL) was added to 2-phenyl-1,3,4-oxadiazole (20 mmol) and heated at 190 °C for 12 h. The resulting oil or crystalline product was washed 5–6 times with petroleum ether (50 mL). Then 10% aqueous solution of potassium hydroxide was added and the mixture was stirred for 15–20 min. The resulting crystalline precipitate was filtered off, washed with water, then with a mixture of petroleum ether–isopropanol (3 : 1) and petroleum ether and dried. The obtained products were recrystallized from benzene.

3-Phenyl-4-(2,4,6-trimethylphenyl)-1,2,4-triazole (**6b**)

Yield 2.63 g, 50%, mp 125–126 °C (benzene). δ_{H} (400 MHz; CDCl_3 ; Me_4Si): 1.86 (s, 6H, *o*- CH_3 C), 2.31 (s, 3H, *p*- CH_3 C), 7.09s (2H, Ar), 7.39s (5H, Ar), 8.73s (1H, C^5HN). Compound **6b** was identical to the product of the same reaction in ionic liquids isolated by column chromatography³³ (mp 125–126 °C).

3-Phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazole (**6c**)

Yield 3.05 g, 50%, mp 98–100 °C (benzene). Found: C, 78.7; H, 7.4; N, 13.9. Calc. for $\text{C}_{20}\text{H}_{23}\text{N}_3$: C, 78.7; H, 7.6; N, 13.8%. δ_{H} (400 MHz; DMSO- d_6 ; Me_4Si): 0.88 (d, 6H, CH_3C , *J* 6.8 Hz), 1.09 (d, 6H, CH_3C , *J* 6.8 Hz), 2.24 (qr, 2H, CHC , *J* 6.8 Hz), 7.25–7.42 (m, 7H, Ar), 7.58 (m, 1H, Ar), 8.88 (s, 1H, C^5HN). δ_{H} (400 MHz; CDCl_3 ; Me_4Si): 1 form: 0.90 (d, 6H, CH_3C , *J* 6.8 Hz), 1.10 (d, 6H, CH_3C , *J* 6.8 Hz), 2.37 (qr, 2H, CHC , *J* 6.8 Hz) (*i*-Pr), 8.10 (s, 1H, C^5HN); 2 form: 1.19 (d, 12H, CH_3C), 3.02 (qr, 2H, CHC , *J* 6.8 Hz) (*i*-Pr); 8.30 (s, 1H, C^5HN); general for both forms: 7.20–7.37 (m, 12H, Ar), 7.45 (d, 3H, Ar, *J* 7.6 Hz), 7.51 (dd, 1H,

Ar, *J* 7.6 Hz). The first form of compound **6c** was identical to the product of the same reaction in ionic liquids isolated by column chromatography³³ (mp 114–115 °C).

1-Alkyl-3,4-diaryl-1,2,4-triazolium perchlorates (7b–d) (general procedure)

The mixture of sodium iodide (50–60 mmol) and 1.5-fold excess of *tert*-butyl chloride in acetic acid (10 mL) was boiled for 1.5 h. One equivalent (to iodide amount) of the respective 3,4-diaryl-1,2,4-triazole **6a,c** was added and boiled for 21 h. Then water (20 mL) and sodium sulfite were added to the obtained mixture to discolor the solution. The mixture was diluted by water (0.5 L) and activated carbon (1–3 g) was added. The subsequent mixture was heated to boiling and filtered hot. Sodium perchlorate was added in an amount exceeding equivalence by 10–20%. The precipitate that formed was filtered off, washed with isopropanol and ether, recrystallized from the respective solvent, and subsequently dried. The salts **7b,d** were obtained similarly using 1-bromoadamantane instead of *tert*-butyl iodide. The heating time was 10–18 h.

1-(1-Adamantyl)-3-phenyl-4-(2,4,6-trimethylphenyl)-1,2,4-triazolium perchlorate (7b)

The heating time was 10 h. Yield 85%, mp > 300 °C (acetic acid). Found: C, 65.0; H, 6.5; Cl, 7.1; N, 8.5. Calc. for C₂₇H₃₂ClN₃O₄: C, 65.1; H, 6.5; Cl, 7.1; N, 8.4%. δ_H (400 MHz; CDCl₃; Me₄Si): 1.79 (s, 6H, Ad), 2.02 (s, 6H, CH₃C), 2.30 (s, 3H, Ad), 2.35 (s, 3H, *p*-CH₃C), 2.36 (s, 6H, *o*-CH₃C), 7.19 (s, 2H, Ar), 7.44 (dd, 2H, Ar, *J* 7.6 Hz), 7.51 (dd, 2H, Ar, *J* 8.0 Hz), 7.62 (dd, 1H, Ar, *J* 7.2 Hz), 10.59 (s, 1H, C⁵HN).

1-*tert*-Butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazolium perchlorate (7c)

Yield 64%, mp 232–234 °C (acetic acid–water, 1 : 1). Found: C, 62.5; H, 6.9; Cl, 7.6; N, 9.2. Calc. for C₂₄H₃₂ClN₃O₄: C, 62.4; H, 7.0; Cl, 7.7; N, 9.1%. δ_H (400 MHz; DMSO-*d*₆; Me₄Si): 0.91 (d, 6H, *J* 6.8 Hz), 1.16 (d, 6H, *J* 6.8 Hz) (CH₃C, *i*-Pr), 1.81 (s, 9H, CH₃C, *t*-Bu), 2.40 (qr, 2H, CHC, *i*-Pr, *J* 6.8 Hz), 7.44 (d, 2H, *J* 7.6 Hz), 7.51 (d, 2H, *J* 7.6 Hz), 7.54 (d, 2H, *J* 7.6 Hz), 7.63 (dd, 1H, *J* 7.6 Hz), 7.73 (dd, 1H, *J* 7.6 Hz) (Ar), 10.91 (s, 1H, C⁵HN).

1-(1-Adamantyl)-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazolium perchlorate (7d)

The heating time was 18 h. Yield 59%, mp 210–212 °C (acetic acid–water, 1 : 1). Found: C, 66.8; H, 7.0; Cl, 6.6; N, 7.7. Calc. for C₃₀H₃₈ClN₃O₄: C, 66.7; H, 7.1; Cl, 6.7; N, 7.8%. δ_H (400 MHz; CDCl₃; Me₄Si): 0.91 (d, 6H, *J* 4.4 Hz), 1.28 (d, 6H, *J* 4.4 Hz) (CH₃C, *i*-Pr), 1.87 (m, 6H, Ad), 2.23 (qr, 2H, CHC, *i*-Pr, *J* 4.4 Hz), 2.42 (s, 3H, Ad), 2.48 (s, 6H, Ad), 7.37 (m, 4H), 7.42 (m, 2H), 7.51 (m, 1H), 7.63 (m, 1H) (Ar), 10.28 (s, 1H, C⁵HN).

1-Alkyl-3,4-diaryl-1,2,4-triazol-5-ylidenes (8b–d) (general procedure)

A suspension of 1-*tert*-butyl-3,4-diaryl-1,2,4-triazolium perchlorate **7a–d** and potassium *tert*-butoxide (95% from the calculated amount) was added to the mixture of anhydrous

isopropanol and toluene (1 : 3) or in THF and stirred for 1.5 h. The solvent was evaporated and the oil was obtained. This oil was then stirred with petroleum ether and the subsequent precipitate that was formed was filtered off and dried. Then, anhydrous toluene (10–20 mL) was added to the precipitate and stirred for 1 h. The resulting inorganic salt was filtered off. The mother liquor was evaporated and the resulting product was crystallized by stirring with petroleum ether. The precipitate was filtered off, washed several times with petroleum ether, dried, and recrystallized from toluene.

1-(1-Adamantyl)-3-phenyl-4-(2,4,6-trimethylphenyl)-1,2,4-triazol-5-ylidene (8b)

Yield 50%, mp 168–169 °C (toluene). Found: C, 81.7; H, 7.8; N, 10.5%. Calc. for C₂₇H₃₁N₃: C, 81.6; H, 7.9; N, 10.6%. δ_H (400 MHz; C₆D₆; Me₄Si): 1.64 (m, 3H, Ad), 1.73 (m, 3H, Ad), 1.97 (s, 6H, *o*-CH₃C), 2.06 (s, 3H, *p*-CH₃C), 2.13 (m, 3H, Ad), 2.66 (m, 6H, Ad), 6.69 (s, 2H, Ar), 6.94 (s, 3H, Ar), 7.61 (d, 2H, Ar, *J* 3.6 Hz). δ_C (100 MHz, C₆D₆, Me₄Si): 18.3 (*o*-CH₃C), 21.0 (*p*-CH₃C), 30.2 (CHC, Ad), 36.7 (CH₂C, Ad), 44.2 (CH₂C, Ad), 59.4 (*ipso*-C, *t*-Bu), 127.7, 128.4, 128.6, 129.5, 130.2 (Ar), 135.5, 136.8, 137.9 (*ipso*-C, Ar), 151.1 (C³), 211.5 (C⁵).

1-*tert*-Butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene (8c)

Yield 70%, mp 112–114 °C (toluene). Found: C, 79.8; H, 8.5; N, 11.7. Calc. for C₂₄H₃₁N₃: C, 79.7; H, 8.6; N, 11.6%. δ_H (400 MHz; C₆D₆; Me₄Si): 0.85 (d, 6H, *J* 7.2 Hz), 1.18 (d, 6H, *J* 7.2 Hz) (CH₃C, *i*-Pr), 1.83 (s, 9H, CH₃C, *t*-Bu), 2.74 (qr, 2H, *J* 7.2 Hz, CHC, *i*-Pr), 6.90 (m, 3H), 7.10 (d, 2H, *J* 7.6 Hz), 7.24 (dd, 1H, *J* 7.6 Hz), 7.61 (m, 2H) (Ar). δ_C (100 MHz; C₆D₆; Me₄Si): 22.5, 24.5 (CH₃C, *i*-Pr), 29.0 (CH, *i*-Pr), 30.9 (CH₃C, *t*-Bu), 59.3 (*ipso*-C, *t*-Bu), 128.2, 128.5, 129.5, 129.6 (Ar), 124.2, 136.6, 146.1 (*ipso*-C, Ar), 151.8 (C³), 213.0 (C⁵).

1-(1-Adamantyl)-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene (8d)

Yield 68%, mp 151–153 °C (toluene). Found: C, 82.1; H, 8.4; N, 9.6. Calc. for C₃₀H₃₇N₃: C, 82.0; H, 8.5; N, 9.6%. δ_H (400 MHz; C₆D₆; Me₄Si): 0.87 (d, 6H, *J* 6.8 Hz), 1.20 (d, 6H, *J* 6.8 Hz) (CH₃C, *i*-Pr), 1.64 (m, 3H), 1.71 (m, 3H), 2.12 (m, 3H), 2.63 (m, 6H) (Ad), 2.76 (qr, 2H, CHC, *i*-Pr, *J* 6.8 Hz), 6.94 (m, 3H), 7.12 (d, 2H, *J* 7.6 Hz), 7.26 (dd, 1H, *J* 7.6 Hz), 7.62 (m, 2H) (Ar). δ_C (100 MHz, C₆D₆, Me₄Si): 22.5, 24.5 (CH₃C, *i*-Pr), 29.4 (CH, *i*-Pr), 30.5 (CHC), 36.6, 44.6 (CH₂C) (Ad), 59.5 (*ipso*-C, Ad), 124.2, 128.1, 128.5, 129.4, 129.5 (Ar), 127.6, 136.8, 146.3 (*ipso*-C, Ar), 151.5 (C³), 211.8 (C⁵N).

trans-trans-Bis(1-*tert*-butyl-3-phenyl-4-*p*-bromophenyl-1,2,4-triazol-5-ylidene)palladium chloride (9a)

A powder of carbene **8a**⁹ (2 g, 0.56 mmol) and anhydrous dimethylsulfoxide (0.03 mL) were added to the suspension of palladium chloride (0.05 g, 0.28 mmol) in THF (5 mL) and stirred for 0.5 h. The suspension of complex **9a** was evaporated to dryness and stirred with petroleum ether. The precipitate was filtered off and dried. Yield 0.25 g (100%), mp 160–163 °C

(subl.) (acetonitrile). Found: C, 48.7; H, 4.0; Br, 18.3; Cl, 8.1; N, 9.4. Calc. for $C_{36}H_{36}Br_2Cl_2N_6Pd$: C, 48.6; H, 4.1; Br, 18.0; Cl, 8.0; N, 9.4%. δ_H (400 MHz; $CDCl_3$; Me_4Si): 1.65 (s, 18H, CH_3C), 7.26 (m, 10H), 7.53 (m, 4H), 7.69 (m, 4H) (Ar). δ_C (100 MHz, $CDCl_3$, Me_4Si): 30.6 (CH_3C), 61.9 (CH_3C), 128.5, 128.6, 130.3, 131.6, 132.9 (Ar), 124.0, 125.1, 136.3 (*ipso-C*), 151.8 (C^3), 171.3 (C^5Pd).

Bis[1-*tert*-butyl-3-phenyl-4-*p*-bromophenyl-1,2,4-triazol-5-ylidene]palladium iodide (9b)

The same procedure that was used for complex **9a** was applied to the synthesis of **9b**. Yield 75%, mp 119–120 °C (acetonitrile). Found: C, 40.4; H, 3.4; Br, 14.8; I, 23.6; N, 7.9. Calc. for $C_{36}H_{36}Br_2I_2N_6Pd$: C, 40.3; H, 3.4; Br, 14.9; I, 23.7; N, 7.8%. The product was a mixture of two isomers *trans-trans*- and presumably *cis-trans*-forms:

For the *trans-trans*-isomer: δ_H (400 MHz, $CDCl_3$, Me_4Si): 1.58 (s, 18H, CH_3C), 7.24 (m, 10H), 7.61 (m, 8H, Ar). δ_C (100 MHz; $CDCl_3$; Me_4Si): 30.9 (CH_3C), 62.2 (CH_3C), 128.6, 128.7, 130.5, 132.4, 132.8 (Ar), 124.0, 125.3, 136.3 (*ipso-C*), 152.5 (C^3), 168.7 (C^5Pd).

For the *cis-trans*-isomer: δ_H (400 MHz; $CDCl_3$; Me_4Si): 2.11 (s, 18H, CH_3C), 7.01 (m, 4H), 7.33 (m, 14H); δ_C (100 MHz; $CDCl_3$; Me_4Si): 31.9 (CH_3C), 63.0 (CH_3C), 128.5, 128.6, 130.3, 131.4, 132.0 (Ar), 123.4, 125.5, 136.1 (*ipso-C*), 152.8 (C^3), 167.7 (C^5Pd).

The *trans-trans*-isomer was isolated from the mixture by additional recrystallization from acetonitrile and identified by X-ray diffraction study.

Bis-[1-(1-adamantyl)-3-phenyl-4-(2,4,6-trimethylphenyl)-1,2,4-triazol-5-ylidene]palladium iodide (9c)

The powder carbene **8b** (0.4 g, 1 mmol) was added to a solution of palladium iodide (0.18 g, 0.5 mmol) in THF (4 mL), the mixture was stirred for 30 h and boiled for 2 h. The precipitate formed was filtered off and dried. Yield 0.42 g (72%), mp > 300 °C (dimethylformamide). Found: C, 56.3; H, 5.3; I, 21.9; N, 7.4%. Calc. for $C_{54}H_{62}I_2N_6Pd$: C, 56.1; H, 5.4; I, 22.0; N, 7.3%. δ_C (100 MHz; solid state; DSS): 21.3, 23.7 (*o-CH_3C*), 24.6 (*p-CH_3C*), 30.9 (CHC, Ad), 37.8, 45.2 (CH_2C , Ad), 64.1 (*ipso-C*, Ad), 127.7, 130.0, 131.2, 133.3, 135.6, 139.8, 140.7 (Ar), 154.8 (C^3), 168.2 (C^5Pd).

Bis[1-*tert*-butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene]palladium iodide (9d)

The powder palladium iodide (0.12 g, 0.34 mmol) and dimethylsulfoxide (0.04 mL) were added to a solution of carbene **8c** (0.25 g, 0.68 mmol) in toluene (5 mL), stirred at room temperature for 8 h and boiled for 2 h. The precipitate that formed was filtered off, washed with petroleum ether, and dried. Yield 0.29 g (78%), mp 260–261 °C (acetonitrile). Found: C, 53.3; H, 5.8; I, 23.3; N, 7.7%. Calc. for $C_{48}H_{62}I_2N_6Pd$: C, 53.2; H, 5.8; I, 23.4; N, 7.8%. δ_C (100 MHz; solid state; DSS): 25.5, 27.1, 27.8 (CH_3C , *i-Pr*), 31.6 (CH_3C , *t-Bu*), 47.4, 48.1 (CHC, *i-Pr*), 64.3 (CH_3C , *t-Bu*), 126.4, 127.2, 128.9, 130.1, 132.1, 132.8 (Ar), 134.0, 146.7, 147.4 (*ipso-C*), 154.9 (C^3), 171.9 (C^5Pd).

[1-*tert*-Butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene]palladium chloride (10a)

A mixture of carbene **8c** (0.25 g, 0.68 mmol) and palladium chloride (0.12 g, 0.68 mmol) in anhydrous dimethylsulfoxide (2.5 mL) was stirred for 3 days. The precipitate that formed was filtered off, washed by petroleum ether, and dried. Dichloromethane (5 mL) and water (5 mL) were added to the mother liquor and stirred again. The phases that formed were then subsequently separated. The dichloromethane phase was then dried by anhydrous sodium sulfate. The solvent was evaporated; the oil that formed was stirred with petroleum ether. The precipitate that formed was filtered off and dried. Yield 0.15 g (41%), mp 196–198 °C (acetonitrile). Found: C, 53.4; H, 5.9; Cl, 13.2; N, 7.7%. Calc. for $C_{24}H_{31}Cl_2N_3Pd$: C, 53.5; H, 5.8; Cl, 13.2; N, 7.8%. δ_H (400 MHz; $CDCl_3$; Me_4Si): 1 form: 0.54 d, 0.63 d, 1.15 d, 1.30 d (12H, CH_3C , *J* 4.8 Hz, *i-Pr*), 2.29 (9H, *t-Bu*), 2.50 (m, 2H, CHC, *i-Pr*); 2 form: 0.57 d, 0.71 d, 1.38 d, 1.53 d (12H, CH_3C , *J* 4.8 Hz, *i-Pr*), 2.18 (9H, *t-Bu*), 2.61 (m, 2H, CHC, *i-Pr*); a general part of two forms: 7.17 (m, 4H), 7.26–7.48 (m, 3H), 7.26 (m, 1H) (Ar). δ_C (100 MHz, $CDCl_3$, Me_4Si): 25.1, 25.4, 28.4, 28.6, 28.7 (CH_3C , *i-Pr*), 31.0 (CH_3C , *t-Bu*), 24.3, 24.5 (CH, *i-Pr*), 64.3 (CH_3C , *t-Bu*), 125.3, 125.7, 127.9, 128.4, 130.9 (Ar), 131.3, 132.1, 146.8, 147.2 (*ipso-C*, Ar), 149.1 (C^3), 153.3 (C^5Pd).

[1-*tert*-Butyl-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene]palladium iodide (10b)

A solution of carbene **8c** (0.25 g, 0.68 mmol) and anhydrous dimethylsulfoxide (0.03 mL) in anhydrous toluene (1 mL) was added to a mixture of palladium iodide (0.246 g, 0.68 mmol) in toluene (5 mL) and stirred for 8 h. Then the mixture was boiled for 2.5 h. The precipitate that formed was filtered off, washed with petroleum ether, and dried. Yield 0.40 g (81%), mp 254–255 °C (acetonitrile). Found: C, 40.0; H, 4.2; I, 35.1; N, 5.9%. Calc. for $C_{24}H_{31}I_2N_3Pd$: C, 39.9; H, 4.3; I, 35.2; N, 5.8%. δ_H (400 MHz; $CDCl_3$; Me_4Si): 1 form: 0.62 d, 1.36 d (12H, CH_3C , *J* 4.8 Hz, *i-Pr*), 2.94 (m, 2H, CHC, *i-Pr*); 2 form: 0.83 d, 1.25 s (12H, CH_3C , *J* 4.8 Hz, *i-Pr*), 4.11 (s, 2H, CHC, *i-Pr*); a general part of two forms: 2.09 (s, 9H, *t-Bu*), 7.19 (m, 2H), 7.30 (m, 2H), 7.39 (m, 3H), 7.60 (m, 1H) (Ar). δ_C (100 MHz; solid state; DSS): 23.7, 25.8, 26.5, 27.2, 29.6 (CH_3C , *i-Pr*), 31.7, 32.3 (CH_3C , *t-Bu*), 47.3 (CHC, *i-Pr*), 64.5 (CH_3C , *t-Bu*), 126.0, 127.7, 128.6, 131.0, 131.5, 132.7, 134.2, 136.3, 137.3 (Ar), 146.6, 147.2, 149.2, 154.2 (*ipso-C*), 156.1, 157.6 (C^3), 167.0, 169.2, 170.0, 171.3 (C^5Pd).

[1-(1-Adamantyl)-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene]palladium chloride (10c)

A solution of carbene **8d** (0.2 g, 0.455 mmol) in toluene (1 mL) and anhydrous dimethylsulfoxide (0.03 mL) was added to a suspension of palladium chloride (0.08 g, 0.455 mmol) in anhydrous toluene (4 mL) for 10 min, and the mixture was stirred for 8 h. The precipitate that formed was filtered off, washed by petroleum ether, and dried. Yield 0.19 g (68%), mp 146 °C (dec.) (acetonitrile). Found: C, 58.3; H, 6.1; Cl, 11.5; N,

6.9. Calc. for $C_{30}H_{37}Cl_2N_3Pd$: C, 58.4; H, 6.0; Cl, 11.5; N, 6.8%. δ_H (400 MHz, $CDCl_3$, Me_4Si): 0.65 s, 13.8 s (12H, CH_3C , *i*-Pr), 2.25 (m, 9H, Ad), 2.62 (m, 6H, Ad; 2H, CHC , *i*-Pr), 7.19–7.60 (m, 8H, Ar). δ_C (100 MHz; $CDCl_3$; Me_4Si): 24.0 (CH_3C , *i*-Pr), 24.4 (CH , *i*-Pr), 28.5 (CH , Ad), 31.3, 41.0 (CH_2 , Ad), 64.2 (*ipso*-C, Ad), 125.4, 125.6, 127.9, 128.5, 130.9 (Ar), 131.4, 132.1, 147.0 (*ipso*-C), 149.1 (C^3), 153.3 (C^5Pd). δ_C (100 MHz; solid state; DSS): 26.2, 27.1, 29.6, 31.0 (CH_3C , *i*-Pr), 31.0, 36.6, 44.5 (CH_2 , Ad), 41.8, 48.6 (CH , *i*-Pr), 65.6 (*ipso*-C, Ad), 126.0, 127.2, 130.5, 131.9, 133.2 (*ipso*-C, Ar), 148.2 (C^3), 154.9 (C^5Pd).

[1-(1-Adamantyl)-3-phenyl-4-(2,6-diisopropylphenyl)-1,2,4-triazol-5-ylidene]palladium iodide (10d)

A mixture of carbene **8d** (0.25 g, 0.55 mmol), palladium iodide (0.19 g, 0.55 mmol) and anhydrous dimethylsulfoxide (0.3 mL) in anhydrous toluene (5 mL) was stirred for 4 h, and then the mixture was boiled for 3 h. The precipitate that formed was filtered off, washed by petroleum ether, and dried. Yield 0.37 g (84%), mp > 310 °C (dimethylformamide). Found: C, 45.0; H, 4.8; I, 31.6; N, 5.2%. Calc. for $C_{30}H_{37}I_2N_3Pd$: C, 45.1; H, 4.7; I, 31.7; N, 5.3%. δ_C (100 MHz; solid state; DSS): 26.9, 27.6, 29.0, 29.9 (CH_3C , *i*-Pr), 31.6, 37.5, 43.3 (CH_2 , Ad), 47.5, 49.1 (CH , *i*-Pr), 66.2 (*ipso*-C, Ad), 125.7, 128.0, 129.0, 130.5, 130.9, 132.0, 133.2 (Ar), 136.3, 146.9, 148.6 (*ipso*-C, Ar), 156.4 (C^3), 169.5 (C^5Pd).

Acknowledgements

We thank the Ukrainian Academy of Sciences for financial support (Grant no. 118, 26.02.2014), and the generous financial support provided by the Robert A. Welch Foundation (Grant F-0003) (A.H.C.).

Notes and references

- 1 Y. Cheng and O. Meth-Cohn, *Chem. Rev.*, 2004, **104**, 2507.
- 2 (a) D. Enders, O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, **107**, 5606; (b) *N-Heterocyclic Carbenes in Synthesis*, ed. S. P. Nolan, Wiley VCH Verlag GmbH & Co KGaA, Weinheim, 2006, p. 68; (c) J. Izquierdo, G. E. Hutson, D. T. Cohen and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2012, **51**, 11686; (d) M. J. Fuchter, *Chem. – Eur. J.*, 2010, **16**, 12286; (e) P.-C. Chiang and J. Y. W. Bode, in *RSC Catalysis Series No. 6: N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*, ed. S. Diez-Gonzalez, Royal Society of Chemistry, 2013, ch. 14, p. 399.
- 3 N. I. Korotkikh and O. P. Shvaika, *Carbene and carbene complex catalysis of organic reactions*, DonNU, Donetsk, Ukraine, 2013, p. 372.
- 4 (a) D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39; (b) M. C. Jahnke and F. E. Hahn, in *RSC Catalysis Series No. 6: N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*, ed. S. Diez-Gonzalez, RSC, 2013, ch. 1, p. 1; (c) D. Martin, M. Melaimi, M. Soleilhavoup and G. Bertrand, *Organometallics*, 2011, **30**, 5304; (d) W. Kirmse, *Angew. Chem., Int. Ed.*, 2010, **49**, 8798.
- 5 D. Enders, K. Breuer, G. Raabe, J. Runsink, J. H. Teles, J. P. Melder, K. Ebel and S. Brode, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**(9), 1021.
- 6 D. Enders, K. Breuer, J. Runsink and J. H. Teles, *Liebigs Ann. Chem.*, 1996, 2019.
- 7 N. I. Korotkikh, G. F. Rayenko, O. P. Shvaika, T. M. Pekhtereva, A. H. Cowley, J. N. Jones and C. L. B. Macdonald, *J. Org. Chem.*, 2003, **68**(14), 5762.
- 8 N. I. Korotkikh, O. P. Shvaika, G. F. Rayenko, A. V. Kiselyov, A. V. Knishevitsky, A. H. Cowley, J. N. Jones and C. L. B. Macdonald, *ARKIVOC*, 2005, **8**, 10.
- 9 N. I. Korotkikh, N. V. Glinyanaya, A. H. Cowley, J. A. Moore, A. V. Knishevitsky, T. M. Pekhtereva and O. P. Shvaika, *ARKIVOC*, 2007, **16**, 156.
- 10 N. I. Korotkikh, A. H. Cowley, J. A. Moore, N. V. Glinyanaya, I. S. Panov, G. F. Rayenko, T. M. Pekhtereva and O. P. Shvaika, *Org. Biomol. Chem.*, 2008, **1**, 195.
- 11 A. V. Knishevitsky, N. I. Korotkikh, A. H. Cowley, J. A. Moore, T. M. Pekhtereva, O. P. Shvaika and G. Reeske, *J. Organomet. Chem.*, 2008, **693**, 1405.
- 12 A. V. Kiselyov, N. I. Korotkikh, A. H. Cowley, J. A. Moore, T. M. Pekhtereva and O. P. Shvaika, *ARKIVOC*, 2008, **15**, 329.
- 13 C. Dash, M. M. Shaikh, R. J. Butcher and P. Ghosh, *Dalton Trans.*, 2010, **39**, 2515.
- 14 V. Hornillos, J. Guerra, A. de Cózar, P. Prieto, S. Merino, M. A. Maestro, E. Díez-Barra and J. Tejada, *Dalton Trans.*, 2011, **40**, 4095.
- 15 J. Turek, I. Panov, M. Horáček, Z. Černošek, Z. Padělková and A. Růžička, *Organometallics*, 2013, **32**, 7234.
- 16 T. I. Gorbunov, V. I. Saloutin and O. N. Chupakhin, *Russ. Chem. Rev.*, 2010, **79**(6), 565.
- 17 C. L. Geiger, K. Carvalho-Knighton, S. Novaes-Card, P. Maloney and R. DeVor, in *Environmental Applications of Nanoscale and Microscale Reactive Metal Particles*, *ACS Symposium Series*, American Chemical Society, Washington, DC, 2010, p. 1–20.
- 18 V. Dichiarante, M. Fagnoni and A. Albini, *Green Chem.*, 2009, **11**, 942.
- 19 K. A. Cannon, M. E. Geuther, C. K. Kelly, S. Lin and A. H. R. MacArthur, *Organometallics*, 2011, **30**, 4067.
- 20 W. M. Czaplik, S. Grupe, M. Mayer and A. J. von Wangelin, *Chem. Commun.*, 2010, **46**, 6350.
- 21 Y. Mitoma, T. Uda and N. Egashira, *Environ. Sci. Technol.*, 2004, **38**, 1216.
- 22 Y. Mitoma, M. Kakeda, A. M. Simion, N. Egashira and C. Simion, *Environ. Sci. Technol.*, 2009, **43**, 5952.
- 23 A. M. Simion, H. Miyata, M. Kakeda, N. Egashira, Y. Mitoma and C. Simion, *Sep. Purif. Technol.*, 2013, **103**, 267.
- 24 Y. Huang, S. Liu, Z. Lin, W. Li, X. Li and R. Cao, *J. Catal.*, 2012, **292**, 111.

- 25 P. Selvam, S. U. Sonavane, S. K. Mohapatra and R. V. Jayaram, *Tetrahedron Lett.*, 2004, **45**, 3071.
- 26 M. S. Viciu, G. A. Grasa and S. P. Nolan, *Organometallics*, 2001, **20**(16), 3607.
- 27 O. Navarro, H. Kaur, P. Mahjoor and S. P. Nolan, *J. Org. Chem.*, 2004, **69**(9), 3173.
- 28 O. Navarro, N. Marion, Y. Oonishi, R. A. Kelly and S. P. Nolan, *J. Org. Chem.*, 2006, **71**(2), 685.
- 29 C. J. E. Davies, M. J. Page, C. E. Ellul, M. F. Mahon and M. K. Whittlesey, *Chem. Commun.*, 2010, **46**, 5151.
- 30 S. Kuhl, R. Schneider and Y. Fort, *Adv. Synth. Catal.*, 2003, **345**(3), 341.
- 31 S. Akzinnay, F. Bisaro and C. S. J. Cazin, *Chem. Commun.*, 2009, 5752.
- 32 N. I. Korotkikh, V. Sh. Saberov, A. V. Kiselev, N. V. Glinyanaya, K. A. Marichev, T. M. Pekhtereva, G. V. Dudarenko, N. A. Bumagin and O. P. Shvaika, *Chem. Heterocycl. Compd.*, 2012, **47**(12), 1551.
- 33 X. Chen, R. Liu, Y. Xu and G. Zou, *Tetrahedron*, 2012, **68**, 4813.
- 34 N. I. Korotkikh, G. F. Raenko, T. M. Pekhtereva, O. P. Shvaika, A. H. Cowley and J. N. Jones, *Russ. J. Org. Chem.*, 2006, **42**(12), 1822.
- 35 (a) G. M. Sheldrick, *SHELXTL-PC, version 5.03*, Siemens Analytical X-ray Instruments, Inc., Madison, WI, 1994. See also; (b) G. M. Sheldrick, *Acta Crystallogr., Sect. A: Fundam. Crystallogr.*, 2008, **64**, 112.
- 36 A. L. Spek, *PLATON, A Multipurpose Crystallographic Tool*, Utrecht University, The Netherlands, 1998.
- 37 N. I. Korotkikh, A. V. Kiselyov, A. V. Knishevitsky, G. F. Rayenko, T. M. Pekhtereva and O. P. Shvaika, *Chem. Heterocycl. Compd.*, 2005, **7**, 1026.