

## REPTILIA: TESTUDINES: EMYDIDAE

*Trachemys nebulosa*

## Catalogue of American Amphibians and Reptiles.

Seidel, M.E. 2010. *Trachemys nebulosa*.

***Trachemys nebulosa* (Van Denburgh)  
Northwestern Mexican Slider**

*Pseudemys ornata*: Yarrow 1882:33.

*Chrysemys nebulosa* Van Denburgh 1895:84. Type-locality, "Los Dolores... Mainland abreast of San Jose Island, Lower California, [Mexico]". Holotype, California Academy of Sciences (CAS) 2244, collected by W.E. Bryant in 1889, date unknown (not examined by author). See **Comment**.

*Chrysemys ornata*: Siebenrock 1909:466 (part).

*Pseudemys nebulosa*: Van Denburgh and Slevin 1921:53.

*Clemmys marmorata*: Linsdale 1932:385 (part).

*Pseudemys scripta*: Carr 1938:135 (part).

*Chrysemys concinna*: Wermuth and Mertens 1961:140 (part).

*Chrysemys scripta*: Smith and Taylor 1966:12 (part).

*Chrysemys gageae*: Weaver and Rose 1967:63 (part).

*Trachemys scripta*: Iverson 1985:4 (part).

*Trachemys dorbigni*: Obst in Wermuth and Mertens 1996:473 (part).

*Trachemys ornata*: Walls 1996:50 (part).

*Trachemys nebulosa*: Wilms in Siebenrock 1999:198. First use of present combination.

• **CONTENT.** Two subspecies are recognized: *Trachemys nebulosa nebulosa* and *T. n. hiltoni*. See **Comment**.

• **DEFINITION.** The carapace of the adult female ranges to 37 cm and that of the male to 33 cm. It is widest at the level of marginal scutes 7–8. The carapace is oval, weakly keeled down the dorsal midline, and not notched or only slightly serrate along the posterior margin. The pleural scutes often have a series of rounded longitudinal ridges. The ground color of the adult carapace is brown to olive with obscured, black-bordered, orange reticulations along the pleural scutes, appearing as distorted "ocelli". The cervical scute underlap (ventral surface) is shallow, ventral length/carapace length < 0.035. The lateral and posterior marginals have dark spots or smudges on the dorsal and ventral surfaces. Markings of hatchlings and juveniles are more prominent.

The large, hingeless, plastron is relatively flat and slightly broader anterior to the bridge. The ground color of the plastron is pale to bright yellow with 2 rows of elongate black (smudge-like) markings variably separated by yellow at the midline. Elongate dark smudges also occur on the bridge.

The head and limbs are olive brown with a variety of yellow stripes. The supratemporal (postorbital) stripe is orange and does not contact the orbit of the eye. A yellow vertical bar branches dorsally from the orbitocervical stripe and crosses the tympanum. The throat is pale yellow, mostly without dark markings or clearly defined stripes. The left and right aspects of




**FIGURE 1.** Juvenile *T. nebulosa nebulosa* from El Corro, Baja California Sur, Mexico (photograph by L. Grismer).



**FIGURE 2.** Head and plastral views of adult female *T. nebulosa hiltoni* from Río Cuchujaqui, Sonora, Mexico (photographs by C. Schwalbe).

the upper jaw form an acute angle where they join at the midline. The tomial edge of the mandible is not distinctly serrate and choanal papillae are present in the roof of the mouth.

In contrast to females, adult males have a longer and wider tail with the vent positioned beyond the posterior edge of the carapace. The snout of males is longer and narrower than in females, but there is no



**MAP.** Circles mark type-localities; dots indicate other records.

marked sexual difference in foreclaw length.

• **DESCRIPTIONS.** General descriptions are presented in Bogert and Oliver (1945), Bonin et al. (1996, 2006), Bour (2003), Carr (1942, 1952), Casas Andreu (1965, 1967), Ernst (1990), Ernst and Barbour (1989), Ernst et al. (2000), Grismer (2002), Legler (1990), Legler and Webb (1970), Obst (1985), Pritchard (1967, 1979), Rogner (1995), Seidel (2002), Siebenrock (1909), Smith and Smith (1980), Smith and Taylor (1966), and Van Denburgh (1922). Other descriptions include: **head pattern** (Legler 1990; Ward 1980), **plastron pattern** (Williams 1956), **skull** (Ward 1980), **choanae** (Parsons 1960), **rostral pores** (Winokur and Legler 1974), **sexual dimorphism** (Gibbons and Lovich 1990), **egg** (Carr 1942), **habitat** (Carr 1952; Grismer 2002; Grismer and McGuire 1993; Hardy and McDiarmid 1969; Pritchard 1979), and **reproduction and nesting site** (Grismer 2002; Hardy and McDiarmid 1969).

• **ILLUSTRATIONS.** Black-and-white drawings are in Smith and Smith (1980), Van Denburgh (1895), Ward (1980), and Williams (1956). Black-and-white photographs are in Carr (1942) and Smith and Smith (1980). Color photographs are in Grismer (2002), Pritchard (1979), Rogner (1995), and Vetter (2004). Drawings of the egg are in Carr (1942) and of the plastron in Williams (1956). A color drawing of a hatchling is in Bour (2003).

• **DISTRIBUTION.** *Trachemys nebulosa* occurs in Mexico in disjunct bodies of fresh water in the southern half of the Baja Peninsula, Baja California Sur, and on the Mexican mainland, where it is restricted to the Río Fuerte drainage system of southern Sonora and northern Sinaloa. Range maps or discussions of the distribution of *T. nebulosa* are presented in Bonin et al. (1996, 2006), Bringsøe (2001), Carr (1952), Ernst (1990), Grismer (2002), Hardy and McDiarmid (1969), Iverson (1986, 1992), Legler (1990), Leviton and Banta (1964), Moll and Legler (1971), Mosauer (1935), Obst (1985), Roberts (1982), Seidel (2002),

Smith and Smith (1980), and Vetter (2004). The subspecies *Trachemys nebulosa nebulosa* is restricted to Baja California Sur, Mexico, south of San Ignacio; the distribution of *Trachemys nebulosa hiltoni* encompasses the watershed of the Río Fuerte in extreme southern Sonora and northern Sinaloa.

Conant (1969) and Murphy (1983) suggested that *T. n. nebulosa* on the Baja Peninsula may have been introduced, presumably from an ancestral population in the Río Fuerte (Smith and Smith 1980). Grismer and McGuire (1993) speculated that most of the populations north of the Cape Region of extreme southern Baja are the result of human introduction. Zweifel and Norris (1955) erroneously extended the range of *T. n. hiltoni* 160 km northwest of the Río Fuerte. They based this on a slider turtle they collected in the Río Yaqui, Sonora, Mexico. That specimen (MVZ 55384, now apparently lost) probably was *T. yaquia*, subsequently described by Legler and Webb (1970).

• **FOSSIL RECORD.** No fossils have been reported. A Recent archeological record was reported from the west coast of Mexico (Voorhies et al. 2002).

• **PERTINENT LITERATURE.** General accounts are in Bonin et al. (1996, 2006), Carr (1952), Casas Andreu (1967), Ditmars (1907, 1936), Ernst et al. (2000), Grismer (2002), Nietzsche (1969), Obst (1985), Pritchard (1979), Rogner (1995), Smith and Smith (1980), and Wermuth and Mertens (1961). Other important references include: **biogeography** (Clavigero 1789; Grismer 2002; Legler 1990; Savage 1960; Smith and Smith 1980), **common name** (Bogert and Oliver 1945; Liner 1994, 2007), **habitat and distribution** (Bogert and Oliver 1945; Bonin et al. 1996; Casas Andreu 1965; Hoard 1964; Mocquard 1899; Neill 1958; Nelson 1922; Roberts 1982; Savage 1960; Schmidt 1922; Smith and Smith 1975; Van Denburgh 1922), **human exploitation** (Grismer 2002), **longevity** (Bowler 1977; Snider and Bowler 1992), **macrosympatry** (Moll and Moll 2004), **morphology** (Parsons 1960; Winokur and Legler 1974), **natural history** (Bogert and Oliver 1945; Casas Andreu 1965; Grismer 2002; Zweifel and Norris 1955), **nesting behavior** (Carr 1952; Grismer 2002), **systematics and taxonomy** (Bickham et al. 2007; Bour 2003; Carr 1942; David 1994; Ernst 1990; Fritz 1981, 1990; Fritz and Havas 2007; Iverson 1986, 1992; Iverson et al. 2007; Legler 1990; Legler and Webb 1970; Obst 1985; Rust 1934; Seidel 2002; Smith and Smith 1980; Stephens and Wiens 2003; Van Denburgh 1895; Van Denburgh and Slevin 1921; Ward 1980; Weaver and Rose 1967; Wermuth and Mertens 1977; Williams 1956), **conservation** (Roberts 1982; Smith and Smith 1980), and **zoo holdings** (Nietzke 1969; Slavens 1976).

• **ETYMOLOGY.** The specific name *nebulosa*, from the Latin for “cloudy”, “misty”, “dark”, or “indefinite”, apparently refers to the obscured dark “ocelli-like” markings on the pleural scutes of the carapace. The subspecific name *hiltoni* is a Latinized patronym hon-



oring John W. Hilton, collector of the type-series.

### 1. *Trachemys nebulosa nebulosa* (Van Denburgh) Baja California Slider

*Chrysemys nebulosa* Van Denburgh 1895:84. See species synonymy.

*Chrysemys scripta elegans*: Mocquard 1899:300 (part).

*Chrysemys ornata nebulosa*: Siebenrock 1909:466.

*Pseudemys ornata nebulosa*: Stejneger and Barbour 1917:120.

*Pseudemys nebulosa*: Van Denburgh and Slevin 1921:53.

*Pseudemys scripta nebulosa*: Carr 1942:1

*Chrysemys scripta nebulosa*: Smith and Taylor 1966:12.

*Trachemys scripta nebulosa*: Iverson 1985:4.

*Chrysemys gaigeae nebulosa*: Obst 1985:7.

*Trachemys dorbigni nebulosa*: Obst in Wermuth and Mertens 1996:473.

*Trachemys ornata nebulosa*: Walls 1996:50.

*Trachemys nebulosa*: Wilms in Siebenrock 1999:198.

*Trachemys nebulosa nebulosa*: Bringsøe 2001:520. First use of present combination.

• **DEFINITION.** Maximum carapace length is greater than 32 cm. The two rows of dark smudges on the plastron are distinctly separated by a yellow midline region. The pygal bone typically extends beyond the marginal-vertebral seam.

### 2. *Trachemys nebulosa hiltoni* (Carr) Río Fuerte Slider

*Pseudemys scripta hiltoni* Carr 1942:1. Type-locality, "...Güirocoba about 28 miles southeast of Alamos, Sonora, Mexico, at an elevation of approximately 1,485 feet." Holotype, American Museum of Natural History (AMNH) 63747, fluid preserved mature male, collected by John W. Hilton in 1941 (examined by author).

*Pseudemys concinna hiltoni*: Wermuth and Mertens 1961:140.

*Chrysemys scripta hiltoni*: Smith and Taylor 1966:11.

*Chrysemys gaigeae hiltoni*: Weaver and Rose 1967:70.

*Trachemys scripta hiltoni*: Iverson 1985:4.

*Trachemys ornata hiltoni*: Walls 1996:50.

*Trachemys nebulosa hiltoni*: Bringsøe 2001:520. First use of present combination.

• **DEFINITION.** Maximum carapace length is less than 32 cm. The two rows of dark smudges on the plastron are frequently connected along the midline. The pygal bone does not extend beyond the marginal-vertebral seam.

• **COMMENT.** The first apparent reference to this turtle was by Yarrow (1882:33) who lists USNM 11622, shell, as *Pseudemys ornata* Gray, 1831, collected in February 1882 by Lyman Belding at La Paz, Baja California Sur, Mexico. The locality was correct-

ed to the San Jose River by Belding (1887:97).

Smith and Smith (1980) indicate that the type-specimen of *T. n. nebulosa* was "destroyed". The existence of the holotype (2244) in the California Academy of Sciences collection was recently confirmed by Robert Drewes and Jens Vindum (pers. comm.). Apparently Smith and Smith (1980) confused the holotype with CAS 2245 (a topotype also collected by W.E. Bryant) which was indeed lost in the earthquake and fire of 1906.

*Trachemys n. hiltoni* and *T. n. nebulosa* were formerly recognized as subspecies of *T. scripta*. Based on a phylogenetic analysis using morphological characters, Seidel (2002) found that these turtles are very similar (sister taxa) and assigned them to subspecies of *T. nebulosa*. Their phylogenetic relationship and natural origins remains somewhat uncertain (Legler 1990; Smith and Smith 1980).

## LITERATURE CITED

- Belding, L. 1887. Reptiles of the Cape Region of Lower California. West Amer. Sci. 3:93–97.
- Bickham, J.W., J.B. Iverson, J.F. Parham, H. Philippen, A.G.J. Rhodin, H.B. Shaffer, P.Q. Spinks, and P.P. Van Dijk. 2007. An annotated list of modern turtle terminal taxa with comments on areas of taxonomic instability and recent change. Chelonian Res. Monogr. (4):173–199.
- Bogert, C.M. and J.A. Oliver. 1945. A preliminary analysis of the herpetofauna of Sonora. Bull. Amer. Mus. Nat. Hist. 83:301–425.
- Bonin, F., B. Devaux, and A. Dupré. 1996. Toutes les Tortues du Monde. Delachaux et Niestlé S.A., Lausanne, Switzerland.
- , –, and –. 2006. Turtles of the World. Johns Hopkins Univ. Press, Baltimore, Maryland.
- Bour, R. 2003. Le genre *Trachemys*: systematique et repartition. Manouria 6:2–9.
- Bowler, J.K. 1977. Longevity of reptiles and amphibians in North American collections as of November, 1975. SSAR Herpetol. Circ. (6):1–32.
- Bringsøe, H. 2001. *Trachemys* Agassiz, 1857–Schmuckschildkröten, p. 517–523. In U. Fritz (ed.), Handbuch der Reptilien und Amphibien Europas. Vol. 3/IIIA: Schildkroten (Testudines) I (Bataguridae, Testudinidae, Emydidae). AULA-Verlag GmbH, Wiebelsheim, Germany.
- Carr, A.F. 1938. Notes on the *Pseudemys scripta* complex. Herpetologica 1:131–135.
- . 1942. A new *Pseudemys* from Sonora, Mexico. Amer. Mus. Nov. (1181):1–4.
- . 1952. Handbook of Turtles. The Turtles of the United States, Canada, and Baja California. Cornell Univ. Press, Ithaca, New York.
- Casas Andreu, G. 1965. Estudio preliminar sobre las tortugas de agua dulce Mexico, D.F. Ann. Inst. Nac. Invest. Biol. Pesq. 1:365–401.
- . 1967. Contribucion al conocimiento de las tortugas dulceauicolas de Mexico. Fac. Cien. Dept. Biol. Univ. Natl. Auton. Mexico.
- Clavigero, F.J. 1789. Storia della California. Venice. Translated and annotated as the history of

- [Lower] California by S.E. Lake and A.A. Gray, 1937. Stanford Univ. Press, Palo Alto, California.
- Conant, R. 1969. A review of the water snakes of the genus *Natrix* in Mexico. *Bull. Amer. Mus. Nat. Hist.* 142:1–140.
- David, P. 1994. Liste des reptiles actuels du Monde. I. Chelonii. *Dumerilia* 1:7–127.
- Ditmars, R.L. 1907. *The Reptile Book*. Doubleday, Page Co. Inc., New York.
- . 1936. *The Reptiles of North America*. Doubleday, Doran Co., Garden City, New York.
- Ernst, C.H. 1990. Systematics, taxonomy, variation, and geographic variation of the slider turtle, p. 57–67. *In* J. W. Gibbons (ed.), *Life History and Ecology of the Slider Turtle*. Smithsonian Inst. Press, Washington D.C.
- , R.G.M. Altenburg, and R.W. Barbour. 2000. *Turtles of the World*. Multimedia interactive CD-ROM. Biodiv. Ctr., Exp. Ctr. Taxon. Ident., Amsterdam. Springer-Verlag, UNESCO, Heidelberg.
- and R.W. Barbour. 1989. *Turtles of the World*. Smithsonian Inst. Press, Washington, D. C.
- Fritz, U. 1981. Zwei interessante mexikanische Schmuckschildkroten der Gattung [sic] *Chrysemys* Gray, 1844. *Herpetofauna (Germany)* 3: 25–32.
- . 1990. Balzverhalten and systematik in der subtribus Nectemydina. 1. Die Gattung *Trachemys*, besonders *Trachemys scripta callirostris* (Gray, 1855). *Salamandra* 26:221–245.
- and P. Havas. 2007. Checklist of chelonians of the World. *Vert. Zool.* 57:149–368.
- Gibbons, J.W. and J.E. Lovich. 1990. Sexual dimorphism in turtles with emphasis on the slider turtle (*Trachemys scripta*). *Herpetol. Monogr.* (4):1–29.
- Gray, J.E. 1831. *Synopsis Reptilium*. Part 1. Cataphracta, Tortoises, Crocodiles, and Enaliosaurians. Treüttel., Würtz and Co., London.
- Grismer, L.L. 2002. *Amphibians and Reptiles of Baja California*. Univ. California Press, Berkeley.
- and J.A. McGuire. 1993. The oases of central Baja California, Mexico, part I: a preliminary account of the relict mesophilic herpetofauna and the status of the oases. *Bull. So. California Acad. Sci.* 92:2–24.
- Hardy, L.M. and R.W. McDiarmid. 1969. The amphibians and reptiles of Sinaloa, Mexico. *Univ. Kansas Publ. Mus. Nat. Hist.* 18:39–252.
- Hoard, H.F. 1964. New Lower California reptile records. *J. Entomol. Zool.* 31:4–5.
- Iverson, J.B. 1985. Checklist of the turtles of the world with English common names. *SSAR Herpetol. Circ.* (14):ii +14 p.
- . 1986. A Checklist with Distribution Maps of the Turtles of the World. Privately printed. Richmond, Indiana.
- . 1992. A Revised Checklist with Distribution Maps of the Turtles of the World. Privately printed. Richmond, Indiana.
- , R.M. Brown, T.S. Akre, T.J. Near, M. Le, R.C. Thomson, and D.E. Starkey. 2007. In search of the tree of life for turtles. *Chelonian Res. Monogr.* (4):85–106.
- Legler, J.M. 1990. The genus *Pseudemys* in Mesoamerica: taxonomy, distribution, and origins, p. 82–105. *In* J.W. Gibbons (ed.), *Life History and Ecology of the Slider Turtle*. Smithsonian Inst. Press, Washington D.C.
- and R.G. Webb. 1970. A new slider turtle (*Pseudemys scripta*) from Sonora, Mexico. *Herpetologica* 26:157–168.
- Leviton, A.E. and B.H. Banta. 1964. Midwinter reconnaissance of the herpetofauna of the Cape region of Baja California, Mexico. *Proc. California Acad. Sci.* 30:127–156.
- Liner, E.A. 1994. Scientific and common names for the amphibians and reptiles of Mexico in English and Spanish. *SSAR Herpetol. Circ.* (23):v +113 p.
- . 2007. A checklist of the amphibians and reptiles of México. *Occas. Pap. Mus. Nat. Sci. Louisiana St. Univ.* (80):1–60.
- Linsdale, J.M. 1932. Amphibians and reptiles from Lower California. *Univ. California Publ. Zool.* 38: 346–386.
- Mocquard, F. 1899. Contribution a la faune herpetologique de la Basse Californie. *Nouv. Arch. Mus. Nat. Hist. Natl.* 4:297–344.
- Moll, D. and E.O. Moll. 2004. *The Ecology, Exploitation and Conservation of River Turtles*. Oxford Univ. Press, New York.
- Moll, E.O. and J.M. Legler. 1971. The life history of a Neotropical slider turtle, *Pseudemys scripta* (Schoepff), in Panama. *Bull. Los Angeles Co. Mus. Nat. Hist. Sci.* (11):1–102.
- Mosauer, W. 1935. The identity of the turtles from San Ignacio, Lower California. *Copeia* 1935:191.
- Murphy, R.W. 1983. Paleobiogeography and genetic differentiation of the Baja California herpetofauna. *Occas. Pap. California Acad. Sci.* (137):1–48.
- Neill, W.T. 1958. The occurrence of amphibians and reptiles in saltwater areas, and a bibliography. *Bull. Marine Sci. Gulf Caribbean.* 8:1–97.
- Nelson, E.W. 1922. Lower California and its natural resources. *Mem. Natl. Acad. Sci. Washington* (16):1–194.
- Nietzke, G. 1969. *Die Terrarierentier* 1. Lechler, Stuttgart, Germany.
- Obst, F.J. 1985. *Schmuckchildkroten. Die Gattung Chrysemys*. A. Ziemsen Verlag, Wittenberg Lutherstadt, Germany.
- Parsons, T.S. 1960. The structure of the choanae of the Emydinae (Testudines: Testudinidae). *Bull. Mus. Comp. Zool.* 123:113–127.
- Pritchard, P.C.H. 1967. *Living Turtles of the World*. T.F.H. Publ., Inc., Jersey City, New Jersey.
- . 1979. *Encyclopedia of Turtles*. T.F.H. Publ., Inc., Neptune, New Jersey.
- Roberts, N. 1982. A preliminary report on the status of Chelydridae, Trionychidae, and Testudinidae in the region of Baja California, Mexico. *Proc. Desert Tortoise Council* 1982:154–161.
- Rogner, M. 1995. *Schildkroten* 1. Heidi Rogner-Verlag, Hurtgenwald, Germany.
- Rust, H.T. 1934. Systematische liste der lebenden schildkroten. *Bl. Aquar. Terrar.* 45:42–45, 59–67.
- Savage, J.M. 1960. Evolution of a peninsular herpe-

- tofauna. Syst. Zool. 9:184–212.
- Schmidt, K.P. 1922. The amphibians and reptiles of Lower California and the neighboring islands. Bull. Amer. Mus. Nat. Hist. 46:607–707.
- Seidel, M.E. 2002. Taxonomic observations on extant species and subspecies of slider turtles, genus *Trachemys*. J. Herpetol. 36:285–292.
- Siebenrock, F. 1909 (reprinted 1999). Synopsis der rezenten Schildkröten, mit Berücksichtigung der in historischer Zeit ausgestorbenen Arten. Zool. Jahrb. Suppl. 10:428–618.
- Slavens, F.L. 1976. The Working Guide to Breeding Potential for Reptiles and Amphibians in the United States Zoos. Woodland Park Zoological Gardens, Seattle, Washington.
- Smith, H.M. and R.B. Smith. 1975. An analysis of the knowledge of the turtle fauna of Mexico. Chelonia 2:3–8.
- and –. 1980 (1979). Synopsis of the Herpetofauna of Mexico. Vol. VI. Guide to Mexican Turtles. Bibliographic Addendum III. John Johnson, North Bennington, Vermont.
- and E.H. Taylor. 1966. Herpetology of Mexico: Annotated Checklists and Keys to the Amphibians and Reptiles. A Reprint of Bulletins 187, 194 and 199 of the U.S. National Museum with a list of Subsequent Taxonomic Innovations. Eric Lundberg, Ashton, Maryland.
- Snider, A.T. and J.K. Bowler. 1992. Longevity of Reptiles and Amphibians in North American Collections, 2nd ed. SSAR Herpetol. Circ. (21):1–40.
- Stejneger, L.H. and T. Barbour. 1917. A Check List of North American Amphibians and Reptiles. Harvard Univ. Press, Cambridge.
- Stephens, P.R. and J.J. Wiens. 2003. Ecological diversification and phylogeny of emydid turtles. Biol. J. Linn. Soc. 79:577–610.
- Van Denburgh, J. 1895. A review of the herpetology of Lower California. Part I - Reptiles. Proc. California Acad. Sci. 2:77–162, plates iv–xiv.
- . 1922. The reptiles of western North America Vol. II. Snakes and Turtles. Occas. Pap. California Acad. Sci. (10):612–1028, pl. 58–128.
- and J.R. Slevin. 1921. A list of amphibians and reptiles of the peninsula of Lower California, with notes on the species in the California Academy. Proc. California Acad. Sci. 11:27–72.
- Vetter, H. 2004. Turtles of the World. Vol. 2, North America. Andreas S. Brahm, Germany.
- Voorhies, B., D.J. Kennett, J.G. Jones, and T.A. Wake. 2002. A Middle Archaic archaeological site on the west coast of Mexico. Latin Amer. Antiquity 13:179–200.
- Walls, J.G. 1996. Cooters, Sliders and Painted Turtles. T.F.H Publ., Neptune City, New Jersey.
- Ward, J.P. 1980. Comparative Cranial Morphology of the Freshwater Turtle Subfamily Emydinae: an Analysis of the Feeding Mechanisms and the Systematics. Ph.D. Diss., North Carolina St. Univ., Raleigh.
- Weaver, W.G. and F.L. Rose. 1967. Systematics, fossil history, and evolution of the genus *Chrysemys*. Tulane Stud. Zool. 14:63–73.
- Wermuth, H. and R. Mertens. 1961. Schildkröten, Krokodile, Brückeneschen. Gustav Fischer, Jena.
- and –. 1977. Liste der rezenten Amphibien und Reptilien. Testudines, Crocodylia, Rhynchocephalia. Das Tierreich (100):xxvii + 174 p.
- and –. 1996. Schildkröten, Krokodile, Brückeneschen. Gustav Fischer, Jena. Reprinted Edition.
- Williams, E.E. 1956. *Pseudemys scripta callirostris* from Venezuela with a general survey of the *scripta* series. Bull. Mus. Comp. Zool. 115:145–160.
- Winokur, R.M. and J.M. Legler. 1974. Rostral pores in turtles. J. Morphol. 143:107–118.
- Yarrow, H.C. 1882. Check list of North American Reptilia and Batrachia, with catalogue of specimens in U.S. National Museum. Bull. U.S. Natl. Mus. (24): v + [i] + 249 p.
- Zweifel, R.G. and K.S. Norris. 1955. Contributions to the herpetology of Sonora, Mexico: descriptions of new subspecies of snakes (*Micruroides euryxanthus* and *Lampropeltis getulus*) and miscellaneous collecting notes. Amer. Midl. Nat. 54:230–249.

---

**Michael E. Seidel**, 4430 Richmond Park Drive East, Jacksonville, Florida 32224 (seidel@marshall.edu).

Primary editor for this account, Andrew H. Price.

Published 15 January 2010 and Copyright © 2010 by the Society for the Study of Amphibians and Reptiles.

---





