



(12) **United States Patent**
Chandrashekar et al.

(10) **Patent No.: US 6,309,644 B1**
(45) **Date of Patent: Oct. 30, 2001**

(54) **PARASITIC NEMATODE
TRANSGLUTAMINASE PROTEINS AND
USES THEREOF**

(75) Inventors: **Ramaswamy Chandrashekar**, Fort
Collins, CO (US); **Kapil Mehta**,
Houston, TX (US)

(73) Assignees: **Heska Corporation**, Fort Collins, CO
(US); **The Board of Regents, The
University of Texas System**, Austin,
TX (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **08/874,102**

(22) Filed: **Jun. 12, 1997**

Related U.S. Application Data

(63) Continuation-in-part of application No. 08/781,420, filed on
Dec. 3, 1996, now Pat. No. 6,248,872.

(51) **Int. Cl.⁷** **A61K 39/00**; A61K 39/02;
C12N 9/00; C07K 1/00

(52) **U.S. Cl.** **424/191.1**; 435/7.1; 435/183;
530/350

(58) **Field of Search** 530/350; 424/191.1;
435/7.1, 183

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,970,297 11/1990 Castelhana et al. .
5,124,358 6/1992 Kapil et al. .
5,686,080 * 11/1997 Tripp et al. .

FOREIGN PATENT DOCUMENTS

WO 95/00636 * 1/1995 (WO) .

OTHER PUBLICATIONS

Bain et al (Int. J. for Parasitology vol. 29 pp. 185–191,
1999.*

Hoetz et al (Emerging Infectious Diseases vol. 3(3) pp.
303–310), Jul.–Sep. 1997.*

Bourdi et al., “cDNA Cloning and Baculovirus Expression
of the Human Liver Endoplasmic Reticulum P58: Charac-
terization as a Protein Disulfide Isomerase Isoform, but Not
as a Protease or a Carnitine Acyltransferase,” 1995, pp.
397–403, *Arch Biochem Biophys* 323:2.

Cariello et al., “A new transglutaminase–like from the
ascidian *Ciona intestinalis*,” 1997, pp. 171–176, *FEBS
Letters* 408.

Edman et al., “Sequence of protein disulphide isomerase and
implications of its relationship to thioredoxin,” 1985, pp.
267–270, *Nature* 317.

Finken et al., “Characterization of the Complete Protein
Disulfide Isomerase Gene of *Schistosoma mansoni* and
identification of the Tissues of its Expression,” 1994, pp.
135–144, *Molecular and Biochemical Parasitology* 640.

Freedman et al., “Protein Disulphide Isomerase: Building
Bridges in Protein Folding,” 1994, pp. 331–336, *TIBS* 19.

Freedman, Robert B., “Protein Disulfide Isomerase: Mul-
tiple Roles in the Modification of Nascent Secretory Pro-
teins,” 1989, pp. 1069–1072, *Cell* 57.

Gething et al., “Protein folding in the cell,” 1992, pp. 33–45,
Nature 355.

Holmgren, Arne, “Thioredoxin,” 1985, pp. 237–271, *Annu.
Rev. Biochem* 54.

Lustigman, S., “Molting Enzymes and New Targets for
Chemotherapy of *Onchocerca volvulus*,” 1993, pp.
294–297, *Parasitology Today* 9(8).

Lustigman, “Transglutaminase–Catalyzed Reaction is
Important for Molting of *Onchocerca volvulus* Third–Stage
Larvae,” 1995, pp. 1913–1919, *Antimicrobial Agents and
Chemotherapy* 39(9).

Mehta et al., “Identification of a Novel Transglutaminase
from the Filarial Parasite *Brugia malayi* and its Role in
Growth and Development,” 1992, pp. 1–16, *Molecular and
Biochemical Parasitology* 53.

Mehta et al., “Significance of Transglutaminase–Catalyzed
Reactions in Growth and Development of Filarial Parasite,
Brugia malayi,” 1990, pp. 1051–1057, *Biochemical and
Biophysical Research Communications* 173(3).

Mehta et al., “Transglutaminase–Catalyzed Incorporation of
Host Proteins in *Brugia malayi* Microfilariae,” 1996, pp.
105–114, *Molecular and Biochemical Parasitology* 76.

Noiva et al., “Protein Disulfide Isomerase,” 1992, pp.
3553–3556, *The Journal of Biological Chemistry* 267:6.

Rao et al., “*Brugia malayi* and *Acanthocheilonema viteae*:
Antifilarial Activity of Transglutaminase Inhibitors In
Vitro,” 1991, pp. 2219–2224, *Antimicrobial Agents and
Chemotherapy* 35(11).

Singh et al., “Purification and Characterization of a Novel
Transglutaminase from Filarial Nematode *Brugia malayi*,”
1994, pp. 625–634, *Eur. J. Biochem.* 225.

(List continued on next page.)

Primary Examiner—Mark Navarro

(74) *Attorney, Agent, or Firm*—Heska Corporation

(57) **ABSTRACT**

The present invention relates to parasitic nematode trans-
glutaminase proteins; to parasitic nematode transglutami-
nase nucleic acid molecules, including those that encode
such transglutaminase proteins; to antibodies raised against
such transglutaminase proteins; and to compounds that
inhibit parasitic nematode transglutaminase activity. The
present invention also includes methods to obtain such
proteins, nucleic acid molecules, antibodies, and inhibitory
compounds. Also included in the present invention are
therapeutic compositions comprising such proteins, nucleic
acid molecules, antibodies and/or inhibitory compounds as
well as the use of such therapeutic compositions to protect
animals from diseases caused by parasitic nematodes. This
invention also relates to the surprising discovery that para-
sitic nematode transglutaminase proteins have protein dis-
ulfide isomerase activity. Accordingly, this invention relates
further to inhibitors of the protein disulfide isomerase activ-
ity of said transglutaminases.

18 Claims, No Drawings

OTHER PUBLICATIONS

Singh et al., "Purification and Partial Characterization of a Transglutaminase from Dog Filarial Parasite, *Dirofilaria immitis*," 1995, pp. 1285-1291, *Int. J. Biochem. Cell Biol.* 27(12).

Van et al., "CaBP2 is a rat homolog of ERp72 with protein disulfide isomerase activity," 1993, pp. 789-795, *Eur J. Biochem* 213.

Wilson et al., "The *Onchocerca volvulus* Homologue of the Multifunctional Polypeptide Protein Disulfide Isomerase," 1994, pp. 103-117, *Molecular and Biochemical Parasitology* 68.

Bennett et al., 1988, *Nature*, vol. 334, pp. 268-270.

Hempel et al., 1991, *J. of Immunol*, vol. 146, pp. 3713-3720.

Koivunen et al., 1997, *Genomics*, vol. 42, pp. 397-404.

* cited by examiner

1

**PARASITIC NEMATODE
TRANSGLUTAMINASE PROTEINS AND
USES THEREOF**

CROSS REFERENCE TO RELATED
APPLICATIONS

This application is a continuation-in-part of application U.S. Ser. No. 08/781,420, filed Dec. 3, 1996, now U.S. Pat. No. 6,248,872, entitled "PARASITIC NEMATODE TRANSGLUTAMINASE PROTEINS, NUCLEIC ACID MOLECULES, AND USES THEREOF".

FIELD OF THE INVENTION

The present invention relates to parasitic nematode transglutaminase nucleic acid molecules, proteins encoded by such nucleic acid molecules, antibodies raised against such proteins, and inhibitors of such proteins. The present invention also includes therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies, inhibitors, and combinations thereof, as well as the use of these compositions to protect animals from diseases caused by parasitic nematodes.

BACKGROUND OF THE INVENTION

Parasitic nematode infections in animals, including humans, are typically treated by chemical drugs. One disadvantage with chemical drugs is that they must be administered often. For example, dogs susceptible to heartworm are typically treated monthly. Repeated administration of drugs, however, often leads to the development of resistant nematode strains that no longer respond to treatment. Furthermore, many of the chemical drugs cause harmful side effects in the animals being treated, and as larger doses become required due to the build up of resistance, the side effects become even greater. Moreover, a number of drugs only treat symptoms of a parasitic disease but are unable to prevent infection by the parasitic nematode.

An alternative method to prevent parasitic nematode infection includes administering a vaccine against a parasitic nematode. Although many investigators have tried to develop vaccines based on specific antigens, it is well understood that the ability of an antigen to stimulate antibody production does not necessarily correlate with the ability of the antigen to stimulate an immune response capable of protecting an animal from infection, particularly in the case of parasitic nematodes. Although a number of prominent antigens have been identified in several parasitic nematodes, including in *Dirofilaria*, there is yet to be a commercially available vaccine developed for any parasitic nematode.

The life cycle of parasitic nematodes generally includes development through four molts, the last two molts taking place in the host animal. Molting is a complex process involving a variety of different mechanisms. However, a lack of understanding of the basic biology, metabolism and biochemistry of parasitic nematodes has resulted in the identification of few targets for chemotherapy or vaccines.

As an example of the complexity of parasitic nematodes, the life cycle of *D. immitis*, the nematode that causes heartworm, includes a variety of life forms, each of which presents different targets, and challenges, for immunization. Adult forms of the parasite are quite large and preferentially inhabit the heart and pulmonary arteries of an animal. Sexually mature adults, after mating, produce microfilariae which traverse capillary beds and circulate in the vascular

2

system of the dog. One method of demonstrating infection in the dog is to detect the circulating microfilariae. If a dog is maintained in an insect-free environment, the life cycle of the parasite cannot progress. However, when microfilariae are ingested by a female mosquito during blood feeding on an infected dog, subsequent development of the microfilariae into larvae occurs in the mosquito. The microfilariae go through two larval stages (L1 and L2) and finally become mature third stage larvae (L3) which can then be transmitted back to the dog through the bite of the mosquito. It is this L3 stage, therefore, that accounts for the initial infection. As early as three days after infection, the L3 molt to the fourth larval (LA) stage, and subsequently to the fifth stage, or immature adults. The immature adults migrate to the heart and pulmonary arteries, where they mature and reproduce, thus producing the microfilariae in the blood. "Occult" infection with heartworm in dogs is defined as that wherein no microfilariae can be detected, but the existence of the adult heartworms can be determined through thoracic examination.

Heartworm not only is a major problem in dogs, which typically cannot even develop immunity upon infection (i.e., dogs can become reinfected even after being cured by chemotherapy), but is also becoming increasingly widespread in other companion animals, such as cats and ferrets. Heartworm infections have also been reported in humans. Other parasitic nematode infections are also widespread, and all require better treatment, including a preventative vaccine program. *O. volvulus*, for example, causes onchocerciasis (also known as river blindness) in humans. Up to 50 million people throughout the world are reported to be infected with *O. volvulus*, with over a million being blinded due to infection.

Although many investigators have tried to develop vaccines based on specific antigens, it is well understood that the ability of an antigen to stimulate antibody production does not necessarily correlate with the ability of the antigen to stimulate an immune response capable of protecting an animal from infection, particularly in the case of parasitic nematodes. Although a number of prominent antigens have been identified in several parasitic nematodes, including in *Dirofilaria* and *Onchocerca*, there is yet to be an effective vaccine developed for any parasitic nematode.

In just the past few years, there has developed an interest in the identification of larval stage-specific enzymes as potential targets for treatment or prevention of nematode diseases. Nematode transglutaminase-catalyzed reactions have recently been identified as possibly important for the growth, development and survival of nematodes, including *Acanthocheilonema vitae*, *Brugia malayi*, and *Onchocerca volvulus*. See, for example, Mehta, 1992, *Mol. Biochem. Parasitol.*, 53, 1-16; Lustigman, 1995, *Antimicrobial Agents and Chemother.*, 39:9, 1913-1919; Lustigman, 1993, *Parasitology Today*, 9:8, 294-297. However, until now, no compounds or methods based on specific known targets in parasitic nematode development have been designed for treating or preventing parasitic nematode disease.

There remains a need to identify an efficacious composition that protects animals against diseases caused by parasitic nematodes and that, preferably, also protects animals from infection by such nematodes.

SUMMARY OF THE INVENTION

The present invention relates to novel products and processes for prevention and treatment of parasitic nematode infection. According to the present invention there are

provided parasitic nematode transglutaminase proteins and mimetopes thereof; nematode nucleic acid molecules, including those that encode such proteins; antibodies raised against parasitic nematode transglutaminase proteins (i.e., anti-parasitic nematode transglutaminase antibodies); and other compounds that inhibit parasitic nematode transglutaminase activity or the protein disulfide isomerase activity of parasitic nematode transglutaminase (i.e., inhibitory compounds or inhibitors).

The present invention also includes methods to obtain the proteins, mimetopes, nucleic acid molecules, antibodies and inhibitory compounds herein described. Also included in the present invention are therapeutic compositions comprising such proteins, mimetopes, nucleic acid molecules, antibodies, inhibitory compounds, or mixtures thereof, as well as the use of such therapeutic compositions to protect animals from diseases caused by parasitic nematodes.

One embodiment of the present invention is an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a parasitic nematode transglutaminase gene. Preferred parasitic nematode transglutaminase genes of the present invention are transglutaminase genes from *Brugia malayi*, *Dirofilaria immitis*, and *Onchocerca volvulus*. Such nucleic acid molecules are referred to as nematode transglutaminase nucleic acid molecules. A parasitic nematode transglutaminase gene preferably includes at least one of the following nucleic acid sequences: SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59.

Another embodiment of the present invention is an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Dirofilaria immitis* (*D. immitis*) transglutaminase gene such nucleic acid molecules are referred to as *Dirofilaria immitis* (or *D. immitis*) transglutaminase nucleic acid molecules. A *D. immitis* transglutaminase gene preferably includes one of the following nucleic acid sequences: SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54 or SEQ ID NO:56. A preferred *D. immitis* transglutaminase protein includes at least a portion of a protein represented by SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28 or SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:52 or SEQ ID NO:55.

The present invention also relates to recombinant molecules, recombinant viruses and recombinant cells that include a transglutaminase nucleic acid molecule of the present invention. Also included are methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

Another embodiment of the present invention includes a non-native nematode transglutaminase protein encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions with a parasitic nematode transglutaminase gene. A preferred nematode transglutaminase protein is

capable of eliciting an immune response when administered to an animal and/or of having parasitic nematode transglutaminase or protein disulfide isomerase (PDI) activity, or both. A preferred nematode transglutaminase protein is encoded by a nucleic acid molecule that hybridizes under stringent conditions with a nucleic acid molecule including either SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56 or SEQ ID NO:59. A preferred nematode transglutaminase protein includes at least a portion of a protein represented by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58.

Another embodiment of the present invention includes a non-native *D. immitis* transglutaminase protein encoded by a nucleic acid molecule that hybridizes under stringent conditions with a *D. immitis* transglutaminase gene. A preferred *D. immitis* transglutaminase protein is capable of eliciting an immune response when administered to an animal and/or of having parasitic nematode transglutaminase activity. A preferred *D. immitis* transglutaminase protein is encoded by a nucleic acid molecule that hybridizes under stringent conditions with a nucleic acid molecule including either SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:53 or SEQ ID NO:56. A preferred *D. immitis* transglutaminase protein includes at least a portion of a protein represented by SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:52 or SEQ ID NO:55.

The present invention also relates to mimetopes of parasitic nematode transglutaminase proteins as well as to isolated antibodies that selectively bind to parasitic nematode transglutaminase proteins or mimetopes thereof. Also included are methods, including recombinant methods, to produce proteins, mimetopes and antibodies of the present invention.

Another embodiment of the present invention is a method to identify a compound capable of inhibiting nematode transglutaminase activity. The method includes the steps of: (a) contacting an isolated nematode transglutaminase protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has nematode transglutaminase activity; and (b) determining if the putative inhibitory compound inhibits the nematode transglutaminase activity. Also included in the present invention is a test kit to identify a compound capable of inhibiting nematode transglutaminase activity. Such a test kit includes an isolated nematode transglutaminase protein having nematode transglutaminase activity and a means for determining the extent of inhibition of the nematode transglutaminase activity in the presence of a putative inhibitory compound.

The present invention also includes an inhibitor of nematode transglutaminase activity identified by its ability to inhibit the activity of a nematode transglutaminase and by its inability to substantially inhibit mammalian transglutaminase. Examples of such inhibitors are substrate analogs of nematode transglutaminase, active site inhibitors of nematode transglutaminase, and antibodies that specifically recognize nematode transglutaminase.

Another embodiment of the present invention is a method to identify a compound capable of inhibiting protein disul-

fide isomerase (PDI) activity. The method includes the steps of: (a) contacting an isolated nematode transglutaminase protein with a putative PDI inhibitory compound under conditions in which, in the absence of the compound, the protein has nematode PDI activity; and (b) determining if the putative inhibitory compound inhibits the nematode PDI activity. Also included in the present invention is a test kit to identify a compound capable of inhibiting nematode PDI activity. Such a test kit includes an isolated nematode PDI protein having nematode PDI activity and a means for determining the extent of inhibition of the nematode PDI activity in the presence of a putative inhibitory compound.

The present invention also includes an inhibitor of nematode PDI activity identified by its ability to inhibit the PDI activity of a nematode transglutaminase protein and by its inability to substantially inhibit mammalian PDI. Examples of such inhibitors are substrate analogs of nematode PDI, active site inhibitors of nematode PDI, and antibodies that specifically recognize parasitic nematode transglutaminase.

Yet another embodiment of the present invention is a therapeutic composition that is capable of protecting an animal from disease caused by a parasitic nematode. Such a therapeutic composition includes an excipient and one or more of the following protective compounds: an isolated nematode transglutaminase protein or a mimotope thereof; an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a nematode transglutaminase gene; an isolated antibody that selectively binds to a nematode transglutaminase protein; an inhibitor of nematode transglutaminase protein activity identified by its ability to (a) inhibit nematode transglutaminase activity, and (b) not substantially inhibit mammalian transglutaminase activity; an inhibitor of nematode PDI protein activity identified by its ability to (a) inhibit nematode PDI activity, and (b) not substantially inhibit mammalian PDI activity; or any combinations thereof. A preferred therapeutic composition of the present invention also includes an adjuvant, a carrier, or both. Preferred nematode transglutaminase nucleic acid molecule compounds of the present invention include naked nucleic acid vaccines recombinant virus vaccines and recombinant cell vaccines. Also included in the present invention is a method to protect an animal from disease caused by a parasitic nematode comprising the step of administering to the animal at least one protective compound of the present invention.

Yet another embodiment of the present invention is a method to produce a transglutaminase protein, the method comprising culturing a cell transformed with a nucleic acid molecule that hybridizes under stringent hybridization conditions with a nematode transglutaminase gene.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides for isolated parasitic nematode transglutaminase proteins, isolated parasitic nematode transglutaminase nucleic acid molecules, antibodies directed against parasitic nematode transglutaminase proteins, and other inhibitors of nematode transglutaminase activity. As used herein, the terms isolated parasitic nematode transglutaminase proteins and isolated parasitic nematode transglutaminase nucleic acid molecules refer to nematode transglutaminase proteins and nematode transglutaminase nucleic acid molecules derived from parasitic nematodes. The proteins and nucleic acid molecules of the present invention can be obtained from their natural source, or they can be produced using, for example, recom-

binant nucleic acid technology (also referred to herein as recombinant DNA technology) or chemical synthesis. The terms non-native parasitic nematode transglutaminase protein or non-native parasitic nematode protein, as used herein, refer to a parasitic nematode transglutaminase protein which is produced either synthetically or by transcribing a molecularly cloned or chemically synthesized parasitic nematode transglutaminase or nucleic acid molecule of the present invention (in other words, by recombinant DNA technology). Also included in the present invention is the use of these proteins, nucleic acid molecules, antibodies and other inhibitors as therapeutic compositions to protect animals from parasitic nematode diseases as well as in other applications, such as those disclosed below. An entirely unexpected finding with respect to nematode transglutaminase proteins of the present invention, herein disclosed for the first time, is that a nematode transglutaminase protein has protein disulfide isomerase (PDI) activity.

Parasitic nematode transglutaminase proteins and nucleic acid molecules of the present invention have utility because they represent novel targets for anti-parasite vaccines and drugs. The products and processes of the present invention are advantageous because they enable the inhibition of crucial steps in nematode molting that involve nematode transglutaminase. While not being bound by theory, it is believed that nematode transglutaminase protein activity is essential for successful development of nematode larvae. In addition, the unexpected finding that nematode transglutaminase proteins have PDI activity supports the use of transglutaminase proteins of the present invention having PDI activity as targets for potential anti-parasite vaccines or therapeutics.

One embodiment of the present invention is an isolated protein comprising a *D. immitis* transglutaminase protein. It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, a protein refers to one or more proteins or at least one protein. The terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. Furthermore, a compound "selected from the group consisting of" refers to one or more of the compounds in the list that follows, including mixtures (i.e., combinations) of two or more of the compounds.

According to the present invention, an isolated, or biologically pure, protein, is a protein that has been removed from its natural milieu. Accordingly, "isolated" and "biologically pure" do not necessarily reflect the extent to which the protein has been purified. An isolated protein of the present invention can be obtained from its natural source, can be produced using recombinant DNA technology or can be produced by chemical synthesis. When an isolated protein of the present invention is produced using recombinant DNA technology or produced by chemical synthesis, the protein is referred to herein as either an isolated protein or as a non-native protein.

As used herein, an isolated parasitic nematode transglutaminase protein can be a full-length protein or any homolog of such a protein. An isolated protein of the present invention, including a homolog, can be identified in a straightforward manner by the protein's ability to elicit an immune response against parasitic nematode transglutaminase proteins, to exhibit transglutaminase activity, or to have any combination of these characteristics. Examples of parasitic nematode transglutaminase homologs include parasitic nematode transglutaminase proteins in which amino acids have been deleted (e.g., a truncated version of the protein,

such as a peptide), inserted, inverted, substituted, derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation, addition of glycerophosphatidyl inositol), or any combination of the above, so that the homolog includes at least one epitope capable of eliciting an immune response against a parasitic nematode transglutaminase protein. In other words, when the homolog is administered to an animal as an immunogen, using techniques known to those skilled in the art, the animal will produce an immune response against at least one epitope of a natural parasitic nematode transglutaminase protein. The ability of a protein to effect an immune response can be measured using techniques known to those skilled in the art. Techniques to measure parasitic nematode transglutaminase activity are also known to those skilled in the art, and are described in the Examples.

Parasitic nematode transglutaminase protein homologs can be the result of natural allelic variation or natural mutation. Nematode transglutaminase protein homologs of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein or modifications to the gene encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

Isolated proteins of the present invention have the further characteristic of being encoded by nucleic acid molecules that hybridize under stringent hybridization conditions to a gene encoding a *D. immitis*, a *B. malayi* or an *O. volvulus* nematode transglutaminase protein (i.e., to a *D. immitis*, a *B. malayi* or an *O. volvulus* nematode transglutaminase gene). As used herein, stringent hybridization conditions refer to standard hybridization conditions under which nucleic acid molecules, including oligonucleotides, are used to identify molecules having similar nucleic acid sequences. Stringent hybridization conditions typically permit isolation of nucleic acid molecules having at least about 70% nucleic acid sequence identity with the nucleic acid molecule being used as a probe in the hybridization reaction. Standard conditions are disclosed, for example, in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press. The reference Sambrook et al., *ibid.*, is incorporated by reference herein in its entirety. Formulae to calculate the appropriate hybridization and wash conditions to achieve hybridization permitting 30% or less mismatch of nucleotides are disclosed, for example, in Meinkoth et al., 1984, *Anal. Biochem.* 138, 267-284; Meinkoth et al., *ibid.*, is incorporated by reference herein in its entirety.

As used herein, a parasitic nematode transglutaminase gene includes all nucleic acid sequences related to a natural parasitic nematode transglutaminase gene including, for example, regulatory regions that control production of the nematode transglutaminase protein encoded by that gene (such as, but not limited to, transcription, translation or post-translation control regions), as well as the coding region itself. A *D. immitis* gene can include nDiTG₇₀₇, nDiTG₁₄₇₂, nDiTG₁₄₃, nDiTG₁₄₀₇, nDiTG₁₈₈₁ or nDiTG₁₄₉₄; a *B. malayi* gene can include nBmTG₅₃₇ or nBmTG₄₄₀, and an *O. volvulus* gene can include nOvTG₅₃₇. In one embodiment, a parasitic nematode transglutaminase gene of the present invention includes the nucleic acid sequence SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:54 or SEQ ID NO:57, as well as the complement of these sequences (i.e. SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:29,

SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56 or SEQ ID NO:59, respectively). The nucleic acid sequence SEQ ID NO:8 represents the deduced sequence of the coding strand of the apparent coding region of a cDNA (complementary DNA) molecule denoted herein as nDiTG₇₀₅, the production of which is disclosed in the Examples. The complement of SEQ ID NO:8 (represented herein by SEQ ID NO:9) refers to the nucleic acid sequence of the strand complementary to the strand having SEQ ID NO:8, the sequence of which can easily be determined from SEQ ID NO:8 by those skilled in the art. Likewise, a nucleic acid sequence complement of any nucleic acid sequence of the present invention refers to the nucleic acid sequence of the nucleic acid strand that is complementary to (i.e., can form a double helix with) the strand for which the sequence is cited.

Nucleic acid sequence SEQ ID NO:13 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nDiTG₁₁₀₇, the production of which is disclosed in the Examples. The complement of SEQ ID NO:13 is represented herein by SEQ ID NO:14.

It should be noted that because nucleic acid and amino acid sequencing technology is not entirely error-free, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:54 and SEQ ID NO:57, as well as other nucleic acid and protein sequences presented herein, represent apparent nucleic acid sequences of the nucleic acid molecules encoding a parasitic nematode transglutaminase protein of the present invention, and apparent amino acid sequences of the proteins of the present invention, respectively.

In another embodiment, a nematode transglutaminase gene can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59. An allelic variant of a nematode transglutaminase gene including SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59, is a gene that occurs at essentially the same locus (or loci) in the genome as the gene including SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59, but which, due to natural variations caused by, for

example, mutation or recombination, has a similar but not identical sequence. Allelic variants typically encode proteins having similar activity to that of the protein encoded by the gene to which they are being compared. Allelic variants can also comprise alterations in the 5' or 3' untranslated regions of the gene (e.g., in regulatory control regions). Allelic variants are well known to those skilled in the art and would be expected to be found within a given parasitic nematode, or among a group of two or more parasitic nematodes, because of the diploid nematode genome.

The minimum size of a nematode transglutaminase protein homolog of the present invention is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid (i.e., hybridize under stringent hybridization conditions) with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. The size of the nucleic acid molecule encoding such a protein homolog is dependent on nucleic acid composition and percent homology between the nucleic acid molecule and a complementary sequence. It should also be noted that the extent of homology required to form a stable hybrid can vary depending on whether the homologous sequences are interspersed throughout the nucleic acid molecules or are clustered (i.e., localized) in distinct regions on the nucleic acid molecules. The minimum size of nucleic acid molecules that can form stable hybrids under standard hybridization conditions is typically at least about 12 to about 15 nucleotides in length if the nucleic acid molecules are GC-rich, and at least about 15 to about 17 bases in length if they are AT-rich. Therefore, the minimum size of a nucleic acid molecule used to encode a nematode transglutaminase protein homolog of the present invention is from about 12 to about 18 nucleotides in length. Accordingly, the minimum size of a nematode transglutaminase protein homolog of the present invention is from about 4 to about 6 amino acids in length. There is no limit, other than a practical limit, on the maximum size of a nucleic acid molecule of the present invention because nucleic acid molecules of the present invention can include a portion of a gene, an entire gene, multiple genes, or portions thereof. The preferred size of a protein encoded by a nucleic acid molecule of the present invention depends on whether a full-length, fusion, multivalent, or functional portion of a protein is desired.

Suitable parasitic nematodes from which to isolate nematode transglutaminase proteins of the present invention (including isolation of the natural protein or production of the natural or non-native protein by recombinant or synthetic techniques) include filarioid, ancylostomatoid, ascaridoid, dioctophymatoid, dracunculoid, metastrongyloid, oxyuroid, physalopteroid, rhabditoid, spiruroid, strongyloid, thelazoid, trichinelloid, and trichostrongyloid nematodes. Particularly preferred nematodes are those of the genera *Dirofilaria*, *Onchocerca*, *Brugia*, *Acanthocheilonema*, *Aelurostrongylus*, *Ancylostoma*, *Angiostrongylus*, *Ascaris*, *Bunostomum*, *Capillaria*, *Chabertia*, *Cooperia*, *Crenosoma*, *Dictyocaulus*, *Dioctophyme*, *Dipetalonema*, *Dracunculus*, *Enterobius*, *Filaroides*, *Haemonchus*, *Lagochilascaris*, *Loa*, *Mansonella*, *Muellerius*, *Necator*, *Nematodirus*, *Oesophagostomum*, *Ostertagia*, *Parafilaria*, *Parascaris*, *Physaloptera*, *Protostrongylus*, *Setaria*, *Spirocerca*, *Stephanofilaria*, *Strongyloides*, *Strongylus*, *Thelazia*, *Toxascaris*, *Toxocara*, *Trichinella*, *Trichostrongylus*, *Trichuris*, *Uncinaria*, and *Wuchereria*. Particularly preferred are filarioid nematodes including *Dirofilaria*, *Onchocerca*, *Brugia*, *Acanthocheilonema*, *Dipetalonema*, *Loa*, *Mansonella*, *Parafilaria*, *Setaria*, *Stephanofilaria*, and *Wuchereria*, with *D. immitis*, *B. malayi*, *O. volvulus* and *T.*

canis being even more preferred, and *D. immitis* being particularly preferred.

A preferred parasitic nematode transglutaminase protein of the present invention is a compound that is not substantially toxic to host animals (that is, does not substantially inhibit host animal transglutaminase; the term, "does not substantially inhibit" as used herein can be used interchangeably with the term, "inability to substantially interfere"; a compound that does not substantially inhibit host animal transglutaminase activity is one that, when administered to a host animal, the host animal shows no significant adverse effects attributable to the compound) and which, when administered to an animal in an effective manner, is capable of protecting that animal from disease caused by a parasitic nematode. In accordance with the present invention, the ability of a nematode transglutaminase protein of the present invention to protect an animal from disease by a parasitic nematode refers to the ability of that protein to, for example, treat, ameliorate or prevent disease caused by parasitic nematodes. In particular, the phrase, "protect an animal from disease by a parasitic nematode," refers to reducing the potential for parasitic nematode population expansion in the host animal by inhibiting parasitic nematode molting and subsequent growth. Nematode molting is an essential step in the life cycle and development of all nematodes, and characterizes the progression of the nematode larvae through the development of larval stages to the adult. A host animal, as used herein, is an animal in which a parasitic nematode can live and multiply. In one embodiment, a nematode transglutaminase protein of the present invention can elicit an immune response (including a humoral or cellular immune response, or both) against a parasitic nematode.

Suitable nematodes to target with therapeutic compounds of the present invention include any nematodes that are essentially incapable of molting, or are inhibited in the ability to molt, in a host animal when a nematode transglutaminase protein of the present invention, or inhibitor of such a protein, has been administered to that animal. Accordingly, a nematode to target includes any nematode that produces a protein having one or more epitopes that can be neutralized by either a humoral or a cellular immune response, or both, elicited by a nematode transglutaminase protein of the present invention, or that produces a protein that can be targeted by a compound that otherwise inhibits nematode transglutaminase activity, thereby resulting in the decreased ability of the nematode to cause disease in an animal. Preferred nematodes to target include parasitic nematodes disclosed herein as being useful in the production or isolation of parasitic nematode transglutaminase proteins of the present invention.

The present invention also includes mimetopes of parasitic nematode transglutaminase proteins of the present invention. As used herein, a mimetope of a parasitic nematode transglutaminase protein of the present invention refers to any compound that is able to mimic the activity of such a parasitic nematode transglutaminase protein (e.g., has the ability to elicit an immune response against a parasitic nematode transglutaminase protein of the present invention or ability to inhibit parasitic nematode transglutaminase activity). The ability to mimic the activity of a parasitic nematode transglutaminase protein is likely to be the result of a structural similarity between the parasitic nematode transglutaminase protein and the mimetope. It is to be noted, however, that the mimetope need not have a structure similar to a parasitic nematode transglutaminase protein as long as the mimetope functionally mimics the protein. Mimetopes

can be, but are not limited to: peptides that have been modified to decrease their susceptibility to degradation; anti-idiotypic and/or catalytic antibodies, or fragments thereof; non-proteinaceous immunogenic portions of an isolated protein (e.g., carbohydrate structures); synthetic or natural organic or inorganic molecules, including nucleic acids; and/or any other peptidomimetic compounds. Mimetopes of the present invention can be designed using computer-generated structures of parasitic nematode transglutaminase proteins of the present invention. Mimetopes can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides or other organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner, (e.g., an anti-parasitic nematode transglutaminase antibody). A preferred mimetope is a peptidomimetic compound that is structurally and/or functionally similar to a parasitic nematode transglutaminase protein of the present invention, particularly to the active site of the parasitic nematode transglutaminase protein.

One embodiment of a parasitic nematode transglutaminase protein of the present invention is a fusion protein that includes a parasitic nematode transglutaminase protein-containing domain attached to one or more fusion segments. Suitable fusion segments for use with the present invention include, but are not limited to, segments that can: enhance a protein's stability; act as an immunopotentiator to enhance an immune response against a nematode transglutaminase protein; assist in purification of a nematode transglutaminase protein (e.g., by affinity chromatography); or any combination of the above listed functions. A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, imparts increased immunogenicity to a protein, or simplifies purification of a protein). Fusion segments can be joined to amino or carboxyl termini, or both, of the nematode transglutaminase-containing domain of the protein and can be susceptible to cleavage in order to enable straightforward recovery of a nematode transglutaminase protein. Fusion proteins are preferably produced by culturing a recombinant cell transformed with a fusion nucleic acid molecule that encodes a protein including the fusion segment attached to either or both of the carboxyl or amino terminal ends of a nematode transglutaminase-containing domain. Preferred fusion segments include a metal binding domain (e.g., a poly-histidine segment); an immunoglobulin binding domain (e.g., Protein A; Protein G; T cell; B cell; Fc receptor; or complement protein antibody-binding domains); a sugar binding domain (e.g., a maltose-binding domain); a "tag" domain (e.g., in at least a portion of β -galactoside, a strep tag peptide, other domains that can be purified using compounds that bind to the domain, such as monoclonal antibodies), or any combination of the above listed fusion segments. More preferred fusion segments include metal binding domains, such as a poly-histidine segment; a maltose-binding domain; a strep tag peptide, such as that available from Biometra in Tampa, Fla.; and an S10 peptide.

In another embodiment, a parasitic nematode transglutaminase protein of the present invention also includes at least one additional protein segment that is capable of protecting an animal from one or more diseases such as a multivalent protective protein can be produced by culturing a cell transformed with a nucleic acid molecule comprising two or more nucleic acid domains joined together in such a manner that the resulting nucleic acid molecule is expressed as a multivalent protective compound containing at least two protective compounds, or portions thereof, capable of pro-

tecting an animal from diseases caused, for example, by at least one other infectious agent.

Examples of multivalent protective compounds include, but are not limited to, a parasitic nematode transglutaminase protein of the present invention attached to one or more compounds protective against one or more other infectious agents, particularly an agent that infects humans, cats, dogs, cattle, sheep pigs, goats or horses, such as, but not limited to: viruses (e.g., adenoviruses, caliciviruses, coronaviruses, distemper viruses, hepatitis viruses, herpesviruses, immunodeficiency viruses, infectious peritonitis viruses, leukemia viruses, oncogenic viruses, panleukopenia viruses, papilloma viruses, parainfluenza viruses, parvoviruses, rabies viruses, and reoviruses, as well as other cancer-causing or cancer-related viruses); bacteria (e.g., Actinomyces, Bacillus, Bacteroides, Bordetella, Bartonella, Brucella, Campylobacter, Capnocytophaga, Clostridium, Corynebacterium, Coxiella, Dermatophilus, Enterococcus, Ehrlichia, Escherichia, Francisella, Fusobacterium, Haemobartonella, Helicobacter, Klebsiella, L-form bacteria, Leptospira, Listeria, Mycobacteria, Mycoplasma, Neorickettsia, Nocardia, Pasteurella, Peptococcus, Peptostreptococcus, Proteus, Pseudomonas, Rickettsia, Rochalimaea, Salmonella, Shigella, Staphylococcus, Streptococcus, and Yersinia; fungi and fungal-related microorganisms (e.g., Absidia, Acremonium, Alternaria, Aspergillus, Basidiobolus, Bipolaris, Blastomyces, Candida, Chlamydia, Coccidioides, Conidiobolus, Cryptococcus, Curvularia, Epidermophyton, Exophiala, Geotrichum, Histoplasma, Madurella, Malassezia, Microsporum, Moniliella, Mortierella, Mucor, Paecilomyces, Penicillium, Phialemonium, Phialophora, Prototheca, Pseudallescheria, Pseudomicrodochium, Pythium, Rhinosporidium, Rhizopus, Scolecobaidium, Sporothrix, Stemphylium, Trichophyton, Trichosporon, and Xylohypha); and other parasites (e.g., Babesia, Balantidium, Besnoitia, Cryptosporidium, Eimeria, Encephalitozoon, Entamoeba, Giardia, Hammondia, Hepatozoon, Isospora, Leishmania, Microsporidia, Nosepora, Nosema, Pentatrichomonas, Plasmodium, Pneumocystis, Sarcocystis, Schistosoma, Theileria, Toxoplasma, and Trypanosoma, as well as other nematode parasites, including, but not limited to those disclosed herein). In one embodiment, a parasitic nematode transglutaminase protein of the present invention is attached to one or more additional compounds protective against parasitic nematode disease. In another embodiment one or more protective compounds, such as those listed above, can be included in a multivalent vaccine comprising a parasitic nematode transglutaminase protein of the present invention and one or more other protective molecules as separate compounds.

A preferred isolated protein of the present invention is a protein encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nDiTG₇₀₇, nDiTG₇₀₅, nDiTG₁₄₇₂, nDiTG₁₁₀₇, nDiTG₁₄₃, nDiTG₄₅, nDiTG₁₂₀, nDiTG₁₄₀₇, nDiTG₁₈₈₁, nDiTG₁₄₉₄, nDiTG₁₄₁₆, nBmTG₄₄₀, nBmTG₄₁₇, nBmTG₃₃₉, nBmTG₅₃₇ or nOvTG₁₃₇. A further preferred isolated protein is encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56 or SEQ ID NO:59.

Translation of SEQ ID NO:5 suggests that nucleic acid molecule nDiTG₇₀₇ encodes a partial-length parasitic nema-

tode transglutaminase protein of about 235 amino acids, referred to herein as PDiTG₂₃₅, represented by SEQ ID NO:6, assuming an open reading frame having a first codon spanning from about nucleotide 1 through about nucleotide 3 of SEQ ID NO:5. The coding region encoding PDiTG₂₃₅ is represented by nucleic acid molecule nDiTG₇₀₅, having the nucleic acid sequence represented by SEQ ID NO:8 (the coding strand) and SEQ ID NO:9 (the complementary strand). The deduced amino acid sequence (represented by SEQ ID NO:6) suggests a protein having a molecular weight of about 27.2 kilodaltons (kD) and an estimated pI of about 5.07.

The amino acid sequence of PDiTG₂₃₅ includes a thioredoxin family active site from residues about 24 to 30. Thioredoxins participate in various redox reactions through the reversible oxidation of an active center disulfide bond Holmgren, A., 1985 *Annual Review of Biochemistry*, 54, 237–271. A number of eukaryotic proteins contain domains evolutionarily related to thioredoxin.

Translation of SEQ ID NO:10 suggests that nucleic acid molecule nDiTG₄₇₂ encodes a partial-length parasitic nematode transglutaminase protein of about 368 amino acids, referred to herein as PDiTG₃₆₈, represented by SEQ ID NO:11, assuming an open reading frame having a first codon spanning from about nucleotide 2 through about nucleotide 4 of SEQ ID NO:10, and a putative stop codon spanning from about nucleotide 1105 through nucleotide 1107 of SEQ ID NO:10. The coding region encoding PDiTG₃₆₈ (including a putative stop codon) is represented by nucleic acid molecule nDiTG₁₁₀₇, having the nucleic acid sequence represented by SEQ ID NO:13 (the coding strand) and SEQ ID NO:14 (the complementary strand). The deduced amino acid sequence SEQ ID NO:11 suggests a protein having a molecular weight of about 42.6 kD and an estimated pI of about 5.71.

The amino acid sequence of PDiTG₃₆₈ (i.e., SEQ ID NO:11) includes: i) a thioredoxin family active site detected from residues 268 to 274; ii) an endoplasmic reticulum (ER) targeting sequence from residues 365 to 368 (KEEL) (proteins that permanently reside in the lumen of ER seem to be distinguished from newly synthesized secretory proteins by the presence of the C-terminal sequence Lys-Asp-Glu-Leu (KDEL); see, for example, Munro et al., 1987, *Cell* 48,899–907; Pelham, 1990, *Trends in Biochemical Sciences*, 15,483–486; and iii) a tachykinin family signature from residues 186 to 202 (tachykinins are a group of biologically active peptides that excite neurons, evoke behavioral responses, are potent vasodilators, and contract many smooth muscles; see, for example, Maggio, 1988, *Annual Review of Neurosciences*, 11,13–28).

More preferred parasitic nematode transglutaminase proteins of the present invention include proteins comprising amino acid sequences that are at least about 80%, preferably at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to amino acid sequence SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, or SEQ ID NO:4. These sequences are described in the Examples. Even more preferred parasitic nematode transglutaminase proteins of the present invention include proteins comprising amino acid sequences that are at least about 50%, preferably at least about 60%, more preferably at least about 70%, more preferably at least about 80%, more preferably at least about 90%, and even more preferably at least about 95% identical to amino acid sequence SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58.

More preferred parasitic nematode transglutaminase proteins of the present invention include proteins encoded by a nucleic acid molecule comprising at least a portion of nDiTG₇₀₇, nDiTG₇₀₅, nDiTG₁₄₇₂, nDiTG₁₁₀₇, nDiTG₁₄₃, nDiTG₄₅, nDiTG₁₂₀, nDiTG₁₄₀₇, nDiTG₁₈₈₁, nDiTG₁₄₉₄, nDiTG₁₄₁₆, nBmTG₄₄₀, nBmTG₄₁₇, nBmTG₃₃₉, nBmTG₅₃₇ and nOvTG₅₃₇, or at least a portion of allelic variants of these nucleic acid molecules. In one embodiment, a preferred nematode transglutaminase protein of the present invention is encoded by at least a portion of SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:10, or SEQ ID NO:13, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:54 or SEQ ID NO:57, and has an amino acid sequence that includes at least a portion of SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58. Also preferred is a protein encoded by an allelic variant of a nucleic acid molecule comprising at least a portion of SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:10, or SEQ ID NO:13, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:54 or SEQ ID NO:57. Particularly preferred proteins of the present invention are those comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58, and proteins encoded by an allelic variant of a nucleic acid molecule encoding any of these amino acid sequences. Also preferred is an isolated *D. immitis* transglutaminase protein. An isolated *D. immitis* transglutaminase protein of the present invention can be either native or can be chemically synthesized or produced in a cell transformed with a nucleic acid molecule encoding a *D. immitis* transglutaminase protein.

Another embodiment of the present invention is an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a parasitic nematode transglutaminase gene, and particularly with a *D. immitis*, *B. malayi* or *O. volvulus* transglutaminase gene. The identifying characteristics of such a gene are herein described. A nucleic acid molecule of the present invention can include an isolated natural parasitic nematode transglutaminase gene or a homolog thereof, the latter of which is described in more detail below. A nucleic acid molecule of the present invention can include one or more regulatory regions, full-length or partial coding regions, or any combinations thereof. The minimum size of a nucleic acid molecule of the present invention is the minimum size that can form a stable hybrid with a parasitic nematode transglutaminase gene under stringent hybridization conditions. Suitable and preferred parasitic nematodes are disclosed above.

In accordance with the present invention, an isolated nucleic acid molecule is a nucleic acid molecule that has been removed from its natural milieu (i.e., that has been subjected to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA. As used herein, the term, “isolated,” does not reflect the extent to which the nucleic acid molecule has been purified. An isolated parasitic nematode transglutaminase nucleic acid molecule of the present invention can be isolated from its natural source or can be produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning) or

chemical synthesis. Isolated nematode transglutaminase nucleic acid molecules can include, for example, natural allelic variants and nucleic acid molecules modified by nucleotide insertions, deletions, substitutions, inversions, variants created during PCR amplification, or any combination of the above modifications. According to the present invention, acceptable modifications to nematode transglutaminase nucleic acid molecules do not substantially interfere with the nucleic acid molecule's ability to encode a nematode transglutaminase protein of the present invention or to form stable hybrids under stringent conditions with natural parasitic nematode transglutaminase gene isolates.

A parasitic nematode transglutaminase nucleic acid molecule homolog can be produced using a number of methods known to those skilled in the art (see, for example, Sambrook et al., *ibid.*). For example, nucleic acid molecules can be modified using a variety of techniques including, but not limited to, classic mutagenesis and recombinant DNA techniques (e.g., site-directed mutagenesis, chemical treatment, restriction enzyme cleavage, ligation of nucleic acid fragments, and PCR amplification), or synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules and combinations thereof. Nucleic acid molecule homologs can be selected by hybridization with a nematode transglutaminase gene or by screening expression products of the nematode transglutaminase nucleic acid molecule homologs for the function of a protein encoded by the nucleic acid molecule (e.g., the ability to elicit an immune response against at least one epitope of a parasitic nematode transglutaminase protein or parasitic nematode transglutaminase activity).

An isolated nucleic acid molecule of the present invention can include a nucleic acid sequence that encodes at least one parasitic nematode transglutaminase protein of the present invention; examples of such proteins are herein disclosed. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides in the nucleic acid molecule, the two phrases can be used interchangeably, especially with respect to a nucleic acid molecule, or a nucleic acid sequence, being capable of encoding a parasitic nematode transglutaminase protein.

A preferred nucleic acid molecule of the present invention, when administered to an animal, is substantially not toxic to the animal and is capable of protecting that animal from disease caused by a parasitic nematode. As will be disclosed in more detail below, such a nucleic acid molecule can be, or encode, an antisense RNA, a molecule capable of triple helix formation, a ribozyme, or other nucleic acid-based drug compound. In additional embodiments, a nucleic acid molecule of the present invention can encode a protective protein (e.g., a nematode transglutaminase protein of the present invention), the nucleic acid molecule being delivered to the animal, for example, by direct injection (i.e., as a composition comprising a naked nucleic acid molecule of the present invention) or in a vehicle such as a recombinant virus vaccine or a recombinant cell vaccine.

One embodiment of the present invention is a parasitic nematode transglutaminase nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nDiTG₇₀₇, and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:5 or SEQ ID NO:7. Such a nucleic acid molecule would also hybridize with nDiTG₇₀₅, and thus would also hybridize with SEQ ID NO:8 or SEQ ID NO:9. Comparison of nucleic acid sequence SEQ ID NO:5 (i.e., the nucleic acid

sequence of the coding strand of nDiTG₇₀₇) with nucleic acid sequences reported in GenBank™ indicates that SEQ ID NO:5 showed the most homology (i.e., about 37% identity) with human clone PA3 (GenBank™ accession number J05016), a protein disulfide isomerase related to protein (Erp72) mRNA.

Another embodiment of the present invention is a parasitic nematode transglutaminase nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nDiTG₁₄₂₇, and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:10 or SEQ ID NO:12. Such a nucleic acid molecule would also hybridize with nDiTG₁₁₀₇, and thus would also hybridize with SEQ ID NO:13 or SEQ ID NO:14. Comparison of nucleic acid sequence SEQ ID NO:10 (i.e., the nucleic acid sequence of the coding strand of nDiTG₁₄₂₇) with nucleic acid sequences reported in GenBank™ indicates that SEQ ID NO:10 showed the most homology (i.e., about 63% sequence identity) with a human epithelial cell mRNA for ER-60 protease (GenBank™ accession number D83485), spanning from nucleotide about 1143 to about 1458 of the ER-60 protease.

Preferred parasitic nematode transglutaminase nucleic acid molecules include nucleic acid molecules having a nucleic acid sequence that is at least about 70%, preferably at least about 75%, more preferably at least about 80%, more preferably at least about 85%, even more preferably at least about 90% and even more preferably at least about 95% identical to nucleic acid sequence SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59.

Another preferred embodiment of the present invention includes at least a portion of a nucleic acid sequence SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59 that is capable of hybridizing with a nematode transglutaminase gene of the present invention, as well as allelic variants thereof. Such nucleic acid molecules can include nucleotides in addition to those included in the sequences listed above, such as, but not limited to, a full-length gene, a full-length coding region, a nucleic acid molecule encoding a fusion protein, or a nucleic acid molecule encoding a multivalent protective compound. Particularly preferred nucleic acid molecules include nDiTG₇₀₇, nDiTG₇₀₅, nDiTG₁₄₇₂, nDiTG₁₁₀₇, nDiTG₁₄₃, nDiTG₁₂₀, nDiTG₁₄₀₇, nDiTG₁₈₈₁, nDiTG₁₄₉₄, nDiTG₁₄₁₆, nBmTG₄₄₀, nBmTG₄₁₇, nBmTG₃₃₉, nBmTG₅₃₇ or nOvTG₅₃₇, and allelic variants of these nucleic acid molecules. Also particularly preferred nucleic acid molecules include those including SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35,

SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57 or SEQ ID NO:59, and allelic variants of these preferred nucleic acid molecules.

The present invention also includes a nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58, including allelic variants of these sequences and nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed. Particularly preferred are nucleic acid molecules that encode amino acid sequences including those represented by SEQ ID NO:6, SEQ ID NO:11, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:52, SEQ ID NO:55 or SEQ ID NO:58, and allelic variants of these nucleic acid molecules.

Knowing the nucleic acid sequences of certain parasitic nematode transglutaminase nucleic acid molecules of the present invention allows one skilled in the art to, for example, (a) make copies of those nucleic acid molecules, (b) obtain nucleic acid molecules including at least a portion of such nucleic acid molecules (e.g., nucleic acid molecules including full-length genes, full-length coding regions, regulatory control sequences, truncated coding regions), and (c) obtain parasitic nematode transglutaminase nucleic acid molecules from other parasitic nematodes. Such nucleic acid molecules can be obtained in a variety of ways including screening appropriate expression libraries with antibodies of the present invention; traditional cloning techniques using oligonucleotide probes of the present invention to screen appropriate libraries DNA, or RNA; and PCR amplification of appropriate libraries, DNA, or RNA using oligonucleotide primers of the present invention. Preferred libraries to screen or from which to amplify nucleic acid molecule include adult and larval stage parasitic nematode cDNA libraries as well as genomic DNA libraries. Similarly, preferred DNA or RNA sources to screen or from which to amplify nucleic acid molecules include adult and larval stage parasitic nematode cDNA, adult and larval mRNA, and genomic DNA. Techniques to clone and amplify genes are disclosed, for example, in Sambrook et al., *ibid*, as well as in the Examples section.

The present invention also includes nucleic acid molecules that are oligonucleotides capable of hybridizing, under stringent hybridization conditions, with complementary regions of other, preferably longer, nucleic acid molecules of the present invention such as those comprising parasitic nematode transglutaminase genes or other parasitic nematode transglutaminase nucleic acid molecules. Oligonucleotides of the present invention can be RNA, DNA, or derivatives of either. The minimum size of such oligonucleotides is the size required for formation of a stable hybrid between an oligonucleotide and a complementary sequence on a nucleic acid molecule of the present invention. Minimal size characteristics are disclosed herein. The present invention includes oligonucleotides that can be used as, for example, probes to identify nucleic acid molecules, primers to produce nucleic acid molecules or therapeutic reagents to inhibit nematode transglutaminase protein production or activity (e.g., as antisense-, triplex formation-, ribozyme- and/or RNA drug-based reagents). The present invention also includes the use of such oligonucleotides to protect

animals from disease using one or more of such technologies. Appropriate oligonucleotide-containing therapeutic compositions can be administered to an animal using techniques known to those skilled in the art.

One embodiment of the present invention includes a recombinant vector, which includes at least one isolated nucleic acid molecule of the present invention, inserted into any vector capable of delivering the nucleic acid molecule into a cell. Such a vector contains heterologous nucleic acid sequences, that is nucleic acid sequences that are not naturally found adjacent to nucleic acid molecules of the present invention and that preferably are derived from a species other than the species from which the nucleic acid molecule (s) are derived. The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a virus or a plasmid. Recombinant vectors can be used in the cloning, sequencing, and/or otherwise manipulation of parasitic nematode transglutaminase nucleic acid molecules of the present invention.

One type of recombinant vector, referred to herein as a recombinant molecule, comprises a nucleic acid molecule of the present invention operatively linked to an expression vector. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a cell. As used herein, an expression vector is a DNA or RNA vector that is capable of transforming a cell and of effecting expression of a specified nucleic acid molecule. Preferably, the expression vector is also capable of replicating within the cell. Expression vectors can be either prokaryotic or eukaryotic, and are typically viruses (including viral genomes) or plasmids. Expression vectors of the present invention include any vectors that function (i.e., direct gene expression) in recombinant cells of the present invention, including in bacterial, fungal, endoparasite, insect, other animal, and plant cells. Preferred expression vectors of the present invention can direct gene expression in bacterial, yeast, insect and mammalian cells and more preferably in the cell types disclosed herein.

In particular, expression vectors of the present invention contain regulatory sequences such as transcription control sequences, translation control sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell and that control the expression of nucleic acid molecules of the present invention. In particular, recombinant molecules of the present invention include transcription control sequences. Transcription control sequences are sequences that control the initiation, elongation, and termination of transcription. Particularly important transcription control sequences are those that control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in at least one of the recombinant cells of the present invention. A variety of such transcription control sequences are known to those skilled in the art. Preferred transcription control sequences include those that function in bacterial, yeast, insect and mammalian cells, such as, but not limited to, *tac*, *lac*, *trp*, *trc*, *oxy-pro*, *omp/lpp*, *rrnB*, bacteriophage lambda (such as lambda p_L and lambda p_R and fusions that include such promoters), bacteriophage T7, T7lac, bacteriophage T3, bacteriophage SP6, bacteriophage SP01, metallothionein, alpha-mating factor, *Pichia* alcohol oxidase, alphavirus subgenomic promoters (such as Sindbis virus subgenomic promoters), antibiotic resistance gene, baculovirus, *Heliothis zea* insect virus, vaccinia virus, herpesvirus, raccoon poxvirus, other

poxvirus, adenovirus, cytomegalovirus (such as intermediate early promoters), simian virus 40, retrovirus, actin, retroviral long terminal repeat, Rous sarcoma virus, heat shock, phosphate and nitrate transcription control sequences as well as other sequences capable of controlling gene expression in prokaryotic or eukaryotic cells. Additional suitable transcription control sequences include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters (e.g., promoters inducible by interferons or interleukins). Transcription control sequences of the present invention can also include naturally occurring transcription control sequences naturally associated with parasitic nematodes, for example *D. immitis*.

Suitable and preferred nucleic acid molecules to include in recombinant vectors of the present invention are as disclosed herein. Particularly preferred nucleic acid molecules to include in recombinant vectors, and particularly in recombinant molecules, include nDiTG₇₀₇, nDiTG₇₀₅, nDiTG₁₄₇₂, nDiTG₁₁₀₇, nDiTG₁₄₃, nDiTG₁₂₀, nDiTG₄₅, nDiTG₁₄₀₇, nDiTG₁₈₈₁, nDiTG₁₄₉₄, nDiTG₁₄₁₆, nBmTG₄₄₀, nBmTG₄₁₇, nBmTG₃₃₉, nBmTG₅₃₇ or nOvTG₅₃₇.

Recombinant molecules of the present invention may also (a) contain secretory signals (i.e., signal segment nucleic acid sequences) to enable an expressed nematode transglutaminase protein of the present invention to be secreted from the cell that produces the protein and/or (b) contain fusion sequences that lead to the expression of nucleic acid molecules of the present invention as fusion proteins. Examples of suitable signal segments include any signal segment capable of directing the secretion of a protein of the present invention. Preferred signal segments include, but are not limited to, tissue plasminogen activator (t-PA), interferon, interleukin, growth hormone, histocompatibility and viral envelope glycoprotein signal segments, as well as natural signal segments. Suitable fusion segments encoded by fusion segment nucleic acids are disclosed herein. In addition, a nucleic acid molecule of the present invention can be joined to a fusion segment that directs the encoded protein to the proteasome, such as a ubiquitin fusion segment. Recombinant molecules may also include intervening and/or untranslated sequences surrounding and/or within the nucleic acid sequences of nucleic acid molecules of the present invention. An example of a preferred intervening sequence for eukaryotic gene expression is cytomegalovirus intron A.

Another embodiment of the present invention includes a recombinant cell transformed with one or more recombinant molecules of the present invention. Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. A nucleic acid molecule of the present invention that has been transformed into a cell is referred to herein as a transformed nucleic acid molecule. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or more sites within a chromosome of the transformed (i.e., recombinant) cell in such a manner that their ability to be expressed is retained. Preferred nucleic acid molecules with which to transform a cell include parasitic nematode transglutaminase nucleic acid molecules disclosed herein. Particularly preferred nucleic acid molecules with which to transform a cell include nDiTG₇₀₅, nDiTG₁₁₀₇, nDiTG₁₂₀,

nDiTG₁₄₀₇, nDiTG₁₄₉₄, nBmTG₄₁₇, nBmTG₅₃₇ or nOvTG₅₃₇. Also preferred are nDiTG₁₈₈₁, nDiTG₇₀₇, nDiTG₁₄₇₂, nDiTG₁₄₃, nDiTG₄₅, nDiTG₁₄₁₆, nBmTG₃₃₉ and nBmTG₄₄₀.

Suitable cells to transform include any cell that can be transformed with a nucleic acid molecule of the present invention. A transformed cell of the present invention is also herein referred to as a recombinant cell. Suitable cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule (e.g., nucleic acid molecules encoding one or more proteins of the present invention, other proteins useful in the production of multivalent vaccines, or a combination thereof). Suitable cells for transformation according to the present invention can be either a) endogenously (i.e., naturally) capable of producing parasitic nematode transglutaminase proteins of the present invention, or b) capable of producing such proteins after transformation with at least one nucleic acid molecule of the present invention. Cells of the present invention can be any cell capable of producing at least one protein of the present invention, and include bacterial, fungal (including yeast), insect, other nematode, other and plant cells. Preferred cells for transformation by nucleic acid molecules of the present invention include bacterial, mycobacterial, yeast, parasite, insect and mammalian cells. More preferred cells include *Salmonella*, *Escherichia*, *Bacillus*, *Listeria*, *Saccharomyces*, *Spodoptera*, *Mycobacteria*, *Trichoplusia*, BHK (baby hamster kidney) cells, MDCK cells (normal dog kidney cell line for canine herpesvirus cultivation), CRFK cells (normal cat kidney cell line for feline herpesvirus cultivation), CV-1 cells (African monkey kidney cell line used, for example, to culture raccoon poxvirus), COS (e.g., COS-7) cells, and Vero cells. Particularly preferred cells for transformation are *Escherichia coli*, including *E. coli* K-12 derivatives; *Salmonella typhi*; *Salmonella typhimurium*, including attenuated strains such as UK-1_{x3987} and SR-11_{x4072}; *Spodoptera frugiperda*; *Trichoplusia ni*; BHK cells; MDCK cells; CRFK cells; CV-1 cells; COS cells; Vero cells; and non-tumorigenic mouse myoblast G8 cells (e.g., ATCC CRL 1246). Additional appropriate mammalian cells suitable for transformation by nucleic acid molecules of the present invention include other kidney cell lines, other fibroblast cell lines (e.g., human, murine or chicken embryo fibroblast cell lines), myeloma cell lines, Chinese hamster ovary cells, mouse NIH3T3 cells, LMTIK³¹ cells and/or HeLa cells. In one embodiment, the proteins may be expressed as heterologous proteins in myeloma cell lines employing immunoglobulin promoters.

A recombinant cell is preferably produced by transforming a suitable cell with one or more recombinant molecules, each comprising one or more nucleic acid molecules of the present invention operatively linked to an expression vector containing one or more transcription control sequences. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a suitable cell as described above.

A recombinant molecule of the present invention is a molecule that can include at least one of any parasitic nematode transglutaminase nucleic acid molecule herein described, operatively linked to at least one of any transcription control sequence capable of effectively regulating expression of the nucleic acid molecule(s) in the cell suitable for transformation, examples of which are disclosed herein.

A recombinant cell of the present invention includes any cell transformed with at least one of any nucleic acid molecule of the present invention. Suitable and preferred

nucleic acid molecules as well as suitable and preferred recombinant molecules with which to transform cells are disclosed herein.

Recombinant cells of the present invention can also be co-transformed with one or more recombinant molecules including parasitic nematode transglutaminase nucleic acid molecules encoding one or more proteins of the present invention and one or more other nucleic acid molecules encoding other protective compounds, as disclosed herein (e.g., to produce multivalent vaccines).

Recombinant DNA technologies can be used to improve expression of transformed nucleic acid molecules by manipulating, for example, the number of copies of the nucleic acid molecules within a transformed cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Recombinant techniques useful for increasing the expression of nucleic acid molecules of the present invention include, but are not limited to, operatively linking nucleic acid molecules of the present invention to nucleic acid molecules that direct the production of a high-copy number of plasmids, integration of the nucleic acid molecules into one or more chromosomes in the transformed cell, addition of vector stability sequences to plasmids containing nucleic acid sequences of the present invention, substitutions or modifications of transcription control signals (e.g., promoters, operators, enhancers), substitutions or modifications of translational control signals (e.g., ribosome binding sites, Shine-Dalgarno sequences), modification of nucleic acid molecules of the present invention to correspond to the codon usage of the transformed cell, deletion of sequences that destabilize transcripts, and use of control signals that temporally separate recombinant cell growth from recombinant enzyme production during fermentation. The activity of an expressed recombinant protein of the present invention may be improved by fragmenting, modifying, or derivatizing nucleic acid molecules encoding such a protein.

Isolated nematode transglutaminase proteins of the present invention can be produced in a variety of ways, including production and recovery of natural proteins, production and recovery of recombinant proteins, and chemical synthesis of the proteins. In one embodiment, an isolated parasitic nematode transglutaminase protein of the present invention is produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering the protein. A preferred cell to culture is a recombinant cell of the present invention. Effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. An effective medium refers to any medium in which a cell is cultured to produce a parasitic nematode transglutaminase protein of the present invention. Such medium typically comprises an aqueous medium having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art.

Depending on the vector and transformed cell system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be

secreted into the fermentation medium; be secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or be retained on the outer surface of a cell or viral membrane. The phrase "recovering the protein", as well as similar phrases, can refer to collecting the whole fermentation medium containing the protein and need not imply additional steps of separation or purification. Proteins of the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, immunoaffinity chromatography, thermoprecipitation, ammonium sulphate precipitation, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization. Proteins of the present invention are preferably retrieved in "substantially pure" form. As used herein, "substantially pure" refers to a purity that allows for the effective use of the protein as a therapeutic composition or diagnostic. A therapeutic composition for animals, for example, should exhibit no substantial toxicity and preferably should be capable of stimulating the production of antibodies in a treated animal.

The present invention also includes isolated (i.e., removed from their natural Milieu) antibodies that selectively bind to a parasitic nematode transglutaminase protein of the present invention or a mimotope thereof (e.g., anti-parasitic nematode transglutaminase antibodies). As used herein, the term "selectively binds to" a nematode transglutaminase protein refers to the ability of antibodies of the present invention to preferentially bind to specified proteins and mimitopes thereof of the present invention. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays (e.g., ELISA), immunoblot assays, etc.; see, for example, Sambrook et al., *ibid.* An anti-parasitic nematode transglutaminase antibody preferably selectively binds to a parasitic nematode transglutaminase protein in such a way as to reduce the activity of that protein.

Isolated antibodies of the present invention can include antibodies in serum, or antibodies that have been purified to varying degrees. Antibodies of the present invention can be polyclonal or monoclonal, or can be functional equivalents such as antibody fragments and genetically-engineered antibodies, including single chain antibodies or chimeric antibodies that can bind to more than one epitope.

A preferred method to produce antibodies of the present invention includes (a) administering to an animal an effective amount of a protein (ranging in size from a peptide to a full length protein) or mimotope thereof of the present invention to produce the antibodies and (b) recovering the antibodies. In another method, antibodies of the present invention are produced recombinantly using techniques as disclosed to produce parasitic nematode transglutaminase proteins of the present invention. Antibodies raised against defined proteins or mimitopes can be advantageous because such antibodies are not substantially contaminated with antibodies against other substances that might otherwise cause interference in a diagnostic assay or side effects if used in a therapeutic composition.

Antibodies of the present invention have a variety of potential uses that are within the scope of the present invention. For example, such antibodies can be used (a) as therapeutic compounds to passively immunize an animal in order to protect the animal from parasitic nematodes susceptible to treatment by such antibodies, (b) as reagents in assays to detect infection by such nematodes, (c) as tools to screen expression libraries, (d) as tools to recover desired

proteins of the present invention from a mixture of proteins and other contaminants, and (e) for any combination of the above listed uses. Furthermore, antibodies of the present invention can be used to target cytotoxic agents to parasitic nematodes of the present invention in order to directly kill such nematodes. Targeting can be accomplished by conjugating (i.e., stably joining) such antibodies to the cytotoxic agents using techniques known to those skilled in the art. Suitable cytotoxic agents are known to those skilled in the art.

One embodiment of the present invention is a therapeutic composition that, when administered to an animal in an effective manner, is capable of protecting that animal from disease caused by a parasitic nematode. Therapeutic compositions of the present invention include an excipient and at least one of the following protective compounds, an isolated native nematode transglutaminase protein; an isolated non-native nematode transglutaminase protein; a mimotope of a nematode transglutaminase protein; an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a nematode transglutaminase gene; an isolated antibody that selectively binds to a nematode transglutaminase protein, an inhibitor of nematode transglutaminase protein activity identified by its ability to inhibit nematode transglutaminase activity and its inability to substantially interfere with host animal transglutaminase activity, or a mixture thereof (i.e., combination of at least two of the compounds). The term "inability to substantially interfere with" host animal transglutaminase activity, as used herein, refers to the failure of a nematode transglutaminase inhibitor compound to inhibit host animal transglutaminase activity to such a degree that such an inhibitor is not substantially toxic to a host animal when it is administered to that animal. The inability to interfere with host animal transglutaminase activity can be identified by transglutaminase assay *in vitro*, as described in the Examples section. Candidate inhibitors can also be tested for toxicity in standard animal studies. Preferred parasitic nematodes to target are herein disclosed. Examples of proteins, nucleic acid molecules, antibodies and inhibitors of the present invention are disclosed herein.

The present invention also includes a therapeutic composition comprising at least one nematode transglutaminase-based compound of the present invention in combination with at least one additional compound protective against one or more infectious agents. Examples of such compounds and infectious agents are disclosed herein.

Therapeutic compositions of the present invention can be administered to any animal susceptible to such therapy, preferably to mammals and birds, and more preferably to dogs, cats, humans, ferrets, horses, cattle, sheep, goats and pigs as well as other pets, food animals, work animals or zoo animals. Preferred animals to protect against parasitic nematode disease include dogs, cats, humans and ferrets, with dogs, cats and humans being particularly preferred.

Suitable inhibitors of nematode transglutaminase activity include compounds that interact directly with a nematode transglutaminase protein active site, thereby inhibiting transglutaminase activity, usually by binding to or otherwise interacting with or otherwise modifying the nematode transglutaminase active site. Nematode transglutaminase inhibitors can also interact with other regions of the nematode transglutaminase protein to inhibit transglutaminase activity, for example, by allosteric interaction. Inhibitors of nematode transglutaminase can be relatively small compounds, or they can be quite large, as in the case of anti-parasitic nematode transglutaminase antibodies. Preferably, a nematode transglutaminase inhibitor of the present invention is identified

by its ability to inhibit the activity of a nematode transglutaminase, and further by its failure to substantially inhibit the activity of host animal transglutaminase. Methods for measuring inhibition of transglutaminase activity, useful for determining inhibition of either nematode or host animal transglutaminase activity, are described in the Examples section.

Inhibitors of a nematode transglutaminase can be used directly as compounds in compositions of the present invention to treat host animals, provided that such compounds do not substantially inhibit the activity of the host animal transglutaminase.

Inhibitors of a nematode transglutaminase protein can also be used to identify preferred types of nematode transglutaminase proteins to target using compositions of the present invention, for example by affinity chromatography. For example, an inhibitor of the present invention could be bound to a gel or a filter, or another substrate, and larval or adult nematode extracts could be contacted with the bound inhibitor. Those compounds in either larval or adult nematode extracts that bound to or otherwise interacted with the inhibitor could then be separated from the bound inhibitor and further analyzed for nematode transglutaminase activity.

Preferred inhibitors of a nematode transglutaminase of the present invention include, but are not limited to, nematode transglutaminase substrate analogs and other molecules that bind to a nematode transglutaminase (e.g., to an allosteric site) in such a manner that nematode transglutaminase activity of the nematode transglutaminase is inhibited. A nematode transglutaminase substrate analog refers to a compound that interacts with (e.g., binds to, associates with, modifies) the active site of a nematode transglutaminase protein. A preferred nematode transglutaminase substrate analog inhibits nematode transglutaminase activity. Nematode transglutaminase substrate analogs can be any inorganic or organic composition, and can be, but are not limited to, peptides, nucleic acids, and peptidomimetic compounds. Nematode transglutaminase substrate analogs can be, but need not be, structurally similar to a nematode transglutaminase protein's natural substrate provided they can interact with the active site of that nematode transglutaminase protein. Nematode transglutaminase substrate analogs can be designed using computer-generated structures of nematode transglutaminase proteins of the present invention or computer structures of nematode transglutaminase proteins' natural substrates. Substrate analogs can also be obtained by generating random samples of molecules (for example, oligonucleotides, peptides, peptidomimetic compounds, or other inorganic or organic molecules), and screening such samples by affinity chromatography techniques using the corresponding binding partner, (e.g., a nematode transglutaminase or anti-nematode transglutaminase substrate antibody). A preferred nematode transglutaminase substrate analog is a peptidomimetic compound (i.e., a compound that is structurally or functionally similar to a natural substrate of a nematode transglutaminase of the present invention, particularly to the region of the substrate that interacts with the nematode transglutaminase active site, but that inhibits nematode transglutaminase activity upon interacting with the nematode transglutaminase active site).

Parasitic nematode transglutaminase peptides, mimetopes and substrate analogs, as well as other protective compounds (nucleic acid molecules, proteins, antibodies, for example), can be used directly as compounds in compositions of the present invention to treat animals as long as such compounds are not harmful to the animals being treated. Methods to test the safety of such compounds are disclosed herein.

In accordance with the present invention, a host animal (i.e., an animal that is infected with or is capable of being infected by a parasitic nematode) is treated by administering to the animal a therapeutic composition of the present invention in such a manner that the composition itself (e.g., an inhibitor of a nematode transglutaminase protein, mimetope, a nematode transglutaminase synthesis suppressor (i.e., a compound that decreases the production of nematode transglutaminase in the nematode), a nematode transglutaminase mimetope or an anti-parasitic nematode transglutaminase antibody) or a product generated by the animal in response to administration of the composition (e.g., antibodies produced in response to a parasitic nematode transglutaminase protein or nucleic acid molecule vaccine, or conversion of an inactive inhibitor "prodrug" to an active inhibitor of a nematode transglutaminase protein) contacts the nematode, thereby reducing transglutaminase activity in the nematode. A host animal is preferably treated in such a way that the compound or product thereof enters the bodily fluids (e.g., blood and lymph systems) and/or tissues of the animal. Parasitic nematodes are then exposed to the composition or product when they are present in the host animal. For example, nematode transglutaminase protein inhibitors administered to an animal are, administered in such a way that the inhibitors enter the blood and tissues of the animal where parasitic nematodes will come in contact with the inhibitors. In another embodiment, when a parasitic nematode transglutaminase protein, mimetopes or nucleic acid molecule vaccine is administered to a host animal, the treated animal mounts an immune response resulting in the production of antibodies against the parasitic nematode transglutaminase protein (i.e., anti-parasitic nematode transglutaminase antibodies) that circulate in the animal's blood stream and/or other bodily fluids thereby coming into contact with parasitic nematodes.

In order to protect an animal from disease caused by a parasitic nematode of the present invention, a therapeutic composition of the present invention is administered to the animal in an effective manner such that the composition is capable of protecting that animal from a disease caused by a parasitic nematode. Therapeutic compositions of the present invention can be administered to animals prior to infection in order to prevent infection (i.e., as a preventative vaccine), or can be administered to animals after infection in order to treat disease caused by the parasitic nematode (i.e., as a therapeutic vaccine), or both techniques may be used.

Therapeutic compositions of the present invention preferably are formulated in an excipient that the animal to be treated can tolerate. Examples of such excipients include water, saline, Ringer's solution, dextrose solution, Hank's solution, and other aqueous physiologically balanced salt solutions. Nonaqueous vehicles, such as fixed oils, sesame oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include suspensions containing viscosity enhancing agents, such as sodium carboxymethylcellulose, sorbitol, or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer and Tris buffer, while examples of preservatives include, m- or o-cresol, formalin and benzyl alcohol. Standard formulations can either be liquid injectables or solids that can be taken up in a suitable liquid as a suspension or solution for injection. Thus, in a non-liquid formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

In one embodiment of the present invention, a therapeutic composition can include an adjuvant. Adjuvants are agents that are capable of enhancing the immune response of an animal to a specific antigen. Suitable adjuvants include, but are not limited to, cytokines, chemokines, and compounds that induce the production of cytokines and chemokines (e.g., granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interleukin 12 (IL-12), interferon gamma, interferon gamma inducing factor I (IGIF), transforming growth factor beta, RANTES (regulated upon activation, normal T-cell expressed and presumably secreted), macrophage inflammatory proteins (e.g., MIP-1 alpha and MIP-1 beta), and leishmania elongation initiating factor (LEIF)); bacterial components (e.g., endotoxing, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum proteins, viral coat proteins; block copolymer adjuvants (e.g., Hunter's Titermax™ adjuvant (Vaxcel™, Inc. Norcross, Ga.), Ribi adjuvants (Ribi ImmunoChem Research, Inc., Hamilton, Mont.); and saponins and their derivatives (e.g., Quil A (Superfos Biosector A/S, Denmark). Protein adjuvants of the present invention can be delivered in the form of the protein themselves or of nucleic acid molecules encoding such proteins using the methods described herein. In addition to the foregoing adjuvants, when an isolated nucleic acid molecule of the present invention is used as a protective compound in the therapeutic composition, one or more DNA adjuvants can be operatively linked to that nucleic acid molecule using molecular biology techniques known to those skilled in the art.

In one embodiment of the present invention, a therapeutic composition can include a carrier. Carriers include compounds that increase the half-life of a therapeutic composition in the treated animal. Suitable carriers include, but are not limited to, polymeric controlled release vehicles, biodegradable implants, liposomes, bacteria, viruses, other cells, oils, esters, and glycols.

One embodiment of the present invention is a controlled release formulation that is capable of slowly releasing a composition of the present invention into an animal. As used herein, a controlled release formulation comprises a composition of the present invention in a controlled release vehicle. Suitable Controlled release vehicles include, but are not limited to, biocompatible polymers, other polymeric matrices, capsules, microcapsules, microparticles, gels (including hydrogels), bolus preparations, osmotic pumps, diffusion devices, liposomes, lipospheres, and transdermal delivery systems. Other controlled release formulations of the present invention include liquids that, upon administration to an animal, form a solid or a gel in situ. Preferred controlled release formulations are biodegradable (i.e., bioerodible).

A preferred controlled release formulation of the present invention is capable of releasing a composition of the present invention into the blood of the treated animal at a constant rate sufficient to attain dose levels of the composition effective to protect an animal from disease caused by parasitic nematodes. The therapeutic composition is preferably released over a period of time ranging from about 1 to about 12 months. A controlled release formulation of the present invention is capable of effecting a treatment prefer-

ably for at least about 1 month, more preferably for at least about 3 months, even more preferably for at least about 6 months, even more preferably for at least about 9 months, and even more preferably for at least about 12 months.

Acceptable protocols to administer therapeutic compositions in an effective manner include the specification of individual dose size, number of doses, frequency of dose administration, and mode of administration. Determination of such protocols can be accomplished by those skilled in the art. A suitable single dose is a dose that is capable of protecting an animal from disease when administered one or more times over a suitable time period. For example, a preferred single dose of a protein, mimetope or antibody therapeutic composition is from about 1 microgram (μg) to about 10 milligrams (mg) of the therapeutic composition per kilogram body weight of the animal. Booster vaccinations can be administered from about 2 weeks to several years after the original administration. Booster administrations preferably are administered when the immune response of the animal becomes insufficient to protect the animal from disease. A preferred administration schedule is one in which from about 10 μg to about 1 mg of the therapeutic composition per kg body weight of the animal is administered from about one to about two times over a time period of from about 2 weeks to about 12 months. Modes of administration can include, but are not limited to, subcutaneous, intradermal, intravenous, intranasal, oral, intraocular, transdermal and intramuscular routes.

According to one embodiment, a nucleic acid molecule of the present invention can be administered to a host animal in a fashion enabling expression of that nucleic acid molecule into a protective protein or protective RNA (e.g., antisense RNA, ribozyme, triple helix forms or RNA drug) in the host animal. Nucleic acid molecules can be delivered to an animal using a variety of methods including, but not limited to, (a) administering a naked (i.e., not packaged in a viral coat or cellular membrane) nucleic acid vaccine (e.g., as naked DNA or RNA molecules, such as is taught, for example in Wolff et al., 1990, *Science* 247, 1465-1468) or (b) administering a nucleic acid molecule packaged as a recombinant virus vaccine or as a recombinant cell vaccine (i.e., the nucleic acid molecule is delivered by a viral or cellular vehicle)).

A naked nucleic acid vaccine of the present invention includes a nucleic acid molecule of the present invention and preferably includes a recombinant molecule of the present invention that preferably is replication, or otherwise amplification, competent. A naked nucleic acid vaccine of the present invention can comprise one or more nucleic acid molecules of the present invention in the form of, for example, a bicistronic recombinant molecule, having, for example, one or more internal entry sites. Preferred naked nucleic acid vaccines include at least a portion of a viral genome (i.e., a viral vector). Preferred viral vectors include those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, and retroviruses, with those based on alphaviruses (such as Sindbis or Semliki virus), species-specific herpesviruses and species-specific poxviruses being particularly preferred. Any suitable transcription control sequence can be used, including those disclosed as suitable for protein production. Particularly preferred transcription control sequence include cytomegalovirus intermediate early (preferably in conjunction with intron-A), Rous Sarcoma Virus long terminal repeat, and tissue-specific transcription control sequences, as well as transcription control sequences endogenous to viral vectors if viral vectors are used. The incorporation of "strong" poly(A) sequences are also preferred.

Naked nucleic acid vaccines of the present invention can be administered in a variety of ways, with intramuscular, subcutaneous, intradermal, transdermal, intranasal, intraocular and oral routes of administration being preferred. A preferred single dose of a naked nucleic acid vaccines ranges from about 1 nanogram (ng) to about 100 μg , depending on the route of administration and method of delivery, as can be determined by those skilled in the art. Suitable delivery methods include, for example, by injection, as drops, by aerosolization and by topical application. Naked DNA of the present invention can be contained in an aqueous excipient (e.g., phosphate buffered saline) alone or in a carrier (e.g., lipid-based vehicles).

A recombinant virus vaccine of the present invention includes a recombinant molecule of the present invention that is packaged in a viral coat and that can be expressed in an animal after administration. Preferably, the recombinant molecule is packaging-deficient, encodes an attenuated virus, or both. A number of recombinant viruses can be used including, but not limited to, those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, and retroviruses. Preferred recombinant virus vaccines are those based on alphaviruses (such as Sindbis virus), raccoon poxviruses, species-specific herpesviruses and species-specific poxviruses. An example of methods to produce and use alphavirus recombinant virus vaccines is disclosed in PCT Publication No. WO 94/17813, by Xiong et al., published Aug. 18, 1994, which is incorporated by reference herein in its entirety. An example of methods to produce and use racoon poxvirus recombinant virus vaccines is disclosed in U.S. Pat. No. 5,266,314, to Esposito, et al., issued November 30, 1993, which is incorporated by reference herein in its entirety.

When administered to an animal, a recombinant virus vaccine of the present invention infects cells within the immunized animal and directs the production of a protective protein or RNA nucleic acid molecule that is capable of protecting the animal from disease caused by a parasitic nematode as disclosed herein. For example, a recombinant virus vaccine comprising a parasitic nematode transglutaminase nucleic acid molecule of the present invention is administered according to a protocol that results in the animal producing a sufficient immune response to protect itself from disease caused by a parasitic nematode. A preferred single dose of a recombinant virus vaccine of the present invention is from about 1×10^4 to about 1×10^7 virus plaque forming units (pfu) per kilogram body weight of the animal. Administration protocols are similar to those described herein for protein-based vaccines, with subcutaneous, intramuscular, intranasal and oral administration routes being preferred.

A recombinant cell vaccine of the present invention includes recombinant cells of the present invention that express at least one protein of the present invention. Preferred recombinant cells for this embodiment include *Salmonella*, *E. coli*, *Ligteria*, *Mycobacterium*, *S. frugiperda*, yeast, (including *Saccharomyces cerevisiae*), BHK, CV-1, myoblast G8, COS (e.g., COS-7), Vero, MDCK and CRFK recombinant cells. Recombinant cell vaccines of the present invention can be administered in a variety of ways but have the advantage that they can be administered orally, preferably at doses ranging from about 10^8 to about 10^{12} cells per kilogram body weight. Administration protocols are similar to those described herein for protein-based vaccines. Recombinant cell vaccines can comprise whole cells, cells stripped of cell walls or cell lysates.

The efficacy of a therapeutic composition of the present invention to protect an animal from disease caused by a

parasitic nematode can be tested in a variety of ways including, but not limited to, detection of protective antibodies (using, for example, proteins or mimetopes of the present invention), detection of cellular immunity within the treated animal, or challenge of the treated animal with the parasitic nematode to determine whether the treated animal is resistant to disease. Challenge studies can include implan-

5 tation of chambers including parasitic nematode larvae into the treated animal, or direct administration of larvae to the treated animal, or both. In one embodiment, therapeutic compositions can be tested in animal models such as mice. Such techniques are known to those skilled in the art.

One preferred embodiment of the present invention is the use of parasitic nematode transglutaminase proteins, nucleic acid molecules, antibodies and inhibitory compounds of the present invention, to protect an animal from heartworm. It is particularly preferred to prevent L3 that are delivered to the animal by the mosquito intermediate host from maturing into adult worms. Preferred therapeutic compositions are those that are able to inhibit at least one step in the portion of the parasite's development cycle that includes L3, third molt, L4, fourth molt, immature adult prior to entering the host animal's tissues or circulatory system. In dogs, this portion of the development cycle is about 70 days. Particularly preferred therapeutic compositions include nematode transglutaminase-based therapeutic compositions of the present invention, particularly because *D. immitis* transglutaminase is necessary for *D. immitis* larval molting and development, as disclosed herein. These preferred therapeutic compositions include nematode transglutaminase nucleic acid molecules, nematode transglutaminase proteins and mimetopes thereof, anti-nematode transglutaminase antibodies, and inhibitors of nematode transglutaminase activity that fail to substantially inhibit host animal transglutaminase activity. Particularly preferred are *D. immitis* forms of any of the therapeutic compositions of the present invention. Therapeutic compositions are administered to animals in a manner effective to protect the animals heartworm. Additional protection may be obtained by administering additional protective compounds, including other nematode proteins, nucleic acid molecules, antibodies and inhibitory compounds, as disclosed herein and elsewhere.

One therapeutic composition of the present invention includes an inhibitor of nematode transglutaminase activity that does not substantially inhibit host animal transglutaminase activity. In other words, in one embodiment, a therapeutic composition of the present invention includes a compound capable of substantially interfering with the function of a nematode transglutaminase susceptible to inhibition by an inhibitor of nematode transglutaminase activity. The term, "substantially interfering with the function of nematode transglutaminase," as used herein, refers to the ability of an inhibitor compound to interfere with a nematode transglutaminase activity to such a degree that development of heartworm in a host animal is impaired. For example, an isolated protein or mimetope thereof, is administered in an amount and manner that elicits (i.e., stimulates) an immune response that is sufficient to protect the animal from the disease. Similarly, an antibody of the present invention, when administered to an animal in an effective manner, is administered in an amount so as to be present in the animal at a titer that is sufficient to protect the animal from the disease, at least temporarily. Oligonucleotide nucleic acid molecules of the present invention can also be administered in an effective manner, thereby reducing expression of parasitic nematode transglutaminase proteins in order to interfere with development of parasitic nematodes targeted in accordance with the present invention.

An inhibitor of nematode transglutaminase activity can be identified using parasitic nematode transglutaminase proteins of the present invention. One embodiment of the present invention is a method to identify a compound that is capable of inhibiting nematode transglutaminase activity, but that does not substantially inhibit host animal transglutaminase activity. Such a method includes the steps of (a) contacting (e.g., combining, mixing) an isolated nematode transglutaminase protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has nematode transglutaminase activity; (b) determining if the putative inhibitory compound inhibits the nematode transglutaminase activity; and (c) repeating steps (a) and (b), but substituting host animal transglutaminase for nematode transglutaminase. Putative inhibitory compounds to screen for include small organic molecules, antibodies (including fragments and mimetopes thereof) and substrate analogs. Methods to determine nematode and host animal transglutaminase activities are known to those skilled in the art; see, for example, citations in the background section and references included therein.

The present invention also includes a test kit to identify a compound capable of inhibiting nematode transglutaminase activity of a parasitic nematode. Such a test kit includes an isolated nematode transglutaminase protein having transglutaminase activity and a means for determining the extent of inhibition of transglutaminase activity in the presence of (i.e., effected by) a putative inhibitory compound. Compounds determined to inhibit nematode transglutaminase activity are also screened to identify those that are not substantially toxic to host animals.

Nematode transglutaminase inhibitors isolated by the method or by the test kit described, or by both, can be used to inhibit any nematode transglutaminase that is susceptible to such an inhibitor. Preferred parasitic nematode transglutaminase proteins to inhibit are those produced by *D. immitis*, *B. malayi* or *O. volvulus*. A particularly preferred transglutaminase inhibitor of the present invention is capable of protecting an animal from heartworm. Effective amounts and dosing regimens can be determined using techniques known to those skilled in the art.

Another therapeutic composition of the present invention includes an inhibitor of nematode PDI activity that does not substantially inhibit host animal PDI activity. In other words, in one embodiment, a therapeutic composition of the present invention includes a compound capable of substantially interfering with the function of nematode transglutaminase PDI activity susceptible to inhibition by an inhibitor of nematode PDI activity. The term, "substantially interfering with the function of nematode transglutaminase PDI activity," as used herein, refers to the ability of an inhibitor compound to interfere with a nematode PDI activity to such a degree that development of heartworm in a host animal is impaired.

An inhibitor of nematode PDI activity can be identified using parasitic nematode transglutaminase proteins of the present invention. One embodiment of the present invention is a method to identify a compound that is capable of inhibiting nematode PDI activity, but that does not substantially inhibit host animal PDI activity. Such a method includes the steps of (a) contacting (e.g., combining, mixing) an isolated nematode transglutaminase protein (having PDI activity) with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has nematode PDI activity; (b) determining if the putative inhibitory compound inhibits the nematode PDI activity; and (c) repeating steps (a) and (b), but substituting a host

animal PDI for nematode PDI. Putative inhibitory compounds to screen for include small organic molecules, antibodies (including fragments and mimetopes thereof) and substrate analogs. Methods to determine nematode and host animal PDI activities are known to those skilled in the art; see, for example, citations in the background section and references included therein.

The present invention also includes a test kit to identify a compound capable of inhibiting PDI activity of a parasitic nematode. Such a test kit includes an isolated nematode transglutaminase protein having PDI activity and a means for determining the extent of inhibition of PDI activity in the presence of (i.e., effected by) a putative inhibitory compound. Compounds determined to inhibit nematode PDI activity are also screened to identify those that are not substantially toxic to host animals.

Nematode PDI inhibitors isolated by the method or by the test kit described, or by both, can be used to inhibit any nematode PDI that is susceptible to such an inhibitor. Preferred parasitic transglutaminase proteins to inhibit are those produced by *D. immitis*, *B. malayi* or *O. volvulus*. A particularly preferred PDI inhibitor of the present invention is capable of protecting an animal from heartworm. Effective amounts and dosing regimens can be determined using techniques known to those skilled in the art.

It is also within the scope of the present invention to use isolated proteins, mimetopes, nucleic acid molecules and antibodies of the present invention as diagnostic reagents to detect infection by parasitic nematodes. Such diagnostic reagents can be supplemented with additional compounds that can detect other phases of the parasite's life cycle. Methods to use such diagnostic reagents to diagnose parasitic nematode infection are well known to those skilled in the art. Suitable and preferred parasitic nematodes to detect are those to which therapeutic compositions of the present invention are targeted. A particularly preferred parasitic nematode to detect using diagnostic reagents of the present invention is *D. immitis*.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

EXAMPLES

It is to be noted that these Examples include a number of molecular biology, microbiology, immunology and biochemistry techniques familiar to those skilled in the art. Disclosure of such techniques can be found, for example, in Sambrook et al., *ibid.*, Ausubel, et al., 1993, *Current Protocols in Molecular Biology*, Greene/Wiley Interscience, New York, N.Y., and related references. Ausubel, et al., *ibid.* is incorporated by reference herein in its entirety. DNA and protein sequence analyses were carried out using the PC/GENE™ sequence analysis program (available from Intelligenetics, Inc., Mountainview, Calif.) and the Wisconsin Package™ Version 9.0 (available from the Genetics Computer Group (GCG), Madison, Wis.). It should also be noted that because nucleic acid sequencing technology, and in particular the sequencing of PCR products, is not entirely error-free, that the nucleic acid and deduced protein sequences presented herein represent apparent nucleic acid sequences of the nucleic acid molecules encoding *D. immitis*, *B. malayi*, and *O. volvulus* transglutaminase proteins of the present invention.

Example 1

This example describes a novel N-terminal amino acid sequence of a transglutaminase protein purified from *Brugia malayi*. This example further describes the use of a protein

encoded by that sequence to purify and partially characterize a rare and novel transglutaminase protein from *D. immitis*.

Purification and partial characterization of a novel transglutaminase protein from *B. malayi* has been previously described See, Singh, et al., 1994, *Eur J. Biochem.*, 225, 625-634 (incorporated herein by reference). The amino acid sequence of this protein, referred to as SEQ ID NO:1, is herein disclosed for the first time as follows:

(D)(G)DVMKFTDADFKE(G)IK(X)(Y)(D)

The amino acids in brackets are the most probable amino acids at those positions, and the amino acid (X) at position 18 could not be detected.

A protein molecule corresponding to the N-terminal sequence of the previously described 56-kD transglutaminase of *B. malayi* was synthesized commercially and is herein denoted as PBmTG₂₀. The amino acid sequence of this protein represents amino acids from about position 3 (amino acid residue A, or aspartic acid) through about position 17 (amino acid residue K, or lysine). A cysteine residue was added to the N-terminus of the synthetic peptide (immediately before the aspartic acid residue at about position 3) for the convenience of its conjugation with the carrier protein keyhole limpet hemocyanin (KLH) via maleimidobenzoyl-N-hydroxysuccinimide ester (MBS), as follows. 5.0 mg of KLH in 50 mM phosphate buffer, pH 8.0, was reacted with MBS (dissolved in dimethyl sulfoxide) at a molar ratio of 1 KLH:40 MBS. The solution was stirred for 30 min at room temperature. The unreacted MBS was removed by gel filtration, and 5.0 mg of peptide hapten was added to the MBS-activated KLH in 50 mM phosphate buffer, pH 7.5. The solution was stirred at room temperature for 3 hr. Unconjugated peptide was removed by gel filtration. The conjugation efficiency was 40%.

Anti-*B. malayi* transglutaminase peptide PBmTG₂₀ antiserum was produced as follows. A rabbit was immunized subcutaneously, first with approximately 150 μg of the conjugated peptide mixed with Complete Freund's Adjuvant, and then with five subsequent immunizations of the same dose mixed in Incomplete Freund's Adjuvant. Bleeding and immunization were performed at alternate weeks. Unused antisera were preserved in 0.1% sodium azide at 4° C. For immobilizing the anti-peptide antibodies on Affigel-10 (available from Bio-Rad Laboratories, Hercules, Calif.), the immunoglobulin G (IgG) fraction from this antiserum was collected by 40% ammonium sulfate precipitation. Ammonium ions were removed on a NAP-25 column (Sephadex G-25 available from Pharmacia Biotechnology, Piscataway, N.J.) preequilibrated with 100 mM (3-[N-morpholino]propanesulfonic acid) (MOPS) buffer, pH 7.5 (buffer A) to obtain a desalted IgG fraction.

A crude *D. immitis* extract preparation was prepared as follows. All operations were performed at 4° C. unless otherwise mentioned. Thirty-two frozen adult female worms of *D. immitis* (available from TRA laboratories, Athens, Ga.) were homogenized twice in a Pyrex homogenizer in 20 mM Tris-HCl (pH 8.5) containing 0.1% Triton X-100, 2 mM 1,4-dithiothreitol, 1 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), and 0.1 mM N-tosyl-L-lysine chloromethane. The resulting 40 ml of homogenate was sonicated as described in Singh et al., *ibid.* The extract was frozen and thawed between sonications to maximize the solubilization of membrane-bound enzyme. The extract was centrifuged at 15,000 g for 20 min, and the supernatant (36 ml) was collected for further purification.

Anti-*B. malayi* peptide PmTG₂₀ antiserum produced as described above, was found to react with a 56-kD protein band in a western blot of a *D. immitis* extract. This reactivity of anti-PBmTG₂₀ antiserum could be completely inhibited in the presence of excess synthetic peptide. In order to monitor the progress of the purification process, Western

blots were performed on samples of the extract after each major step in the purification process as follows: Sodium dodecyl sulfate (SDS)-polyacrylamide (10%) gel electrophoresis was performed according to the method of Laemmli (1970). Western blotting was performed by transferring the protein bands to the nitrocellulose paper (0.47 μ M, available from Schleicher & Schuell, Keene, N.H.) using a Semiphor dry blot apparatus (available from Hoefer Scientific Instruments, San Francisco, Calif.). All solutions used for membrane processing were made in phosphate buffered saline (PBS), and incubations were done at room temperature unless otherwise noted. The membrane was blocked with 5% nonfat dry milk for 1 hr. and incubated for 1 hr with 1000-fold diluted anti-PBmTG₂₀ antiserum in 5% nonfat dry milk. After two washes with 100 ml of PBST (PBS containing 0.1% Tween 20) for 20 min each, the membrane was incubated for 1 hr with 5000-fold diluted alkaline phosphatase-linked anti-rabbit IgG (available from Kirkegaard and Perry Laboratories, Gaithersburg, Md.) in 5% nonfat dry milk. After two washes in 100 ml of PBST for 20 min each, the membrane was treated with alkaline phosphatase color development reagent (available from Bio-Rad Laboratories, Hercules, Calif.) as per manufacturer's instructions.

Following crude extract preparation, the first step in *D. immitis* transglutaminase protein purification was thermo-precipitation and ammonium sulfate precipitation as follows. The crude extract from adult female worms was subjected to thermo-precipitation at 55° C. in a water bath for 10 min with constant shaking. The precipitate was discarded by centrifugation at 15,000 g for 20 min, and the supernatant (31 ml) was precipitated at a 60% ammonium sulfate cutoff. The precipitate was collected by centrifugation (15,000 g for 30 min) and was dissolved in 2.5 ml of buffer A. The ammonium ions in the preparation were removed by passing the preparation through an NAP-25 column preequilibrated with buffer A. The final volume of the *D. immitis* preparation obtained from the NAP-25 column was 3.5 ml.

The next step in *D. immitis* transglutaminase protein purification, immunoaffinity chromatography, was accomplished as follows. The immunoglobulin fraction was conjugated to Affigel-10 according to the manufacturer's instructions. Specifically, 3.5 ml of the desalted IgG fraction containing anti-*B. malayi* transglutaminase PBmTG₂₀ (containing 17.5 mg of protein), obtained as described above in this Example, was added to 1 ml of Affigel-10 that was previously washed with cold deionized water. The suspension was incubated and rotated overnight at 4° C. Next day, the unbound IgG was removed by repeated washing with buffer A.

The 3.5 ml *D. immitis* preparation obtained after ammonium sulfate precipitation and desalting was incubated and rotated with the IgG-bound Affigel-10 overnight at 4° C. The slurry was then packed in a column, and the gel was washed extensively with buffer A. Nonspecifically bound proteins were removed by washing the gel with 0.5% Triton X-100 in buffer A to remove the nonspecific hydrophobic interactions. This step was necessary before the specific elution of transglutaminase at pH 2.8. The gel was washed again with buffer A. *D. immitis* transglutaminase was eluted with 3 ml of 100 mM glycine-HCl buffer (pH 2.8) with a flow rate of 10 ml/hr. The pH of the *D. immitis* transglutaminase-containing collected fraction was immediately adjusted to pH 8.0 by adding 300 μ l of 1 M sodium bicarbonate; the collected fraction was then subjected to overnight dialysis against 100 mM Tris-HCl buffer (pH 8.5). The dialyzed fraction was concentrated to 0.5 ml in a Centricon-10 tube (available from Pharmacia Biotechnology Piscataway, N.J.), and used for further characterization.

The eluted protein was enzymatically active and gave a single major band of 56-kD when subjected to electrophoresis

under denaturing conditions. The same 56 kD band was detected by western blot analysis (described below) when the anti-*B. malayi* transglutaminase peptide PBmTG₂₀ antibody was used to detect protein.

A summary of the steps used in the purification of transglutaminase from *D. immitis* is shown in Table 1. The starting transglutaminase activity in 224 mg of initial soluble protein obtained from 32 adult female worms was extremely low. The specific activity in the crude extract obtained from *D. immitis* was at least 5 times lower than that previously reported for *B. malayi* transglutaminase preparation (Singh et al., *ibid.*). Transglutaminase activity was determined in a microtiter plate assay according to a recently published procedure; see, Slaughter et al., 1992, *Anal. Biochem.* 205, 166-171 (incorporated herein by reference). One milliunit (mU) transglutaminase activity is defined as the V_{max} ($\Delta A_{405}/\text{min}$) generated in a microtiter plate assay by 0.74 μ g of purified guinea pig liver transglutaminase (available from Sigma Chemical Co., T-5398). The effects of pH and temperature on the transglutaminase activity and stability as well as the effects of inhibitors, metal ions and other cofactors on the enzyme activity were determined as described by Singh et al., *ibid.* The amount of protein was estimated according to the Bradford method (see Bradford, 1976, *Anal. Biochem.*, 72, 248-254), using reagents available from Bio-Rad Laboratories, Hercules, Calif.

TABLE 1

Summary of steps used for the purification of transglutaminase from *D. immitis* adult worms.

Steps	Total protein (μ g)	Total volume (ml)	Total activity (mU)	Specific activity (mU/mg)	Cumulative fold purification	Yield (%)
1. Crude extract	224,200	36.0	42.6	0.19	1.0	100
2. Thermo-precipitation	129,634	31.0	35.0	0.27	1.4	82
3. (NH ₄) ₂ SO ₄ precipitation	11,157	3.5	26.8	2.33	12.2	63
4. Immunoaffinity chromatography	2.1	0.5	4.2	2032.0	10,694.0	10

The *D. immitis* transglutaminase protein preparation protocol presented herein resulted in a high degree of purification of *D. immitis* transglutaminase protein. The final product was approximately 5 times purer than that previously reported for *B. malayi* transglutaminase purified by the lengthy conventional protocol of Singh et al., *ibid.* The specific activity of the purified *D. immitis* enzyme was 2.0 U/mg protein, and is very close to that previously reported for *B. malayi* transglutaminase. Although the enzyme was stable over a wide pH range (data not shown), it was most active in the basic pH range, between pH 8 and pH 9.5, as are the other known transglutaminases. In contrast to mammalian transglutaminases, the *D. immitis* enzyme, like the transglutaminase isolated from *B. malayi* (see, Singh, et al., *ibid.*) was active and stable at high temperatures (data not shown).

The effects of various reagents on the activity of the transglutaminase purified from adult *D. immitis* worms are shown in Table 2. The enzyme required calcium for its activity, and chelating agents like EGTA and EDTA completely blocked the activity. Dithiothreitol and mercaptoethanol increased the enzyme activity substantially, whereas iodoacetamide decreased the activity drastically, suggesting that the enzyme requires at least one cysteine residue at the

active site, like most of the transglutaminases; see, for example, Folk et al., 1977, *Adv. Protein Chem.* 31, 1–133. The effect of iodoacetamide was severe when the enzyme was pretreated with calcium ions, suggesting that calcium ions open the active site for high molecular weight substrates like casein. The enzyme was inhibited competitively by amine donor substrate analogues like monodansyl cadaverine and putrescine, and by the active-site inhibitor cystamine. High concentrations of sodium and potassium ions, Tris and the end product of the reaction, ammonia, reversibly inhibited the enzyme. The observation that Cbz-Gln-Gly affects the enzyme activity only slightly (Table 2) suggests that this compound is a poor amine acceptor substrate for the enzyme. In contrast to mammalian tissue type transglutaminase (Folk et al., *ibid.*; Bergamini et al., 1987, *Biochim. Biophys. Acta* 916, 149–151; Achyuthan et al., 1987, *J. Biol. Chem.* 262, 1901–1906; Bergamini, 1988, *FEBS Lett.* 239, 255–258; Lee et al., 1989, *Biochem. Biophys. Res. Commun.* 162, 1370–1375), this enzyme was not affected adversely by micromolar concentrations of GTP. This suggests that GTP is not involved in the regulation of this enzyme as in nematode, transglutaminase from *B. malayi* (Singh et al., *ibid.*) and *Limulus* hemocyte transglutaminase (Tokunaga et al., 1993, *J. Biol. Chem.* 268, 252–261).

TABLE 2

Effect of ions, inhibitors and other reagents on <i>D. immitis</i> transglutaminase activity		
Reagent*	Concentration (mM)	Transglutaminase activity† (% of control)
Control‡	—	100
NaCl	500	48
KCl	500	54
(NH ₄) ₂ SO ₄	10	0
EDTA	10	0
EGTA	10	0
Iodoacetamide (–Ca ²⁺)§	10	61
Iodoacetamide (+Ca ²⁺)¶	10	27
Tris-HCl (pH 8.5)	250	66
Tris-HCl (pH 8.5)	500	40
N α -CBZ-Gln-Gly	10	92
Monodansyl cadaverine	1	9
Putrescine	1	14
Cystamine	1	52
GTP	0.1	100
GTP	1	95

*The effect of metals, ions and other reagents on transglutaminase activity was determined in the presence of CaCl₂ and dithiothreitol.

†The results shown are the average values from two independent experiments each performed in triplicate. Standard deviation from the mean was less than 5%.

‡Control tubes contained 10 mM CaCl₂ and 10 mM dithiothreitol.

§Iodoacetamide was preincubated with the enzyme in the absence of calcium overnight at 4° C., and the activity was determined in the presence of 10 mM each of calcium and dithiothreitol after removal of iodoacetamide by dialysis.

¶Iodoacetamide was preincubated with the enzyme in the presence of 10 mM calcium overnight at 4° C., and the activity was determined in the presence of 10 mM each of calcium and dithiothreitol after removal of iodoacetamide by dialysis.

Example 2

This Example evaluated the effect of a number of transglutaminase inhibitors on *D. immitis* larval viability in an in vitro larval culture system.

The following transglutaminase inhibitors were tested at the indicated final concentrations in the culture system:

- (a) Monodansyl cadaverine (MDC), a known high affinity substrate analog, was tested at concentrations of 25, 50, 75, 85 and 100 μ M;

- (b) Cystamine, a transglutaminase active site inhibitor, was tested at concentrations of 25, 50, 75, 85 and 100 μ M;

- (c) Iodoacetamide was tested at concentrations of 2.5, 5 and 10 μ M.

All inhibitors are available from Sigma Chemical Co, St. Louis, Mo. Inhibitors were made in NI media (50% NCTC+50% IMDM, available from GibcoBRL, Gaithersburg, Md.) containing antibiotics and 20% SeruMax (available from Sigma Chemical Co.).

The general protocol for the larval viability assays was as follows: Briefly, 50 *D. immitis* L₃ larva were cultured for 6 days in vitro in 1 ml of NI media containing antibiotics and 20% SeruMax. In some assays, transglutaminase inhibitors were added on different days of culture, and in other assays the inhibitors were present for only 24 hours of culture. The cultures were examined microscopically every 24 hours until day 6 when the cultures were terminated. The number of larvae that molted were determined by counting shed cuticles.

Results of these studies are presented below in Tables 3, 4, and 5. All transglutaminase inhibitors tested in the present study reduced in a dose-dependent manner the molting of *D. immitis* L₃ larvae to L₄ larvae (Table 3).

TABLE 3

Effect of TGase inhibitors on molting of <i>D. immitis</i> L ₃		
Inhibitor	Concentration (μ M)	Percent molted
Monodansylcadaverine (MDC)	0	84
	25	75
	50	63
	75	28
	85	2.5
Cystamine	100	0
	0	84
	25	61
	50	62
	75	17
Iodoacetamide	85	1
	100	0
	0	84
	2.5	65
	5	3
	10	0

MDC and cystamine at 100 μ M concentration completely inhibited the molting process; Iodoacetamide at a final concentration of 10 μ M was able to inhibit the molting of L₃ to L₄. In each case, complete inhibition of *D. immitis* molting required the presence of inhibitors during the first 24–48 hr of the molting process (Tables 4 and 5). The transglutaminase active site inhibitor (cystamine) was a very effective inhibitor of larval molting even when added on day 2 during the culture (Table 5).

TABLE 4

Presence of TGase inhibitors during first 24 hr of <i>D. immitis</i> L ₃ culture - Effect on molting		
Inhibitor	Concentration (μ M)	Percent molted
Monodansylcadaverine (MDC)	0	68
	50	64
	100	41
Cystamine	0	68
	50	68
	100	13
Iodoacetamide	0	68
	10	3

TABLE 5

Molting of <i>D. immitis</i> L ₃ in presence of TGase inhibitors added to culture on different days.		
Inhibitor	Day added	Percent molted
Monodansylcadaverine* (MDC)	0	0
	1	18
	2	78
Cystamine*	0	0
	1	0
	2	10
Iodoacetamide†	0	0
	1	0
	2	82
None		70

*MDC and Cystamine were used at a concentration of 100 μ M

†Iodoacetamide was used at a 10 μ M concentration

Example 3

This Example demonstrates that soluble adult and larval *D. immitis* parasite extracts contain transglutaminase activity.

Larval and adult male and female heartworm parasites were separately homogenized in buffer B (20 mM Tris/HCl pH 8.5, containing 2 mM 1,4-dithiothreitol, 2 mM EDTA, 1 mM phenylmethylsulfonyl fluoride, 0.1 mM N-tosyl-L-lysine chloromethane and 0.1 mM N-tosyl-L-phenylalanine chloromethane; all available from Sigma) for 20 min on ice. The crude extracts thus obtained were sonicated continuously for three 1-min periods, with 5-min intervals between each sonication, using a pre-chilled small probe of the W-380 Ultrasonic Processor (available from Heat Systems-Ultrasonics, Farmingdale, N.Y.). The third sonication was done in the presence of 0.1% Triton X-100. The suspensions were centrifuged at 15,000 \times g for 20 min. The supernatant thus obtained (referred to herein as the parasite extracts, or crude parasite extracts) was used to assay for transglutaminase activity.

Transglutaminase activity was determined in a microtiter plate assay as described above in Example 1. In brief, the microtiter plates were coated with 1% dimethylcasein (available from Sigma) at room temperature overnight; uncoated sites were blocked with 1% nonfat dry milk. The reaction mixtures contained in total volumes of 200 μ l each: 100 mM Tris/HCl pH 8.5, 10 mM CaCl₂, 10 mM dithiothreitol, 1 mM amine donor substrate 5(biotinamido) pentylamine (BPT), (available from Sigma), and crude parasite extracts. The reactions were performed at 37° C. for 2 hours and transglutaminase-catalyzed conjugation of BPT into dimethylcasein was determined by streptavidin-peroxidase and orthophenyldiamine as a reporter system. The enzyme activity (expressed as mU) in extracts was determined relative to the activity of purified guinea pig liver transglutaminase (available from Sigma) tested in the same microtiter plate. The results of this assay are given in Table 6. There was detectable transglutaminase activity both in larval and adult extracts. The activity in males was lower than in females for the same amount of protein tested.

TABLE 6

Transglutaminase enzyme activity in <i>D. immitis</i> larvae and adults		
Parasite stage	Amount used	Total activity (mU)
0 hr L ₃	100 L ₃	38.9
48 hr L ₃	100 L ₃	42.3
6 day L ₄	100 L ₄	27.6
Male adult	60 μ g	9.0
Female adult	60 μ g	50.0

Example 4

This Example describes the identification of native *D. immitis* transglutaminase (DiTG) by immunoblot analysis. Rabbit anti-*B. malayi* transglutaminase peptide PBmTG₂₀ antisera, produced as described in Example 1, was used to identify a native *D. immitis* transglutaminase protein in *D. immitis* extracts as follows.

The materials in crude extracts from *D. immitis* larvae and adult male and female worms were separated by running 5 μ g protein per lane on a 12-well 10% Tris-glycine SDS-PAGE gel at 200 volts for 1 hour, and then transferred to a nitrocellulose membrane by standard methods. After transfer, the membrane was blocked in 5% dry milk for 1 hr at 37° C. The membrane was then incubated with rabbit anti-*B. malayi* transglutaminase peptide PBmTG₂₀ antibody at a dilution of 1:2500 in Tris buffered saline. After 1 hr incubation at room temperature, the blot was washed, and antibody binding resolved using a peroxidase-labeled rabbit IgG secondary antibody and the substrate nitroblue tetrazolium chloride, 5-bromo-4-chloro-3-indolylphosphate p-toluidine salt (NBT/BCIP) (available from Gibco BRL, Gaithersburg, Md). Using this antibody, immunoblot analysis of *D. immitis* adult male, female and larval extracts identified a 56 kD native *D. immitis* protein (DiTG) similar to the size of native Brugia protein (Singh et al., *ibid.*).

Example 5

This Example describes the amino acid sequence analysis of the 56 kD *D. immitis* transglutaminase.

The native 56 kD *D. immitis* transglutaminase protein from adult female *D. immitis* parasite extracts was separated by two dimensional SDS-PAGE. The first dimension was an isoelectric focusing gel using a non-equilibrium pH gradient containing ampholines of pI 5-8 (available from Pharmacia Biotech, Uppsala, Sweden). The second dimension was run on an 8% Tris-glycine gel; the resulting protein spots were transferred to PVDF membrane, and the spot corresponding to *D. immitis* transglutaminase was excised. 17 such spots were then used for N-terminal sequence analysis using an automated protein sequencer (ABI437A, available from Applied Biosystems, Inc., Foster City, Calif.).

For internal amino acid sequence analysis, spots containing *D. immitis* transglutaminase were excised from Coomassie blue stained preparative two dimensional SDS-PAGE gels of female *D. immitis* parasite extract. 48 such spots were pooled and then subjected to trypsin digestion in the gel. The digested protein sample was then separated using high pressure liquid chromatography (HPLC). Digested proteins were then sequenced as described above. Preparation and sequencing of the internal protein fragments were performed by the Harvard Microchemistry Facility, Cambridge, Mass.

The results of the amino acid sequence analysis of *D. immitis* transglutaminase are given below. A partial

N-terminal amino acid sequence of about 29 amino acids was determined and is represented herein as SEQ ID NO:2:
DGDVMKFTDADFKEGKPYDVLVVKFYAP

A homology search of a non-redundant protein sequence database was performed on this amino acid sequence through the National Center for Biotechnology Information (NCBI) (National Library of Medicine, National Institute of Health, Baltimore, Md.) using the BLAST network. This database includes SwissProt+PIR+SPupdate+GenPept+GPUupdate+PDB databases. The search was performed using SEQ ID NO:2 and showed significant homology to probable protein disulfide isomerases (PDIs) spanning from amino acid residue 1 through 29 of SEQ ID NO:2. The highest scoring match of the homology search at the amino acid level was GenBank™ accession number Z37139, *Caenorhabditis elegans* clone C14B1.1. SEQ ID NO:2 showed about 44% identity to residues 24 to 50 of the clone C14B1.1. SEQ ID NO:2 also showed a near sequence identity to the *B. malayi* peptide, PBmTG₂₀, SEQ ID NO:1.

The two internal *D. immitis* transglutaminase amino acid sequences obtained as described above were characterized as follows: A partial internal amino acids sequence of about 14 amino acids was determined and is represented herein as SEQ ID NO:3:

YQYDLLPMFVVYGK

A homology search of a non-redundant protein sequence database was performed on SEQ ID NO:3 through the NCBI using the BLAST network as described above. This database includes SwissProt+PIR+SPupdate+GenPept+GPUupdates+PDB. Results of the search showed no significant homology of SEQ ID NO:3 to other proteins in the database.

Another partial internal amino acid sequence of about 19 amino acids was determined and is represented herein as SEQ ID NO:4:

MDATANDVPPPFVQVGFPT

A homology search of a non-redundant protein sequence database was performed on this amino acid sequence using the BLAST network through the NCBI, as described above. The search was performed using SEQ ID NO:4 and showed significant homology to probable PDIs spanning from amino acid residue 1 through 19 of SEQ ID NO:4. The highest scoring match of the homology search at the amino acid level was GenBank™ accession number PC1298 (chicken nuclear matrix 57 K protein). SEQ ID NO:4 showed about 78% identity to residues 42 to 60 of this chicken nuclear matrix protein sequence.

Example 6

This Example describes the isolation and sequencing of a nucleic acid molecule encoding a *D. immitis* transglutaminase protein.

A *D. immitis* transglutaminase nucleic acid molecule of about 707 nucleotides, denoted nDiTG₇₀₇, was identified by polymerase chain reaction (PCR) amplification from *D. immitis* first strand cDNA reverse transcribed from adult female mRNA as follows. The following primers were used to PCR amplify the *D. immitis* transglutaminase nucleic acid molecule from the cDNA template: A sense primer spanning nucleotides encoding amino acid residue number about 5 through amino acid residue number about 15 of SEQ ID NO:2, and having the nucleic acid sequence 5' ATGAARTYACNGAYGCGAYTTTAAARGARGG 3' (denoted herein as SEQ ID NO:15); and an anti-sense primer spanning nucleotides encoding amino acid residue number about 8 through amino acid residue number about 14 of SEQ ID NO:3 and having the nucleic acid sequence 5' TTNCCR-TANACNACRAACAT 3' (denoted herein as SEQ ID NO:16).

The PCR amplified *D. immitis* transglutaminase nucleic acid molecule, referred to herein as nDiTG₇₀₇, was separated from the rest of the PCR reaction products on a 1% agarose gel at 60 v for 2 hr. After separation of the PCR products, the band of interest was excised from the agarose gel. The gel slice was then processed to release the DNA using the QIAquick kit (available from Qiagen, Chatsworth, Calif.) as per manufacturer's instructions. The purified DNA was then cloned into TA cloning vector (available from Invitrogen, San Diego, Calif.) as per the manufacturer's instructions and submitted for automated sequence analysis. The sequences of the two complementary strands of nDiTG₇₀₇ are presented as SEQ ID NO:5 and SEQ ID NO:7.

Translation of SEQ ID NO:5 yields a protein of about 235 amino acids, denoted PDiTG₂₃₅, the amino acid sequence of which is presented in SEQ ID NO:6. The nucleic acid molecule encoding PDiTG₂₃₅ is referred to herein as nDiTG₇₀₅, the nucleic acid sequence of which is represented in SEQ ID NO:8 (the coding strand) and SEQ ID NO:9 (the complementary strand). Based on its amino acid sequence, PDiTG₂₃₅ has a predicted molecular weight of about 27.2 kD and an estimated pI of about 5.07.

Amino acid sequence of PDiTG₂₃₅ (i.e. SEQ ID NO:6) was analyzed using the PC/GENE™ (available from Intelligenetics, Inc., Mountainview, Calif.) sequence analysis program for sites and signatures. A thioredoxin family active site was detected from residues about 24 to 30. Thioredoxins participate in various redox reactions through the reversible oxidation of an active center disulfide bond; see, for example, Holmgren, *ibid.* A number of eukaryotic proteins contain similar domains evolutionarily related to thioredoxin.

A homology search of a non-redundant protein sequence database was performed through the NCBI using the BLAST network, as described above. The search performed using SEQ ID NO:6 showed that this sequence has significant homology to protein disulfide isomerases (PDI), and PDI-related proteins, of eukaryotic origin. The homology spans from about amino acid 1 through about amino acid 235 of SEQ ID NO:6. The highest scoring match of the homology search at the amino acid level was GenBank™ accession number P38658, *Schistosoma mansoni*, probable PDI ER-60 precursor. SEQ ID NO:6 showed about 37% identity to P38658. At the nucleotide level, the coding regions represented in SEQ ID NO:8, from nucleotide 7 to 246, were similar to that of the human clone PA3 (GenBank™ accession number J05016), PDI-related protein (Erp72) mRNA. SEQ ID NO:8 showed about 59% nucleic acid identity spanning from nucleotide 589 to 828 of clone PA3.

Example 7

The following experiment was performed in order to confirm the *D. immitis* origin of the isolated DiTG cDNA nucleic acid molecule nDiTG₇₀₇, and in order to identify he genomic restriction fragments corresponding to nDiTG₇₀₇. A Southern blot containing about 10 μg of EcoRI and XhoI restricted *D. immitis* genomic DNA was hybridized under stringent conditions with nDiTG₇₀₇ DNA labeled with a chemiluminescent label (ECL labeling kit, available from Amersham, Arlington Heights, Ill.). The probe detected a single band of about 11.7 kilobase pairs (kb) in the genomic DNA digested with XhoI, where as in EcoRI digested genomic DNA, the probe detected three bands at about 9.5, 1.07 and 0.43 kb, respectively.

Example 8

This Example describes the isolation and characterization of transglutaminase nucleic acid molecules of the present invention from a *D. immitis* L₄ cDNA library.

D. immitis transglutaminase nucleic acid molecules were cloned from a cDNA library by nucleic acid screening using *D. immitis* transglutaminase nucleic acid molecules of the present invention as probes. Specifically, a 6 day *D. immitis* larval cDNA library was constructed from 140,420 L₄ as follows. *D. immitis* 6 day, L₄ larvae were cultured in NCTC 135:IMDM media and 20% Seru-MaX™ for 48 hours. Larvae were settled by gravity at 37° C. culture media were removed and larvae were disrupted in 4 M guanidinium thiocyanate, 1.5% sarkosyl, 0.5 M 2-mercaptoethanol. Total RNA (290 µg) was recovered by the acid guanidinium-thiocyanate-phenol-chloroform procedure (Chomczynski, et al., 1987, *Anal. Biochem.* 162, pp. 156–159). Poly A⁺ mRNA (6.954 µg) was isolated with oligo(dT) cellulose using the RiboSep Mini mRNA Isolation Kit (available from Collaborative Research, Inc., Bedford, Mass.). The ZAP-cDNA@ Synthesis Kit (available from Stratagene, La Jolla, Calif.) was used to synthesize cDNA, which was then ligated into the Uni-ZAP XR vector (Stratagene), packaged and amplified to produce the L₄ CDNA library. The nucleic acid molecule, nDiTG₇₀₇, (represented herein by the sequences SEQ ID NO:5 and SEQ ID NO:7) was labeled with a chemiluminescent label as described in Example 7, and used as a DNA probe to screen the L₄ cDNA expression library. A clone containing a *D. immitis* transglutaminase nucleic acid molecule referred to herein as nDiTG₁₄₇₂ was plaque-purified from the expression library using standard methods, and then sequenced. The following nucleotide primers were used to sequence this clone: a) two pBluescript™ vector primers consisting of a sense T₃X primer (denoted herein as SEQ ID NO:17) having the nucleic acid sequence 5' AAT-TAACCTCACTAAAGGG 3'; and an antisense T₇X primer (denoted herein as SEQ ID NO:18) having the nucleotide sequence, 5' GTAATACGACTCACTATAGGGC 3'; and b) three internal primers including a sense primer having the nucleic acid sequence 5' GAAAACCGTTAT-CAGTATGATCT 3' (denoted herein as SEQ ID NO:19), and two antisense primers having the nucleic acid sequences 5' CTGTGGAATGATTTAAATATTTATCC 3' (denoted herein as SEQ ID NO:20) and 5' GTCCATTTTTGCAATAACAA-CACC 3' (denoted herein as SEQ ID NO:21), respectively. The resulting nucleic acid sequences of the two complementary DNA strands of nDiTG₁₄₇₂ are referred to herein as SEQ ID NO:10 and SEQ ID NO:12. The sense primer represented by SEQ ID NO:19 spans nucleotides about nucleotide 359 to about nucleotide 381 of SEQ ID NO:10; the antisense primer represented by SEQ ID NO:20 spans about nucleotide 1171 to about nucleotide 1192 of SEQ ID NO:10; and the antisense primer SEQ ID NO:21 spans from about nucleotide 878 to about nucleotide 901 of SEQ ID NO:10.

Translation of SEQ ID NO:10 yields a protein of about 368 amino acids, denoted as PDiTG₃₆₈, the amino acid sequence of which is presented in SEQ ID NO:11. The nucleic acid molecule encoding PDiTG₃₆₈ is referred to herein as nDiTG₁₁₀₇, the nucleic acid sequence of which is represented in SEQ ID NO:13 (the coding strand) and SEQ ID NO:14 (the complementary strand) assuming that the first codon spans from about nucleotide 2 through about nucleotide 5, and a putative stop codon spans from about nucleotide 1106 to about nucleotide 1108 (the stop codon included in nDiTG₁₁₀₇). The amino acid sequence of *D. immitis* PDiTG₃₆₈ (i.e., SEQ ID NO:11) predicts that PDiTG₃₆₈ has an estimated molecular weight of about 42.6 kD and an estimated pI of 5.71.

The amino acid sequence of PDiTG₃₆₈ (i.e., SEQ ID NO:11) was analyzed using the PC/GENE™ program to

identify sites and signatures. A number of interesting sites were detected. They include: i) a thioredoxin family active site detected from residues 268 to 274; ii) an endoplasmic reticulum (ER) targeting sequence from residues 365 to 368 (KEEL); proteins that permanently reside in the lumen of ER seem to be distinguished from newly synthesized secretory proteins by the presence of the C-terminal sequence Lys-Asp-Glu-Leu (KDEL), see Munro et al., *ibid. Cell* 48, 899–907; Pelham, *ibid.*; and iii) a tachykinin family signature from residues 186 to 202 (tachykinins are a group of biologically active peptides that excite neurons, evoke behavioral responses, are potent vasodilators, and contract many smooth muscles; see, Maggio, 1988, *Annual Review of Neurosciences* 11, 13–28).

A homology search of a non-redundant protein sequence database was performed on SEQ ID NO:11 using the BLAST network through the NCBI, as described above. The search showed significant homology to PDI, and PDI-related proteins of eukaryotic origins, spanning from about amino acid 1 through about amino acid 368 of SEQ ID NO:11. The highest scoring match of the homology search at the amino acid level was to GenBank™ accession number D16234 (from amino acid residues 130 to 505), a human phospholipase C- α clone. This match revealed about 47% identity spanning amino acid residues about 3 to about 368 of SEQ ID NO:11. The nucleic acid coding region represented in SEQ ID NO:13, from about nucleotide 717 to about nucleotide 1032, was similar to that of human epithelial cell mRNA for ER-60 protease (GenBank™ accession number D83485), being about 63% identical to nucleotides 1143 through 1458 of the ER-60 protease sequence.

Example 9

This Example describes the identification of *D. immitis* poly(A)⁺ RNA transcripts corresponding to nDiTG₇₀₇.

A Northern blot was performed as follows: *D. immitis* adult female and male total RNA (8 µg) and adult female and male poly(A)⁺ RNA (0.5 µg) were electrophoresed on a 1% formaldehyde gel and transferred to a N+ nylon membrane (available from Amersham). The RNA was cross-linked to the membrane using the Stratalinker (available from Stratagene). The Northern blot was then hybridized with peroxidase-labeled nDiTG₇₀₇ cDNA using the ECL direct nucleic acid labeling and detection system (available from Amersham) as per the manufacturer's instructions. In each of the four samples, the nDiTG₇₀₇ cDNA probe hybridized to a single band of approximately 2613 nucleotides as calculated by MacVector's mobility program.

Example 10

This Example describes the PCR amplification and subsequent isolation of a 5 nematode transglutaminase nucleic acid molecule from *D. immitis* cDNA using the nematode 22 nucleotide splice leader sequence as the primer.

Nematode transglutaminase nucleic acid molecules were PCR amplified from *D. immitis* female adult cDNA using a primer corresponding to the sequence of a nematode splice leader (SL). Most, but not all nematode messenger RNAs have the SL sequence at their 5' ends, and the presence of the 5' SL sequence is indicative of an apparent full length cDNA. See, for example Blaxter and Liu, 1996, *Int. J. Parasitol.* 26, 1025–1033, which is incorporated herein by reference. The two primers used in the PCR amplification reaction were a sense primer representing the SL sequence, having the nucleotide sequence 5' GGTTAATTAC-CCAAGTTTGAG 3' (herein denoted as SEQ ID NO:22)

and an antisense primer having the sequence 5' TCCCTC-CTTGAAGTCCGCATCTGTAAATTTTCAT 3' (herein denoted as SEQ ID NO:23; SEQ ID NO:23 represents nucleotides from about, nucleotide 673 to about nucleotide 705 of SEQ ID NO:9) PCR amplification of adult female cDNA using these primers resulted in the production of an 143 bp nucleic acid molecule (herein denoted as nDiTG₁₄₃).

Nucleic acid molecule nDiTG₁₄₃ was gel purified, cloned into TA cloning vector (available from Invitrogen, Carlsbad, Calif.) and sequenced using an automated DNA sequencer. Sequence analysis of the nDiTG₁₄₃ coding strand (herein denoted as SEQ ID NO:27) and the complementary strand (herein denoted as SEQ ID NO:29) demonstrated that nDiTG₁₄₃ had the SL sequence at its 5' end. Translation of SEQ ID NO:27 yields a protein of about 40 amino acids, herein denoted as PDiTG₄₀, the amino acid sequence of which is presented in SEQ ID NO:28. The nucleic acid molecule encoding PDiTG₄₀ is referred to herein as nDiTG₁₂₀, the nucleic acid sequence of which is represented in SEQ ID NO:30 (the coding strand) and SEQ ID NO:31 (the complementary strand). The amino acid sequence of *D. immitis* PDiTG₄₀ (i.e., SEQ ID NO:28) predicts that PDiTG₄₀ has an estimated molecular weight of about 4.5 kD and an estimated pI of about 4.6. PC/GENE™ sequence analysis of SEQ ID NO:27 predicts a translation product having an N-terminal hydrophobic signal sequence spanning from about amino acid residue 1 through about amino acid residue 25 of SEQ ID NO:28, and having a predicted cleavage site between about amino acid residue 25 and about amino acid residue 26. The nucleic acid sequence of the predicted mature protein product of PDiTG₄₀ (after cleavage at the predicted cleavage site) is an approximately 15 amino acid protein herein denoted as PDiTG₁₅ (the amino acid sequence of which is represented by SEQ ID NO:52), encoded by a nucleic acid molecule spanning from about nucleotide 99 to about nucleotide 143 of SEQ ID NO:27 (the coding and complementary sequences of which are herein designated as SEQ ID NO:51, and SEQ ID NO:53, respectively).

Example 11

This Example describes the amplification and subsequent isolation of a nematode transglutaminase nucleic acid molecule of the present invention from *D. immitis* female adult cDNA using primers designed for protein expression in the pTrcHisB vector (available from Invitrogen).

Nematode transglutaminase nucleic acid molecules were PCR amplified from *D. immitis* female adult cDNA using the following two primers in the PCR amplification reaction: a sense primer (DiTG-XhoI) with the sequence, 5' CCGAGCTCGAGAATGAAATTTACAGATGCGGAC 3' (herein denoted as SEQ ID NO:24, XhoI site in bold; nucleic acid residues 13 through 33 of this primer represent sequence from about position 1 to about position 21 of SEQ ID NO:5, while the remainder of the primer was designed to include the restriction endonuclease cleavage site) and an antisense primer (DiTG-HindIII) 5' CAGCCAAGCTTCT-TACAATCTTCTTCTTCTTCGGTTTTC 3' (herein denoted as SEQ ID NO:25; HindIII site in bold) for PCR amplification. PCR amplification of *D. immitis* adult female cDNA using these primers resulted in the production of a 1407 bp nucleic acid molecule (herein denoted as nDiTG₁₄₀₇).

The nucleic acid molecule designated nDiTG₁₄₀₇, was gel purified, cloned into a TA cloning vector (available from Invitrogen) and sequenced using an automated DNA

sequencer. The nucleic acid sequence of the coding strand of nDiTG₁₄₀₇ is herein denoted as SEQ ID NO:32, and the complementary strand is herein denoted as SEQ ID NO:34. Translation of SEQ ID NO:32 yields a protein of about 468 amino acids, herein denoted as PDiTG₄₆₈, the amino acid sequence of which is presented in SEQ ID NO:33. The amino acid sequence of *D. immitis* PDiTG₄₆₈ (i.e., SEQ ID NO:33) predicts that PDiTG₄₆₈ has an estimated molecular weight of about 54.3 kD and an estimated pI of about 5.6.

The sequence of nDiTG₁₄₀₇ overlaps with that of nDiTG₁₄₃ and nDiTG₁₄₇₂, allowing for the construction of a composite transglutaminase sequence representing a full-length nematode transglutaminase gene. The nucleic acid molecule represented by this sequence (herein denoted as nDiTG₁₈₈₁) includes both the nematode splice leader sequence at the 5' end of the molecule, and the poly(A)⁺ sequence at the 3' end of the molecule. The nucleic acid sequence of the coding and complementary strands of nDiTG₁₈₈₁ are herein represented by SEQ ID NO:46 and SEQ ID NO:48, respectively. Translation of nDiTG₁₈₈₁ yields an approximately 497 amino acid protein herein denoted as PDiTG₄₉₇, the amino acid sequence of which is presented in SEQ ID NO:47. The nucleic acid molecule encoding PDiTG₄₉₇ is referred to herein as nDiTG₁₄₉₄, the nucleic acid sequence of which is represented in SEQ ID NO:49 (the coding strand) and SEQ ID NO:50 (the complementary strand).

Sequence analysis of SEQ ID NO:27 (the origin of the sequence of the 5' end of nDiTG₁₈₈₁, i.e., SEQ ID NO:46) predicts a translation product having an N-terminal hydrophobic signal sequence spanning from about amino acid residue 1 through about amino acid residue 25 of SEQ ID NO:46, and having a predicted cleavage site between about amino acid residue 25 and about amino acid residue 26. The nucleic acid sequence of the predicted mature protein product of nDiTG₁₈₈₁ (after cleavage at the predicted cleavage site) is an approximately 472 amino acid protein herein denoted as PDiTG₄₇₂ (the amino acid sequence of which is represented by SEQ ID NO:55), encoded by a nucleic acid molecule spanning from about nucleotide 99 to about nucleotide 1514 of SEQ ID NO:46 (the coding and complementary sequences of which are herein designated as SEQ ID NO:54, and SEQ ID NO:56, respectively).

Example 12

This Example discloses the production of a recombinant molecule and a recombinant cell of the present invention.

Recombinant molecule pTrc-nDiTG₁₄₀₇, containing a *D. immitis* transglutaminase nucleic acid molecule represented by nucleotides from about 1 through about 1407 of SEQ ID NO:32 operatively linked to trc transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidine residues, was produced in the following manner. An about 1407 base nucleic acid molecule including nucleotides spanning from about nucleotides 1 through 1407 of SEQ ID NO:32 was PCR amplified from nDiTG₁₄₀₇ using the primers DiTG-XhoI sense primer 5' CCGAGCTCGAGAATGAAATTTACAGATGCGGAC 3' (herein denoted as SEQ ID NO:24; XhoI site in bold) and DiTG-HindIII antisense primer 5' CAGCCAAGCTTCTTACAATCTTCTTCTTCTTCGGTTTTC 3' (herein denoted as SEQ ID NO:25; KpnI site in bold). Recombinant molecule PHis-DiTG₁₄₀₇ was produced by digesting the nDiTG₁₄₀₇-containing PCR product with XhoI and HindIII restriction endonucleases, gel purifying the resulting fragment and directionally subclon-

ing it into the expression vector pTrcHisB (available from Invitrogen) that had been cleaved with XhoI and HindIII and gel purified.

Recombinant molecule pTrc-nDiTG₁₄₀₇ was transformed into *E. coli* to form recombinant cell *E. coli*:pTrc-nDiTG₁₄₀₇, using standard techniques as disclosed, for example, in Sambrook et al., *ibid*.

Example 13

This Example describes the production of a nematode transglutaminase protein of the present invention in a prokaryotic cell as well as studies to characterize that protein.

Recombinant cell *E. coli*:pTrc-nDiTG₁₄₀₇ was cultured in shake-flasks containing an enriched bacterial growth medium containing 0.1 mg/ml ampicillin at about 37° C. When the cells reached an OD₆₀₀ of about 0.5, expression of *D. immitis* pTrc-nDiTG₁₄₀₇ was induced by addition of about 0.5 mM isopropyl-β-D-thiogalactoside (IPTG), and the cells were cultured for about 3 hr. at about 37° C. Protein production was monitored by SDS-PAGE of recombinant cell lysates, followed by Coomassie blue staining, using standard techniques. Recombinant cell *E. coli*:pTrc-nDiTG₁₄₀₇ produced a fusion protein, herein denoted as PHIS-PDiTG₄₆₈, that migrated with an apparent molecular weight of about 60 kD.

Immunoblot analysis of recombinant cell *E. coli*:pTrc-nDiTG₁₄₀₇ lysates indicated that the about 60 kD protein was able to bind to a T₇ Tag® monoclonal antibody (available from Novagen, Inc., Madison, Wis.) directed against the fusion portion of the recombinant PHIS-PDiTG₄₆₈ fusion protein.

The PHIS-PDiTG₄₆₈ histidine fusion protein was separated from *E. coli* proteins by cobalt chelation chromatography with an imidazole gradient elution. Immunoblot analysis of the *E. coli*:pTrc-nDiTG₁₄₀₇ lysates, column eluate and column void volume indicated that the approximately 60 kD PHIS-PDiTG₄₆₈ protein isolated using cobalt column chromatography was able to selectively bind to a T₇ Tag® monoclonal antibody. The fusion peptide expressed in pTrcHisB contributes approximately 4 kD of vector-encoded amino acid sequence to the recombinant protein; and thus nDiTG₁₄₀₇ encodes a protein of approximately 55kD.

Example 14

This Example discloses the purification of a nematode transglutaminase fusion protein of the present invention from total cell lysates. Also described is the production of anti-DiTG antibodies of the present invention.

Nematode transglutaminase fusion protein PHIS-PDiTG₄₆₈ was separated from *E. coli* proteins by Talon™ Metal Affinity Resin Chromatography (available from CLONTECH Laboratories, Inc., Palo Alto, Calif.) according to the manufacturer's instructions. The nematode transglutaminase fusion protein was eluted using an imidazole gradient, pooled and dialyzed against 1xPBS to produce cobalt column-purified PHIS-PDiTG₄₆₈. The dialyzed protein was concentrated using a 10K molecular weight cut off Centrifugal Ultra-free® concentrator (available from Millipore Corporation, Bedford, Mass.). The protein content of the fusion protein was determined using a MicroBCA™ Protein Assay (available from Pierce, Rockford, Ill.). The purified protein was tested for its purity by SDS PAGE and immunoblot analysis as described in Example 3.

Anti-PHIS-PDiTG₄₆₈ (anti-DiTG) antisera was produced as follows: A rabbit was immunized subcutaneously, first with approximately 75 μg of the purified PHIS-PDiTG₄₆₈ (see above) protein with complete Freund's Adjuvant (available from Sigma, St. Louis, Mo.), and then with three subsequent immunizations of the same dose of PHIS-PDiTG₄₆₈ mixed in Incomplete Freund's Adjuvant. Bleeding and immunization were performed at alternate weeks. Sera were separated and stored at -70° C. until use.

The immunoglobulin G (IgG) fraction (anti-DiTG IgG) from anti-DiTG antisera was collected by 50% ammonium sulfate precipitation. Ammonium ions were removed by extensive dialysis in 0.1 M PBS, pH 7.2. The IgG content was determined by measuring absorbance at OD₂₈₀ and comparing absorbance with that of a blank PBS control. The anti-DiTG IgG had a titer of 1:124,000 as determined by ELISA.

Example 15

This Example describes the PCR amplification and subsequent isolation and characterization of transglutaminase nucleic acid molecules from other related filarial Nematode transglutaminase nucleic acid molecules from *Brugia malayi* and *Onchocerca volvulus* were identified using standard PCR technology and methods as follows. In brief, nematode transglutaminase nucleic acid molecules were PCR amplified from *B. malayi* female adult cDNA using two primers, a sense primer representing the SL sequence, having the nucleotide sequence, 5' GGTTTAATTAC-CCAAGTTTGAG 3' (herein denoted as SEQ ID NO:22) and an antisense primer 5' GCTGATGGACCTGCCTGTC-CACGC 3' (herein denoted as SEQ ID NO:26). PCR amplification of *B. malayi* adult female cDNA using these primers resulted in the production of a 440-bp nucleic acid molecule (herein denoted as nBmTG₄₄₀). Nematode transglutaminase nucleic acid molecules were also amplified from *B. malayi* female adult cDNA using primers corresponding to internal sequences in the cDNA sequence of nDiTG₁₄₇₂ (SEQ ID NO:10). The two primers used were: i) a sense primer spanning from about nucleotide 359 to about nucleotide 371 of SEQ ID NO:10, and having the nucleotide sequence 5' GAAAACCGTTATCAGTATGATCT 3' (SEQ ID NO:19); and ii) an antisense primer spanning from about nucleotides 878 to 901 of SEQ ID NO:10, and having the nucleotide sequence 5' GTCCATTTTTGCAATAACAACACC 3' (SEQ ID NO:21). PCR amplification of adult female cDNA using these primers resulted in the production of a 537 bp nucleic acid molecule (herein denoted as nBmTG₅₃₇; this nucleic acid molecule was previously identified in U.S. patent application Ser. No. 08/781,420 as nBmTG₅₄₂).

Transglutaminase nucleic acid molecules were also PCR amplified from an *O. volvulus* larval cDNA library using two primers that represent internal sequences of nDiTG₁₄₇₂: a sense primer spanning from about nucleotide 359 to about nucleotide 371 of SEQ ID NO:10, and having the nucleotide sequence 5' GAAAACCGTTATCAGTATGATCT 3' (SEQ ID NO:19), and an antisense primer spanning from about nucleotide 878 to about nucleotide 901 of SEQ ID NO:10, and having the nucleotide sequence GTCCATTTG-CAATAACAACACC 3' (SEQ ID NO:21). PCR amplification of adult female cDNA using these primers resulted in the production of a 537 bp nucleic acid molecule (herein denoted as nOvTG₅₃₇; this nucleic acid molecule was, previously identified in U.S. patent application Ser. No. 08/781,420 as nOvTG₅₄₂).

Nucleic acid molecules nBmTG₄₄₀ (the sequences of the coding and complementary strands herein denoted as SEQ

ID NO:35 and SEQ ID NO:37, respectively), nBmTG₅₃₇ (the sequences of the coding and complementary strands herein denoted as SEQ ID NO:42 and SEQ ID NO:44, respectively), and nOvTG₅₃₇ (the sequences of the coding and complementary strands herein denoted as SEQ ID NO:43 and SEQ ID NO:45, respectively) were gel purified, cloned into TA cloning vector and sequenced using an automated DNA sequencer. Sequence analysis of nBmTG₄₄₀ coding and noncoding strands (represented by SEQ ID NO:35 and SEQ ID NO:37, respectively) showed that nBmTG₄₄₀ included the SL sequence at its 5' end. Furthermore, the entire sequences of nOvTG₅₃₇ and nBmTG₅₃₇ were, respectively, 98.9% and 87.7% identical in the region spanning from about nucleotide 359 to about nucleotide 901 of nDiTG₁₄₇₂ (SEQ ID NO:10).

Translation of SEQ ID NO:35 yields a protein of about 139 amino acids, herein denoted as PBmTG₁₃₉, the amino acid sequence of which is presented in SEQ ID NO:36. The nucleic acid molecule encoding PBmTG₁₃₉ is herein referred to as nBmTG₄₁₇, the nucleic acid sequence of which is represented in SEQ ID NO:38 (the coding strand) and SEQ ID NO:39 (the complementary strand). The amino acid sequence of *D. immitis* PBmTG₁₃₉ (i.e., SEQ ID NO:36) predicts that PBmTG₁₃₉ has an estimated molecular weight of about 14.1 kD and an estimated pI of about 5.1. PC/GENE™ sequence analysis of SEQ ID NO:35 predicts a translation product having an N-terminal hydrophobic signal sequence spanning from about amino acid residue 1 through about amino acid residue 26 of SEQ ID NO:36, and having a predicted cleavage site between about amino acid residue 26 and about amino acid residue 27. The nucleic acid sequence of the predicted mature protein product of nBmTG₄₁₇ (after cleavage at the predicted cleavage site) is an approximately 113 amino acid protein herein denoted as PBmTG₅₃₉ (the amino acid sequence of which is represented by SEQ ID NO:58), encoded by a nucleic acid molecule spanning from about nucleotide 102 to about nucleotide 440 of SEQ ID NO:35 (the coding and complementary sequences of which are herein designated as SEQ ID NO:57, and SEQ ID NO:59, respectively).

Translation of SEQ ID NO:42 yields a protein of about 179 amino acids, herein denoted as PBmTG₁₇₉, the amino acid sequence of which is presented in SEQ ID NO:43. The amino acid sequence of *D. immitis* PBmTG₁₇₉ predicts that PBmTG₁₇₉ has an estimated molecular weight of about 20.8 kD and an estimated pI of about 4.6. Translation of SEQ ID NO:43 yields a protein of about 179 amino acids, herein denoted as POvTG₁₇₉, the amino acid sequence of which is presented in SEQ ID NO: 48. The amino acid sequence of *D. immitis* POvTG₁₇₉ (i.e., SEQ ID NO:44) predicts that POvTG₁₇₉ has an estimated molecular weight of about 20.8 kD and an estimated pI of about 4.6.

Example 16

This Example demonstrates that proteins of the present invention possess transglutaminase activity. The transglutaminase activity of column-purified nematode transglutaminase (PIS-PDiTG₄₆₈) was determined in a microtiter plate assay essentially as previously herein described. In brief, microtiter wells were coated with 1% dimethylcasein, (available from Sigma) at room temperature overnight; uncoated sites were blocked with 1% (w/v) nonfat dry milk. The following reaction mixtures were contained in a total volume of 200 μ l: 100 mM Tris/HCl pH 8.5, 10 mM CaCl₂ (except where otherwise indicated), 10 mM dithiothreitol, 1 mM amine donor substrate 5(biotinamido) pentylamine (BPT), (available from Sigma), and PHIS-PDiTG₄₆₈. Reac-

tions were performed at 55° C. (or as herein indicated) for 2 hours and transglutaminase-catalyzed conjugation of BPT into dimethylcasein was determined by streptavidin-peroxidase and orthophenyldiamine as a reporter system. The enzyme activity (expressed as mU) of extracts was determined relative to the activity of purified guinea pig liver transglutaminase (available from Sigma) tested in the same microtiter plate. Several biochemical factors required for transglutaminase activity of PHIS-PDiTG₄₆₈ were also investigated. The results of these assays are given in Tables 7–12.

TABLE 7

Effect of enzyme concentration on PHIS-PDiTG₄₆₈ activity

Concentration of PHIS-PDiTG ₄₆₈ (μ g/ml)	Activity (mU \pm SEM)
5.0	14.4 \pm 0.6
10.0	42.1 \pm 4.4
15.0	74.4 \pm 7.1

TABLE 8

Effect of Ca²⁺ on transglutaminase activity of PHIS-PDiTG₄₆₈ (15 μ g/ml)

Concentration of Ca ²⁺ (mM)	Activity (mU \pm SEM)
0.0	2.1 \pm 1.0
0.5	65.7 \pm 0.3
2.0	110.6 \pm 1.0
4.0	125.8 \pm 0.0
8.0	125.8 \pm 0.0

TABLE 9

Effect of EDTA on transglutaminase activity of PHIS-PDiTG₄₆₈ (15 μ g/ml)

Concentration of EDTA (mM)	Activity (mU \pm SEM)
0.0	79.8 \pm 9.6
0.5	34.1 \pm 2.8
2.0	6.6 \pm 0.2
4.0	4.8 \pm 2.3
8.0	0.69 \pm 0.0

TABLE 10

Effect of temperature on transglutaminase activity of PHIS-PDiTG₄₆₈ (15 μ g/ml)

Temperature (° C.)	Activity (mU \pm SEM)
37	8.2 \pm 3.1
45	14.3 \pm 3.1
55	52.6 \pm 13
65	30.5 \pm 4.0

TABLE 11

Effect of dithiothreitol (DTT) on transglutaminase activity of PHIS-PDiTG ₄₆₈ (15 μ g/ml)		
Concentration of DTT (μ g/ml)	Activity (mU \pm SEM)	
	+DTT	-DTT
10.0	55.0 \pm 8.0	98.6 \pm 2.0
15.0	83.0 \pm 6.6	149.0 \pm 5.7
20.0	123.0 \pm 3.0	161.0 \pm 3.0

TABLE 12

Effect of inhibitors on transglutaminase activity of PHIS-PDiTG ₄₆₈ (15 μ g/ml)		
Inhibitor	Concentration (mM)	Activity (mU \pm SEM)
None		81.2 \pm 3.3
Monodansylcadaverine	2.0	48.3 \pm 4.0
	4.0	44.6 \pm 5.1
Putrescine	2.0	36.5 \pm 3.9
	4.0	35.3 \pm 4.3
Cystamine	0.5	46.4 \pm 4.6
	1.0	28.3 \pm 2.2

As can be seen from the results presented above, PHIS-PDiTG₄₆₈ was able to cross-link BPT to dimethylcasein and the rate of cross-linking activity was concentration dependent (Table 7). Furthermore, the transglutaminase activity of PHIS-PDiTG₄₆₈ was Ca²⁺-dependent (Table 8) and was inhibited by EDTA (Table 9) and EGTA (data not shown). Interestingly, the PHIS-PDiTG₄₆₈ was found to be highly thermostable with optimum activity observed at 55° C. (Table 10). Dithiothreitol (DTT, available from Sigma) was not absolutely essential for transglutaminase activity. In fact, PHIS-PDiTG₄₆₈ was more active in the absence of DTT (Table 11). The known inhibitors of transglutaminase such as monodansylcadaverine, putrescine, cystamine and iodoacetamide inhibited the transglutaminase activity of PHIS-PDiTG₄₆₈ (Table 12).

Example 17

This Example demonstrates the transglutaminase activity for bovine protein disulfide isomerase (PDI).

Sequence analysis of nDiTG₄₀₇ showed significant homology between the protein encoded by nDiTG and known PDIs. Therefore, bovine PDI was tested to see if it has transglutaminase activity. Transglutaminase activity of PDI (Bovine PDI, available from Sigma) was determined in a microtiter plate assay as described above. The results of the assays are given in Tables 13, 14 and 15.

TABLE 13

Transglutaminase activity of bovine protein disulfide isomerase (PDI)	
Concentration of PDI (μ g/ml)	Transglutaminase activity (mU \pm SEM)
2.5	57.4 \pm 3.0
5.0	93.1 \pm 7.3
10.0	119.9 \pm 7.1
15.0	173.0 \pm 4.5

TABLE 14

Effect of EDTA (10 mM) on transglutaminase activity of bovine protein disulfide isomerase (PDI)	
Concentration of PDI (μ g/ml)	Transglutaminase activity (mU \pm SEM)
2.5	1.5 \pm 0.1
5.0	0.0 \pm 0.0
10.0	0.0 \pm 0.0
15.0	0.0 \pm 0.0

TABLE 15

Effect of temperature on transglutaminase activity of PDI (15 μ g/ml)	
Temperature (° C.)	Activity (mU \pm SEM)
37	30.3 \pm 1.0
45	47.2 \pm 2.5
55	92.1 \pm 4.3
65	67.0 \pm 2.9

The data presented in these Tables demonstrate that, surprisingly, bovine PDI was able to cross-link BPT to dimethylcasein and that the rate of cross-linking activity was concentration dependent (Table 13). The transglutaminase activity of bovine PDI was Ca²⁺-dependent and was inhibited EDTA (Tables 14). Like PHIS-PDiTG₄₆₈, bovine PDI had optimum transglutaminase activity at 55° C. (Table 15).

Example 18

This Example discloses a novel protein disulfide isomerase (PDI) activity of a nematode transglutaminase protein of the present invention.

The protein disulfide isomerase activity of PHIS-PDiTG₄₆₈ was determined essentially as described by Lambert and Freedman (Biochem. J 213:235, 1983). Briefly, known amounts of purified bovine liver PDI and PHIS-PDiTG₄₆₈ were incubated for 2 min. at 30° C. in 0.1 ml of sodium-phosphate buffer (50 mM, pH 7.5) containing 10 μ M DTT and 2.5 mM EDTA. 10 μ l of "scrambled" RNase (5 mg/ml stock, available from Sigma) was then to the above mixture and the incubation was continued for an additional 10 min. 10 μ l samples were drawn from each reaction mixture, and added immediately to 3 ml of pre-chilled TKM buffer (50 mM Tris HCl/25 mM KCl/5 MM MgCl₂, pH 7.5) and assayed for RNase activity at 30° C. in the presence of 0.25 mg yeast RNA (Sigma) by measuring the increase in A₂₆₀ for 2 min. in a Beckman DU-600 spectrophotometer. The results of these assays are presented in Tables 16 and 17.

TABLE 16

PDI activity of PHIS-PDiTG ₄₆₈ and purified PDI from bovine liver		
Protein concentration (μ g/ml)	PDI activity* (Mean \pm SD)	
	PDI	PHIS-PDiTG ₄₆₈
0.5	10.90 \pm 1.1	2.84 \pm 0.3
1.0	9.84 \pm 0.9	22.84 \pm 0.9
2.5	27.36 \pm 2.8	34.36 \pm 3.5
5.0	39.70 \pm 0.4	33.60 \pm 5.2
10.0	58.20 \pm 7.3	43.50 \pm 1.2

TABLE 16-continued

PDI activity of PHIS-PDiTG ₄₆₈ and purified PDI from bovine liver		
Protein concentration ($\mu\text{g/ml}$)	PDI activity* (Mean \pm SD)	
	PDI	PHIS-PDiTG ₄₆₈

*The PDI activity was determined as described (Lambert, N. and Freedman, R.B., 1983, Biochem. J 213, pp. 235-243), and was expressed as the change in A_{260} relative to A_{280} min^{-1} [$\Delta A \text{ min}^{-1}$], measured using a Beckman DU-600 dual wavelength spectrophotometer.

TABLE 17

Effect of transglutaminase inhibitors on PDI activity		
Inhibitor (1 mM)	PDI Activity* (Mean \pm SD)	
	PDI	PHIS-PDiTG ₄₆₈
None	27.4 \pm 2.8	34.4 \pm 3.5
MDC	24.0 \pm 2.0	41.2 \pm 2.9
Cystamine	26.4 \pm 2.9	33.3 \pm 1.8

*The PDI activity was determined as described (Lambert, N. and Freedman, R. B., 1983, Biochem. J. 213, pp. 235-243), and was expressed as the change in A_{260} relative to A_{280} min^{-1} [$\Delta A \text{ min}^{-1}$], measured using a Beckman DU-600 dual wavelength spectrophotometer.

PHIS-PDiTG₄₆₈ was capable of reactivating "scrambled" RNase, and this effect was time- and dose-dependent (Table 16). However, the PDI activity of PHIS-PDiTG₄₆₈ was not inhibited by transglutaminase inhibitors (Table 17).

Example 19

This Example describes ultra-structural studies of molting inhibition of *D. immitis* L₃ by the transglutaminase pseudo substrate monodansylcadaverine (MDC).

As herein described, MDC at a final concentration of 100 μM completely inhibits the molting of *D. immitis* L₃ to L₄. The ultra structural events in larval molting inhibition were studied by culturing *D. immitis* larvae in the presence of MDC and observing the effects on development. Larvae cultured in the presence of MDC were collected every 24 hr. for 6 days, fixed using standard procedures and embedded in resin. Ultra-microtome sections of larvae were prepared and then examined by electron microscopy. The L₃ cuticle in untreated controls started separating from the new L₄ cuticle after 24 hr. in culture and molted by 72 hr. In contrast, the MDC treated L₃ failed to show any separation between the L₃ and L₄ cuticles. In addition, the L₄ cuticle and the accompanying hypodermis were much thinner in MDC treated worms than in controls. Finally, the MDC treated larvae failed to molt even on day 6 whereas most of the larvae in control cultures had molted to L₄ by day six.

Example 20

This Example describes the identification of native *D. immitis* transglutaminase protein by immunoblot analysis.

Rabbit anti-DiTG IgG was used to identify a native *D. immitis* transglutaminase protein in *D. immitis* extracts as follows. The material in crude extracts from *D. immitis* larvae, adult male and female worms, and excretory-secretory (E-S) products from larvae and adults were separated by separating 5 μg protein per lane on a 10-well, 4-20% gradient Tris-glycine SDS-PAGE gel at 200 volts for 1 hour. The separated proteins were then transferred to a nitrocellulose membrane by standard methods. After transfer, the membrane was blocked in 5% (w/v) nonfat dry

milk for 1 hr. at 37° C. The membrane was incubated with rabbit anti-DiTG IgG at a dilution of 1:2500 in Tris buffered saline. After 1 hr. incubation at room temperature, the blot was washed and antibody binding resolved using a peroxidase-labeled goat anti-rabbit IgG secondary antibody (available from Kirkegaard and Perry Laboratories) and the substrate NBT/BCIP (available from Sigma). Rabbit anti-DiTG IgG recognized a 56 kD native *D. immitis* protein in *D. immitis* adult male, female and larval extracts. In addition, a 57 kD *D. immitis* protein was identified in the larval E-S products, but not in the adult E-S.

Example 21

This Example describes immunoblot analysis of bovine protein disulfide isomerase using rabbit anti-DiTG IgG as the primary antibody.

Antibodies raised against PHIS-PDiTG₄₆₈ were analyzed for cross-reactivity with bovine PDI as follows. Bovine PDI (100 ng) was separated by SDS-PAGE and transferred to a nitrocellulose filter essentially as previously described. The nitrocellulose filter containing bovine PDI was probed with rabbit anti-DiTG. Rabbit anti-DiTG failed to react with bovine PDI.

Example 22

This Example describes the immuno-localization of native antigen encoded by nDiTG₁₄₀₇ by light microscopy.

Adult male and female *D. immitis* worms were fixed in 4% paraformaldehyde (available from Sigma) in 0.1 M phosphate buffer, pH 7.2 overnight at 4° C. Fixed worms were cut into 1-cm pieces, dehydrated and embedded in paraffin. Thin sections (about 7 microns) of the worm were then prepared using a microtome. The sections on glass slides were de-paraffinized and dehydrated using graded series of alcohol. The sections were then rehydrated in PBS, and treated for 1 hr. in 0.7% of 30% H₂O₂ in PBS containing 10% ethanol in order to block endogenous peroxidases. For immuno-localization, the slides were blocked in PBS containing 10% fetal calf serum (available from Sigma) and 3% bovine serum albumin (available from Sigma) (PBS/FCS/BSA) for 1 hr. at room temperature. The slides were then flooded with a 1:1000 dilution of anti-DiTG IgG in PBS/BSA, and incubated overnight at 4° C. The slides were then rinsed thoroughly with PBS and the antibody binding resolved using a peroxidase-labeled goat anti-rabbit IgG secondary antibody, and the substrate 3',3'-diaminobenzidine tetrahydrochloride (SigmaFast™ tablets, available from Sigma). After color development, the slides were dehydrated in graded series of alcohol and cleared in xylene. The slides were then covered with cover slips and observed under a Nikon MicroPhot-FXA™ microscope (available from Nikon Corporation, Japan). Using anti-DiTG IgG antibody, the native antigen corresponding to nematode transglutaminase was localized mainly in the contents and the walls of the male reproductive system. In the females, reaction products were seen in the gut epithelium and in the channels in the hypodermis. In addition, labeling was seen in the afibrillar muscle cells in males and in some areas of uterine walls in females.

Example 23

This Example demonstrates that *D. immitis*-infected cats, *D. immitis*-infected dogs and immune dogs generate antibodies that recognize a nematode transglutaminase protein of the present invention.

Recombinant antigen PHIS-PDiTG₄₆₈ (100 μl /well; 1.0 $\mu\text{g/ml}$ in 0.06 M carbonate buffer, pH 9.6) was incubated in

Immulon® 2 microtiter plates (Dynatech Laboratories, Alexandria, Va.) overnight at 4° C. Plates were blocked with 0.01 M PBS (pH 7.4) with 0.05% Tween 20 (available from Sigma) and 5% fetal calf serum (PBS/T/FCS) for 1 hr. at 37° C. Sera from infected and immune animals, diluted 1:25 in PBS/T/FCS, were added to the first row of the ELISA plates and two-fold dilution was carried out. After 1 hr. incubation at 37° C., the plates were washed with PBS/T and a peroxidase-conjugated anti-dog IgG antibody (1:5000) (available from Sigma) was added to detect binding of the primary antibody. After 1 hr. incubation, the plates were washed and substrate was added (o-phenyldiamine, available from Amresco®, Solon, Ohio) with H₂O₂. The enzyme reaction was stopped after 5 min. at room temperature with 4M H₂SO₄. Optical density (OD) was compared with a PBS blank at 490 nm using a SpectraMax™ 250 ELISA reader (available from Molecular Devices, Sunnyvale, Calif.).

Immune dogs (n=4) (immune dogs are defined as described in PCT Publication No. WO 94/15593, published Jul. 21, 1994, by Grieve et al.), D., infected dog (n=8), and infected cats (n=6) had detectable levels of IgG antibodies to PHIS-PDiTG₄₆₈. In infected dogs and cats, the mean antibody levels were significantly higher at days 140–160 days post infection than antibody levels earlier in the infection. Specific antibody response to PHIS-PDiTG₄₆₈ coincided with the onset of maturity of developing worms in the host.

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.

 SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 59

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME/KEY: Xaa = Unknown
 (B) LOCATION: 18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile
 1 5 10 15

Lys Xaa Tyr Asp
 20

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile
 1 5 10 15

Lys Pro Tyr Asp Val Leu Val Lys Phe Tyr Ala Pro
 20 25

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val Tyr Gly Lys
 1 5 10

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Asp Ala Thr Ala Asn Asp Val Pro Pro Phe Gln Val Gln Gly
 1 5 10 15

Phe Pro Thr

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 707 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1...705

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG AAA TTT ACA GAT GCG GAC TTC AAG GAG GGA ATT AAA CCA TAT GAT	48
Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile Lys Pro Tyr Asp	
1 5 10 15	
GTA TTA CTT GTG AAA TTT TAT GCA CCA TGG TGC GGA CAC TGC AAA AAG	96
Val Leu Leu Val Lys Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Lys	
20 25 30	
ATA GCA CCA GAA TTT GAA AAA GCA GCA ACC AAA CTT TTA CAG AAT GAT	144
Ile Ala Pro Glu Phe Glu Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp	
35 40 45	
CCG CCT ATT CAT TTA GCA GAG GTT GAC TGT ACG GAG GAG AAG AAA ACT	192
Pro Pro Ile His Leu Ala Glu Val Asp Cys Thr Glu Glu Lys Lys Thr	
50 55 60	
TGC GAT GAA TAC GGT GTT AGT GGC TTC CCG ACT TTG AAA ATT TTC CGT	240
Cys Asp Glu Tyr Gly Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg	
65 70 75 80	
AAG GGA GAA CTA GCA CAG GAT TAT GAT GGT CCG AGA GTA GCA GAA GGT	288
Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala Glu Gly	
85 90 95	
ATT GTG AAA TAT ATG CGT GGA CAG GCA GGT CCA TCA GCT ACA GAA ATT	336
Ile Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu Ile	
100 105 110	
AAT ACA CAA CAA GAA TTC GAA AAA ATG TTG CAA GCC GAT GAC GTT ACT	384
Asn Thr Gln Gln Glu Phe Glu Lys Met Leu Gln Ala Asp Asp Val Thr	
115 120 125	
ATT TGT GGA TTT TTC GAA GAG AAC AGC AAG TTA AAA GAC TCA TTC TTA	432
Ile Cys Gly Phe Phe Glu Glu Asn Ser Lys Leu Lys Asp Ser Phe Leu	
130 135 140	
AAA GTT GCG GAT ACA GAA AGA GAT CGT TTT AAG TTT GTG TGG ACA TCA	480
Lys Val Ala Asp Thr Glu Arg Asp Arg Phe Lys Phe Val Trp Thr Ser	
145 150 155 160	

-continued

AAT AAA CAA ATT CTG GAA TCA AGG GGA TAC AAT GAT GAT ATC GTC GCA	528
Asn Lys Gln Ile Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala	
165 170 175	
TAT CAA CCG AAG AAA TTT CAT AAT AAA TTT GAA CCA AAT GAA TTC AAG	576
Tyr Gln Pro Lys Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys	
180 185 190	
TAT GAT GGA AAT TAC GAC ACA GAC AAG ATT AAA GAA TTT CTC CTA CAC	624
Tyr Asp Gly Asn Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His	
195 200 205	
GAA ACA AAT GGG CTT GTT GGT ATA CGA ACG GCC GAA AAC CGT TAT CAG	672
Glu Thr Asn Gly Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr Gln	
210 215 220	
TAT GAT CTA CTT CCG ATG TTC GTC GTC TAT GGC AA	707
Tyr Asp Leu Leu Pro Met Phe Val Val Tyr Gly	
225 230 235	

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 235 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile Lys Pro Tyr Asp	
1 5 10 15	
Val Leu Leu Val Lys Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Lys	
20 25 30	
Ile Ala Pro Glu Phe Glu Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp	
35 40 45	
Pro Pro Ile His Leu Ala Glu Val Asp Cys Thr Glu Glu Lys Lys Thr	
50 55 60	
Cys Asp Glu Tyr Gly Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg	
65 70 75 80	
Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala Glu Gly	
85 90 95	
Ile Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu Ile	
100 105 110	
Asn Thr Gln Gln Glu Phe Glu Lys Met Leu Gln Ala Asp Asp Val Thr	
115 120 125	
Ile Cys Gly Phe Phe Glu Glu Asn Ser Lys Leu Lys Asp Ser Phe Leu	
130 135 140	
Lys Val Ala Asp Thr Glu Arg Asp Arg Phe Lys Phe Val Trp Thr Ser	
145 150 155 160	
Asn Lys Gln Ile Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala	
165 170 175	
Tyr Gln Pro Lys Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys	
180 185 190	
Tyr Asp Gly Asn Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His	
195 200 205	
Glu Thr Asn Gly Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr Gln	
210 215 220	
Tyr Asp Leu Leu Pro Met Phe Val Val Tyr Gly	
225 230 235	

(2) INFORMATION FOR SEQ ID NO:7:

-continued

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 707 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

TTGCCATAGA CGACGAACAT CGGAAGTAGA TCATACTGAT AACGGTTTTT GCCTGTCGT      60
ATACCAACAA GCCCATTTGT TTCGTGTAGG AGAAATCTT TAATCTTGTC TGTGTCGTAA      120
TTTCCATCAT ACTTGAATTC ATTTGGTTCA AATTTATTAT GAAATTTCTT CGGTTGATAT      180
GCGACGATAT CATCATGTGA TCCCCTTGAT TCCAGAATTT GTTTATTTGA TGTCCACACA      240
AACTTAAAAC GATCTCTTTC TGTATCCGCA ACTTTTAAGA ATGAGTCTTT TAACTTGCTG      300
TTCTCTTCGA AAAATCCACA AATAGTAACG TCATCGGCTT GCAACATTTT TTCGAATTCT      360
TGTTGTGTAT TAATTTCTGT AGCTGATGGA CCTGCCTGTC CACGCATATA TTTCACAATA      420
CCTTCTGCTA CTCTCGGACC ATCATAATCC TGTGCTAGTT CTCCCTTACG GAAAATTTTC      480
AAAGTCGGGA AGCCACTAAC ACCGTATTCA TCGCAAGTTT TCTTCTCCTC CGTACAGTCA      540
ACCTCTGCTA AATGAATAGG CGGATCATTC TGTAAGAGTT TGTTGCTGTC TTTTCAAAT      600
TCTGGTGCTA TCTTTTTCGA GTGTCCGCAC CATGGTGCAT AAAATTTTAC AAGTAATACA      660
TCATATGGTT TAATCCCTC CTTGAAGTCC GCATCTGTAA ATTTTCAT      707

```

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 705 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

ATGAAATTTA CAGATCGGA CTTCAAGGAG GGAATTAAC CATATGATGT ATTACTTGTG      60
AAATTTTATG CACCATGGTG CGGACACTGC AAAAAGATAG CACCAGAATT TGAAAAGCA      120
GCAACCAAAC TTTTACAGAA TGATCCGCCT ATTCATTTAG CAGAGGTTGA CTGTACGGAG      180
GAGAAGAAAA CTTGCGATGA ATACGGTGT AGTGGCTTCC CGACTTTGAA AATTTTCCGT      240
AAGGGAGAAC TAGCACAGGA TTATGATGGT CCGAGAGTAG CAGAAGGTAT TGTGAAATAT      300
ATGCGTGGAC AGGCAGGTCC ATCAGCTACA GAAATTAATA CACAACAAGA ATTGCAAAAA      360
ATGTTGCAAG CCGATGACGT TACTATTTGT GGATTTTTCG AAGAGAACAG CAAGTTAAAA      420
GACTCATTCT TAAAAGTTGC GGATACAGAA AGAGATCGTT TTAAGTTTGT GTGGACATCA      480
AATAAACAAA TTCTGGAATC AAGGGGATAC AATGATGATA TCGTCGCATA TCAACCGAAG      540
AAATTTTATA ATAAATTTGA ACCAAATGAA TTCAAGTATG ATGGAATTA CGACACAGAC      600
AAGATTAAAG AATTTCTCCT ACACGAAACA AATGGGCTTG TTGGTATAC AACGGCCGAA      660
AACCGTTATC AGTATGATCT ACTTCCGATG TTCGTCGTCT ATGGC      705

```

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 705 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

GCCATAGACG ACGAACATCG GAAGTAGATC ATACTGATAA CGGTTTTTCGG CCGTTCGTAT      60
ACCAACAAGC CCATTTGTTT CGTGTAGGAG AAATCTTTA ATCTTGCTCG TGTCGTAATT      120
TCCATCATACTTGAATTCAT TTGGTTCAAA TTTATTATGA AATTTCTTCG GTTGATATGC      180
GACGATATCA TCATTGTATC CCCTTGATTC CAGAATTTGT TTATTTGATG TCCACACAAA      240
CTTAAAACGA TCTCTTCTCG TATCCGCAAC TTTTAAGAAT GAGTCTTTTA ACTTGCTGTT      300
CTCTTCGAAA AATCCACAAA TAGTAACGTC ATCGGCTTGC AACATTTTTT CGAATTCCTG      360
TTGTGTATTA ATTTCTGTAG CTGATGGACC TGCCTGTCCA CGCATATATT TCACAATACC      420
TTCTGCTACT CTCGGACCAT CATAATCTCG TGCTAGTTCT CCCTTACGGA AAATTTTCAA      480
AGTCGGGAAG CCACTAACAC CGTATTCATC GCAAGTTTTT TTCTCCTCCG TACAGTCAAC      540
CTCTGCTAAA TGAATAGGCG GATCATTCTG TAAAAGTTTG GTTGCTGCTT TTTCAAATTC      600
TGGTGCTATC TTTTTCAGT GTCCGCACCA TGGTGCATAA AATTTACAA GTAATACATC      660
ATATGGTTTA ATCCCTCCT TGAAGTCCGC ATCTGTAAAT TTCAT                          705

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1472 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 2..1105

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

T ATG CGT GGA CAG GCA GGT CCA TCA GCT ACA GAA ATT AAT ACA CAA CAA      49
Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu Ile Asn Thr Gln Gln
   1             5             10             15

GAA TTC GAA AAA ATG TTG CAA GCC GAT GAC GTT ACT ATT TGT GGA TTT      97
Glu Phe Glu Lys Met Leu Gln Ala Asp Asp Val Thr Ile Cys Gly Phe
   20             25             30

TTC GAA GAG AAC AGC AAG TTA AAA GAC TCA TTC TTA AAA GTT GCG GAT      145
Phe Glu Glu Asn Ser Lys Leu Lys Asp Ser Phe Leu Lys Val Ala Asp
   35             40             45

ACA GAA AGA GAT CGT TTT AAG TTT GTG TGG ACA TCA AAT AAA CAA ATT      193
Thr Glu Arg Asp Arg Phe Lys Phe Val Trp Thr Ser Asn Lys Gln Ile
   50             55             60

CTG GAA TCA AGG GGA TAC AAT GAT GAT ATC GTC GCA TAT CAA CCG AAG      241
Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln Pro Lys
   65             70             75             80

AAA TTT CAT AAT AAA TTT GAA CCA AAT GAA TTC AAG TAT GAT GGA AAT      289
Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys Tyr Asp Gly Asn
   85             90             95

TAC GAC ACA GAC AAG ATT AAA GAA TTT CTC CTA CAC GAA ACA AAT GGG      337
Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His Glu Thr Asn Gly
   100            105            110

CTT GTT GGT ATA CGA ACG GCC GAA AAC CGT TAT CAG TAT GAT CTA CTT      385
Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr Gln Tyr Asp Leu Leu
   115            120            125

```

-continued

CCG ATG TTT GTT GTG TAT GGC AAG GTT GAC TAT GAA TTG GAT CCA AAA	433
Pro Met Phe Val Val Tyr Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys	
130 135 140	
GGT TCC AAC TAT TGG CGA AAT CGT GTT CTT ATG GTT GCA AAA GAT TAC	481
Gly Ser Asn Tyr Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr	
145 150 155 160	
AAA AGG AAA GCA AAT TTT GCT ATG AGT AAC AAA GAA GAC TTC TCT TTT	529
Lys Arg Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser Phe	
165 170 175	
GAT CTT GAT GAA TTT GGC TTA GCT AAT CGT AAA GAT ACC AAG CCG CTT	577
Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys Pro Leu	
180 185 190	
GTT GCA GCA CGT AGC AAA AAA GGC AAA TTC TTT ATG AAA GAA GAA TTC	625
Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe Met Lys Glu Glu Phe	
195 200 205	
AGT TTT AGC GTG GAA AAT TTG AAA AAA TTT GTC GAA GAT GTT ATT GGT	673
Ser Phe Ser Val Glu Asn Leu Lys Lys Phe Val Glu Asp Val Ile Gly	
210 215 220	
GAT AGA TTA GAA CCG TAT ATG AAG AGC GAA GAA GCA CCT GAA GAT CAG	721
Asp Arg Leu Glu Pro Tyr Met Lys Ser Glu Glu Ala Pro Glu Asp Gln	
225 230 235 240	
GGT GAT GTT AAG GTC GTT GTT GCT AAG ACA TTC CAA GAA ATG ATC ATG	769
Gly Asp Val Lys Val Val Ala Lys Thr Phe Gln Glu Met Ile Met	
245 250 255	
AAT GTG GAA AAG GAT GTT TTA ATC GAA TTT TAT GCT CCA TGG TGT GGC	817
Asn Val Glu Lys Asp Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly	
260 265 270	
CAC TGC AAA GCA CTC GCA CCG AAA TAT GAT GAA TTA GGC CAG AAA TTA	865
His Cys Lys Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu	
275 280 285	
TCC GGT GAA CCA GGT GTT GTT ATT GCA AAA ATG GAC GCA ACA GCG AAT	913
Ser Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp Ala Thr Ala Asn	
290 295 300	
GAT GTC CCA CCA CCA TTC CAA GTA CAA GGA TTT CCA ACT CTT TAC TGG	961
Asp Val Pro Pro Pro Phe Gln Val Gln Gly Phe Pro Thr Leu Tyr Trp	
305 310 315 320	
GTA CCG AAG AAT AAA AAA GAC AAA CCA GAG CCA TAC TCT GGT GGT CGA	1009
Val Pro Lys Asn Lys Lys Asp Lys Pro Glu Pro Tyr Ser Gly Gly Arg	
325 330 335	
GAA GTG GAT GAT TTT ATT AAA TAC ATC GCG AAG CAT GCA ACG GAA GAA	1057
Glu Val Asp Asp Phe Ile Lys Tyr Ile Ala Lys His Ala Thr Glu Glu	
340 345 350	
CTG AAG GGA TAC AAG AGA GAT GGA AAA CCG AAG AAG AAG GAA GAA TTG	1105
Leu Lys Gly Tyr Lys Arg Asp Gly Lys Pro Lys Lys Lys Glu Glu Leu	
355 360 365	
TAAAGGGTAA TAATGATGAA TTTTAAATTT GATGTGAACC CAAACACCT CAGTTGCTTA	1165
TTGTTGGATA AATATTAAA TCATCCACA GAGCTGTGAT ATGAATTTTC AAATATGTTT	1225
TTTTTTGGTT TATTTTGATA AATTCATATT TTAAGTTGTT ATTTTTTAGT GCCTTAGGCT	1285
GTTTCATCAG TTGCCTTAGG CTATTTTGTC AGTTCGGAAT GTTTATTCGG TTAGCTTAGG	1345
CTTTTTTTTG TTTACCTTAT GTTACTGTTG TTATTGTATT ACTATTTTGC CCTTGTTTTT	1405
TAAATTTTAA ATAAATTTTT TTTGGAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA	1465
AAAAAAA	1472

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 368 amino acids

-continued

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu Ile Asn Thr Gln Gln
 1           5           10           15
Glu Phe Glu Lys Met Leu Gln Ala Asp Val Thr Ile Cys Gly Phe
          20           25           30
Phe Glu Glu Asn Ser Lys Leu Lys Asp Ser Phe Leu Lys Val Ala Asp
          35           40           45
Thr Glu Arg Asp Arg Phe Lys Phe Val Trp Thr Ser Asn Lys Gln Ile
          50           55           60
Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln Pro Lys
          65           70           75           80
Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys Tyr Asp Gly Asn
          85           90           95
Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His Glu Thr Asn Gly
          100          105          110
Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr Gln Tyr Asp Leu Leu
          115          120          125
Pro Met Phe Val Val Tyr Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys
          130          135          140
Gly Ser Asn Tyr Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr
          145          150          155          160
Lys Arg Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser Phe
          165          170          175
Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys Pro Leu
          180          185          190
Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe Met Lys Glu Glu Phe
          195          200          205
Ser Phe Ser Val Glu Asn Leu Lys Lys Phe Val Glu Asp Val Ile Gly
          210          215          220
Asp Arg Leu Glu Pro Tyr Met Lys Ser Glu Glu Ala Pro Glu Asp Gln
          225          230          235          240
Gly Asp Val Lys Val Val Val Ala Lys Thr Phe Gln Glu Met Ile Met
          245          250          255
Asn Val Glu Lys Asp Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly
          260          265          270
His Cys Lys Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu
          275          280          285
Ser Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp Ala Thr Ala Asn
          290          295          300
Asp Val Pro Pro Pro Phe Gln Val Gln Gly Phe Pro Thr Leu Tyr Trp
          305          310          315          320
Val Pro Lys Asn Lys Lys Asp Lys Pro Glu Pro Tyr Ser Gly Gly Arg
          325          330          335
Glu Val Asp Asp Phe Ile Lys Tyr Ile Ala Lys His Ala Thr Glu Glu
          340          345          350
Leu Lys Gly Tyr Lys Arg Asp Gly Lys Pro Lys Lys Lys Glu Glu Leu
          355          360          365

```

(2) INFORMATION FOR SEQ ID NO:12:

-continued

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1472 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTCCAAAAAA AATTTATTTA      60
AAATTTAAAA AACAAGGGCA AAATAGTAAT ACAATAACAA CAGTAACATA AGGTAACAA      120
AAAAAAGCCT AAGCTAACGG AATAAACATT CCGAACTGAC AAAATAGCCT AAGGCAACTG      180
ATGAAACAGC CTAAGGCACT AAAAAATAAC AACTTAAAT ATGAATTTAT CAAAATAAAC      240
CAAAAAAAA CATATTTGAA AATTCATATC ACAGCTCTGT GGAATGATTT AAATATTTAT      300
CCACCAATAA GCAACTGAGG TTGTTTGGGT TCACATCAA TTA AAAATTC ATCATTATTA      360
CCCTTTACAA TTCTTCCTTC TTCTTCGGTT TTCCATCTCT CTTGTATCCC TTCAGTTCTT      420
CCGTTGCATG CTTGCGATG TATTTAATAA AATCATCCAC TTCTCGACCA CCAGAGTATG      480
GCTCTGGTTT GTCTTTTTTA TTCTTCGGTA CCCAGTAAAG AGTTGGAAT CCTTGTACTT      540
GGAATGGTGG TGGGACATCA TTCGCTGTTG CGTCCATTTT TGCAATAACA AAACCTGGTT      600
CACCGATAA TTTCTGGCCT AATTCATCAT ATTTGCGTGC GAGTGCTTTG CAGTGGCCAC      660
ACCATGGAGC ATAAAATTCG ATTA AACAT CCTTTTCCAC ATTCATGATC ATTTCTTGGA      720
ATGTCCTAGC AACACGACC TTAACATCAC CCTGATCTTC AGGTGCTTCT TCGCTCTTCA      780
TATACGGTTC TAATCTATCA CCAATAACAT CTTCGACAAA TTTTTTCAA TTTTCCACGC      840
TAAAAC TGAA TTCTTCTTTC ATAAAGAATT TGCCTTTTTT GCTACGTGCT GCAACAAGCG      900
GCTTGGTATC TTTACGATTA GCTAAGCCAA ATTCATCAAG ATCAAAGAG AAGTCTTCTT      960
TGTTACTCAT AGCAAATTT GCTTTCCTTT TGTAATCTTT TGCAACCATA AGAACAGGAT      1020
TTCGCCAATA GTTGAACCT TTTGGATCCA ATTCATAGTC AACCTTGCCA TACACAACAA      1080
ACATCGGAAG TAGATCATA TATAACGGT TTTGCGCCGT TCGTATACCA ACAAGCCCAT      1140
TTGTTTCGTG TAGGAGAAAT TCTTTAATCT TGTCTGTGTC GTAATTTCCA TCATACTTGA      1200
ATTCATTTGG TTCAAATTTA TATGAAATT TCTTCGGTTG ATATGCGACG ATATCATCAT      1260
TGTATCCCTT TGATTCAGA ATTTGTTTAT TTGATGTCCA CACAACTTA AAACGATCTC      1320
TTTCTGTATC CGCAACTTTT AAGAATGAGT CTTTTAATT GCTGTCTCTC TCGAAAATC      1380
CACAAATAGT AACGTCATCG GCTTGCAACA TTTTTTCGAA TTCTTGTGT GTATTAATTT      1440
CTGTAGCTGA TGGACCTGCC TGTCCACGCA TA      1472

```

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1107 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

ATGCGTGGAC AGGCAGGTCC ATCAGCTACA GAAATTAATA CACAACAAGA ATTCGAAAAA      60
ATGTTGCAAG CCGATGACGT TACTATTTGT GGATTTTTCG AAGAGAACAG CAAGTTAAAA      120
GACTCATTCT TAAAAGTTGC GGATACAGAA AGAGATCGTT TTAAGTTTGT GTGGACATCA      180

```

-continued

AATAAACAAA TTCTGGAATC AAGGGGATAC AATGATGATA TCGTCGCATA TCAACCGAAG	240
AAATTTTATA ATAAATTTGA ACCAAATGAA TTCAAGTATG ATGGAATTA CGACACAGAC	300
AAGATTAAG AATTTCTCCT ACACGAAACA AATGGGCTTG TTGGTATACG AACGGCCGAA	360
AACCGTTATC AGTATGATCT ACTTCCGATG TTTGTTGTGT ATGGCAAGGT TGAATATGAA	420
TTGGATCCAA AAGGTTCCAA CTATTGGCGA AATCGTGTTT TTATGGTTGC AAAAGATTAC	480
AAAAGGAAAG CAAATTTTGC TATGAGTAAC AAAGAAGACT TCTCTTTTGA TCTTGATGAA	540
TTTGGCTTAG CTAATCGTAA AGATACCAAG CCGCTTGTG CAGCACGTAG CAAAAAGGC	600
AAATCTTTA TGAAAGAAGA ATTCAGTTT AGCGTGAAA ATTTGAAAA ATTTGTCGAA	660
GATGTTATTG GTGATAGATT AGAACCGTAT ATGAAGAGCG AAGAAGCACC TGAAGATCAG	720
GGTATGTTA AGGTCGTTGT TGCTAAGACA TTCCAAGAAA TGATCATGAA TGTGAAAAG	780
GATGTTTTAA TCGAATTTTA TGCTCCATGG TGTGGCCACT GCAAAGCACT CGCACCGAAA	840
TATGATGAAT TAGGCCAGAA ATTATCCGGT GAACCAGGTG TTGTTATGTC AAAAATGGAC	900
GCAACAGCGA ATGATGTCCC ACCACCATTC CAAGTACAAG GATTTCCAAC TCTTTACTGG	960
GTACCGAAGA ATAAAAAGA CAAACCAGAG CCATACTCTG GTGGTCGAGA AGTGGATGAT	1020
TTTATTAAAT ACATCGCGAA GCATGCAACG GAAGAAGTGA AGGGATACAA GAGAGATGGA	1080
AAACCGAAGA AGAAGGAAGA ATTTGTAA	1107

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1107 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TTACAATTCT TCCTTCTTCT TCGGTTTTCC ATCTCTCTTG TATCCCTTCA GTTCTTCCGT	60
TGCATGCTTC GCGATGTATT TAATAAAATC ATCCACTTCT CGACCACCAG AGTATGGCTC	120
TGGTTTGTCT TTTTTATTCT TCGGTACCCA GTAAAGAGTT GAAATCCTT TACTTTGGAA	180
TGGTGGTGGG ACATCATTTC CTGTTGCGTC CATTTTTGCA ATAACAACAC CTGGTTCACC	240
GGATAATTC TGGCCTAAT CATCATATTT CGGTGCGAGT GCTTTGCAGT GGCCACACCA	300
TGGAGCATAA AATTCGATTA AAACATCCTT TTCCACATTC ATGATCATTT CTGGGAATGT	360
CTTAGCAACA ACGACCTTAA CATCACCTG ATCTTCAGGT GCTTCTTCGC TCTTCATATA	420
CGGTTCTAAT CTATCACCAA TAACATCTTC GACAAATTTT TTCAAATTTT CCACGCTAAA	480
ACTGAATTCT TCTTTCATAA AGAATTTGCC TTTTTTGCTA CGTGCTGCAA CAAGCGGCTT	540
GGTATCTTTA CGATTAGCTA AGCCAAATTC ATCAAGATCA AAAGAGAAGT CTTCTTTGTT	600
ACTCATAGCA AAATTTGCTT TCCTTTTGTG ATCTTTTGTG ACCATAAGAA CACGATTTTCG	660
CCAATAGTTG GAACCTTTTG GATCCAATTC ATAGTCAACC TTGCCATACA CAACAAACAT	720
CGGAAGTAGA TCATACTGAT AACGGTTTTT GGCCGTTTCG ATACCAACAA GCCCATTGTT	780
TTCGTGTAGG AGAAATCTTT TAATCTTGTG TGTGTCGTAA TTTCCATCAT ACTTGAATTC	840
ATTTGGTTCA AATTTATTAT GAAATTTCTT CGGTTGATAT GCGACGATAT CATCATTGTA	900
TCCCTTGAT TCCAGAATTT GTTTATTTGA TGTCCACACA AACTTAAAC GATCTCTTTC	960
TGTATCCGCA ACTTTTAAGA ATGAGTCTTT TAACTTGCTG TTCTCTTCGA AAAATCCACA	1020

-continued

AATAGTAACG TCATCGGCTT GCAACATTTT TTCGAATTCT TGTTGTGTAT TAATTTCTGT 1080
 AGCTGATGGA CCTGCCTGTC CACGCAT 1107

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATGAARTTYA CNGAYGCNGA YTTYAARGAR GG 32

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TTNCCRTANA CNACRAACAT 20

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AATTAACCCT CACTAAAGGG 20

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTAATACGAC TCACTATAGG GC 22

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAAAACCGTT ATCAGTATGA TCT 23

-continued

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CTGTGGAATG ATTTAAATAT TTATCC

26

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GTCCATTTTT GCAATAACAA CACC

24

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GGTTAATTA CCAAGTTG AG

22

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TCCCTCCTTG AAGTCCGCAT CTGTAAATTT CAT

33

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCGAGCTCGA GAATGAAATT TACAGATGCG GAC

33

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 42 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:
 CAGCCAAGCT TCTTACAATT CTCCTTCTT CTCGGTTTT CC 42

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:
 GCTGATGGAC CTGCCTGTCC ACGC 24

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 143 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 24..143

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:
 GGTTTAATTA CCCAAGTTG AGG ATG ACA CTG GTG AGG TTG TTT 44
 Met Thr Leu Val Arg Leu Phe
 1 5

GAT GCT TCG ATT TTT AAA TTA TTC TTG TTT CTG ATA TTG CCA 86
 Asp Ala Ser Ile Phe Lys Leu Phe Leu Phe Leu Ile Leu Pro
 10 15 20

TTA ACG AAT GCC GAT GGC GAT GTG ATG AAA TTT ACA GAT GCG 128
 Leu Thr Asn Ala Asp Gly Asp Val Met Lys Phe Thr Asp Ala
 25 30 35

GAC TTC AAG GAG GGA 143
 Asp Phe Lys Glu Gly
 40

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 40 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:
 Met Thr Leu Val Arg Leu Phe Asp Ala Ser Ile Phe Lys Leu
 1 5 10

Phe Leu Phe Leu Ile Leu Pro Leu Thr Asn Ala Asp Gly Asp
 15 20 25

Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly
 30 35 40

-continued

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 143 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

```
TCCCTCCTTG AAGTCCGCAT CTGTAAATTT CATCACATCG CCATCGGCAT      50
TCGTTAATGG CAATATCAGA AACAAGAATA ATTTAAAAAT CGAAGCATCA      100
AACAACTCA CCAGTGCAT CCTCAAATT GGGTAATTAA ACC                 143
```

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 120 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

```
ATGACACTGG TGAGGTGTT TGATGCTTCG ATTTTAAAT TATTCTTGTT      50
TCTGATATTG CCATTAACGA ATGCCGATGG CGATGTGATG AAATTTACAG      100
ATGCGGACTT CAAGGAGGGA                                       120
```

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 120 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

```
TCCCTCCTTG AAGTCCGCAT CTGTAAATTT CATCACATCG CCATCGGCAT      50
TCGTTAATGG CAATATCAGA AACAAGAATA ATTTAAAAAT CGAAGCATCA      100
AACAACTCA CCAGTGCAT                                       120
```

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1407 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1404

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

```
ATG AAA TTT ACA GAT GCG GAC TTC AAG GAG GGA ATT AAA CCA      42
Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile Lys Pro
1           5           10
```

-continued

TAT GAT GTA TTA CTT GTG AAA TTT TAT GCA CCA TGG TGC GGA	84
Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala Pro Trp Cys Gly	
15 20 25	
CAC TGC AAA AAG ATA GCA CCA GAA TTT GAA AAA GCA GCA ACC	126
His Cys Lys Lys Ile Ala Pro Glu Phe Glu Lys Ala Ala Thr	
30 35 40	
AAA CTT TTA CAG AAT GAT CCG CCT ATT CAT TTA GCA GAG GTT	168
Lys Leu Leu Gln Asn Asp Pro Pro Ile His Leu Ala Glu Val	
45 50 55	
GAC TGT ACG GAG GAG AAG AAA ACT TGC GAT GAA TAC GGT GTT	210
Asp Cys Thr Glu Lys Lys Thr Cys Asp Glu Tyr Gly Val	
60 65 70	
AGT GGC TTC CCG ACT TTG AAA ATT TTC CGT AAG GGA GAA CTA	252
Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg Lys Gly Glu Leu	
75 80	
GCA CAG GAT TAT GAT GGT CCG AGA GTA GCA GAA GGT ATT GTG	294
Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala Glu Gly Ile Val	
85 90 95	
AAA TAT ATG CGT GGA CAG GCA GGT CCA TCA GCT ACA GAA ATT	336
Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu Ile	
100 105 110	
AAT ACA CAA CAA GAA TTC GAA AAA ATG TTG CAA GCC GAT GAC	378
Asn Thr Gln Gln Glu Phe Glu Lys Met Leu Gln Ala Asp Asp	
115 120 125	
GTT ACT ATT TGT GGA TTT TTC GAA GAG AAC AGC AAG TTA AAA	420
Val Thr Ile Cys Gly Phe Phe Glu Glu Asn Ser Lys Leu Lys	
130 135 140	
GAC TCA TTC TTA AAA GTT GCG GAT ACA GAA AGA GAT CGT TTT	462
Asp Ser Phe Leu Lys Val Ala Asp Thr Glu Arg Asp Arg Phe	
145 150	
AAG TTT GTG TGG ACA TCA AAT AAA CAA ATT CTG GAA TCA AGG	504
Lys Phe Val Trp Thr Ser Asn Lys Gln Ile Leu Glu Ser Arg	
155 160 165	
GGA TAC AAT GAT GAT ATC GTC GCA TAT CAA CCG AAG AAA TTT	546
Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln Pro Lys Lys Phe	
170 175 180	
CAT AAT AAA TTT GAA CCA AAT GAA TTC AAG TAT GAT GGA AAT	588
His Asn Lys Phe Glu Pro Asn Glu Phe Lys Tyr Asp Gly Asn	
185 190 195	
TAC GAC ACA GAC AAG ATT AAA GAA TTT CTC CTA CAC GAA ACA	630
Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His Glu Thr	
200 205 210	
AAT GGG CTT GTT GGT ATA CGA ACG GCC GAA AAC CGT TAT CAG	672
Asn Gly Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr Gln	
215 220	
TAT GAT CTA CTT CCG ATG TTC GTC GTC TAT GGC AAG GTT GAC	714
Tyr Asp Leu Leu Pro Met Phe Val Val Tyr Gly Lys Val Asp	
225 230 235	
TAT GAA TTG GAT CCA AAA GGT TCC AAC TAT TGG CGA AAT CGT	756
Tyr Glu Leu Asp Pro Lys Gly Ser Asn Tyr Trp Arg Asn Arg	
240 245 250	
GTT CTT ATG GTT GCA AAA GAT TAC AAA AGG AAA GCA AAT TTT	798
Val Leu Met Val Ala Lys Asp Tyr Lys Arg Lys Ala Asn Phe	
255 260 265	
GCT ATG AGT AAC AAA GAA GAC TTC TCT TTT GAT CTT GAT GAA	840
Ala Met Ser Asn Lys Glu Asp Phe Ser Phe Asp Leu Asp Glu	
270 275 280	
TTT GGC TTA GCT AAT CGT AAA GAT ACC AAG CCG CTT GTT GCA	882
Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys Pro Leu Val Ala	
285 290	

-continued

GCA CGT AGC AAA AAA GGC AAA TTC TTT ATG AAA GAA GAA TTC Ala Arg Ser Lys Lys Gly Lys Phe Phe Met Lys Glu Glu Phe 295 300 305	924
AGT TTT AGC GTG GAA AAT TTG AAA AAA TTT GTC GAA GAT GTT Ser Phe Ser Val Glu Asn Leu Lys Lys Phe Val Glu Asp Val 310 315 320	966
ATT GGT GAT AGA TTA GAA CCG TAT ATG AAG AGC GAA GAA GCA Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys Ser Glu Glu Ala 325 330 335	1008
CCT GAA GAT CAG GGT GAT GTT AAG GTC GTT GTT GCT AAG ACA Pro Glu Asp Gln Gly Asp Val Lys Val Val Val Ala Lys Thr 340 345 350	1050
TTC CAA GAA ATG ATC ATG AAT GTG GAA AAG GAT GTT TTA ATC Phe Gln Glu Met Ile Met Asn Val Glu Lys Asp Val Leu Ile 355 360	1092
GAA TTT TAT GCT CCA TGG TGT GGC CAC TGC AAA GCA CTC GCA Glu Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Ala Leu Ala 365 370 375	1134
CCG AAA TAT GAT GAA TTA GGC CAG AAA TTA TCC GGT GAA CCA Pro Lys Tyr Asp Glu Leu Gly Lys Leu Ser Gly Glu Pro 380 385 390	1176
GGT GTT GTT ATT GCA AAA ATG GAC GCA ACA GCG AAT GAT GTC Gly Val Val Ile Ala Lys Met Asp Ala Thr Ala Asn Asp Val 395 400 405	1218
CCA CCA CCA TTC CAA GTA CAA GGA TTT CCA ACT CTT TAC TGG Pro Pro Pro Phe Gln Val Gln Gly Phe Pro Thr Leu Tyr Trp 410 415 420	1260
GTA CCG AAG AAT AAA AAA GAC AAA CCA GAG CCA TAC TCT GGT Val Pro Lys Asn Lys Lys Asp Lys Pro Glu Pro Tyr Ser Gly 425 430	1302
GGT CGA GAA GTG GAT GAT TTT ATT AAA TAC ATC GCG AAG CAT Gly Arg Glu Val Asp Asp Phe Ile Lys Tyr Ile Ala Lys His 435 440 445	1344
GCA ACG GAA GAA CTG AAG GGA TAC AAG AGA GAT GGA AAA CCG Ala Thr Glu Glu Leu Lys Gly Tyr Lys Arg Asp Gly Lys Pro 450 455 460	1386
AAG AAG AAG GAA GAA TTG TAA Lys Lys Lys Glu Glu Leu 465	1407

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 468 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile Lys Pro 1 5 10
Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala Pro Trp Cys Gly 15 20 25
His Cys Lys Lys Ile Ala Pro Glu Phe Glu Lys Ala Ala Thr 30 35 40
Lys Leu Leu Gln Asn Asp Pro Pro Ile His Leu Ala Glu Val 45 50 55
Asp Cys Thr Glu Glu Lys Lys Thr Cys Asp Glu Tyr Gly Val 60 65 70

-continued

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 440 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 24..440

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

```

GGTTTAATTA CCCAAGTTTG AGG ATG GCG CAG TTG AGG CTG TTT      44
      Met Ala Gln Leu Arg Leu Phe
      1                               5

AAT CAT GCT TCG GTT TTG AAT TTA TTC TTA TTA CTG GTA TTG      86
Asn His Ala Ser Val Leu Asn Leu Phe Leu Leu Leu Val Leu
      10                             15                             20

CCG GTA GCA AAT GGC GAT GGT GAT GTG ATG AAA TTC ACA GAT      128
Pro Val Ala Asn Gly Asp Gly Asp Val Met Lys Phe Thr Asp
      25                             30                             35

GCT GAT TTT AAG GAA GGA ATC AAA TCA TAT GAT GTA TTA CTT      170
Ala Asp Phe Lys Glu Gly Ile Lys Ser Tyr Asp Val Leu Leu
      40                             45

GTG AAA TTT TAT GCA CCA TGG TGT GGG CAC TGC AAG AAA CTG      212
Val Lys Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Lys Leu
      50                             55                             60

GCC CCA GAA TTT GAG AAG GCA GCA ACA AAA CTT TTA CAA AAT      254
Ala Pro Glu Phe Glu Lys Ala Ala Thr Lys Leu Leu Gln Asn
      65                             70                             75

GAT CCA CCT ATT CAT TTA GCA GAT GTC GAT TGC ACA GAG GAA      296
Asp Pro Pro Ile His Leu Ala Asp Val Asp Cys Thr Glu Glu
      80                             85                             90

AAG AAA ATT TGC GAT GAA TTC AGT GTT AGT GGT TTT CCG ACT      338
Lys Lys Ile Cys Asp Glu Phe Ser Val Ser Gly Phe Pro Thr
      95                             100                            105

TTA AAA ATT TTC CGT AAG GGT GAA CTG GCT CAG GAT TAT GAT      380
Leu Lys Ile Phe Arg Lys Gly Glu Leu Ala Gln Asp Tyr Asp
      110                            115

GGC CCA CGA GTT GCA GAA GGT ATT GTT AAA TAT ATG CGT GGA      422
Gly Pro Arg Val Ala Glu Gly Ile Val Lys Tyr Met Arg Gly
      120                            125                            130

CAG GCA GGT CCA TCA GCT      440
Gln Ala Gly Pro Ser Ala
      135

```

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 139 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

```

Met Ala Gln Leu Arg Leu Phe Asn His Ala Ser Val Leu Asn
  1                               5                               10

Leu Phe Leu Leu Leu Val Leu Pro Val Ala Asn Gly Asp Gly

```

-continued

15	20	25										
Asp Val Met Lys Phe Thr	Asp Ala Asp Phe Lys	Glu Gly Ile										
30	35	40										
Lys Ser Tyr Asp Val Leu Leu	Val Lys Phe Tyr Ala	Pro Trp										
45	50	55										
Cys Gly His Cys Lys Lys Leu	Ala Pro Glu Phe Glu	Lys Ala										
60	65	70										
Ala Thr Lys Leu Leu Gln Asn	Asp Pro Pro Ile His	Leu Ala										
75	80											
Asp Val Asp Cys Thr Glu Glu	Lys Lys Ile Cys Asp	Glu Phe										
85	90	95										
Ser Val Ser Gly Phe Pro Thr	Leu Lys Ile Phe Arg	Lys Gly										
100	105	110										
Glu Leu Ala Gln Asp Tyr Asp	Gly Pro Arg Val Ala	Glu Gly										
115	120	125										
Ile Val Lys Tyr Met Arg Gly	Gln Ala Gly Pro Ser	Ala										
130	135											

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 440 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

AGCTGATGGA CCTGCCTGTC CACGCATATA TTTAACAATA CCTTCTGCAA	50
CTCGTGGGCC ATCATAATCC TGAGCCAGTT CACCCTTACG GAAAATTTTT	100
AAAGTCGGAA AACCACTAAC ACTGAATTC TCGCAAATTT TCTTTTCCTC	150
TGTGCAATCG ACATCTGCTA AATGAATAGG TGGATCATT TGTAAAAGTT	200
TTGTTGCTGC CTTCTCAAAT TCTGGGGCCA GTTCTTTCGCA GTGCCACAC	250
CATGGTGCAT AAAATTTTAC AAGTAATACA TCATATGATT TGATTCCTTC	300
CTTAAATCA GCATCTGTGA ATTTTCATCAC ATCACCATCG CCATTTGCTA	350
CCGGCAATAC CAGTAATAAG AATAAATTC AAACCGAAGC ATGATTAAAC	400
AGCCTCAACT GCGCCATCCT CAAACTTGGG TAATTAAACC	440

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 417 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

ATGGCGCAGT TGAGGCTGTT TAATCATGCT TCGGTTTTGA ATTTATTCTT	50
ATTACTGGTA TTGCCGGTAG CAAATGGCGA TGGTGATGTG ATGAAATTC	100
CAGATGCTGA TTTTAAGGAA GGAATCAAAT CATATGATGT ATTACTTGTG	150
AAATTTTATG CACCATGGTG TGGGCACTGC AAGAACTGG CCCGAGAATT	200
TGAGAAGGCA GCAACAAAAC TTTTACAAA TGATCCACCT ATTCATTTAG	250

-continued

CAGATGTCGA TTGCACAGAG GAAAAGAAAA TTTGCGATGA ATTCAGTGTT	300
AGTGGTTTTTC CGACTTTAAA AATTTTCCGT AAGGGTGAAC TGGCTCAGGA	350
TTATGATGGC CCACGAGTTG CAGAAGGTAT TGTTAAATAT ATGCGTGGAC	400
AGGCAGGTCC ATCAGCT	417

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 417 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

AGCTGATGGA CTGCCTGTC CACGCATATA TTAAACAATA CCTTCTGCAA	50
CTCGTGGGCC ATCATAATCC TGAGCCAGTT CACCCTTACG GAAAATTTTT	100
AAAGTCGGAA AACCACTAAC ACTGAATTCA TCGCAAATTT TCTTTTCCCTC	150
TGTGCAATCG ACATCTGCTA AATGAATAGG TGGATCATT TGTAAAAGTT	200
TTGTTGCTGC CTTCTCAAT TCTGGGCCA GTTCTTGCA GTGCCACAC	250
CATGGTGCAT AAAATTTCAC AAGTAATACA TCATATGATT TGATTCCCTC	300
CTTAAATCA GCATCTGTGA ATTTTCATCAC ATCACCATCG CCATTTGCTA	350
CCGGCAATAC CAGTAATAAG AATAAATCA AAACCGAAGC ATGATTA AAC	400
AGCCTCAACT GCGCCAT	417

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..537

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GAA AAC CGT TAT CAG TAT GAT CTG CTC CCA ATG TTT GTT GTG Glu Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val 1 5 10	42
TAC AGC AAG ATT GAC TAT GAA TTG GAT CCA AAA GGG TCC AAT Tyr Ser Lys Ile Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn 15 20 25	84
TAT TGG AGA AAT CGT GTT CTT ACA GTT GCA AAG GAT TAC AGA Tyr Trp Arg Asn Arg Val Leu Thr Val Ala Lys Asp Tyr Arg 30 35 40	126
AGA AAA GCA TAT TTT GCT ATA AGT AAT AAG GAC GAT TTC TCA Arg Lys Ala Tyr Phe Ala Ile Ser Asn Lys Asp Asp Phe Ser 45 50 55	168
TTT GAC CTT GAT GAA TTT GGC TTA GCT GGT CGT AAA GAT ACT Phe Asp Leu Asp Glu Phe Gly Leu Ala Gly Arg Lys Asp Thr 60 65 70	210
AAA CCG CTT GTT GCA GCT CGT AGT AAG AAA GGC AAA TTC TTC Lys Pro Leu Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe	252

-continued

75	80	
ATG AAA GAA GAG TTC AGC GTG GAA AAT TTG AGA AAA TTT GTC		294
Met Lys Glu Glu Phe Ser Val Glu Asn Leu Arg Lys Phe Val		
85	90	95
GAA GAC GTT ATA AAT GAT AGA TTA GAA CCA CAT ATG AAA AGC		336
Glu Asp Val Ile Asn Asp Arg Leu Glu Pro His Met Lys Ser		
100	105	110
GAG GAA CCA CCG GAA GAA CAG GGC GAT GTT AAG GTT GTT GTT		378
Glu Glu Pro Pro Glu Glu Gln Gly Asp Val Lys Val Val Val		
115	120	125
GCT AAA ACA TTC CAA GAG ATG GTT GTT GAT GTG GAA AAG GAT		420
Ala Lys Thr Phe Gln Glu Met Val Val Asp Val Glu Lys Asp		
130	135	140
GTC CTG ATC GAA TTC TAT GCT CCA TGG TGT GGA CAT TGC AAG		462
Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys Lys		
145	150	
GCA CTA GCA CCT AAA TAT GAT GAA TTA GGC CAG AAA TTA TCC		504
Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu Ser		
155	160	165
GGC GAA CCA GGT GTT GTT ATT GCA AAA ATG GAC		537
Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp		
170	175	

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 179 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Glu	Asn	Arg	Tyr	Gln	Tyr	Asp	Leu	Leu	Pro	Met	Phe	Val	Val
1				5					10				
Tyr	Ser	Lys	Ile	Asp	Tyr	Glu	Leu	Asp	Pro	Lys	Gly	Ser	Asn
15				20					25				
Tyr	Trp	Arg	Asn	Arg	Val	Leu	Thr	Val	Ala	Lys	Asp	Tyr	Arg
30					35				40				
Arg	Lys	Ala	Tyr	Phe	Ala	Ile	Ser	Asn	Lys	Asp	Asp	Phe	Ser
45					50				55				
Phe	Asp	Leu	Asp	Glu	Phe	Gly	Leu	Ala	Gly	Arg	Lys	Asp	Thr
60						65			70				
Lys	Pro	Leu	Val	Ala	Ala	Arg	Ser	Lys	Lys	Gly	Lys	Phe	Phe
75						80							
Met	Lys	Glu	Glu	Phe	Ser	Val	Glu	Asn	Leu	Arg	Lys	Phe	Val
85					90				95				
Glu	Asp	Val	Ile	Asn	Asp	Arg	Leu	Glu	Pro	His	Met	Lys	Ser
100					105				110				
Glu	Glu	Pro	Pro	Glu	Glu	Gln	Gly	Asp	Val	Lys	Val	Val	Val
115						120			125				
Ala	Lys	Thr	Phe	Gln	Glu	Met	Val	Val	Asp	Val	Glu	Lys	Asp
130						135			140				
Val	Leu	Ile	Glu	Phe	Tyr	Ala	Pro	Trp	Cys	Gly	His	Cys	Lys
145						150							
Ala	Leu	Ala	Pro	Lys	Tyr	Asp	Glu	Leu	Gly	Gln	Lys	Leu	Ser
155					160				165				
Gly	Glu	Pro	Gly	Val	Val	Ile	Ala	Lys	Met	Asp			

-continued

170

175

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GTCACATTTTT GCAATAACAA CACCTGGTTC ACCGGATAAT TTCTGGCCTA	50
ATTTCATCATA TTTCGGTGCG AGTGTCTTGC AGTGCCACA CCATGGAGCA	100
TAAAATTCGA TTTAAACATC CTTTTCCACA TTCATGATCA TTTCTTGAA	150
TGTCTTAGCA ACAACGACCT TAGCATCACC CTGATCTTCA GGTGCTTCTT	200
CGCTCTTCAT ATACGGTCTT AATCTATCAC CAATAACATC TTCGACAAAT	250
TTTTCCAAAT TTCCACGCT GAATCTTCTT TTCATAAAGA ATTTGCCTTT	300
TTTGCTACGT GCTGCAACAA GCGGCTTGGT ATCTTTACGA TTAGCTAAGC	350
CAAATTCATC AAGATCAAAA GAGAAGTCTT CTTTGTTACT CATAGCAAAA	400
TTTGCTTTCC TTTTGTAATC TTTTGCAACC ATAAGAACAC GATTTGCGCA	450
ATAGTTGGAA CCTTTTGGAT CCAATTCATA GTCAACCTTG CCATACACAA	500
CAAACATCGG AAGTAGATCA TACTGATAAC GGTTTTC	537

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..537

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GAA AAC CGT TAT CAG TAT GAT CTA CTT CCG ATG TTT GTT GTG	42
Glu Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val	
1 5 10	
TAT GGC AAG GTT GAC TAT GAA TTG GAT CCA AAA GGT TCC AAC	84
Tyr Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn	
15 20 25	
TAT TGG CGA AAT CGT GTT CTT ATG GTT GCA AAA GAT TAC AAA	126
Tyr Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr Lys	
30 35 40	
AGG AAA GCA AAT TTT GCT ATG AGT AAC AAA GAA GAC TTC TCT	168
Arg Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser	
45 50 55	
TTT GAT CTT GAT GAA TTT GGC TTA GCT AAT CGT AAA GAT ACC	210
Phe Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr	
60 65 70	
AAG CCG CTT GTT GCA GCA CGT AGC AAA AAA GGC AAA TTC TTT	252
Lys Pro Leu Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe	
75 80	
ATG AAA GAA GAA TTC AGC GTG GAA AAT TTG GAA AAA TTT GTC	294

-continued

Met Lys Glu Glu Phe Ser Val Glu Asn Leu Glu Lys Phe Val	
85 90 95	
GAA GAT GTT ATT GGT GAT AGA TTA GAA CCG TAT ATG AAG AGC	336
Glu Asp Val Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys Ser	
100 105 110	
GAA GAA GCA CCT GAA GAT CAG GGT GAT GCT AAG GTC GTT GTT	378
Glu Glu Ala Pro Glu Asp Gln Gly Asp Ala Lys Val Val Val	
115 120 125	
GCT AAG ACA TTC CAA GAA ATG ATC ATG AAT GTG GAA AAG GAT	420
Ala Lys Thr Phe Gln Glu Met Ile Met Asn Val Glu Lys Asp	
130 135 140	
GTT TTA ATC GAA TTT TAT GCT CCA TGG TGT GGC CAC TGC AAA	462
Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys Lys	
145 150	
GCA CTC GCA CCG AAA TAT GAT GAA TTA GGC CAG AAA TTA TCC	504
Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu Ser	
155 160 165	
GGT GAA CCA GGT GTT GTT ATT GCA AAA ATG GAC	537
Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp	
170 175	

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 179 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Glu Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val	
1 5 10	
Tyr Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn	
15 20 25	
Tyr Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr Lys	
30 35 40	
Arg Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser	
45 50 55	
Phe Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr	
60 65 70	
Lys Pro Leu Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe	
75 80	
Met Lys Glu Glu Phe Ser Val Glu Asn Leu Glu Lys Phe Val	
85 90 95	
Glu Asp Val Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys Ser	
100 105 110	
Glu Glu Ala Pro Glu Asp Gln Gly Asp Ala Lys Val Val Val	
115 120 125	
Ala Lys Thr Phe Gln Glu Met Ile Met Asn Val Glu Lys Asp	
130 135 140	
Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys Lys	
145 150	
Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu Ser	
155 160 165	
Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp	
170 175	

-continued

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

```

GTCACATTTT GCAATAACAA CACCTGGTTC ACCGGATAAT TTCTGGCCTA      50
ATTCATCATA TTTCGGTGCG AGTGTCTTGC AGTGGCCACA CCATGGAGCA      100
TAAAATTCGA TAAAAACATC CTTTTCACA TTCATGATCA TTTCTTGAA      150
TGTCTTAGCA ACAACGACCT TAGCATCACC CTGATCTTCA GGTGCTTCTT      200
CGCTCTTCAT ATACGGTCTT AATCTATCAC CAATAACATC TTCGACAAAT      250
TTTTCCAAAT TTCCACGCT GAATTCTTCT TTCATAAAGA ATTTGCCTTT      300
TTTGCTACGT GCTGCAACAA GCGGCTTGGT ATCTTTACGA TTAGCTAAGC      350
CAAATTCATC AAGATCAAAA GAGAAGTCTT CTTTGTTACT CATAGCAAAA      400
TTTGCTTTCC TTTTGTAATC TTTTGCAACC ATAAGAACAC GATTTGCGCA      450
ATAGTTGGAA CCTTTTGGAT CCAATTCATA GTCAACCTTG CCATACACAA      500
CAAACATCGG AAGTAGATCA TACTGATAAC GGTTTTC      537

```

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1881 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 24..1514

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

```

GGTTTAATTA CCCAAGTTTG AGG ATG ACA CTG GTG AGG TTG TTT      44
      Met Thr Leu Val Arg Leu Phe
      1                5
GAT GCT TCG ATT TTT AAA TTA TTC TTG TTT CTG ATA TTG CCA      86
Asp Ala Ser Ile Phe Lys Leu Phe Leu Phe Leu Ile Leu Pro
      10                15                20
TTA ACG AAT GCC GAT GGC GAT GTG ATG AAA TTT ACA GAT GCG      128
Leu Thr Asn Ala Asp Gly Asp Val Met Lys Phe Thr Asp Ala
      25                30                35
GAC TTC AAG GAG GGA ATT AAA CCA TAT GAT GTA TTA CTT GTG      170
Asp Phe Lys Glu Gly Ile Lys Pro Tyr Asp Val Leu Leu Val
      40                45
AAA TTT TAT GCA CCA TGG TGC GGA CAC TGC AAA AAG ATA GCA      212
Lys Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Lys Ile Ala
      50                55                60
CCA GAA TTT GAA AAA GCA GCA ACC AAA CTT TTA CAG AAT GAT      254
Pro Glu Phe Glu Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp
      65                70                75
CCG CCT ATT CAT TTA GCA GAG GTT GAC TGT ACG GAG GAG AAG      296
Pro Pro Ile His Leu Ala Glu Val Asp Cys Thr Glu Glu Lys
      80                85                90

```

-continued

AAA ACT TGC GAT GAA TAC GGT GTT AGT GGC TTC CCG ACT TTG Lys Thr Cys Asp Glu Tyr Gly Val Ser Gly Phe Pro Thr Leu 95 100 105	338
AAA ATT TTC CGT AAG GGA GAA CTA GCA CAG GAT TAT GAT GGT Lys Ile Phe Arg Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly 110 115	380
CCG AGA GTA GCA GAA GGT ATT GTG AAA TAT ATG CGT GGA CAG Pro Arg Val Ala Glu Gly Ile Val Lys Tyr Met Arg Gly Gln 120 125 130	422
GCA GGT CCA TCA GCT ACA GAA ATT AAT ACA CAA CAA GAA TTC Ala Gly Pro Ser Ala Thr Glu Ile Asn Thr Gln Gln Glu Phe 135 140 145	464
GAA AAA ATG TTG CAA GCC GAT GAC GTT ACT ATT TGT GGA TTT Glu Lys Met Leu Gln Ala Asp Val Thr Ile Cys Gly Phe 150 155 160	506
TTC GAA GAG AAC AGC AAG TTA AAA GAC TCA TTC TTA AAA GTT Phe Glu Glu Asn Ser Lys Leu Lys Asp Ser Phe Leu Lys Val 165 170 175	548
GCG GAT ACA GAA AGA GAT CGT TTT AAG TTT GTG TGG ACA TCA Ala Asp Thr Glu Arg Asp Arg Phe Lys Phe Val Trp Thr Ser 180 185	590
AAT AAA CAA ATT CTG GAA TCA AGG GGA TAC AAT GAT GAT ATC Asn Lys Gln Ile Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile 190 195 200	632
GTC GCA TAT CAA CCG AAG AAA TTT CAT AAT AAA TTT GAA CCA Val Ala Tyr Gln Pro Lys Lys Phe His Asn Lys Phe Glu Pro 205 210 215	674
AAT GAA TTC AAG TAT GAT GGA AAT TAC GAC ACA GAC AAG ATT Asn Glu Phe Lys Tyr Asp Gly Asn Tyr Asp Thr Asp Lys Ile 220 225 230	716
AAA GAA TTT CTC CTA CAC GAA ACA AAT GGG CTT GTT GGT ATA Lys Glu Phe Leu Leu His Glu Thr Asn Gly Leu Val Gly Ile 235 240 245	758
CGA ACG GCC GAA AAC CGT TAT CAG TAT GAT CTA CTT CCG ATG Arg Thr Ala Glu Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met 250 255	800
TTC GTC GTC TAT GGC AAG GTT GAC TAT GAA TTG GAT CCA AAA Phe Val Val Tyr Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys 260 265 270	842
GGT TCC AAC TAT TGG CGA AAT CGT GTT CTT ATG GTT GCA AAA Gly Ser Asn Tyr Trp Arg Asn Arg Val Leu Met Val Ala Lys 275 280 285	884
GAT TAC AAA AGG AAA GCA AAT TTT GCT ATG AGT AAC AAA GAA Asp Tyr Lys Arg Lys Ala Asn Phe Ala Met Ser Asn Lys Glu 290 295 300	926
GAC TTC TCT TTT GAT CTT GAT GAA TTT GGC TTA GCT AAT CGT Asp Phe Ser Phe Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg 305 310 315	968
AAA GAT ACC AAG CCG CTT GTT GCA GCA CGT AGC AAA AAA GGC Lys Asp Thr Lys Pro Leu Val Ala Ala Arg Ser Lys Lys Gly 320 325	1010
AAA TTC TTT ATG AAA GAA GAA TTC AGT TTT AGC GTG GAA AAT Lys Phe Phe Met Lys Glu Glu Phe Ser Phe Ser Val Glu Asn 330 335 340	1052
TTG AAA AAA TTT GTC GAA GAT GTT ATT GGT GAT AGA TTA GAA Leu Lys Lys Phe Val Glu Asp Val Ile Gly Asp Arg Leu Glu 345 350 355	1094
CCG TAT ATG AAG AGC GAA GAA GCA CCT GAA GAT CAG GGT GAT Pro Tyr Met Lys Ser Glu Glu Ala Pro Glu Asp Gln Gly Asp 360 365 370	1136

-continued

GTT AAG GTC GTT GTT GCT AAG ACA TTC CAA GAA ATG ATC ATG Val Lys Val Val Val Ala Lys Thr Phe Gln Glu Met Ile Met 375 380 385	1178
AAT GTG GAA AAG GAT GTT TTA ATC GAA TTT TAT GCT CCA TGG Asn Val Glu Lys Asp Val Leu Ile Glu Phe Tyr Ala Pro Trp 390 395	1220
TGT GGC CAC TGC AAA GCA CTC GCA CCG AAA TAT GAT GAA TTA Cys Gly His Cys Lys Ala Leu Ala Pro Lys Tyr Asp Glu Leu 400 405 410	1262
GGC CAG AAA TTA TCC GGT GAA CCA GGT GTT GTT ATT GCA AAA Gly Gln Lys Leu Ser Gly Glu Pro Gly Val Val Ile Ala Lys 415 420 425	1304
ATG GAC GCA ACA GCG AAT GAT GTC CCA CCA CCA TTC CAA GTA Met Asp Ala Thr Ala Asn Asp Val Pro Pro Pro Phe Gln Val 430 435 440	1346
CAA GGA TTT CCA ACT CTT TAC TGG GTA CCG AAG AAT AAA AAA Gln Gly Phe Pro Thr Leu Tyr Trp Val Pro Lys Asn Lys Lys 445 450 455	1388
GAC AAA CCA GAG CCA TAC TCT GGT GGT CGA GAA GTG GAT GAT Asp Lys Pro Glu Pro Tyr Ser Gly Gly Arg Glu Val Asp Asp 460 465	1430
TTT ATT AAA TAC ATC GCG AAG CAT GCA ACG GAA GAA CTG AAG Phe Ile Lys Tyr Ile Ala Lys His Ala Thr Glu Glu Leu Lys 470 475 480	1472
GGA TAC AAG AGA GAT GGA AAA CCG AAG AAG AAG GAA GAA TTG Gly Tyr Lys Arg Asp Gly Lys Pro Lys Lys Lys Glu Glu Leu 485 490 495	1514
TAAAGGGTAA TAATGATGAA TTTTAAATTT GATGTGAACC CAAACAACCT	1564
CAGTTGCTTA TTGGTGGATA AATATTTAAA TCATTCCACA GAGCTGTGAT	1614
ATGAATTTTC AAATATGTTT TTTTTGGTT TATTTTGATA AATTCATATT	1664
TTAAGTTGTT ATTTTTTAGT GCCTTAGGCT GTTTCATCAG TTGCCTTAGG	1714
CTATTTTGTG AGTTCGGAAT GTTTATTCGG TTAGCTTAGG CTTTTTTTTG	1764
TTTACCTTAT GTTACTGTTG TTATTGTATT ACTATTTTGC CCTTGTTTTT	1814
TAAATTTTAA ATAAATTTTT TTTGGAAAAA AAAAAAAAAA AAAAAAAAAA	1864
AAAAAAAAAA AAAAAAA	1881

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 497 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Met Thr Leu Val Arg Leu Phe Asp Ala Ser Ile Phe Lys Leu
1 5 10

Phe Leu Phe Leu Ile Leu Pro Leu Thr Asn Ala Asp Gly Asp
15 20 25

Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu Gly Ile Lys
30 35 40

Pro Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala Pro Trp Cys
45 50 55

Gly His Cys Lys Lys Ile Ala Pro Glu Phe Glu Lys Ala Ala
60 65 70

-continued

Thr Lys Leu Leu Gln Asn Asp Pro Pro Ile His Leu Ala Glu
 75 80
 Val Asp Cys Thr Glu Glu Lys Lys Thr Cys Asp Glu Tyr Gly
 85 90 95
 Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg Lys Gly Glu
 100 105 110
 Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala Glu Gly Ile
 115 120 125
 Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser Ala Thr Glu
 130 135 140
 Ile Asn Thr Gln Gln Glu Phe Glu Lys Met Leu Gln Ala Asp
 145 150
 Asp Val Thr Ile Cys Gly Phe Phe Glu Glu Asn Ser Lys Leu
 155 160 165
 Lys Asp Ser Phe Leu Lys Val Ala Asp Thr Glu Arg Asp Arg
 170 175 180
 Phe Lys Phe Val Trp Thr Ser Asn Lys Gln Ile Leu Glu Ser
 185 190 195
 Arg Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln Pro Lys Lys
 200 205 210
 Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys Tyr Asp Gly
 215 220
 Asn Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu Leu His Glu
 225 230 235
 Thr Asn Gly Leu Val Gly Ile Arg Thr Ala Glu Asn Arg Tyr
 240 245 250
 Gln Tyr Asp Leu Leu Pro Met Phe Val Val Tyr Gly Lys Val
 255 260 265
 Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn Tyr Trp Arg Asn
 270 275 280
 Arg Val Leu Met Val Ala Lys Asp Tyr Lys Arg Lys Ala Asn
 285 290
 Phe Ala Met Ser Asn Lys Glu Asp Phe Ser Phe Asp Leu Asp
 295 300 305
 Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys Pro Leu Val
 310 315 320
 Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe Met Lys Glu Glu
 325 330 335
 Phe Ser Phe Ser Val Glu Asn Leu Lys Lys Phe Val Glu Asp
 340 345 350
 Val Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys Ser Glu Glu
 355 360
 Ala Pro Glu Asp Gln Gly Asp Val Lys Val Val Val Ala Lys
 365 370 375
 Thr Phe Gln Glu Met Ile Met Asn Val Glu Lys Asp Val Leu
 380 385 390
 Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys Lys Ala Leu
 395 400 405
 Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu Ser Gly Glu
 410 415 420
 Pro Gly Val Val Ile Ala Lys Met Asp Ala Thr Ala Asn Asp
 425 430

-continued

Val Pro Pro Pro Phe Gln Val Gln Gly Phe Pro Thr Leu Tyr
 435 440 445
 Trp Val Pro Lys Asn Lys Lys Asp Lys Pro Glu Pro Tyr Ser
 450 455 460
 Gly Gly Arg Glu Val Asp Asp Phe Ile Lys Tyr Ile Ala Lys
 465 470 475
 His Ala Thr Glu Glu Leu Lys Gly Tyr Lys Arg Asp Gly Lys
 480 485 490
 Pro Lys Lys Lys Glu Glu Leu
 495

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1881 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTTTTTTTTT TTCCAAAAA	50
AATTTATTTA AAATTTAAAA AACAAAGGCA AAATAGTAAT ACAATAACAA	100
CAGTAACATA AGGTAAACAA AAAAAAGCCT AAGCTAACGG AATAAACATT	150
CCGAACGTAC AAAATAGCCT AAGGCAACTG ATGAAACAGC CTAAGGCACT	200
AAAAAATAAC AACTTAAAAAT ATGAATTTAT CAAAATAAAC CAAAAAATAA	250
CATATTTGAA AATTCATATC ACAGCTCTGT GGAATGATTT AAATATTTAT	300
CCACCAATAA GCAACTGAGG TTGTTTGGGT TCACATCAA TTAATAATTC	350
ATCATTATTA CCCTTTACAA TTCTTCCTTC TTCTTCGGTT TTCCATCTCT	400
CTTGATCCC TTCAGTCTT CCGTTGCATG CTTCGCGATG TATTTAATAA	450
AATCATCCAC TTCTCGACCA CCAGAGTATG GCTCTGGTTT GTCTTTTTTA	500
TTCTTCGGTA CCCAGTAAAG AGTTGGAAT CCTTGACTT GGAATGGTGG	550
TGGGACATCA TTCGCTGTTG CGTCCATTTT TGCAATAACA ACACCTGGTT	600
CACCGGATAA TTTCTGGCCT AATTCATCAT ATTTCCGGTGC GAGTGCCTTG	650
CAGTGGCCAC ACCATGGAGC ATAAAAATCG ATTA AACAT CCTTTCCAC	700
ATTCATGATC ATTTCTTGA ATGCTTAGC AACACGACC TTAACATCAC	750
CCTGATCTC AGGTGCTTCT TCGCTCTTCA TATACGGTTC TAATCTATCA	800
CCAATAACAT CTTCGACAAA TTTTTTCAA TTTCCACGC TAAAACTGAA	850
TTCTTCTTTC ATAAAGAATT TGCCTTTTTT GCTACGTGCT GCAACAAGCG	900
GCTTGGTATC TTTACGATTA GCTAAGCCAA ATTCATCAAG ATCAAAAGAG	950
AAGCTTCTT TGTACTCAT AGCAAAATTT GCTTTCCTTT TGTAACTTTT	1000
TGCAACCATA AGAACACGAT TTCGCCAATA GTTGGAACCT TTTGGATCCA	1050
ATTCATAGTC AACCTTGCCA TAGACGACGA ACATCGGAAG TAGATCATAC	1100
TGATAACGGT TTTTCGGCCGT TCGTATACCA ACAAGCCCAT TTGTTTCGTG	1150
TAGGAGAAAT TCTTTAATCT TGTCTGTGTC GTAATTTCCA TCATACTGA	1200
ATTCATTTGG TTCAAATTTA TTATGAAATT TCTTCGGTTG ATATGCGACG	1250
ATATCATCAT TGTATCCCCT TGATCCAGA ATTTGTTTAT TTGATGTCCA	1300

-continued

CACAACTTA AACGATCTC TTTCTGTATC CGCAACTTTT AAGAATGAGT	1350
CTTTTAACTT GCTGTTCTCT TCGAAAAATC CACAAATAGT AACGTCATCG	1400
GCTTGCAACA TTTTTCGAA TTCTTGTTGT GTATTAATTT CTGTAGCTGA	1450
TGGACCTGCC TGTCCACGCA TATATTTAC AATACCTTCT GCTACTCTCG	1500
GACCATCATA ATCCTGTGCT AGTTCTCCCT TACGGAAAAAT TTTCAAAGTC	1550
GGGAAGCCAC TAACACCGTA TTCATCGCAA GTTTTCTTCT CCTCCGTACA	1600
GTCAACCTCT GCTAAATGAA TAGGCGGATC ATTCTGTAAA AGTTTGTTG	1650
CTGCTTTTTC AAATTCGGT GCTATCTTTT TGCAGTGTC GCACCATGGT	1700
GCATAAAATT TCACAAGTAA TACATCATAT GGTTTAATTC CCTCCTTGAA	1750
GTCCGCATCT GTAAATTTCA TCACATCGCC ATCGGCATTC GTTAATGGCA	1800
ATATCAGAAA CAAGAATAAT TTAATAATCG AAGCATCAA CAACCTCACC	1850
AGTGTCTCC TCAAACCTGG GTAATTAAC C	1881

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1494 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

ATGACACTGG TGAGGTTGTT TGATGCTTCG ATTTTAAAT TATTCTTGTT	50
TCTGATATTG CCATTAACGA ATGCCGATGG CGATGTGATG AAATTTACAG	100
ATGCGGACTT CAAGGAGGGA ATTAACCAT ATGATGTATT ACTTGTGAAA	150
TTTTATGCAC CATGGTGGG ACACTGCAAA AAGATAGCAC CAGAATTTGA	200
AAAAGCAGCA ACCAACTTT TACAGAATGA TCCGCCTATT CATTAGCAG	250
AGGTTGACTG TACGGAGGAG AAGAAAATT GCGATGAATA CGGTGTTAGT	300
GGCTTCCCGA CTTTGAAAAT TTTCCGTAAG GGAGAACTAG CACAGGATTA	350
TGATGGTCCG AGAGTAGCAG AAGGTATTGT GAAATATATG CGTGGACAGG	400
CAGGTCCATC AGCTACAGAA ATTAATACAC AACAGAATT CGAAAAATG	450
TTGCAAGCCG ATGACGTTAC TATTTGTGGA TTTTTCGAAG AGAACAGCAA	500
GTTAAAAGAC TCATTCTTAA AAGTTGCGGA TACAGAAAGA GATCGTTTTA	550
AGTTTGTGTG GACATCAAAT AAACAAATTC TGGAATCAAG GGGATACAAT	600
GATGATATCG TCGCATATCA ACCGAAGAAA TTTCATAATA AATTTGAACC	650
AAATGAATTC AAGTATGATG GAAATTACGA CACAGACAAG ATTAAGAAT	700
TTCTCCTACA CGAAACAAAT GGGCTTGTG GTATACGAAC GGCCGAAAAC	750
CGTTATCAGT ATGATCTACT TCCGATGTTT GTCGTCTATG GCAAGGTTGA	800
CTATGAATTG GATCCAAAAG GTTCCAATA TTGGCGAAAT CGTGTCTTTA	850
TGGTTGCAAA AGATTACAAA AGGAAAGCAA ATTTTGCTAT GAGTAACAAA	900
GAAGACTTCT CTTTGTGATCT TGATGAATTT GGCTTAGCTA ATCGTAAAGA	950
TACCAAGCCG CTTGTTGCAG CACGTAGCAA AAAAGGCAA TTCTTTATGA	1000
AAGAAGAATT CAGTTTGTAG GTGAAAAATT TGAAAAAATT TGTCGAAGAT	1050

-continued

GTTATTGGTG ATAGATTAGA ACCGTATATG AAGAGCGAAG AAGCACCTGA	1100
AGATCAGGGT GATGTTAAGG TCGTTGTTGC TAAGACATTC CAAGAAATGA	1150
TCATGAATGT GGAAAAGGAT GTTTTAATCG AATTTTATGC TCCATGGTGT	1200
GGCCACTGCA AAGCACTCGC ACCGAAATAT GATGAATTAG GCCAGAAATT	1250
ATCCGGTGAA CCAGGTGTTG TTTATTGCAA AATGGACGCA ACAGCGAATG	1300
ATGTCCCACC ACCATTCCAA GTACAAGGAT TTCCAACCTT TTAAGGGTA	1350
CCGAAGAATA AAAAAGACAA ACCAGAGCCA TACTCTGGTG GTCGAGAAGT	1400
GGATGATTTT ATTAATAACA TCGCGAAGCA TGCAACGGAA GAACTGAAGG	1450
GATACAAGAG AGATGGAAAA CCGAAGAAGA AGGAAGAATT GTAA	1494

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1494 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

TTACAATTCT TCCTTCTTCT TCGGTTTCC ATCTCTCTTG TATCCCTTCA	50
GTTCTTCCGT TGCATGCTTC GCGATGTATT TAATAAAATC ATCCACTTCT	100
CGACCACCAG AGTATGGCTC TGGTTTGTCT TTTTATCTCT TCGGTACCCA	150
GTAAAGAGTT GGAATCCTT TACTTTGGAA TGGTGGTGGG ACATCATTCG	200
CTGTTGCGTC CATTTTGGCA ATAACAACAC CTGGTTCACC GGATAATTC	250
TGGCCTAATT CATCATATTT CGGTGCGAGT GCTTTGCACT GGCCACACCA	300
TGGAGCATAA AATTCGATTA AAACATCCTT TTCCACATTC ATGATCATTT	350
CTTGGAATGT CTTAGCAACA ACGACCTTAA CATCACCTG ATCTTCAGGT	400
GCTTCTTCGC TCTTCATATA CGGTTCTAAT CTATCACCAA TAACATCTTC	450
GACAAATTTT TTCAAATTTT CCACGCTAAA ACTGAATCTT TCTTTCATAA	500
AGAATTTGCC TTTTTTGCTA CGTGCTGCAA CAAGCGGCTT GGTATCTTTA	550
CGATTAGCTA AGCCAAATTC ATCAAGATCA AAAGAGAAGT CTTCTTTGTT	600
ACTCATAGCA AAATTTGCTT TCCTTTTGTA ATCTTTTGCA ACCATAAGAA	650
CACGATTCG CCAATAGTTG GAACCTTTTG GATCCAATTC ATAGTCAACC	700
TTGCCATAGA CGACGAACAT CGGAAGTAGA TCATACTGAT AACGGTTTTT	750
GGCCGTTTCT ATACCAACAA GCCCATTTGT TTCGTGTAGG AGAAATCTTT	800
TAATCTTGTC TGTGTCGTAA TTTCCATCAT ACTTGAATTC ATTTGGTTCA	850
AATTTATTAT GAAATTTCTT CGGTTGATAT GCGACGATAT CATCATTTGA	900
TCCCCTTGAT TCCAGAATTT GTTTATTTGA TGTCCACACA AACTTAAAAC	950
GATCTCTTTC TGTATCCGCA ACTTTTAAGA ATGAGTCTTT TAACTTGCTG	1000
TTCTCTTCGA AAAATCCACA AATAGTAACG TCATCGGCTT GCAACATTTT	1050
TTCGAATTTT TGTGTTGAT TAATTTCTGT AGCTGATGGA CCTGCCTGTC	1100
CACGCATATA TTTACAATA CCTTCTGCTA CTCCTGGACC ATCATAATCC	1150
TGTGCTAGTT CTCCCTTACG GAAAATTTTC AAAGTCGGGA AGCCACTAAC	1200

-continued

ACCGTATTCA TCGCAAGTTT TCTTCTCCTC CGTACAGTCA ACCTCTGCTA 1250
 AATGAATAGG CGGATCATTC TGTA AAAAGTT TGGTTGCTGC TTTTCAAAT 1300
 TCTGGTGCTA TCTTTTGGCA GTGTCCGCAC CATGGTGCAT AAAATTTTAC 1350
 AAGTAATACA TCATATGGTT TAATCCCTC CTTGAAGTCC GCATCTGTAA 1400
 ATTTCATCAC ATCGCCATCG GCATTGCTTA ATGGCAATAT CAGAAACAAG 1450
 AATAATTTAA AAATCGAAGC ATCAAACAAC CTCACCAGTG TCAT 1494

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 45 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 1..45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

GAT GGC GAT GTG ATG AAA TTT ACA GAT GCG GAC TTC AAG GAG 42
 Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu
 1 5 10
 GGA 45
 Gly
 15

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu
 1 5 10
 Gly
 15

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 45 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

TCCCTCCTTG AAGTCCGCAT CTGTAAATTT CATCACATCG CCATC 45

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1416 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-continued

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1416

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GAT GGC GAT GTG ATG AAA TTT ACA GAT GCG GAC TTC AAG GAG Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu 1 5 10	42
GGA ATT AAA CCA TAT GAT GTA TTA CTT GTG AAA TTT TAT GCA Gly Ile Lys Pro Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala 15 20 25	84
CCA TGG TGC GGA CAC TGC AAA AAG ATA GCA CCA GAA TTT GAA Pro Trp Cys Gly His Cys Lys Lys Ile Ala Pro Glu Phe Glu 30 35 40	126
AAA GCA GCA ACC AAA CTT TTA CAG AAT GAT CCG CCT ATT CAT Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp Pro Pro Ile His 45 50 55	168
TTA GCA GAG GTT GAC TGT ACG GAG GAG AAG AAA ACT TGC GAT Leu Ala Glu Val Asp Cys Thr Glu Glu Lys Lys Thr Cys Asp 60 65 70	210
GAA TAC GGT GTT AGT GGC TTC CCG ACT TTG AAA ATT TTC CGT Glu Tyr Gly Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg 75 80	252
AAG GGA GAA CTA GCA CAG GAT TAT GAT GGT CCG AGA GTA GCA Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala 85 90 95	294
GAA GGT ATT GTG AAA TAT ATG CGT GGA CAG GCA GGT CCA TCA Glu Gly Ile Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser 100 105 110	336
GCT ACA GAA ATT AAT ACA CAA CAA GAA TTC GAA AAA ATG TTG Ala Thr Glu Ile Asn Thr Gln Gln Glu Phe Glu Lys Met Leu 115 120 125	378
CAA GCC GAT GAC GTT ACT ATT TGT GGA TTT TTC GAA GAG AAC Gln Ala Asp Asp Val Thr Ile Cys Gly Phe Phe Glu Glu Asn 130 135 140	420
AGC AAG TTA AAA GAC TCA TTC TTA AAA GTT GCG GAT ACA GAA Ser Lys Leu Lys Asp Ser Phe Leu Lys Val Ala Asp Thr Glu 145 150	462
AGA GAT CGT TTT AAG TTT GTG TGG ACA TCA AAT AAA CAA ATT Arg Asp Arg Phe Lys Phe Val Trp Thr Ser Asn Lys Gln Ile 155 160 165	504
CTG GAA TCA AGG GGA TAC AAT GAT GAT ATC GTC GCA TAT CAA Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln 170 175 180	546
CCG AAG AAA TTT CAT AAT AAA TTT GAA CCA AAT GAA TTC AAG Pro Lys Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys 185 190 195	588
TAT GAT GGA AAT TAC GAC ACA GAC AAG ATT AAA GAA TTT CTC Tyr Asp Gly Asn Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu 200 205 210	630
CTA CAC GAA ACA AAT GGG CTT GTT GGT ATA CGA ACG GCC GAA Leu His Glu Thr Asn Gly Leu Val Gly Ile Arg Thr Ala Glu 215 220	672
AAC CGT TAT CAG TAT GAT CTA CTT CCG ATG TTC GTC GTC TAT Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val Tyr 225 230 235	714
GGC AAG GTT GAC TAT GAA TTG GAT CCA AAA GGT TCC AAC TAT Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn Tyr 240 245 250	756

-continued

TGG CGA AAT CGT GTT CTT ATG GTT GCA AAA GAT TAC AAA AGG	798
Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr Lys Arg	
255 260 265	
AAA GCA AAT TTT GCT ATG AGT AAC AAA GAA GAC TTC TCT TTT	840
Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser Phe	
270 275 280	
GAT CTT GAT GAA TTT GGC TTA GCT AAT CGT AAA GAT ACC AAG	882
Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys	
285 290	
CCG CTT GTT GCA GCA CGT AGC AAA AAA GGC AAA TTC TTT ATG	924
Pro Leu Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe Met	
295 300 305	
AAA GAA GAA TTC AGT TTT AGC GTG GAA AAT TTG AAA AAA TTT	966
Lys Glu Phe Ser Phe Ser Val Glu Asn Leu Lys Lys Phe	
310 315 320	
GTC GAA GAT GTT ATT GGT GAT AGA TTA GAA CCG TAT ATG AAG	1008
Val Glu Asp Val Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys	
325 330 335	
AGC GAA GAA GCA CCT GAA GAT CAG GGT GAT GTT AAG GTC GTT	1050
Ser Glu Glu Ala Pro Glu Asp Gln Gly Asp Val Lys Val Val	
340 345 350	
GTT GCT AAG ACA TTC CAA GAA ATG ATC ATG AAT GTG GAA AAG	1092
Val Ala Lys Thr Phe Gln Glu Met Ile Met Asn Val Glu Lys	
355 360	
GAT GTT TTA ATC GAA TTT TAT GCT CCA TGG TGT GGC CAC TGC	1134
Asp Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys	
365 370 375	
AAA GCA CTC GCA CCG AAA TAT GAT GAA TTA GGC CAG AAA TTA	1176
Lys Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu	
380 385 390	
TCC GGT GAA CCA GGT GTT GTT ATT GCA AAA ATG GAC GCA ACA	1218
Ser Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp Ala Thr	
395 400 405	
GCG AAT GAT GTC CCA CCA CCA TTC CAA GTA CAA GGA TTT CCA	1260
Ala Asn Asp Val Pro Pro Pro Phe Gln Val Gln Gly Phe Pro	
410 415 420	
ACT CTT TAC TGG GTA CCG AAG AAT AAA AAA GAC AAA CCA GAG	1302
Thr Leu Tyr Trp Val Pro Lys Asn Lys Lys Asp Lys Pro Glu	
425 430	
CCA TAC TCT GGT GGT CGA GAA GTG GAT GAT TTT ATT AAA TAC	1344
Pro Tyr Ser Gly Gly Arg Glu Val Asp Asp Phe Ile Lys Tyr	
435 440 445	
ATC GCG AAG CAT GCA ACG GAA GAA CTG AAG GGA TAC AAG AGA	1386
Ile Ala Lys His Ala Thr Glu Glu Leu Lys Gly Tyr Lys Arg	
450 455 460	
GAT GGA AAA CCG AAG AAG AAG GAA GAA TTG	1416
Asp Gly Lys Pro Lys Lys Lys Glu Glu Leu	
465 470	

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 472 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu
 1 5 10

-continued

Gly Ile Lys Pro Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala
 15 20 25
 Pro Trp Cys Gly His Cys Lys Lys Ile Ala Pro Glu Phe Glu
 30 35 40
 Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp Pro Pro Ile His
 45 50 55
 Leu Ala Glu Val Asp Cys Thr Glu Glu Lys Lys Thr Cys Asp
 60 65 70
 Glu Tyr Gly Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg
 75 80
 Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala
 85 90 95
 Glu Gly Ile Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser
 100 105 110
 Ala Thr Glu Ile Asn Thr Gln Gln Glu Phe Glu Lys Met Leu
 115 120 125
 Gln Ala Asp Asp Val Thr Ile Cys Gly Phe Phe Glu Glu Asn
 130 135 140
 Ser Lys Leu Lys Asp Ser Phe Leu Lys Val Ala Asp Thr Glu
 145 150
 Arg Asp Arg Phe Lys Phe Val Trp Thr Ser Asn Lys Gln Ile
 155 160 165
 Leu Glu Ser Arg Gly Tyr Asn Asp Asp Ile Val Ala Tyr Gln
 170 175 180
 Pro Lys Lys Phe His Asn Lys Phe Glu Pro Asn Glu Phe Lys
 185 190 195
 Tyr Asp Gly Asn Tyr Asp Thr Asp Lys Ile Lys Glu Phe Leu
 200 205 210
 Leu His Glu Thr Asn Gly Leu Val Gly Ile Arg Thr Ala Glu
 215 220
 Asn Arg Tyr Gln Tyr Asp Leu Leu Pro Met Phe Val Val Tyr
 225 230 235
 Gly Lys Val Asp Tyr Glu Leu Asp Pro Lys Gly Ser Asn Tyr
 240 245 250
 Trp Arg Asn Arg Val Leu Met Val Ala Lys Asp Tyr Lys Arg
 255 260 265
 Lys Ala Asn Phe Ala Met Ser Asn Lys Glu Asp Phe Ser Phe
 270 275 280
 Asp Leu Asp Glu Phe Gly Leu Ala Asn Arg Lys Asp Thr Lys
 285 290
 Pro Leu Val Ala Ala Arg Ser Lys Lys Gly Lys Phe Phe Met
 295 300 305
 Lys Glu Glu Phe Ser Phe Ser Val Glu Asn Leu Lys Lys Phe
 310 315 320
 Val Glu Asp Val Ile Gly Asp Arg Leu Glu Pro Tyr Met Lys
 325 330 335
 Ser Glu Glu Ala Pro Glu Asp Gln Gly Asp Val Lys Val Val
 340 345 350
 Val Ala Lys Thr Phe Gln Glu Met Ile Met Asn Val Glu Lys
 355 360
 Asp Val Leu Ile Glu Phe Tyr Ala Pro Trp Cys Gly His Cys
 365 370 375
 Lys Ala Leu Ala Pro Lys Tyr Asp Glu Leu Gly Gln Lys Leu

-continued

380	385	390
Ser Gly Glu Pro Gly Val Val Ile Ala Lys Met Asp Ala Thr		
395	400	405
Ala Asn Asp Val Pro Pro Pro Phe Gln Val Gln Gly Phe Pro		
410	415	420
Thr Leu Tyr Trp Val Pro Lys Asn Lys Lys Asp Lys Pro Glu		
425	430	
Pro Tyr Ser Gly Gly Arg Glu Val Asp Asp Phe Ile Lys Tyr		
435	440	445
Ile Ala Lys His Ala Thr Glu Glu Leu Lys Gly Tyr Lys Arg		
450	455	460
Asp Gly Lys Pro Lys Lys Lys Glu Glu Leu		
465	470	

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1419 nucleotides
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

TTACAATTCT TCCTTCTTCT TCGGTTTTCC ATCTCTCTTG TATCCCTTCA	50
GTTCTTCCGT TGCATGCTTC GCGATGTATT TAATAAAATC ATCCACTTCT	100
CGACCACCAG AGTATGGCTC TGGTTTGTCT TTTTATTCT TCGGTACCCA	150
GTAAGAGATT GAAATCCTT GTAAGTGGAA TGGTGGTGGG ACATCATTCG	200
CTGTTGCGTC CTTTTTGCA ATAACAACAC CTGGTTCACC GGATAATTC	250
TGGCCTAATT CATCATATTT CGGTGCGAGT GCTTTGCAGT GGCCACACCA	300
TGGAGCATAA AATTCGATTA AAACATCCTT TTCCACATTC ATGATCATTT	350
CTTGAATGT CTTAGCAACA ACGACCTTAA CATCACCTG ATCTTCAGGT	400
GCTTCTTCGC TCTTCATATA CGGTTCTAAT CTATCACCAA TAACATCTTC	450
GACAAATTTT TTCAAATTTT CCACGCTAAA ACTGAATCT TCTTTCATAA	500
AGAATTTGCC TTTTTTGCTA CGTGTGCAA CAAGCGGCTT GGTATCTTTA	550
CGATTAGCTA AGCCAAATTC ATCAAGATCA AAAGAGAAGT CTTCTTTGTT	600
ACTCATAGCA AAATTTGCTT TCCTTTTGTA ATCTTTTGCA ACCATAAGAA	650
CACGATTCG CCAATAGTTG GAACCTTTTG GATCCAATTC ATAGTCAACC	700
TTGCCATAGA CGACGAACAT CGGAAGTAGA TCATACTGAT AACGGTTTTC	750
GGCCGTTTCG ATACCAACAA GCCCATTTGT TTCGTGTAGG AGAAATCTTT	800
TAATCTTGTC TGTGTCGTAA TTTCCATCAT ACTTGAATTC ATTTGGTTCA	850
AATTTATTAT GAAATTTCTT CGGTTGATAT GCGACGATAT CATCATTGTA	900
TCCCTTGAT TCCAGAATTT GTTTATTTGA TGTCCACACA AACTTAAAC	950
GATCTCTTTC TGTATCCGCA ACTTTTAAGA ATGAGTCTTT TAACTTGCTG	1000
TTCTCTTCGA AAAATCCACA AATAGTAACG TCATCGGCTT GCAACATTTT	1050
TTGCAATCTT TGTGTGTAT TAATTTCTGT AGCTGATGGA CCTGCCTGTC	1100
CACGCATATA TTTCACAATA CCTTCTGCTA CTCTCGGACC ATCATAATCC	1150

-continued

TGTGCTAGTT CTCCTTACG GAAAATTTTC AAAGTCGGGA AGCCACTAAC	1200
ACCGTATTCA TCGCAAGTTT TCTTCTCCTC CGTACAGTCA ACCTCTGCTA	1250
AATGAATAGG CGGATCATTC TGTAAGAGTT TGGTTGCTGC TTTTCAAAT	1300
TCTGGTGCTA TCTTTTGGCA GTGTCCGCAC CATGGTGCAT AAAATTTTAC	1350
AAGTAATACA TCATATGGTT TAATCCCTC CTTGAAGTCC GCATCTGTAA	1400
ATTTTCATCAC ATGCCCATC	1419

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 339 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..339

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

GAT GGT GAT GTG ATG AAA TTC ACA GAT GCT GAT TTT AAG GAA	42
Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu	
1 5 10	
GGA ATC AAA TCA TAT GAT GTA TTA CTT GTG AAA TTT TAT GCA	84
Gly Ile Lys Ser Tyr Asp Val Leu Leu Val Lys Phe Tyr Ala	
15 20 25	
CCA TGG TGT GGG CAC TGC AAG AAA CTG GCC CCA GAA TTT GAG	126
Pro Trp Cys Gly His Cys Lys Lys Leu Ala Pro Glu Phe Glu	
30 35 40	
AAG GCA GCA ACA AAA CTT TTA CAA AAT GAT CCA CCT ATT CAT	168
Lys Ala Ala Thr Lys Leu Leu Gln Asn Asp Pro Pro Ile His	
45 50 55	
TTA GCA GAT GTC GAT TGC ACA GAG GAA AAG AAA ATT TGC GAT	210
Leu Ala Asp Val Asp Cys Thr Glu Glu Lys Lys Ile Cys Asp	
60 65 70	
GAA TTC AGT GTT AGT GGT TTT CCG ACT TTA AAA ATT TTC CGT	252
Glu Phe Ser Val Ser Gly Phe Pro Thr Leu Lys Ile Phe Arg	
75 80	
AAG GGT GAA CTG GCT CAG GAT TAT GAT GGC CCA CGA GTT GCA	294
Lys Gly Glu Leu Ala Gln Asp Tyr Asp Gly Pro Arg Val Ala	
85 90 95	
GAA GGT ATT GTT AAA TAT ATG CGT GGA CAG GCA GGT CCA TCA	336
Glu Gly Ile Val Lys Tyr Met Arg Gly Gln Ala Gly Pro Ser	
100 105 110	
GCT	339
Ala	

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 113 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Asp Gly Asp Val Met Lys Phe Thr Asp Ala Asp Phe Lys Glu
1 5 10

NO:28, SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:52, and SEQ ID NO:55 with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has transglutaminase activity; and (b) determining if said putative inhibitory compound inhibits said activity.

16. The method of claim 15, said method further comprising contacting an isolated host animal transglutaminase protein with a putative nematode transglutaminase inhibitory compound under conditions in which, in the absence of said compound, said host animal transglutaminase protein has transglutaminase activity; and determining if said putative inhibitory compound inhibits the host animal transglutaminase activity.

17. A method to identify a compound capable of inhibiting protein disulfide isomerase activity, said method comprising (a) contacting an isolated protein comprising an amino acid sequence selected from the group consisting of SEQ ID

NO:28, SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:52, and SEQ ID NO:55 with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has protein disulfide isomerase activity; and (b) determining if said putative inhibitory compound inhibits said activity.

18. The method of claim 17, said method further comprising contacting an isolated host animal protein disulfide isomerase protein with a putative nematode protein disulfide isomerase inhibitory compound under conditions in which, in the absence of said compound, said host animal protein disulfide isomerase protein has protein disulfide isomerase activity; and determining if said putative inhibitory compound inhibits the host animal protein disulfide isomerase activity.

* * * * *