

J-ISOSS.
2015 Vol. 1(1), 01-11

IMPROVING STATISTICAL INFERENCE WITH UNCERTAIN NON-SAMPLE PRIOR INFORMATION

**ShahjahanKhan¹, Muhammed Ashraf Memon²
Budi Pratikno³ and Rossita M Yunus⁴**

Email address of corresponding author: Shahjahan.Khan@usq.edu.au

¹School of Agric., Comput. and Environmental Sciences
International Centre for Applied Climate Sciences, Centre for Health
Sciences Research, University of Southern Queensland, Toowoomba,
Australia

²Sunnybank Obesity Centre &SEQS, Suite 9, 259 McCullough Street
Sunnybank, Queensland, AUSTRALIA Mayne Medical School, University
of Queensland, Australia Faculty of Health Sciences & Medicine, Bond
University, Australia

³Department of Mathematics and Natural Science Jenderal Soedirman
University, Purwokerto, Indonesia

⁴Institute of Mathematical Sciences, University of Malaya Kuala Lumpur,
Malaysia

ABSTRACT

In the classical inference, the observed sample data is the only source of information. The Bayesian inferential methods assume prior distribution of the underlying model parameters to combine with sample data. Often non-sample prior information (NSPI) on the value of the model parameters is available from previous studies or expert knowledge which could be used along with the sample data to improve the quality of statistical inference. Obviously the NSPI is not always correct and hence there is uncertainty in the suspected value of the parameter. Any such uncertainty can be removed by conducting an appropriate statistical test, and the quality of statistical inference can be improved by including the outcome of the test in the inferential procedure. This paper provides the underlying methodology to illustrate the process and include an example to demonstrate its application.

KEYWORDS AND PHRASES: Regression model; uncertain non-sample prior information; restricted, preliminary test and shrinkage estimators; bias, relative efficiency, M and score tests, testing after pre-test, power of tests, correlated non-central bivariate chi-square distribution.

2010 Mathematics Subject Classification: Primary 62F03, 62F30, 62J05; Secondary 62H12 and 62F10.

1 INTRODUCTION

Statistical inference uses both sample and non-sample information. Classical inference uses only the sample data for estimation and test of hypotheses. Bayesian methods uses sample data and prior distribution of the model parameters. The notion of inclusion of non-sample prior information (NSPI) on the value of model parameters has been introduced to 'improve' the quality of statistical inference. The natural expectation is that the inclusion of additional information would result in a better estimator and test with relevant statistical properties. In some cases this may be true, but in many other cases the risk of worse consequences can not be ruled out.

A number of estimators have been introduced in the literature that uses NSPI and, under particular situation, over performs the traditional exclusive sample information based unbiased estimators when judged by criteria such as the mean square error and squared error loss function.

In many studies the researchers estimate the slope parameter of the regression model. However, the estimation of the intercept parameter is more difficult than that of the slope parameter. This is because the estimator of the slope parameter is required in the estimation of the intercept parameter. Khan et al. (2002) studied the improved estimation of the slope parameter for the linear regression model. They introduced the coefficient of distrust on the belief of the null hypothesis, and incorporated this coefficient in the definition and analysis of the estimators.

In recent time (eg Khan and Pratikno, 2013; Yunus and Khan, 2008, 2010, 2011a,b) several studies used NSPI on the slope of a regression model to test the intercept parameter. Yunus (2010) applied the NSPI in the testing regime using M-test along the line of Humber's M-estimation. Pratikno (2012) studied the parametric test for the intercept parameter using NSPI information on the slope of different regression models.

In general the NSPI on the slope is uncertain and may fall into one of the following three categories: (i) unspecified, no information available, (ii) specified, correct value known, and (iii) specified with uncertainty.

This paper provides alternative estimators and tests of the intercept parameter when NSPI on the slope of the simple linear regression model is available. This include the unrestricted (UE), restricted (RE), preliminary test (PTE) estimators as well as the unrestricted (UT), restricted (RT) and pre-test (PTT) tests of the intercept parameter. Statistical properties of these estimators and tests are investigated both analytically and graphically. Motivation for a real life application of test for the intercept is found in Kent (2009).

Studies in the area of the estimation include Bancroft (1944), Han and Bancroft (1968), Sclove et al. (1972), Saleh and Sen (1978, 1985), Judge and Bock (1978), Stein (1981), Khan (1998, 2003, 2008), Chiouand Saleh (2002), Saleh (2006), Khan and Saleh (1997, 2001, 2005), Salel (2006), Khan et al. (2002, 2005), Hoque et al. (2009). The testing problem has been investigated by Tamura (1965), Saleh and Sen (1978, 1985), Yunus (2010), Yunus and Khan (2008, 2011a,b), Pratikno (2012) and Khan and Pratikno (2013).

The next section introduces the model and definition of the unrestricted estimators of θ and σ^2 . The three alternative estimators are defined in Section 3 along with their properties. The three tests and their power analyses are provided in Section 4. Some concluding remarks are given in section 5.

2 THE MODEL AND SOME PRELIMINARIES

The n independently and identically distributed responses from a linear regression model can be expressed by the equation

$$y = \theta \mathbf{1}_n + \beta x + e, \quad (2.1)$$

where y and x are the column vectors of response and explanatory variables respectively, $\mathbf{1}_n = (1, \dots, 1)'$ - a vector of n -tuple of 1's, θ and β are the unknown intercept and slope parameters respectively and $e = (e_1, \dots, e_n)'$ is a vector of errors with independent components which is distributed as $N_n(0, \sigma^2 I_n)$. So that $E(e) = 0$ and $E(ee') = \sigma^2 I_n$ where σ^2 is the variance of each of the error component in e and I_n is the identity matrix of order n .

Assume that uncertain NSPI on the value of β is available, either from previous study or from practical experience of the researchers or experts. Let the NSPI be expressed in the form of $H_0: \beta = 0$ which may be true, but not sure. We wish to incorporate both the sample information and the uncertain NSPI in estimating and testing the intercept θ . Following Khan et al (2002) we assign a coefficient of distrust, $0 \leq d \leq 1$, for the NSPI, that represents the degree of distrust in the null hypothesis.

The *unrestricted* mle of the slope β and intercept θ are given by

$$\tilde{\beta} = (x'x)^{-1} x'y \quad \text{and} \quad \tilde{\theta} = \bar{y} - \tilde{\beta}\bar{x}, \quad (2.2)$$

where $\bar{x} = \frac{1}{n} \sum_{j=1}^n x_j$ and $\bar{y} = \frac{1}{n} \sum_{j=1}^n y_j$. The mle of σ^2 is $S_n^{*2} = \frac{1}{n} (y - \hat{y})' (y - \hat{y})$, where

$\hat{y} = \tilde{\theta} \mathbf{1}_n + \tilde{\beta} x$. This estimator is biased for σ^2 . However, $S_n^2 = \frac{1}{n-2} (y - \hat{y})' (y - \hat{y})$ is

unbiased for σ^2 . To remove the uncertainty from the NSPI, we perform an appropriate statistical test on $H_0: \beta = \beta_0$ against $H_a: \beta \neq \beta_0$. Here the appropriate test is given by

$L_v = S_n^{-1} S_{xx}^{\frac{1}{2}} (\tilde{\beta} - \beta_0)$. Under the H_a , L_v , follows a non-central Student-t distribution with $v = (n-2)$ df and non-centrality parameter $\Delta^2 = \sigma^{-2} S_{xx} (\beta - \beta_0)^2$.

3 ALTERNATIVE ESTIMATORS OF INTERCEPT

In this section we define the alternative estimators of the intercept and investigate its properties.

3.1 The Estimators

The UE, RE and PTE of θ are given by

$$\tilde{\theta}^{\text{UE}} = \bar{y} - \tilde{\beta}\bar{x} \quad (3.1)$$

$$\hat{\theta}^{\text{RE}}(d) = d\tilde{\theta} + (1-d)\hat{\theta}, \quad 0 \leq d \leq 1 \quad (3.2)$$

$$\begin{aligned} \hat{\theta}^{\text{PTE}}(d) &= \hat{\theta}^{\text{RE}}(d)I(F < F_\alpha) + \tilde{\theta}I(F \geq F_\alpha) \\ &= \tilde{\theta} + \tilde{\beta}\bar{x}(1-d)I(F < F_\alpha). \end{aligned} \quad (3.3)$$

The bias of the estimators are obtained as (cf Hoque et al. 2006)

$$B_1[\tilde{\theta}^{\text{UE}}(d)] = 0 \quad (3.4)$$

$$B_2[\hat{\theta}^{\text{RE}}(d)] = S_{xx}^{-1/2}\bar{x}\sigma(1-d)\Delta \quad (3.5)$$

$$B_3[\hat{\theta}^{\text{PTE}}(d)] = (1-d)\bar{x}\beta G_{3,v}(3^{-1}F_\alpha; \Delta^2), \quad (3.6)$$

where $G_{n_1, n_2}(\cdot; \Delta^2)$ is the c.d.f. of a non-central F-distribution with (n_1, n_2) df and non-centrality parameter Δ^2 which is the *departure constant* from the null-hypothesis. Among the three estimators, the UE is the only unbiased estimator.

The mean squared errors (MSE) of the estimators become

$$M_1[\tilde{\theta}^{\text{UE}}] = \sigma^2 H \quad (3.7)$$

$$M_2[\hat{\theta}^{\text{RE}}(d)] = \sigma^2 \left[d^2 H + (1-d)^2 S_{xx}^{-1} \bar{x}^2 \Delta^2 \right] \quad (3.8)$$

$$\begin{aligned} M_3[\hat{\theta}^{\text{PTE}}(d)] &= \sigma^2 H + S_{xx}^{-1} \sigma^2 \bar{x}^2 \left[\Delta^2 \left\{ 2(1-d)G_{3,v}(3^{-1}F_\alpha; \Delta^2) \right. \right. \\ &\quad \left. \left. - (1-d^2)G_{5,v}(5^{-1}F_\alpha; \Delta^2) \right\} - (1-d^2)G_{3,v}(3^{-1}F_\alpha; \Delta^2) \right], \end{aligned} \quad (3.9)$$

where $H = \left\{ n^{-1} + S_{xx}^{-1} \bar{x}^2 \right\}$.

The relative efficiency of the PTE relative to the UE and RE is

$$\text{RE}[\hat{\theta}^{\text{PTE}}(d) : \tilde{\theta}^{\text{UE}}] = H \left[H + S_{xx}^{-1} \bar{x}^2 \sigma^2 g(\Delta^2) \right]^{-1} \quad (3.10)$$

and

$$\begin{aligned} \text{RE}[\hat{\theta}^{\text{PTE}}(d) : \hat{\theta}^{\text{RE}}(d)] &= \left[d^2 H + (1-d)^2 \Delta^2 S_{xx}^{-1} \bar{x}^2 \right] \\ &\quad \times \left[H + S_{xx}^{-1} \bar{x}^2 g(\Delta^2) \right]^{-1} \end{aligned} \quad (3.11)$$

respectively, where

$$\begin{aligned} g(\Delta^2) &= \Delta^2 \left\{ 2(1-d)G_{3,v}(3^{-1}F_\alpha; \Delta^2) - (1-d^2)G_{5,v}(5^{-1}F_\alpha; \Delta^2) \right\} \\ &\quad - (1-d^2)G_{3,v}(3^{-1}F_\alpha; \Delta^2). \end{aligned} \quad (3.12)$$

The bias, MSE and relative efficiency functions of the estimators can be analysed for different values of d and Δ and the performances of the estimators could be compared accordingly. Graphs of the relative efficiency of the PTE is given in Figure 1.




Figure 1: Graph of the relative efficiency of PTE relative to UE and RE against Δ^2 .

4 THREE TESTS OF INTERCEPT

In this section we define three alternative tests of the intercept and investigate their properties.

To remove the uncertainty in the NSPI on β , we perform a pretest (PT) on $H_0^* : \beta = \beta_0$ before testing on the intercept. Let ϕ^{PT} be the test function for pretesting $H_0^* : \beta = \beta_0$ (a suspected constant) against $H_a^* : \beta \neq \beta_0$. If the H_0^* is rejected in the PT, then the UT is used to test the intercept, otherwise the RT is used. The appropriate test statistic for the PT is $T^{PT} = S_n^{-1}(\beta - \beta_0) \sqrt{S_{xx}} \sim t_{n-2}$.

4.1 Three Test Statistics

Under the three scenarios on β the UT, RT and PTT for testing $H_0 : \theta = \theta_0$ (known constant) against $H_a : \theta \neq \theta_0$ are defined as follows:

- (i) ϕ^{UT} = test function and T^{UT} is the test statistic when β is unspecified,
- (ii) ϕ^{RT} = test function and T^{RT} is the test statistic when $\beta = \beta_0$ is specified and
- (iii) ϕ^{PTT} = test function and T^{PTT} is the test statistic following a PT on H_0^* when $\beta = \beta_0$ is uncertain.

The test statistics are obtained as

$$T^{UT} = (\tilde{\theta} - \theta_0) / SE(\tilde{\theta}) = \sqrt{n}(\bar{Y} - \tilde{\beta}\bar{X} - \theta_0) \left[S_n^2 (1 + S_{xx}^{-1} n \bar{X}^2) \right]^{-1/2} \quad (4.1)$$

$$T^{RT} = (\hat{\theta} - \theta_0) / SE(\hat{\theta})\% = \frac{(\hat{\theta} - \theta_0)}{s_y / \sqrt{n}} = s_y^{-1} \sqrt{n}(\bar{Y} - \theta_0) \sim t_{n-1}, \quad (4.2)$$

where $T^{UT} \sim t_{n-2}$, and $s_y^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$. Let us choose a positive number

α_j , ($0 < \alpha_j < 1$, for $j=1, 2, 3$) then let t_{n-2, α_j} be such that $P(T^{UT} > t_{n-2, \alpha_1} \hat{U}\theta = \theta_0) = \alpha_1$,

$P(T^{RT} > t_{n-1, \alpha_2} \hat{U}\theta = \theta_0) = \alpha_2$, and

$P(T^{PT} > t_{n-2, \alpha_3} \hat{U}\beta = \beta_0) = \alpha_3$. Then, the PTT for testing $H_0 : \theta = \theta_0$ when $\beta = \beta_0$ is uncertain is given by the test function

$$\Phi^{PTT} = \begin{cases} 1, & \text{if } (T^{PT} \leq t_{n-2, \alpha_3}, T^{RT} > t_{n-1, \alpha_2}) \\ & \text{or } (T^{PT} > t_{n-2, \alpha_3}, T^{UT} > t_{n-2, \alpha_1}); \\ 0, & \text{otherwise.} \end{cases} \quad (4.3)$$

4.2 Properties of the Tests

Let $\{K_n\}$ be a sequence of alternative hypotheses defined as

$$K_n : (\theta - \theta_0, \beta - \beta_0) = \left(\frac{\lambda_1}{\sqrt{n}}, \frac{\lambda_2}{\sqrt{n}} \right) = n^{-1/2} \lambda, \quad (4.4)$$

where $\lambda = (\lambda_1, \lambda_2)$ is a vector of fixed real numbers and θ is the true value of the intercept. Under K_n , $(\theta - \theta_0) \neq 0$ and under H_0 , $(\theta - \theta_0) = 0$.

Note that T^{UT} and T^{PT} are correlated, but T^{RT} and T^{PT} are uncorrelated (but not independent). The joint distribution of the T_1^{UT} and T_3^{PT} is $(T_1^{UT}, T_3^{PT})' \sim t_{n-2}$, a bivariate

Student- t distribution with $(n-2)$ df and correlation coefficient ρ with

$$\text{Cov}(T_1^{UT}, T_3^{PT}) = \frac{(n-2)}{(n-4)} \Sigma \quad (\text{cf., Kotz and Nadarajah, 2004}).$$

The power functions of the tests are given by

$$\begin{aligned} \pi^{UT}(\lambda) &= P(T^{UT} > t_{\alpha_1, n-2} \hat{U}K_n) \\ &= 1 - P(T_1^{UT} \leq t_{\alpha_1, n-2} - \lambda_1 k^{-1}) \end{aligned} \quad (4.5)$$

$$\begin{aligned} \pi^{RT}(\lambda) &= P(T^{RT} > t_{\alpha_1, n-1} \hat{U}K_n) \\ &= P(T_2^{RT} > t_{\alpha_2, n-1} - \sqrt{n}((\theta - \theta_0) + (\beta - \beta_0)\bar{X})s_y^{-1}) \\ &= 1 - P(T_2^{RT} \leq t_{\alpha_2, n-1} - \lambda_1 + \lambda_2 \bar{X}s_y^{-1}) \end{aligned} \quad (4.6)$$

$$\begin{aligned} \pi^{PTT}(\lambda) &= P(T^{PT} \leq t_{n-2, \alpha_3}, T^{RT} > t_{n-1, \alpha_2}) + P(T^{PT} > t_{n-2, \alpha_3}, T^{UT} > t_{n-2, \alpha_1}) \\ &= d_{10} \left\{ t_{n-2, \alpha_3} - \lambda_2 \sqrt{S_{xx}} [S_n^2 n]^{-\frac{1}{2}}, t_{\alpha_2, n-1} - s_y^{-1} (\lambda_1 + \lambda_2 \bar{X}), \rho = 0 \right\} \\ &\quad + d_{2\rho} \left\{ t_{n-2, \alpha_3} - \lambda_2 \sqrt{S_{xx}} [S_n^2 n]^{-\frac{1}{2}}, t_{\alpha_1, n-2} - \lambda_1 k^{-1}, \rho \neq 0 \right\}, \\ &= d_{10} \left\{ t_{n-2, \alpha_3} - \lambda_2 \frac{\sqrt{S_{xx}}}{S_n \sqrt{n}}, \% t_{\alpha_2, n-1} - \frac{(\lambda_1 + \lambda_2 \bar{X})}{s_y}, \rho = 0 \right\} \\ &\quad + d_{2\rho} \left\{ t_{n-2, \alpha_3} - \lambda_2 \frac{\sqrt{S_{xx}}}{S_n \sqrt{n}}, t_{\alpha_1, n-2} - \lambda_1 k^{-1}, \rho \neq 0 \right\}, \end{aligned} \quad (4.7)$$

where $k = S_n \sqrt{(1 + n\bar{X}^2 S_{xx}^{-1})}$, d_{10} and $d_{2\rho}$ are bivariate Student's t probability integrals.

Here d_{10} is defined as $d_{10} = \int_{-\infty}^a \int_c^\infty f(t^{PT}, t^{RT}) dt^{PT} dt^{RT}$,

$$a = \left[t_{n-2, \alpha_3} - \lambda_2 \frac{\sqrt{S_{xx}}}{S_n \sqrt{n}} \right] \text{ and } c = \left[t_{n-1, \alpha_2} - \frac{\lambda_1 + \lambda_2 \bar{X}}{s_y} \right], \text{ and}$$

$d_{2\rho}$ is defined as

$$d_{2\rho}(a, b, \rho) = \frac{\Gamma\left(\frac{v+2}{2}\right)}{\Gamma\left(\frac{v}{2}\right) n \pi \sqrt{1-\rho^2}} \int_a^\infty \int_b^\infty \left[1 + \frac{(x^2 + y^2 - 2\rho xy)}{v(1-\rho^2)} \right]^{-\frac{v+2}{2}} dx dy,$$

in which $-1 < \rho < 1$ is the correlation coefficient between the T^{UT} , T^{PT} and

$$b = \left[t_{\alpha_1, n-2} - \lambda_1 / k \right].$$




Figure 2: Graphs of the power functions of the UE, RE and PTT for various values of λ_1 , and λ_2 with a fixed value of $\rho = 0.1$.

The power curves of the PTT for different values of ρ is provided in Figure 3.




Figure 3: The power curve of the PTT against λ_2 , and its power and size curves against ρ .

5 CONCLUDING REMARKS

In practice, the NSPI is obtained from expert knowledge or previous studies, and hence the value of the parameter available from prior information is expected to be close to its true value and the degree of distrust on the null hypothesis is very likely to be close to 0.

Based on the above analyses, it is evident that the power of the RT is always higher than that of the UT and PTT, and the power of the PTT lies between the power of the RT and UT for all values of λ_1, λ_2 and ρ . The size of the UT is smaller than that of the RT and PTT.

Of the three tests, the RT has the maximum power and size, and the UT has minimum power and size. So none of them is achieving the highest power and lowest size. But the PTT protects against maximum size of the RT and minimum power of the UT. As $\lambda_2 \rightarrow 0$ the difference between the power of the PTT and RT diminishes for all values of $\lambda_2 \rightarrow 0$. That is, if the NSPI is accurate the power of the PTT is about the same as that of the RT. Moreover, the power of the PTT gets closer to that of the RT as $\rho \rightarrow 1$. If $\rho = 1$ then the power of the PTT matches with that of the RT. Thus if there is a high (near 1) correlation between the T^{UT} and T^{PT} the power of the PTT is very close to that of the RT.

ACKNOWLEDGEMENT

The authors are thankful to the referees and editors for their valuable suggestions. A draft version of this paper was presented at the 13th Islamic Countries Conference on Statistical Sciences (ICCS-13) in Bogor, Indonesia,

REFERENCES

1. Bancroft, T.A. (1944). On biases in estimation due to the use of the preliminary tests of significance. *Annals of Mathematical Statistics*, 15, 190-204.
2. Chiou, P., and Saleh, A K Md E. (2002). Preliminary test confidence sets for the mean of a multivariate normal distribution. *Journal of Propagation in Probability and Statistics*, 2, 177-189.
3. Han, C.P., and Bancroft, T.A. (1968). On pooling means when variance is unknown. *Journal of American Statistical Association*, 63, 1333-1342.
4. Hoque, Z., Khan, S., and Wesolowski, J. (2009). Performance of preliminary test estimator under linex loss function. *Communications in Statistics: Theory and Methods*, 38 (2). pp. 252-261.
5. Judge, G.G., and Bock, M.E. (1978). *The Statistical Implications of Pre-test and Stein-rule Estimators in Econometrics*. North-Holland, New York.
6. Kent, R. (2009). Energy miser-know your plants energy, Fingerprint (accessed 23 May 2011). {URL}:<http://www.ptonline.com/articles/know-your-plants-energy-fingerprint>.
7. Khan, S. (2008). Shrinkage estimators of intercept parameters of two simple regression models with suspected equal slopes. *Communications in Statistics - Theory and Methods*, 37, 247-260.
8. Khan, S. (2003). Estimation of the parameters of two parallel regression lines under uncertain prior information. *Biometrical Journal*, 44, 73-90.
9. Khan, S. (1998). On the estimation of the mean vector of Student-t population with uncertain prior information. *Pakistan Journal of Statistics*, 14, 161-175.
10. Khan, S., Hoque, Z., and Saleh, A K Md E. (2005). Estimation of the intercept parameter for linear regression model with uncertain non-sample prior information. *Statistical Papers*, 46 (3). pp. 379-395
11. Khan, S., Hoque, Z., and Saleh, A K Md E. (2002). Improved estimation of the slope parameter for linear regression model with normal errors and uncertain prior information. *Journal of Statistical Research*, 36, 55-73.

12. Khan, S. and Pratikno, B. (2013) Testing base load with non-sample prior information on process load. *Statistical Papers*, 54(3), 605-617
13. Khan, S., and Saleh, A K Md E. (2001). On the comparison of the pre-test and shrinkage estimators for the univariate normal mean. *Statistical Papers*, 42(4), 451-473.
14. Khan, S., and Saleh, A K Md E. (1997). Shrinkage pre-test estimator of the intercept parameter for a regression model with multivariate Student-t errors. *Biometrical Journal*, 39, 1-17.
15. Pratikno, B. (2012). Test of hypotheses for linear regression models with non-sample prior information. Unpublished PhD Thesis, University of Southern Queensland, Australia
16. Saleh, A K Md E. (2006). Theory of Preliminary Test Stein-Type Estimation with Application, Wiley, New York.
17. Saleh, A K Md E., and Sen, P K. (1985). Shrinkage least squares estimation in a general multivariate linear model. *Proceedings of the Fifth Pannonian Symposium on Mathematical Statistics*, 307-325.
18. Saleh, A K Md E., and Sen, P K. (1978). Nonparametric estimation of location parameter after a preliminary test on regression. *Annals of Statistics*, 6, 154-168.
19. Sclove, S.L., Morris, and Rao, C.R. (1972). Non-optimality of preliminary-test estimators for the mean of a multivariate normal distribution. *Ann. Math. Statist.*, 43, 1481-1490.
20. Stein, C. (1956). Inadmissibility of the usual estimator for the mean of a multivariate normal distribution, *Proceedings of the Third Berkeley Symposium on Math. Statist. And Probability*, University of California Press, Berkeley, 1, 197-206.
21. Stein, C. (1981). Estimation of the mean of a multivariate normal distribution. *Annals of Statistics* 9, 1135--1151.
22. Tamura, R. (1965). Nonparametric inferences with a preliminary test. *Bull. Math. Stat.* 11, 38-61.
23. Yunus, R. M. (2010). Increasing power of M-test through pre-testing. Unpublished PhD Thesis, University of Southern Queensland, Australia.
24. Yunus, R. M., and Khan, S. (2008). Test for intercept after pre-testing on slope - a robust method. In: *9th Islamic Countries Conference on Statistical Sciences (ICCS-IX): Statistics in the Contemporary World - Theories, Methods and Applications*, 81-90.
25. Yunus, R. M., and Khan, S. (2011a). Increasing power of the test through pre-test - a robust method. *Communications in Statistics-Theory and Methods*, 40, 581-597.
26. Yunus, R. M., and Khan, S. (2011b). M-tests for multivariate regression model. *Journal of Nonparametric Statistics*, 23, 201-218.