

SPACE LIFE SCIENCES SYMPOSIUM (A1)
Radiation Fields, Effects and Risks in Human Space Missions (5)

Author: Dr. Christa Baumstark-Khan

Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Germany, christa.baumstark-khan@dlr.de

Mr. Bikash Konda

German Aerospace Center (DLR), Germany, bikash.konda@dlr.de

Mr. Luis Fernando Spitta

Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Germany, luis.spitta@gmx.de

Dr. Christine Hellweg

Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Germany, christine.hellweg@dlr.de

RADIATION RESPONSE OF PORCINE LENS EPITHELIAL CELLS AND EYE LENSES IN
ORGAN-CULTURE

Abstract

Astronauts on long-term space missions have a higher risk for the expression of radiation late effects such as cancer or sub-capsular cortical eye lens opacities. This is due to higher dose and different patterns of cellular energy deposition from high-linear-energy-transfer (LET) components of galactic cosmic radiation in space than that of terrestrial low-LET radiation on Earth. The eye lens is a radiation sensitive organ with radiation induced cataract to occur with a threshold absorbed dose of 0.5 Gy (0 - 1 Gy) of sparsely ionizing radiation. Doses perceived by astronauts on the International Space Station (ISS) are in average 150 mSv per year (Cucinotta et al. (2001) Radiat Res. 156:460-466). Radiation-induced lens opacification is assumed to initiate from post irradiation proliferative activity of genetically damaged lens epithelial cells with alterations in cell cycle control, apoptosis, differentiation, and cellular disorganization, or other pathways controlling lens fiber cells' differentiation. As the porcine eye lens is similar to the human lens in size and anatomy, DNA damage response was investigated in ex-vivo porcine lenses in organ culture, in in-vitro cultivated lens epithelial slabs (ES) and in porcine lens epithelial cells (pLEC). Cell survival of proliferative cells was calculated from colony forming ability (CFA) assay. The phosphorylated form of H2AX (γ H2AX) was used as a molecular marker to visualize DNA double strand breaks (DSB) and their repair. Propidium iodide based DNA staining for cellular DNA content marked radiation-induced cell cycle disturbances. In pLEC the cell survival curve of immediate plated cells and after a recovery period of 24 h follow the equation $S=1.40xD+\ln 1.47$ and $S=1.59xD+\ln 1.79$, respectively. DNA DSB are induced in a dose-dependent manner (18 DSB/cell/Gy) and repaired during successive recovery (5 DSB/cell/Gy residual damage after 24 h). For doses ≥ 2 Gy a cell cycle arrest in G2 phase occurred 24 h after X-irradiation and persisted up to 72 h post-irradiation. DNA DSB induction and repair could as well be documented for ES and whole lenses after X-irradiation. In whole lenses, the amount of residual damage (after 24 h and 48 h) was highest in the equatorial zone while in the central epithelial zone DSB repair seemed to proceed with time in a manner comparable to in-vitro cultivated pLEC. Lens organ culture allows cellular metabolism and DNA synthesis in whole lenses. Repair of DNA DSB takes place in the central epithelial layer and is reduced in the equatorial region of cultivated lenses.