



Proceedings of the 7th Nordic Feed Science Conference, Uppsala, Sweden

**Institutionen för husdjurens
utfodring och vård**

**Swedish University of Agricultural Sciences
Department of Animal Nutrition and Management**

**Rapport 293
Report**

Uppsala 2016

ISSN 0347-9838
ISRN SLU-HUV-R-293-SE

Published by:

Organising committee of the 7th Nordic Feed Science Conference
Department of Animal Nutrition and Management
Swedish University of Agricultural Sciences (SLU)
SE- 753 23 Uppsala, Sweden

Copyright © 2016 SLU

All rights reserved. Nothing from this publication may be reproduced, stored in computerised systems or published in any form or any manner, including electronic, mechanical, reprographic or photographic, without prior written permission from the publisher (SLU).

The individual contributions in this publication and any liabilities arising from them remain the responsibility of the authors.

Organising Committee:

Peter Udén
Rolf Spörndly
Bengt-Ove Rustas
Torsten Eriksson

Edited by:

P. Udén
T. Eriksson
B-O. Rustas
R. Danielsson

Printed by:

SLU Repro
SE-75007 Uppsala, Sweden

Distributed by:

Department of Animal Nutrition and Management,
Box 7024,
SE-75323 Uppsala, Sweden

www.slu.se/husdjur-utfodring-vara

A note on sample preparation in the analysis of nitrate and nitrite in forage

R. Spörndly, T. Grönqvist, M. Knicky & B. Ericson

Swedish University of Agricultural Sciences (SLU), Department of Animal Nutrition & Management, Feed Science Division, Box 7024, 750 07 Uppsala.

Correspondence: Rolf.Spörndly@slu.se

Introduction

Nitrate content in forage is of interest as it pose a risk of poisoning for ruminants at levels above 1000 mg NO₃-N per kg DM (Strickland et al, 1995; Undersander et al, 1999). Analysing nitrate in forages by colorimetric methods have been questioned during decades due to interference of coloured substances in the forage (Wiseman & Jacobson, 1962; Wegner, 1972; Anderson & Case, 1999). Nevertheless, the colorimetric method using flow injection analysis (FIA) with a Cu-Cd reduction column has become a standard method in analysing nitrite and nitrate after reduction to nitrite (MacKown & Weik, 2004). The problem of diverging results in nitrate analysis has often appeared as a result of different nitrate extraction techniques, as opposed to differences due to detection methods after extraction (Anderson & Case, 1999). The sample preparation and analytical method in use at the departments of Soil and Environment, Crop Production Ecology and Animal Nutrition and Management at the Swedish University of Agricultural Science has been water extraction of dried samples followed by the nitrite and nitrate analysis according to the ISO method 13395;1996. As nitrate, and in particular nitrite, are unstable compounds, doubts have been raised concerning the influence of the sample preparation step. The present study was made with the aim to compare result of nitrite and nitrate analyses of dried or un-dried samples and extraction with boiling water or room tempered water.

Materials and Methods

Samples of fresh mixed grass-red clover forage were treated as in Figure 1.




Figure 1. Set up of experiment to evaluate the effect of drying and extraction on the level of nitrite and nitrate content in a grass- clover forage crop.

Two kg of the fresh forage crop was wilted to 50% DM and divided in two 1 kg portions. To one portion, 3 ml Safesil[®] (equivalent to 30.435 mg NO₂-N/kg) was added in the form of an aerosol to cover the entire forage surface and mixed inside an inflated plastic bag. Both portions were frozen to -20°C and ground frozen in a meat grinder and thoroughly blended.

These two portions, one blank and one nitrite spiked, were each of them treated as follows. One half was dried at 60°C for 16 h and two 1-g sub-samples were mixed with 100 ml distilled water each. The other half was not dried and 2 samples, equivalent in size on DM basis to the dried samples, were mixed with 100 ml distilled water each. Extraction was performed as follows: one of each parallel water suspension was boiled for 10 minutes and one was shaken for 1 h at room temperature after which they were diluted with distilled water to 250 ml and filtered through filter paper (Whatman 602H¹/₂).

After extraction, all samples were analysed without delay according to the procedure prescribed in the ISO method 13395:1996 as follows: liquid extracts were analysed for nitrite-N by a computerized flow injection analysis (FIA) system (FIAstar™ 5000 Analyzer with the PC software 'soFIA' from FOSS, www.foss.dk). The colorimetric detection was based on nitrite with sulfonilamide forming an azo dye product with N-(1-naphthyl)ethylenediamine (Griess reagent). Absorbance was determined at 540 nm. Nitrate was then reduced to nitrite in a copper-cadmium column and nitrite-N was analysed again. Nitrate was then calculated by difference. Absorbance for all samples was also determined without reagents and deducted from the gross determinations of nitrite. Finally, the nitrate- and nitrite-N were calculated to as per mg/kg DM.

Results were processed by the GLM procedure by SAS using nitrite-treatment, preparation method and extraction method as class variables (SAS ver. 9.4. SAS Institute Inc., Cary, NC, USA).

Results and Discussion

The fresh crop selected was very low in nitrate and nitrite. The values were all very close to the detection limit and are presented in Table 1. Nitrite treatment increased both nitrate and nitrite levels ($p < 0.0028$). No difference could be seen between extraction methods ($p > 0.8156$). However, preparation method made a difference ($p < 0.0035$), as drying the samples resulted in considerably lower nitrate and nitrite levels compared to no drying. Sample drying seemed to reduce nitrate content with by on average 36% and the nitrite content by 90%.

Table 1 Nitrate and nitrite in fresh crop samples with or without added NO₂, with or without drying and extraction in boiling or room temperate water (average ± SEM of two replicates)

Sample	Extraction	NO ₃ -N, mg/kg DM		NO ₂ -N, mg /kg DM	
		Dried	Un-dried	Dried	Un-dried
Fresh crop	Boiled	61 ± 1.0	61 ± 1.0	7 ± 0.5	10 ± 0.5
Fresh crop	Room temp	60 ± 1.0	59 ± 0.5	3 ± 0.0	4 ± 0.0
Fresh crop + NO ₂	Boiled	72 ± 2.0	181 ± 2.0	11 ± 0.0	80 ± 1.5
Fresh crop + NO ₂	Room temp	68 ± 0.0	188 ± 2.0	5 ± 0.0	85 ± 2.5

The low basal nitrate and nitrite content of the crop was not, or only slightly, affected by the drying process. For samples where NaNO₂ was added, the drying process reduced the nitrate and nitrite content ($p < 0.05$).

After evaluation of the effect on fresh crop similar analyses continued with silages. As the extraction method did not affect the result of nitrate and nitrite analyses, it was decided only to evaluate the effect of sample preparation (dried or un-dried) on recovery of nitrate and nitrite in silages. The result from 4 silage samples from the Czech Republic and 7 silages from the USA are presented in Table 2.

Table 2. Nitrate and nitrite in silage samples prepared with or without drying. All samples were extracted in boiling water

	NO ₃ -N, mg/kg DM		NO ₂ -N, mg/kg DM	
	Dried	Un-dried	Dried	Un-dried
Silage Czech republic 1	1155	1134	0.7	0.5
Silage Czech republic 2	892	908	0.7	0.5
Silage Czech republic 3	849	827	0.7	0.6
Silage Czech republic 4	123	95	0.7	0.6
Silage USA 1	182	191	0.7	1.6
Silage USA 2	19	25	3.9	5.6
Silage USA 3	237	270	5.1	9.6
Silage USA 4	253	246	5.6	10.1
Silage USA 5	11	14	4.2	5.5
Silage USA 6	21	32	5.9	8.9
Silage USA 7	23	26	7.1	8.8

No clear difference between dried and non-dried samples in nitrate and nitrite results can be seen in the Czech Republic samples. These samples were maize silages. In the samples from USA, which were alfalfa silages, a tendency for 7% lower nitrate values of the dried compared to un-dried samples were seen. Nitrite was affected to a greater extent by drying showing a 34% reduction.

In this experiment, nitrate and nitrite analysis according to ISO 13395:1996 was performed immediately upon extraction. In a preliminary investigation, we observed that freezing to -18°C, followed by thawing, decreased both nitrate and nitrite content in the sample. This observation, together with the main results in this paper suggest that further detailed studies are needed in order to establish complete recommendation of sample preparation for nitrate and nitrite analysis.

Conclusions

No differences in the nitrate and nitrite levels of fresh forage were observed if the extraction step was made by boiling the sample in water for 10 minutes or shaking in water at room temperature for 1 hour. Effects of drying the samples prior to extraction were however noticed. When NaNO₂ was added to the fresh crop, drying at 60°C for 16 h resulted in nitrate loss of 36% and nitrite loss of 90%. In preparation of silage samples, the drying process resulted in a 13% loss of nitrate and 63% loss of nitrite in some maize and alfalfa silages. It is

recommended not to use drying during sample preparation for nitrate and nitrite analyses. Instead extractions of fresh or frozen samples should be made.

References

- Anderson, K.A. & Case, T.E., 1999. Evaluation of plant nitrate extraction techniques and effect on commonly used analytical methods of detection. *Comm. Soil Sci. Plant Anal.* 30, 1470-1495.
- MacKown, C.T. & Weik, J.C., 2004. Comparison of laboratory and quick-test methods for forage nitrate. *Crop Sci.* 44, 218-226
- Strickland, G., Selk, G., Allen, E & Thedford, T., 1995. Nitrate toxicity in livestock. F-2903. Oklahoma Coop. Ext. Service, Stillwater, OK, USA.
- Undersander, D., Combs, D., Howard, T., Shaver, R., Siemens, M. & Thomas, D., 1999. Nitrate poisoning in cattle, sheep and goats. Univ. Wisconsin Coop. Ext., USA.
- Wiseman, H.G. & Jacobson, W.C., 1965. Determination of nitrate in silages and forages. *J. Agr. Food Chem.* 13, 36-39.
- Wegener, T.N., 1972. Simple and sensitive procedure for determining nitrate and nitrite in mixtures in biological fluids. *J. Dairy Sci.* 55, 642-644.

DISTRIBUTION:
Sveriges Lantbruksuniversitet
Institutionen för husdjurens utfodring och vård
Box 7024
750 07 UPPSALA
Tel.018/672817
Margareta.Norinder@slu.se