

## ORIGINAL ARTICLE

## Airborne cultivable microflora and microbial transfer in farm buildings and rural dwellings

Anne-Cécile Normand,<sup>1</sup> Bertrand Sudre,<sup>2</sup> Mallory Vacheyrou,<sup>2</sup> Martin Depner,<sup>3</sup> Inge M Wouters,<sup>4</sup> Ilka Noss,<sup>4</sup> Dick Heederik,<sup>4</sup> Anne Hyvärinen,<sup>5</sup> Jon Genuneit,<sup>6</sup> Charlotte Braun-Fahrländer,<sup>7</sup> Erika von Mutius,<sup>3</sup> Renaud Piarroux,<sup>1</sup> the GABRIEL-A Study Group

<sup>1</sup>Department of Parasitology and Mycology, Assistance Publique-Hôpitaux de Marseille, Marseille, France

<sup>2</sup>Franche-Comté University, UMR-CNRS 6249

Chrono-Environnement, Mycology Department, Besançon, France

<sup>3</sup>Children's Hospital of the University of Munich, Lindwurmstrasse 4, Munich, Germany

<sup>4</sup>Institute for Risk Assessment Sciences, Division of Environmental Epidemiology, Utrecht University, TD Utrecht, The Netherlands

<sup>5</sup>National Institute for Health and Welfare, Department of Environmental Health, Kuopio, Finland

<sup>6</sup>Institute of Epidemiology, University of Ulm, Ulm, Germany

<sup>7</sup>Swiss Tropical and Public Health Institute, Basel, Switzerland

**Correspondence to**

Anne-Cécile Normand, Laboratoire de Parasitologie/Mycologie, Hôpital de la Timone, 264 rue Saint-Pierre, 13385 Marseille cedex 05, France; [acecilenor@hotmail.com](mailto:acecilenor@hotmail.com)

The GABRIEL-A Study Group consists of the following members of GABRIEL-A study centres: Austria: Katalin Kovacs, Bernhard Morass, Elisabeth Horak; Baden-Württemberg (Germany): Gisela Büchele, Nikolaos Sitaridis, Jon Genuneit; Bavaria (Germany): Juliane Weber, Erika von Mutius; Poland: Anna Debinska, Hanna Danielewicz, Barbara Sozanska; Switzerland: Marco Waser, Charlotte Braun-Fahrländer.

Accepted 2 March 2011  
Published Online First  
24 March 2011

**ABSTRACT**

**Objectives** Exposure to environments rich in microorganisms such as farms has been shown to protect against the development of childhood asthma and allergies. However, it remains unclear where, and how, farm and other rural children are exposed to microbes. Furthermore, the composition of the microbial flora is poorly characterised. We tested the hypothesis that farm children are exposed indoors to substantial levels of viable microbes originating from animal sheds and barns. We also expected that environmental microbial flora on farms and in farm homes would be more complex than in the homes of rural control children.

**Methods** Dust samples were collected using passive samplers in the bedrooms of the following groups of children in rural Bavaria, Germany: (i) those living on farms (n=144), (ii) those regularly exposed to farm environments but not living on farms (n=149) and (iii) those never visiting farms (n=150). For farm children, additional samples were collected in animal sheds and barns. All samples were subjected to fungal and bacterial culturing.

**Results** Detectable levels of microorganisms were more often found in samples taken from farm dwellings than from other homes. Farm dwellings also showed higher microbial levels. Microbial counts of farm dwelling samples correlated with the counts in corresponding animal sheds and barns.

**Conclusions** Microorganisms are transported from animal sheds and barns into farm dwellings. Therefore, children living in these environments are exposed when indoors and when visiting animal sheds and barns. Indoor exposure may also contribute to the protective effect of the farm environment.

**INTRODUCTION**

Exposure to environments rich in microorganisms such as farms has previously been shown to protect against the development of childhood asthma and allergies.<sup>1–7</sup> In particular, frequent contact with farm animals during early childhood seems to be an important determining factor. In farm dwellings, environmental studies have shown increased levels of microbe associated molecular patterns such as endotoxin from Gram-negative bacteria, muramic acid from Gram-positive bacteria, and  $\beta(1-3)$ -D-glucans and extracellular polysaccharides from fungi, as compared to rural non-farm dwellings.

**What this paper adds**

- ▶ The farm environment has been shown to offer protection against asthma and allergies in children.
- ▶ The role of microorganisms has been recognised, but the microbial flora to which farm children are exposed is poorly characterised.
- ▶ This study shows that children living on farms are exposed to many microorganisms and that numerous fungal and bacterial taxa found in farm dwellings are also found in animal sheds and barns.
- ▶ Fewer microorganisms are found in the rural dwellings of non-farmers.

However, little is known about the complex microbial flora which underlies exposure to these microbial markers. We hypothesised that children were exposed to substantial levels of viable microbes not only in animal sheds and barns, but also indoors in the farm dwelling. We also hypothesised that the indoor levels and spectra of microorganisms were determined by the microorganisms present in animal sheds and barns.

Air in animal sheds is highly contaminated with numerous species of moulds and bacteria, including high levels of actinomycetes.<sup>8–11</sup> Individuals working on a farm may transport microorganisms from the animal shed to the dwelling.<sup>12 13</sup> However, the microbial flora to which farm children may be exposed is poorly characterised.

The objectives of the present study were therefore to determine whether the microbes children were exposed to indoors in farm environments originated from animal sheds and barns, and to measure the levels of fungi and bacteria in farm and control rural dwellings. We therefore studied: (i) the microbial flora in the dwellings (bedrooms) of three groups of children living in a rural environment in the German Alpine region: one group living on farms, one group occasionally exposed to farm environments, and one group with no exposure to a farm environment; (ii) the microbial flora collected in the animal sheds and barns of farm dwellings; and (iii) the relationship between the

level of contamination in the farm dwelling and in associated animal sheds and barns.

This project is part of the GABRIEL Advanced Studies, which were designed to identify environmental microbial exposures in farm environments and determine the exposures protecting against the development of asthma and allergic diseases.

## MATERIALS AND METHODS

### Study design


This environmental study is nested in phase III of the GABRIEL Advanced Studies (figure 1). The farms included in this study were mixed traditional family farms, with mainly cows and some small animals such as chickens; some farms had modest numbers of pigs. Crops produced were often grown as animal feed.<sup>14</sup> Three non-overlapping strata were defined: (i) farm children, that is children living on a farm run by their family at the time of assessment; (ii) exposed non-farm children, that is non-farm children regularly visiting animal sheds, barns or consuming cows milk produced on a farm; and (iii) non-exposed rural non-farm children. From the children participating in all phase II study modules, a stratified random sample of 895 subjects using nine dwelling strata (three for health status (those with asthma, those with atopy and controls) and three

for the exposure (non-exposed children, exposed children and farm children)) was invited to participate in phase III in which we performed in-depth environmental sampling. For the nested study presented here, 444 subjects from phase III were selected (approximately 50 subjects per stratum) and data were obtained for 443. For farm children, sampling involved dwellings and associated barns and animal sheds.

### Airborne dust samples

Airborne dust samples were collected by a passive collection method using electrostatic dust fall collectors (EDCs). Details about the use of EDCs in indoor and animal shed environments have been described elsewhere as well as the results of validation pilot studies.<sup>15 16</sup> Briefly, EDCs are plastic sample holders equipped with between two and four electrostatic cloths with a cloth exposure area of 0.0209 m<sup>2</sup> (Zeeman, Utrecht, The Netherlands). When exposed to air, EDCs capture settling dust due to their electrostatic properties. EDCs have been validated against active dust PM<sub>10</sub> sampling methods such as the Harvard impactor.<sup>16</sup> Within each dwelling, one EDC was placed in the child's bedroom and also, if applicable, in whichever barn and/or animal shed in which the child spent most of his or her time, resulting in the collection of 150 EDCs from the non-exposed

**Figure 1** Flow chart of the GABRIEL Advanced Studies in Bavaria, Germany. Samples from 443 dwellings (bedrooms) were included in the current analyses from among the 895 children invited for in-depth exposure assessment within phase III of the GABRIEL Advanced Studies. For all participants, dust was collected in the child's room. For farm children, dust was additionally collected on the farm or the animal shed wherever possible. EDC, electrostatic dust fall collector.



children's bedrooms, 149 from exposed children's bedrooms, 144 from farm children's bedrooms, 111 from animal sheds and 126 from barns. EDCs were placed by a fieldwork team member. After exposure to air for 14 days, the families were asked to close the EDCs and returned them to the study centre in Munich by post, after which they were sent to Utrecht for further processing. The four electrostatic cloths were transferred individually into plastic bags and stored frozen: three cloths were stored at  $-80^{\circ}\text{C}$  and one cloth was stored at  $-20^{\circ}\text{C}$ . Samples stored at  $-20^{\circ}\text{C}$  were used for the analyses described in this manuscript and were sent to the microbiology laboratory in Besançon, France, to be analysed for microbial composition within 4–8 months. Sampling was performed during the 2007–2008 winter, as high humidity levels were conducive to the development of fungi. A total of 680 EDCs were available for analyses.

### Treatment of EDC cloths for microbiological analyses

Microbiological analyses were performed blindly, with the operators unaware of the characteristics of each sample. The cloths were put in a sterile bag with 20 ml of 0.1% Tween 80 and shaken for 10 min in a Stomacher (AES Laboratoire, Combourg, France). Samples of the washing solution were added to different growth media suitable for a range of microorganisms, and the analyses were performed as described below.

### Culturing

Aliquots (100  $\mu\text{l}$ ) of the cloth washing solution were spread on five different agars: (i) Dichloran-Glycerol 18 (Oxoid, Basingstoke, England) with 0.5% chloramphenicol (Sigma-Aldrich, Steinheim, Germany) incubated at  $30^{\circ}\text{C}$  for mesophilic mould isolation, (ii) 3% malt-agar (AES, Bruz, France) with 10% salt and 0.5% chloramphenicol, incubated at room temperature for

osmophilic fungal species, (iii) Difco actinomycetes isolation agar (Becton Dickinson, Le Pont de Claix, France) incubated at  $30^{\circ}\text{C}$  for mesophilic actinomycetes, (iv) R8 medium incubated at  $52^{\circ}\text{C}$  for thermophilic actinomycetes,<sup>17</sup> and (v) CHROMagar Orientation (CHROMagar, Paris, France) incubated at room temperature for bacterial growth and phenotypic differentiation. Because the different samples were blindly analysed and microbial contamination levels were unknown, four dilutions (pure, 10, 100 and 1000-fold dilutions) were used for the analysis of bacteria on the CHROMagar Orientation medium. For fungi and actinomycetes, three dilutions (pure, 10 and 100-fold dilutions) were systematically plated for counting and identification purposes. Microbial counts were then performed on those plates with the more concentrated inoculums that had sufficient colonies.

### Fungal identification

After 7 days of incubation, the colonies were counted and identified using macroscopic and microscopic criteria, following the keys in the *Atlas of Clinical Fungi*.<sup>18</sup> Slow-growing fungi were identified 14 days after incubation, using the same methods. Forty-four fungal species were identified, and taxa were built for use in further medical analyses. We particularly focused on genera and species commonly found in the farm environment and suspected of provoking allergic or pulmonary diseases among farmers, namely *Absidia* spp. (farmer's lung diseases), *Alternaria* spp. (allergy), *Aspergillus* spp. and more particularly *Aspergillus fumigatus* (asthma) and *Aspergillus glaucus* (farmer's lung diseases), *Cladosporium* spp. (allergy), *Penicillium* spp. (farmer's lung diseases and allergy), *Scopulariopsis* sp. (allergy) and *Wallemia sebi* (farmer's lung diseases).<sup>11 19 20</sup> Other fungi were classified into yeasts, other *Mucorales* species and unidentified moulds.

**Table 1** Number and frequency of detected microorganisms, stratified according to exposure strata and location on the farm

	Non-exposed non-farm children dwellings		Exposed non-farm children dwellings		Farm children dwellings		Animal sheds		Barns		Farm vs non-exposed non-farm children dwellings $\chi^2$	Farm vs exposed non-farm children dwellings $\chi^2$
	N	%	N	%	N	%	N	%	N	%		
<i>Absidia</i> spp.	5	3.3	7	4.7	35	24.3	61	55.0	67	53.2	27.5***	22.9***
Other <i>Mucorales</i> spp.	12	8.0	11	7.4	31	21.5	48	43.2	45	35.7	10.8**	11.9**
<i>Aspergillus fumigatus</i>	32	21.3	40	26.9	55	38.2	74	66.7	88	69.8	10.0**	4.3*
Identified <i>Eurotium</i> spp.†	27	18.0	40	26.9	95	66.0	72	64.9	75	59.5	69.6***	45.1***
Other <i>Eurotium</i> spp.	21	14.0	22	14.8	51	35.4	51	46.0	53	42.1	18.2***	16.7***
Other <i>Aspergillus</i> spp.	57	38.0	66	44.3	95	66.0	75	67.6	87	69.1	23.0***	13.9***
<i>Alternaria</i> spp.	8	5.3	11	7.4	19	13.2	35	31.5	44	34.9	5.4*	2.7 (NS)
<i>Cladosporium</i> spp.	77	51.3	73	49.0	116	80.6	82	73.9	104	82.5	27.8***	31.9***
<i>Penicillium</i> spp.	113	75.3	113	75.8	127	88.2	80	72.1	73	57.9	8.1**	7.5**
<i>Scopulariopsis</i> sp.	5	3.3	10	6.7	41	28.5	66	59.5	69	54.8	35.2***	24.1***
<i>Wallemia sebi</i>	18	12.0	27	18.1	69	47.9	50	45.1	78	61.9	45.5***	29.5***
Yeast	12	8.0	19	12.8	44	30.6	72	64.9	48	38.1	24.2***	13.8***
Other fungi	44	29.3	46	30.9	66	45.8	72	64.9	78	61.9	8.5**	6.9**
Mesophilic <i>Actinomycetaceae</i>	38	25.3	45	30.2	111	77.1	109	98.2	117	92.9	78.7***	64.7***
Thermophilic <i>Actinomycetaceae</i>	8	5.3	27	18.1	45	31.3	79	71.2	71	56.4	33.4***	6.8**
Gram negative rods	79	52.7	87	58.4	121	84.0	95	85.6	112	88.9	33.2***	23.4***
Gram positive rods	89	59.3	91	61.1	132	91.7	90	81.1	111	88.1	41.2***	37.7***
Gram negative cocci	41	27.3	44	29.5	67	46.5	38	34.2	30	23.8	11.6**	9.0**
Gram positive cocci	106	70.7	95	63.8	133	92.4	94	84.7	109	86.5	22.7***	34.7***

$\chi^2$  Test statistics representing differences in frequencies between exposure strata (farm vs exposed and farm vs non-exposed) are given.

\* $p < 0.05$ ; \*\* $p < 0.001$ ; \*\*\* $p < 0.0001$ .

† Identified *Eurotium* spp. and their teleomorphs: *Aspergillus hollandicus*, *Aspergillus glaucus*, *Eurotium amstelodami* and *Eurotium herbariorum*.

N, number of positive samples; NS, non-significant.

**Bacterial identification**

Bacterial colonies were selected from CHROMagar Orientation medium according to colony morphotype, subjected to Gram staining and classified. We classified and counted the actinomycetes cultured on Difco and R8 media according to growth temperature. We presented results of fungal identification according to bacterial taxa. We particularly focused on actinomycetes counts because these bacteria are known to be present in high amounts in the farm environment and to be responsible for immuno-allergic diseases such as farmer's lung disease.<sup>11 19</sup> Identification of each bacterial colony at the genus or species level will be continued in a further project on bacterial environmental characterisation as a specimen of every colony was re-isolated by sub-culturing and frozen at  $-80^{\circ}\text{C}$  for analyses. Bacterial and fungal results were expressed in colony forming units (CFU) per plate and further converted to CFU per EDC and  $\text{CFU}/\text{m}^2$  taking into account the dilution steps and the exposure surface area of the EDC ( $0.0209 \text{ m}^2$ ).

**Statistical analyses**

Due to the procedure used, levels below 200 CFU/EDC could not be detected. For calculation of the arithmetical mean, zero values were kept as such. For calculation of the geometric mean, non-detects were replaced by half of the limit of detection ( $1/2 \times 200/0.0209 \text{ CFU}/\text{m}^2 = 4784.69 \text{ CFU}/\text{m}^2$ ). The geometric means and 95% CIs of each microbial category stratified by exposure (non-exposed non-farm children, exposed non-farm children, farm children) and farm location (farm dwelling, animal shed, barn) were calculated. To test for differences in the frequencies of detectable microorganisms between exposure strata, a Pearson's  $\chi^2$  test was used. Spearman's  $r$  was used to calculate

correlations between the corresponding viable microorganisms in dwellings and barns ( $n=125$  matched pairs) and between dwellings and animal sheds ( $n=110$  matched pairs). Statistical significance was set at the 5% level. STATA 10 SE was used for statistical analyses.

**RESULTS**

Numerous fungal and bacterial taxa were detected from the 680 analysed EDCs. However, only one fungal and three bacterial taxa, that is *Penicillium* spp., Gram negative rods, Gram positive rods and Gram positive cocci, were found in more than 50% of samples collected in the different environments (non-exposed dwelling, exposed dwelling, farm dwelling, animal shed and barn). There was a marked trend towards a higher proportion of samples with detectable levels of microorganisms by taxa collected from the dwellings of the farm children than from the dwellings of the non-exposed and exposed children (table 1).

In particular, the proportion of positive samples for each bacterial and fungal taxon was significantly higher in farm dwellings than in exposed or non-exposed dwellings. There was a trend for a higher proportion of detectable fungi and bacteria in exposed dwellings compared to non-exposed dwellings, but the difference was only significant for the thermophilic Actinomycetaceae group (data not shown).

The higher exposure in farm dwellings was evidenced by the proportion of detectable samples and by increased concentrations. As shown in table 2, the geometric mean concentration of detectable samples for every microbial parameter was higher in farm dwellings than in the two other types of dwellings, with the exception of *Scopulariopsis* sp. Moreover, for most taxa, the

**Table 2** Geometric means and GSDs of microorganisms (in  $\text{CFU}/\text{m}^2$ ) by exposure strata and by location on the farm, and Spearman correlation coefficients between microorganism levels in the farm dwelling and in the animal shed or barn

	Non-exposed non-farm dwelling		Exposed non-farm dwelling		Farm dwelling		Animal shed		Barn		Farm dwelling/animal shed (n = 110)		Farm dwelling/barn (n = 125)	
	GM/1000	GSD	GM/1000	GSD	GM/1000	GSD	GM/1000	GSD	GM/1000	GSD	Spearman's r	p Value	Spearman's r	p Value
<i>Absidia</i> spp.	5	0.1	5	0.1	9	1.6	48	5.7	38	6.5	0	***	0	***
Other <i>Mucorales</i> spp.	5	0.1	5	0.3	7	0.9	19	3.4	15	3.6	0	NS	0	*
<i>Aspergillus fumigatus</i>	6	0.6	8	1.2	10	1.9	55	4.1	34	6.0	0	***	0	***
Identified <i>Eurotium</i> spp. †	6	0.6	7	0.8	23	2.6	187	8.7	110	9.8	0	***	0	***
Other <i>Eurotium</i> spp.	6	0.4	6	0.5	11	1.7	69	7.3	39	9.5	0	***	0	NS
Other <i>Aspergillus</i> spp.	9	1.2	10	1.4	20	2.1	131	5.7	78	7.6	0	***	0	***
<i>Alternaria</i> spp.	5	0.1	5	0.1	6	0.3	11	2.2	12	2.5	0	*	0	NS
<i>Cladosporium</i> spp.	10	1.1	9	0.8	23	1.6	189	5.7	316	6.4	0	*	0	NS
<i>Penicillium</i> spp.	22	1.8	28	2.1	48	2.0	122	5.6	57	5.3	0	*	0	*
<i>Scopulariopsis</i> sp.	5	0.0	6	0.4	7	0.8	59	4.3	30	6.6	0.26	**	0.28	**
<i>Wallemia sebi</i>	6	0.3	7	0.8	14	2.0	51	7.6	108	8.1	0	**	0	**
Yeast	5	0.1	6	0.4	10	1.7	126	7.2	34	7.5	0	*	0	*
Other fungi	8	0.8	8	0.9	11	1.6	77	5.9	67	5.8	0	*	0	NS
Mesophilic <i>Actinomycetaceae</i>	6	0.4	7	0.8	41	3.4	1682	5.3	477	4.1	0	***	0	***
Thermophilic <i>Actinomycetaceae</i>	5	0.1	6	0.5	9	1.7	71	4.8	30	5.7	0	***	0	*
Gram negative rods	15	2.0	17	2.7	122	5.2	7022	8.6	2258	12.3	0	**	0	***
Gram positive rods	13	1.4	17	2.9	246	5.3	5604	9.3	2792	15.6	0	**	0	***
Gram negative cocci	9	1.5	9	1.8	21	4.9	68	4.6	15	14.5	0	NS	0	NS
Gram positive cocci	27	3.7	29	4.7	454	6.0	25368	10.0	1841	16.6	0	***	0	***

\* $p < 0.05$ ; \*\* $p < 0.001$ ; \*\*\* $p < 0.0001$ .

† Identified *Eurotium* spp. and their teleomorphs: *Aspergillus hollandicus*, *Aspergillus glaucus*, *Eurotium amstelodami* and *Eurotium herbariorum*.

Mean values are expressed in colony forming unit per square metre ( $\text{CFU}/\text{m}^2$ ). Non-detects have been replaced by half of the limit of detection for GM calculations.

Spearman's  $r$  is shown for farm children for whom electrostatic dust fall collectors from either farm source were returned: child's room and animal shed ( $N=110$ ) or child's room and barn ( $N=125$ ).

GM, geometric mean; GSD, geometric SD; NS, non-significant.



**Table 3** Number of samples with or without specified microorganisms identified in farm dwellings and/or other indoor farm source (animal shed or barn)

	Present in farm dwelling/animal shed or barn				Proportion of +/+ of all farm dwelling positives	Proportion of -/+ of all farm dwelling negatives
	-/-	+/-	-/+	+/+		
<i>Absidia</i> spp.	30	0	39	29	100%	57%
Other <i>Mucorales</i> spp.	36	8	39	15	65%	52%
<i>Aspergillus fumigatus</i>	16	3	46	33	92%	74%
Identified <i>Eurotium</i> spp.*	8	9	26	55	86%	76%
Other <i>Eurotium</i> spp.	25	8	40	25	76%	62%
Other <i>Aspergillus</i> spp.	8	7	26	57	89%	76%
<i>Alternaria</i> spp.	42	3	45	8	73%	52%
<i>Cladosporium</i> spp.	0	4	19	75	95%	100%
<i>Penicillium</i> spp.	1	12	10	75	86%	91%
<i>Scopulariopsis</i> sp.	19	5	49	25	83%	72%
<i>Wallemia sebi</i>	16	8	37	37	82%	70%
Yeast	24	0	42	32	100%	64%
Other fungi	11	5	40	42	89%	78%
Mesophilic <i>Actinomycetaceae</i>	0	0	17	81	100%	100%
Thermophilic <i>Actinomycetaceae</i>	11	2	52	33	94%	83%
Gram negative rods	0	1	17	80	99%	100%
Gram positive rods	2	3	8	85	97%	80%
Gram negative cocci	29	19	24	26	58%	45%
Gram positive cocci	2	3	6	87	97%	75%

N=98 farm children with samples from the child's room, animal shed and barn included.

\*Identified *Eurotium* spp. and their teleomorphs: *Aspergillus hollandicus*, *Aspergillus glaucus*, *Eurotium amstelodami* and *Eurotium herbararium*.

microbial concentrations in the farm dwellings were significantly and positively correlated with the concentrations in the corresponding animal sheds and barns.

The fungal patterns of non-exposed and exposed dwellings were very similar and characterised by the dominant presence of *Penicillium* spp., whereas the microflora from the farm dwellings were more similar to the animal shed microflora with *Eurotium* spp., other *Aspergillus* spp., *Penicillium* spp., *Cladosporium* spp. and Gram positive cocci identified as the dominant groups of microorganisms. Similarities were also found between the bacterial patterns of farm dwellings, animal sheds and barns, and bacterial contamination of non-exposed dwellings was very low compared to that of farm dwellings.

The similarity of the results between animal sheds, barns and farm dwellings is also corroborated by the findings presented in table 3 which show that in a majority of the farm dwellings (58% to 100% depending on the microorganism), a positive sample for a given microorganism in the dwelling corresponded to a positive sample in the animal shed or barn.

As shown in table 4, the concentrations of different microorganisms were lower in farm dwellings compared to animal sheds and barns, with ratios for the arithmetic mean varying from 1:6 (*Penicillium* spp.) to 1:64 (unidentified *Eurotium* spp.) for fungi, and from 1:11 (mesophilic actinomycetes) to 1:171 (Gram negative rods) for bacteria.

## DISCUSSION

Relationships between the presence of fungi in indoor and outdoor air have been studied previously, and the authors found that outdoor factors may influence the presence of viable mould in indoor environments.<sup>21</sup> We sought to describe the association between the patterns and levels of viable microorganisms on the farm and in the indoor environment of children by performing a large and extensive environmental study. The similarities in species found in farm dwellings and barns strongly suggest that microorganisms are transported from animal sheds and barns into farm dwellings. Indeed, there is a marked tendency towards

an increased number of microbial species and higher levels in farm dwellings compared to rural dwellings. Moreover, this strong indication of transport or flow from barns and animal sheds into dwellings clearly modifies the overall composition of airborne microflora in children's bedrooms. While the bedroom indoor environment of non-exposed children is characterised by a predominance of *Penicillium* spp., as previously reported,<sup>22 23</sup> other fungi such as *Aspergillus glaucus* and its teleomorph

**Table 4** Arithmetic means (in 10<sup>3</sup> CFU/m<sup>2</sup>) and ratio of arithmetic mean of level of microorganisms found in animal sheds and farm dwellings

	Arithmetic means (10 <sup>3</sup> CFU/m <sup>2</sup> )		Arithmetic mean ratio (animal shed/farm dwelling)
	Animal shed (N = 111)	Farm dwelling (N = 144)	
<i>Absidia</i> spp.	835	32	26
Other <i>Mucorales</i> spp.	145	14	10
<i>Aspergillus fumigatus</i>	1055	59	18
Identified <i>Eurotium</i> spp.*	4012	177	23
Other <i>Eurotium</i> spp.	3005	47	64
Other <i>Aspergillus</i> spp.	1724	75	23
<i>Alternaria</i> spp.	75	4	19
<i>Cladosporium</i> spp.	2183	51	43
<i>Penicillium</i> spp.	639	109	6
<i>Scopulariopsis</i> sp.	817	13	63
<i>Wallemia sebi</i>	1411	54	26
Yeast	1964	31	63
Other fungi	715	31	23
Mesophilic <i>Actinomycetaceae</i>	6994	624	11
Thermophilic <i>Actinomycetaceae</i>	1073	66	16
Gram negative rods	173 551	1013	171
Gram positive rods	279 564	3270	85
Gram negative cocci	15 733	431 †	37
Gram positive cocci	388 932	6334	61

Arithmetic mean ratio is the animal shed mean/farm dwelling mean.

\*Identified *Eurotium* spp. and their teleomorphs: *Aspergillus hollandicus*, *Aspergillus glaucus*, *Eurotium amstelodami* and *Eurotium herbararium*.

†One missing value.

*Eurotium amstelodami* are also common in farm dwellings. *A. glaucus* is commonly present on farms, especially when hay is distributed to cattle.<sup>24–26</sup> Other fungal species observed in the farm children's bedroom, such as *Absidia* spp. and *Wallemia sebi*, and bacteria such as actinomycetes species, have also been frequently detected in the farm environment<sup>11 27</sup> and are suspected of being involved in farmer's lung diseases. Moreover, our study showed that for numerous taxa, the concentrations of microorganisms in animal sheds and farm dwellings are positively and significantly correlated, corroborating the hypothesis of a noticeable transfer from the animal shed to the farm dwelling. For actinomycetes and most of the fungal groups, the ratio between arithmetic mean exposures in the dwelling compared to the animal shed is between 1:10 and 1:40. For these microorganisms, it can be extrapolated that when a child has spent 10–40 h in his or her bedroom, he or she has been in contact with the same number of microorganisms as if he or she had stayed 1 h in the animal shed. Similarly, bacterial contamination of the bedroom seems influenced by the environment in the animal sheds and barns. This may be due to the propensity of fungal and actinomycetes spores to be easily disseminated in the atmosphere<sup>12</sup> and carried into dwellings on the hands, clothes or shoes of farmers and their children.<sup>13 28 29</sup>

The findings for *Penicillium* spp. were somewhat different. On the one hand, the low arithmetic mean ratio between the air in farm dwellings and the air in animal sheds, and the positive correlation between the numbers of colonies found in these two locations, suggest that the indoor air in a farm dwelling is contaminated by *Penicillium* spp. spores originating from the farm. On the other hand, *Penicillium* species were also the predominant fungal flora in non-farming dwellings, which is concordant with previous studies showing that *Penicillium* species are among the most typical fungi found in the indoor air of homes.<sup>12 30 31</sup> Apart from the farm environment, there are likely to be other sources of *Penicillium* species, as highlighted by studies on mould contamination in indoor air.<sup>22 32–34</sup> More detailed identification of *Penicillium* species would have been of interest to determine if the patterns of individual species found in farm dwellings and non-exposed dwellings are different. However, the identification of *Penicillium* species is particularly difficult even in highly specialised laboratories, as further studies combining molecular, biochemical and physiological approaches are required.

In this study we focused on bacteria and fungi that were easily obtained using standard culture media. Furthermore, we targeted mesophilic and thermophilic fungi and actinomycetes because of their propensity to replicate in indoor environments.<sup>35</sup> We acknowledge that these microorganisms represent only a small fraction of all airborne microorganisms because numerous bacteria and fungi are difficult or even impossible to culture.<sup>35</sup> Other approaches such as microscopy or DNA cloning,<sup>36</sup> that bypass the culturing step, could also be used to assess the microbial diversity in farm dwellings. However, such non-culture based methods, especially the biomolecular approaches, are expensive and difficult to apply to large environmental studies with large series of samples. Both approaches are however complementary and one is not necessarily superior to the other. Nonetheless, the microbial analyses in this study appear to be relevant, since in a separate analysis we have shown that increased diversity of viable microorganisms enhances the protective effect conferred by animal shed fungal and bacterial flora on childhood asthma and atopy.<sup>14</sup>

In conclusion, this study is an important step in the understanding of microbial exposure in farm dwellings because it provides data suggesting transport of microbial species between

animal sheds and farm dwellings. This transport clearly determines children's exposure to microorganisms, as it modifies the composition of airborne microflora in the children's bedrooms.

**Acknowledgements** We thank Dr Wulf Thierfelder and Michael Thamm from the Robert-Koch-Institute, Berlin, Germany for their cooperation and measurements of total and specific IgE that were used in the definition of atopic subjects.

**Funding** This work was supported by the European Commission as part of GABRIEL (a multidisciplinary study to identify the genetic and environmental causes of asthma in the European Community), contract number 018996 under the FP6-LIFESCIHEALTH Integrated Program LSH-2004-1.2.5-1.

**Competing interest** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

1. Braun-Fahrlander C, Gassner M, Grize L, et al. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution. *Clin Exp Allergy* 1999;**29**:28–34.
2. Ernst P, Cormier Y. Relative scarcity of asthma and atopy among rural adolescents raised on a farm. *Am J Respir Crit Care Med* 2000;**161**:1563–6.
3. Kilpelainen M, Terho EO, Helenius H, et al. Farm environment in childhood prevents the development of allergies. *Clin Exp Allergy* 2000;**30**:201–8.
4. Klintberg B, Berglund N, Lijja G, et al. Fewer allergic respiratory disorders among farmers' children in a closed birth cohort from Sweden. *Eur Respir J* 2001;**17**:1151–7.
5. Riedler J, Eder W, Oberfeld G, et al. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy* 2000;**30**:194–200.
6. Von Ehrenstein OS, Von Mutius E, Illi S, et al. Reduced risk of hay fever and asthma among children of farmers. *Clin Exp Allergy* 2000;**30**:187–93.
7. Leynaert B, Neukirch C, Jarvis D, et al. Does living on a farm during childhood protect against asthma, allergic rhinitis, and atopy in adulthood? *Am J Respir Crit Care Med* 2001;**164**:1829–34.
8. American Thoracic Society MSoAIA. Respiratory health hazards in agriculture. *Am J Respir Crit Care Med* 1998;**158**:S1–76.
9. Hanhela R, Louhelainen K, Pasanen AL. Prevalence of microfungi in Finnish cow barns and some aspects of the occurrence of *Wallemia sebi* and *Fusaria*. *Scand J Work Environ Health* 1995;**21**:223–8.
10. Lange L, Thorne P, Kullman GJ. Determinants of culturable bioaerosol concentration in dairy barn. *Ann Agric Environ Med* 1997;**4**:187–94.
11. Reboux G, Piarroux R, Mauny F, et al. Role of molds in farmer's lung disease in Eastern France. *Am J Respir Crit Care Med* 2001;**163**:1534–9.
12. Pasanen AL, Kalliokoski P, Pasanen P, et al. Fungi carried from farmers' work into farm homes. *Am Ind Hyg Assoc J* 1989;**50**:631–3.
13. Lis DO, Mainelis G, Gorny RL. Microbial air contamination in farmhouses—quantitative aspects. *Clean* 2008;**36**:551–5.
14. Ege M, Mayer M, Normand AC, et al. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med* 2011;**364**:701–9.
15. Normand AC, Vacheyrou M, Sudre B, et al. Assessment of dust sampling methods for the study of cultivable-microorganism exposure in stables. *Appl Environ Microbiol* 2009;**75**:7617–23.
16. Noss I, Wouters IM, Visser M, et al. Evaluation of a low-cost electrostatic dust fall collector for indoor air endotoxin exposure assessment. *Appl Environ Microbiol* 2008;**74**:5621–7.
17. Amner W, Edwards C, McCarthy AJ. Improved medium for recovery and enumeration of the farmer's lung organism, *Saccharomonospora viridis*. *Appl Environ Microbiol* 1989;**55**:2669–74.
18. de Hoog GS, Guarro J, Gené J, et al. *Atlas of Clinical Fungi*. 2nd edn. Reus, Spain: Centraalbureau voor Schimmelcultures. Utrecht, The Netherlands / Universitat Rovira I Virgili, 2000:1–1007.
19. Roussel S, Reboux G, Dalphin JC, et al. Farmer's lung disease and microbiological composition of hay: a case-control study. *Mycopathologia* 2005;**160**:273–9.
20. Verhoeff AP, Burge HA. Health risk assessment of fungi in home environments. *Ann Allergy Asthma Immunol* 1997;**78**:544–54; quiz 555–6.
21. Verhoeff AP, van Wijnen JH, Brunekreef B, et al. Presence of viable mould propagules in indoor air in relation to house damp and outdoor air. *Allergy* 1992;**47**:83–91.
22. Reboux G, Bellanger AP, Roussel S, et al. Indoor mold concentration in Eastern France. *Indoor air* 2009;**19**:446–53.
23. Rosas I, Calderon C, Ulloa M, et al. Abundance of airborne *Penicillium* CFU in relation to urbanization in Mexico City. *Appl Environ Microbiol* 1993;**59**:2648–52.
24. Kotimaa MH, Terho EO, Husman K. Airborne moulds and actinomycetes in work environment of farmers. *Eur J Respir Dis Suppl* 1987;**152**:91–100.
25. Kotimaa MH, Oksanen L, Koskela P. Feeding and bedding materials as sources of microbial exposure on dairy farms. *Scand J Work Environ Health* 1991;**17**:117–22.
26. Reboux G, Reiman M, Roussel S, et al. Impact of agricultural practices on microbiology of hay, silage and flour on Finnish and French farms. *Ann Agric Environ Med* 2006;**13**:267–73.

27. **Roussel S**, Reboux G, Dalphin JC, *et al.* Microbiological evolution of hay and relapse in patients with farmer's lung. *Occup Environ Med* 2004;**61**:e3.
28. **Waser M**, Schierl R, von Mutius E, *et al.* Determinants of endotoxin levels in living environments of farmers' children and their peers from rural areas. *Clin Exp Allergy* 2004;**34**:389–97.
29. **Korthals M**, Ege MJ, Tebbe CC, *et al.* Application of PCR-SSCP for molecular epidemiological studies on the exposure of farm children to bacteria in environmental dust. *J Microbiol Methods* 2008;**73**:49–56.
30. **Hyvarinen A**, Reponen T, Husman T, *et al.* Comparison of the indoor air quality in mould damaged and reference buildings in a subarctic climate. *Cent Eur J Public Health* 2001;**9**:133–9.
31. **Ren P**, Jankun TM, Leaderer BP. Comparisons of seasonal fungal prevalence in indoor and outdoor air and in house dusts of dwellings in one Northeast American county. *J Expo Anal Environ Epidemiol* 1999;**9**:560–8.
32. **Roussel S**, Reboux G, Bellanger AP, *et al.* Characteristics of dwellings contaminated by moulds. *J Environ Monit* 2008;**10**:724–9.
33. **Gorny RL**, Dutkiewicz J. Bacterial and fungal aerosols in indoor environment in Central and Eastern European countries. *Ann Agric Environ Med* 2002;**9**:17–23.
34. **Pietarinen VM**, Rintala H, Hyvarinen A, *et al.* Quantitative PCR analysis of fungi and bacteria in building materials and comparison to culture-based analysis. *J Environ Monit* 2008;**10**:655–63.
35. **Eduard W**, Heederik D. Methods for quantitative assessment of airborne levels of noninfectious microorganisms in highly contaminated work environments. *Am Ind Hyg Assoc J* 1998;**59**:113–27.
36. **Pakarinen J**, Hyvarinen A, Salkinoja-Salonen M, *et al.* Predominance of Gram-positive bacteria in house dust in the low-allergy risk Russian Karelia. *Environ Microbiol* 2008;**10**:3317–25.



## Airborne cultivable microflora and microbial transfer in farm buildings and rural dwellings

Anne-Cécile Normand, Bertrand Sudre, Mallory Vacheyrou, Martin Depner, Inge M Wouters, Ilka Noss, Dick Heederik, Anne Hyvärinen, Jon Genuneit, Charlotte Braun-Fahrländer, Erika von Mutius, Renaud Piarroux and the GABRIEL-A Study Group

*Occup Environ Med* 2011 68: 849-855 originally published online March 24, 2011  
doi: 10.1136/oem.2010.061879

---

Updated information and services can be found at:  
<http://oem.bmj.com/content/68/11/849>

---

### References

*These include:*

This article cites 35 articles, 6 of which you can access for free at:  
<http://oem.bmj.com/content/68/11/849#BIBL>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>