

Letter Regarding: Proximal Gastrocnemius Release in the Treatment of Mechanical Metatarsalgia

Foot & Ankle International®
2016, Vol. 37(7) 790–791
© The Author(s) 2016
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1071100716654966
fai.sagepub.com

Dear Editor:

With great interest we read the article by Morales-Muñoz and colleagues.¹⁰ The authors reported a prospective case series of 78 patients suffering mechanical metatarsalgia due to isolated M. gastrocnemius tightness (MGT). All patients were treated by a proximal release of the medial gastrocnemius.³ At 6 months' follow-up, the mean AOFAS score was 84 points (preoperative: 47 points), with nearly 70% of patients being completely satisfied. No major complications were reported.

Morales-Muñoz et al¹⁰ defined impaired ankle dorsiflexion (ADF) as ADF values $\leq 0^\circ$ with the knee extended. Preoperative mean ADF values were -17.5° (range -30° to -5°). Previous studies defined impaired ADF between 0° and 12° with the knee extended.^{4,6,12} In their keystone paper, DiGiovanni et al⁴ conducted a prospective case-control study and reported average ADF values of 4.5° in their patient group (metatarsalgia or related midfoot/forefoot symptoms) and 13.1° in their healthy control group.

Despite similar patient cohorts, the mean ADF values reported by Morales-Muñoz et al¹⁰ and DiGiovanni et al⁴ differ by more than 20° . Although surprising at first glance, these pronounced differences are most likely due to the different testing protocols conducted to assess ADF. Although both authors applied nonweightbearing testing, Morales-Muñoz et al¹⁰ conducted their measurements with the “subtalar joint supinated,” the anatomic landmarks applied were the “shaft of the fibula and lateral border of the foot,” and a goniometer was used. One investigator conducted the measurement, as indicated in Morales-Muñoz et al's Figure 1. DiGiovanni et al⁴, on the contrary, used an equinometer, locked the midfoot, and ensured neutral hindfoot position, applied a constant torque of 10 Nm and used the fibula and plantar aspect of the foot as measurement landmarks. These are just 2 of numerous measurement techniques reported in literature. Testing protocols vary in almost every aspect, the general setup (non-weightbearing,² weightbearing,¹¹ and instrumented⁵), the measurement landmarks used (y-axis: fibula, tibia, or Achilles tendon,^{1,13,14} x-axis: shaft of the fifth metatarsal bone, plantar aspect of the foot, or the floor^{8,9,13}), and the measurement devices used (mobile apps, inclinometers, or goniometers^{1,7,14}). Clearly, each single aspect has a pronounced influence on the ADF values measured. Moreover, some protocols appear more applicable than others. Morales-Muñoz et al,¹⁰ for example, had one investigator controlling foot position, applying maximum ankle dorsiflexion and conducting the

measurements. From our own clinical experience, we find it hard to conduct all of these maneuvers at once.

Despite the promising treatment approach to MGT reported by Morales-Muñoz et al,¹⁰ the examination technique and equinus ADF values reported highlight the major shortcoming in current research on MGT. In order to identify patients suitable for gastrocnemius release and to interpret the results of different procedures in various pathologies, standard techniques to assess ADF should be applied. Currently, various tests are being used to assess ADF. Therefore, the informative value of the studies as well as the comparability between studies suffers considerably. We are convinced that it is essential to first agree on a standardized technique to quantify ADF. Further, using this standardized technique's physiological and pathological norm values should then be defined. These values build the prerequisite for any treatment study. Therefore, we would like to emphasize the need to agree on a standardized, evidence-based examination technique to assess ADF and MGT.

Sebastian F. Baumbach, MD

Mareen Braunstein, MD

Hans Polzer, MD

Munich University Hospital

Department of Trauma Surgery

Nussbaumstr. 20

80336 Munich, Germany

Email: hans.polzer@med.uni-muenchen.de

References

1. Baumbach SF, Brumann M, Binder J, et al. The influence of knee position on ankle dorsiflexion—a biometric study. *BMC Musculoskelet Disord*. 2014;15(1):246. doi:10.1186/1471-2474-15-246.
2. Bolívar YA, Munuera PV, Padillo JP. Relationship between tightness of the posterior muscles of the lower limb and plantar fasciitis. *Foot Ankle Int*. 2013;34(1):42-48. doi:10.1177/1071100712459173.
3. De Los Santos-Real R, Morales-Muñoz P, Payo J, Escalera-Alonso J. Gastrocnemius proximal release with minimal incision: a modified technique. *Foot Ankle Int*. 2012;33(9):750-754. doi:DOI: 10.3113/FAI.2012.0750.
4. DiGiovanni CW, Kuo R, Tejwani N, et al. Isolated gastrocnemius tightness. *J Bone Joint Surg Am*. 2002;84-A(6):962-970.
5. Gatt A, Chockalingam N. Validity and reliability of a new ankle dorsiflexion measurement device. *Prosthet Orthot Int*. 2013;37(4):289-297. doi:10.1177/0309364612465886.

6. Johanson M, Baer J, Hovermale H, Phouthavong P. Subtalar joint position during gastrocnemius stretching and ankle dorsiflexion range of motion. *J Athl Train*. 2008;43(2):172-178. doi:10.4085/1062-6050-43.2.172.
7. Kang M-H, Lee D-K, Park K-H, Oh J-S. Association of ankle kinematics and performance on the y-balance test with inclinometer measurements on the weight-bearing-lunge test. *J Sport Rehabil*. 2015;24(1):62-67. doi:10.1123/jsr.2013-0117.
8. Kim PJ, Peace R, Mieras J, et al. Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used. *J Am Podiatr Med Assoc*. 2011;101(5):407-414.
9. Krause DA, Cloud BA, Forster LA, Schrank JA, Hollman JH. Measurement of ankle dorsiflexion: a comparison of active and passive techniques in multiple positions. *J Sport Rehabil*. 2011;20(3):333-344.
10. Morales-Muñoz P, De Los Santos Real R, Barrio Sanz P, et al. Proximal Gastrocnemius Release in the Treatment of Mechanical Metatarsalgia. *Foot Ankle Int*. 2016;37(7):782-789.
11. Munteanu SE, Strawhorn AB, Landorf KB, Bird AR, Murley GS. A weightbearing technique for the measurement of ankle joint dorsiflexion with the knee extended is reliable. *J Sci Med Sport*. 2009;12(1):54-59. doi:10.1016/j.jsams.2007.06.009.
12. Sobel E, Caselli MA, Velez Z. Effect of persistent toe walking on ankle equinus. Analysis of 60 idiopathic toe walkers. *J Am Podiatr Med Assoc*. 1997;87(1):17-22. doi:10.7547/87507315-87-1-17.
13. Vohralik SL, Bowen AR, Burns J, Hiller CE, Nightingale EJ. Reliability and validity of a smartphone app to measure joint range. *Am J Phys Med Rehabil*. 2015;94(4):325-330. doi:10.1097/PHM.0000000000000221.
14. Williams CM, Caserta AJ, Haines TP. The TiltMeter app is a novel and accurate measurement tool for the weight bearing lunge test. *J Sci Med Sport*. 2013;16(5):392-395. doi:10.1016/j.jsams.2013.02.001.