

BMJ Open Patient trajectories and their impact on mobility, social participation and quality of life in patients with vertigo/dizziness/balance disorders and osteoarthritis (MobilE-TRA): study protocol of an observational, practice-based cohort study

Rebecca Kisch,¹ Antje Bergmann,² Daniela Koller,^{1,3} Reiner Leidl,^{3,4} Ulrich Mansmann,^{1,3} Martin Mueller,⁵ Linda Sanftenberg,⁶ Joerg Schelling,⁶ Leonie Sundmacher,^{3,7} Karen Voigt,² Eva Grill^{1,8}

To cite: Kisch R, Bergmann A, Koller D, *et al.* Patient trajectories and their impact on mobility, social participation and quality of life in patients with vertigo/dizziness/balance disorders and osteoarthritis (MobilE-TRA): study protocol of an observational, practice-based cohort study. *BMJ Open* 2018;**8**:e022970. doi:10.1136/bmjopen-2018-022970

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-022970>).

Received 15 March 2018
Revised 26 March 2018
Accepted 27 March 2018



For numbered affiliations see end of article.

Correspondence to

Dr Rebecca Kisch;
rebecca.kisch@med.uni-muenchen.de

ABSTRACT

Introduction Mobility limitations have a multitude of different negative consequences on elderly patients including decreasing opportunities for social participation, increasing the risk for morbidity and mortality. However, current healthcare has several shortcomings regarding mobility sustainment of older adults, namely a narrow focus on the underlying pathology, fragmentation of care across services and health professions and deficiencies in personalising care based on patients' needs and experiences. A tailored healthcare strategy targeted at mobility of older adults is still missing.

Objective The objective is to develop multiprofessional care pathways targeted at mobility sustainment and social participation in patients with vertigo/dizziness/balance disorders (VDB) and osteoarthritis (OA).

Methods Data regarding quality of life, mobility limitation, pain, stiffness and physical function is collected in a longitudinal observational study between 2017 and 2019. General practitioners (GPs) recruit their patients with VDB or OA. Patients who visited their GP in the last quarter will be identified in the practice software based on VDB and OA-related International Classification of Diseases 10th Revision. Study material will be sent from the practice to patients by mail. Six months and 12 months after baseline, all patients will receive a mail directly from the study team containing the follow-up questionnaire. GPs fill out questionnaires regarding patient diagnostics, therapy and referrals.

Ethics and dissemination The study was approved by the ethical committee of the Ludwig-Maximilians-Universität München and of the Technische Universität Dresden. Results will be published in scientific, peer-reviewed journals and at national and international conferences. Results will be disseminated via newsletters, the project website and a regional conference for representatives of local and national authorities.

Strengths and limitations of this study

- This observational, practice-based prospective cohort study focuses on patients with vertigo/dizziness/balance disorders and/or osteoarthritis, both health conditions with high burden of disability in older adults.
- The study will contribute to identifying determinants of patient trajectories, special concerns in care of specific age groups and significant combinations with comorbidities.
- The study reflects the patients' perspective and treatment trajectories including diagnostic procedures, comorbidities, diagnostic and therapeutic procedures used, tentative and final diagnoses and referrals based on general practitioners' (GPs) information.
- All eligible patients who recently consulted a GP during office hours will be searched in the patient database and contacted for study participation.
- Only those who are able and interested in filling out questionnaires and who are able to go to the physician participate in the study.

INTRODUCTION

With a prevalence of up to 50%, vertigo, dizziness and balance disorders (VDB) are frequent complaints of those patients aged 60 and older. The causes of VDB are often multifactorial. Distinct treatable vestibular disease entities, dizziness caused by medication, cardiovascular disease or diabetes may align with symptoms of the ageing of vestibular, proprioceptive or somatosensory systems. By increasing postural instability, VDB are

among the most obvious and prevalent causes for falls in aged adults¹; they limit mobility and activities of daily life² and restrict social participation.^{2,3}

Hip and knee osteoarthritis (OA) is ranked as the 11th overall out of 291 contributors to global disability.⁴ The risk for OA increases with older age. In Germany, nearly half (40.4%, 95% CI 37.4 to 43.6) of those aged 60–69 are diagnosed as having OA.⁵ Persons with OA have a specific increased risk for cardiovascular and dementia associated mortality, with low levels of physical activity arguably contributing to this excess mortality.⁶ Several evidence-based recommendations and guidelines are available for the management of OA.⁷ Frequently recommended non-surgical treatment options include patient education, exercise, weight loss and the use of assistive devices.

However, in Germany and internationally, appropriate care for patients with VDB and OA is a big challenge in all healthcare settings. Chronological age seems to be one major determinant of inappropriate care for both VDB⁸ and OA.⁹ We have recently shown from the perspective of a tertiary care centre that patients with VDB receive a large range of different unsuccessful diagnostic procedures and treatments prior to referral including extensive and expensive imaging.¹⁰ In OA, the lack of standardised and coordinated approaches to pragmatic management¹¹ has been noted as well. Ultimately, OA regularly results in joint replacement at high economic costs, while benefits are not clear. Large regional variations of surgery rates^{12,13} seem to indicate that to date there are no coordinated care pathways for OA in Germany. Information about the determinants and outcomes of patient trajectories will contribute to the identification of optimal care pathways and the development of targeted interventions. Managing the core causes for disability in old age promotes healthy ageing and a more efficient use of healthcare resources in Germany.

MobilE-Net is a network composed of three projects which aim to develop multiprofessional care pathways targeted at older adults to reduce the burden of disability and to promote healthy ageing, mobility and participation. The subproject MobilE-TRA investigates the status quo of care from the patient perspective with special regard to gateways, decisive transitions and decision interfaces. Patient-relevant participation outcomes will be modelled depending on main potential transition points. Treatment trajectories for patients with VDB and OA include diagnostic procedures, comorbidities, diagnostic and therapeutic procedures used, tentative and final diagnoses and referral practice.

METHODS

Study design

MobilE-TRA is an observational, practice-based prospective cohort study among general practitioner (GP) practices and their patients with VDB or OA. Its longitudinal design with data collection between 2017 and 2019

captures the trajectories and changes to functioning and quality of life as well as resource utilisation over time.

Participants and recruitment

GPs in two German cities, Munich and Dresden, and surrounding areas, recruit their patients presenting with VDB or OA. A recruitment strategy for hard-to-reach individuals will be applied, where eligible patients will be searched in the patient database to increase the number of contacted patients.¹⁴ GPs will check the list of eligible patients before sending the paper-based study material out via mail. Patient information on study contents is done by means of the accompanying letter. After signing the informed consent form, patients are contacted by a researcher of the university team for follow-up. This researcher is not involved in data analysis.

GPs

GP practices will be invited via mail from a network of 262 GP practices associated with the Institute of Primary Care and Family Practice of the Ludwig-Maximilians Universität München (LMU)—as teaching and research partners and about 100 GP practices associated with the Department of General Practice of the University Hospital Carl Gustav Carus of the Technische Universität Dresden.

Patients

Participating practices will identify patients who presented with VDB or OA. Following an established procedure for data collection in GP practices,¹⁴ eligible patients will be identified by searching the practice database. Patients are eligible if in the last quarter they visited the GP for an episode of VDB, or symptomatic hip or knee OA, having statutory health insurance, sufficient command of German language and being 65 years and older. Relevant diagnoses according to the ICD-10 are listed in [table 1](#).

The patient receives the patient information, informed consent and baseline questionnaire along with a postage prepaid return envelope from the GP via mail. In case of consent, the patient fills out these forms and sends them back to the study centre, which is the team at the LMU Institute for Medical Information Processing, Biometrics and Epidemiology. The study is conducted in accordance with the Declaration of Helsinki¹⁵ and was approved by the Ethical Committees of the LMU and Technische Universität Dresden.

For the follow-up surveys (after 6 and 12 months), a cover letter with instructions and the questionnaires will be sent via mail from the study team to the patient's address, with a postage prepaid return envelope.

Patient and public involvement

The final questionnaire was pretested in patients at the University Hospital, LMU Munich, before starting the actual data collection. Patients indicated the time it took to complete the questionnaire and reported any problems in filling out the questionnaire, for example, with the order of the questions or misunderstandings in the instructions.

Table 1 Overview of research outcomes and instruments

Outcome	Instrument	No. of items	Patient with VBD	Patient with OA
Generic quality of life	EuroQol-5D-5L	6	x	x
Mobility limitation	HAQ-DI	44	x	x
Patient satisfaction	PACIC short form	11	x	x
Healthcare utilisation, medication	FIMA	73	x	x
Depression	PHQ-9	9	x	x
Sociodemographic information		16	x	x
Disease-specific measures				
Quality of life	WOMAC	17		x
Self-perceived level of handicap	DHI	25	x	
Activities and participation	VAP	20	x	

Disease-specific scales will be printed on coloured paper. DHI, Dizziness Handicap Inventory; EuroQol-5D-5L, EuroQol Five-Dimensional Five-Level Questionnaire; HAQ-DI, Health Assessment Questionnaire Disability Index; OA, osteoarthritis; PACIC, Patient Assessment of Chronic Illness Care; PHQ-9, Patient Health Questionnaire—Nine questions; VAP, Vestibular Activities and Participation; VBD, vertigo/dizziness balance disorders; WOMAC, Western Ontario and McMaster Universities.

Results of the main study will be disseminated to study participants via the project website.

Study schedule

Outcome measures are applied at baseline at participant level (a continuous inclusion of participants until required sample size is reached) as well as at follow-up 6 month after baseline and 12 months after baseline.

Instruments and outcomes

General practitioners

The GPs are handed out the standardised Questionnaire of Chronic Illness Care in Primary Care.¹⁶ It includes items on the number of patients treated per week, socio-demographic characteristics and GPs professional experience, the share of different patient groups in the practice, structural data of the practice, number of partners and employees in a practice, technical possibilities in the practice, implemented quality management systems and propensity for training.

GPs are also asked to complete a self-developed baseline questionnaire, including information, which might have an influence on diagnostic and referral behaviour.

After patients sent back questionnaires, physicians will complete this questionnaire on comorbidities, diagnostic and therapeutic procedures used, tentative and final diagnoses and referral practice or deliver the respective data out of their practice software.

Patients

Patients receive a self-developed questionnaire including sociodemographic data for example, age, gender, education, health insurance, living situation and difficulties in filling out the questionnaire.

The primary outcomes of the study measured by validated instruments are generic quality of life, mobility limitation, pain, stiffness and physical function in persons with hip and/or knee osteoarthritis and vertigo-specific physical and psychosocial functioning.

1. Generic quality of life is measured through the EuroQol Five-Dimensional Five-Level Questionnaire (EQ-5D-5L) by EuroQol Group.¹⁷ The EQ-5D showed adequate sensitivity to change in a recent randomised trial in unselected patients with vestibular vertigo and dizziness in primary care.^{18 19} For summary valuation, patients' report on the visual analogue scale and, as soon as available, a German value set for utilities based on EQ-5D-5L will be used.
2. Mobility limitation will be measured using the Health Assessment Questionnaire Disability Index.^{20 21} The patients report the amount of difficulty they have in performing activities of daily living.
3. Pain stiffness and physical function in persons with hip and/or knee osteoarthritis is assessed using the German version of the Western Ontario and McMaster Universities Osteoarthritis Index.^{22 23}
4. Patients with vertigo, dizziness and balance disorders will be asked to fill out the Dizziness Handicap Inventory (DHI)²⁴ and the Vestibular Activities and Participation questionnaire.²⁵ The instruments evaluate the effect of the disease on activity limitation, participation restrictions and experienced difficulties. Secondary outcomes are patient satisfaction with healthcare and healthcare utilisation.
 1. The Patient Assessment of Chronic Illness Care will be used as a measure for patient satisfaction.²⁶
 2. Healthcare resource utilisation and medication will be assessed using a validated questionnaire for unit costs for outpatient medical care, inpatient care, informal and formal nursing care and pharmaceuticals (FIMA).^{27 28}
 3. Depression is assessed using the Patient Health Questionnaire.²⁹

An overview of research outcomes, instruments and the number of items is illustrated in [table 2](#).

For non-responder patients to follow-up, a non-responder questionnaire will be provided to explore the reasons for withdrawal and to obtain a crude estimate on health status based on a small set of questions.

Sample size

Based on the recommendations for the minimally important differences of quality of life instruments to be 50% of the SD, we assumed 10.0 points on the Dizziness Handicap Inventory (DHI) as a clinically relevant difference. Assuming an SD of 20,³⁰ a sample size of 48 would allow estimating the mean DHI with a

Table 2 ICD-10 codes of included patients

ICD-10 code	Title
H81	Disorders of vestibular function
H82	Vertiginous syndromes in diseases classified elsewhere
R42	Dizziness and giddiness
A88.1	Epidemic vertigo
E53.8	Deficiency of other specified B group vitamins
F45.8	Other somatoform disorders
G11.8	Other hereditary ataxias
G43.1	Migraine with aura (classical migraine)
G45.0	Vertebro-basilar artery syndrome
G62	Other polyneuropathies
G63	Polyneuropathy in diseases classified elsewhere
H55	Nystagmus and other irregular eye movements
H83.0–2	Other diseases of inner ear
I95.1	Orthostatic hypotension
N95.1	Menopausal and female climacteric states
R26 without R26.1	Abnormalities of gait and mobility (without Paralytic gait)
M16	Coxarthrosis (arthrosis of hip)
M17	Gonarthrosis (arthrosis of knee)

ICD-10, International classification of diseases, 10th Revision.

power of 0.8 while simultaneously adjusting for four CIs ($\alpha=0.0125$).

We assume the loss to follow-up between data collection waves to be 28%.³¹ This means that 100 participants for the baseline data collection will allow for up to three follow-up waves. Considering the design effect with 12 patients on average per cluster (GP practice), and an ICC of 0.017,³² the sample size required is 118. Assuming a similar effect size in patients with OA, the total sample size required is 236, this means we will have to include at least 20 practices. At an estimated response rate of 25% among practices,³³ this means that 80 practices have to be contacted for the inclusion of a total of 236 patients with VDB and/or OA.

Statistical and health economic methods

First, to examine current treatment and referral behaviour of the primary healthcare providers, frequency and distribution of procedures, treatment, medication and referral will be analysed descriptively.

Second, to examine the differences of quality of life, mobility limitation, participation and satisfaction, trajectories will be examined graphically as a function of age stratified by sex.

Differences will be compared using χ^2 tests for categorical variables and analysis of variance for metric variables

which are log transformed if necessary. Mixed-effects regression models will be used with age, gender, socio-demographic variables and comorbidity as independent variables to analyse the determinants of different trajectories of quality of life, mobility limitation, participation and satisfaction. This method is appropriate to investigate longitudinal data with more than one wave of data.³⁴

Lastly, to examine healthcare resource utilisation of current practice, cost utility analysis will be used. Measurement of costs will focus on cost of care and will employ recent German measurement standards.²⁸ Economic assessment will be conducted from a healthcare perspective.^{35 36} Primary endpoint of the economic evaluation is cost per quality-adjusted life year (QALY) at 12 months after inclusion. The incremental cost-effectiveness ratio will be calculated including probabilistic analysis using bootstrapping techniques. To model different trajectories, we will develop a state-transition (Markov) model. This model will characterise the process of treatment through all observable stages and also include both cost and QALY endpoints. Monte-Carlo simulation will be used for probabilistic analysis.³⁷

DISCUSSION

This study aims to investigate the trajectories of healthcare delivery using vertigo, dizziness, balance disorders and joint diseases as a case in point and investigates determinants of healthcare utilisation and outcomes. Chronological age seems to be one major determinant of inappropriate care for both VDB and OA which makes this research particularly relevant.

While there is much variation in the management of VDB and OA in the challenging context of multimorbidity and polypharmacy in elderly patients in primary care, it seems reasonable to assume that there is no 'one-size-fits-all' strategy for optimal outcomes of these complex health conditions in aged adults. This project builds on the idea that care should be stratified based on distinct care pathways for aged adults with VDB or OA. Following Andersen and Newman's Behavioral Model of Healthcare,³⁸ we assume that such pathways are determined by disease-related factors, predisposition and enabling factors. This research will contribute in identifying these determinants, special concerns in care of specific age groups and significant combinations with comorbidities.

A potential limitation is that the study includes only patients who recently consulted a GP during office hours. Although most elderly patients in Germany go to the GP regularly, patients with limited mobility have limited possibility of healthcare utilisation. Thus, those patients who avoid going to a GP are not covered with this study, leading to an underestimation of mobility limitations as only those who are able and interested in filling questionnaires and who are able to go to doctors participate in the study.

The recruitment strategy with a search in the patient database will make the recruitment more efficient. Instead

of physicians inviting each patient face to face during office hours and handing out an envelope containing a personalised cover letter, study information and informed consent form, the questionnaire and a return envelope will be sent via mail. The aim is to (1) reach every patient eligible for the study and (2) reduce GPs' effort of informing each patient about the study.³⁹ The recruitment bears the risk that fewer patients participate in the study compared with when the GPs invite the patients personally during office hours. The feasibility of the recruitment strategy and barriers will be reported.

While the implementation of care pathways in Germany is still at its beginning, there are good practice examples of successful integration.⁴⁰ Using results of MobileE-TRA to develop clinical pathways for patients with VDB and OA can make a significant difference to the practice in primary care. The identification of determinants of VDB and OA will be used to develop individualised treatment strategies to overcome inappropriate care.

ETHICS AND DISSEMINATION

The study was approved by the ethical committee of the Ludwig-Maximilians-Universität München and of the Technische Universität Dresden. Personal data will be pseudonymised before sending the study material out via mail.

Results will be published in scientific, peer-reviewed journals and at national and international conferences. Results will be disseminated via newsletters, the project website and a regional conference for representatives of local and national authorities. The data set will be available for project researchers on request.

Author affiliations

¹Institute for Medical Information Processing, Biometry and Epidemiology, Ludwig-Maximilians-Universität München, Munich, Germany

²Department of General Practice/Medical Clinic III, University Hospital Carl Gustav Carus of the Technische Universität Dresden, Dresden, Germany

³Munich Center of Health Sciences, Ludwig-Maximilians-Universität München, Munich, Germany

⁴Institute of Health Economics and Health Care Management, Helmholtz Zentrum München, Munich, Germany

⁵Faculty of Applied Health and Social Sciences, Hochschule Rosenheim, Rosenheim, Germany

⁶Institute of General Practice and Family Medicine, University Hospital, Ludwig-Maximilians-Universität München, Munich, Germany

⁷Department of Health Services Management, Munich School of Management, Ludwig-Maximilians-Universität München, Munich, Germany

⁸German Centre for Vertigo and Balance Disorders, University Hospital, Ludwig-Maximilians-Universität, Munich, Germany

Acknowledgements The authors acknowledge patients' willingness to participate, support and their valuable feedback in the pilot testing of the questionnaire.

Contributors EG is the principal investigator. She conceived and supervised the project. RK coordinated the study. EG, DK, UM, RL, LSu, JS and MM contributed to the conception of the cohort and design of the project. LSa, JS, AB and KV are responsible for the accusation of the data recruitment. DK is responsible for the study's quality assessment and is, together with EG and UM, in charge of the overall study management. RK and EG drafted the manuscript. All authors approved and critically revised the final manuscript.

Funding This work is part of the project "Munich Network Health Care Research—MobileE-Net" and was supported by the German Federal Ministry of Education and Research grant number 01GY1603A.

Competing interests None declared.

Patient consent Obtained.

Ethics approval The Ethical Committee of the Ludwig-Maximilians-Universität München has approved the study protocol under the number 17-443. The Ethical Committee of the Technische Universität Dresden has approved the study protocol under the number E365092017.

Provenance and peer review Not commissioned; peer reviewed for ethical and funding approval prior to submission.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Agrawal Y, Carey JP, Della Santina CC, *et al.* Disorders of balance and vestibular function in US adults: data from the national health and nutrition examination survey, 2001-2004. *Arch Intern Med* 2009;169:938-44.
2. Mueller M, Strobl R, Jahn K, *et al.* Impact of vertigo and dizziness on self-perceived participation and autonomy in older adults: results from the KORA-Age study. *Qual Life Res* 2014;23:2301-8.
3. Bronstein AM, Golding JF, Gresty MA, *et al.* The social impact of dizziness in London and Siena. *J Neurol* 2010;257:183-90.
4. Cross M, Smith E, Hoy D, *et al.* The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014;73:1323-30.
5. Fuchs J, Rabenberg M, Scheidt-Nave C. [Prevalence of selected musculoskeletal conditions in Germany: results of the German health interview and examination survey for adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:678-86.
6. Nüesch E, Dieppe P, Reichenbach S, *et al.* All cause and disease specific mortality in patients with knee or hip osteoarthritis: population based cohort study. *BMJ* 2011;342:d1165.
7. Nelson AE, Allen KD, Golightly YM, *et al.* A systematic review of recommendations and guidelines for the management of osteoarthritis: the chronic osteoarthritis management initiative of the U.S. bone and joint initiative. *Semin Arthritis Rheum* 2014;43:701-12.
8. Roberts DS, Lin HW, Bhattacharyya N. Health care practice patterns for balance disorders in the elderly. *Laryngoscope* 2013;123:43.
9. Li LC, Sayre EC, Kopec JA, *et al.* Quality of nonpharmacological care in the community for people with knee and hip osteoarthritis. *J Rheumatol* 2011;38:2230-7.
10. Grill E, Strupp M, Müller M, *et al.* Health services utilization of patients with vertigo in primary care: a retrospective cohort study. *J Neurol* 2014;261:1492-8.
11. Smink AJ, Dekker J, Vliet Vlieland TPM, *et al.* Health care use of patients with osteoarthritis of the hip or knee after implementation of a stepped-care strategy: an observational study. *Arthritis Care Res* 2014;66:817-27.
12. OECD. *Geographic variations in health care: what do we know and what can be done to improve health system performance?* Paris: OECD Publishing, 2014.
13. Lüring C, Niethard FU, Günther KP, *et al.* *Faktencheck Gesundheit. Knieoperationen (Endoprothetik)—Regionale Unterschiede und ihre Einflussfaktoren:* Bertelsmann Stiftung, 2013.
14. Brach M, Moschny A, Bucker B, *et al.* Recruiting hard-to-reach subjects for exercise interventions: a multi-centre and multi-stage approach targeting general practitioners and their community-dwelling and mobility-limited patients. *Int J Environ Res Public Health* 2013;10:6611-29.
15. World Medical Organisation. Declaration of helsinki (1964). *Bmj* 1996;313:1448-9.

16. Steinhäuser J, Miksch A, Ose D, *et al.* Questionnaire of chronic illness care in primary care—psychometric properties and test-retest reliability. *BMC Health Serv Res* 2011;11:295.
17. Greene ME, Rader KA, Garellick G, *et al.* The EQ-5D-5L improves on the EQ-5D-3L for health-related quality-of-life assessment in patients undergoing total hip arthroplasty. *Clin Orthop Relat Res* 2015;473:3383–90.
18. Yardley L, Barker F, Muller I, *et al.* Clinical and cost effectiveness of booklet based vestibular rehabilitation for chronic dizziness in primary care: single blind, parallel group, pragmatic, randomised controlled trial. *BMJ* 2012;344:e2237.
19. EuroQol Group. EuroQol—a new facility for the measurement of health-related quality of life. *Health Policy* 1990;16:199–208.
20. Brühlmann P, Stucki G, Michel BA. Evaluation of a German version of the physical dimensions of the Health Assessment Questionnaire in patients with rheumatoid arthritis. *J Rheumatol* 1994;21:1245–9.
21. Fries JF, Spitz PW, Young DY. The dimensions of health outcomes: the health assessment questionnaire, disability and pain scales. *J Rheumatol* 1982;9:789–93.
22. Bellamy N. The WOMAC knee and hip osteoarthritis indices: development, validation, globalization and influence on the development of the AUSCAN hand osteoarthritis indices. *Clin Exp Rheumatol* 2005;23:S148–53.
23. Stucki G, Meier D, Stucki S, *et al.* [Evaluation of a German version of WOMAC (Western Ontario and McMaster Universities) arthrosis index]. *Z Rheumatol* 1996;55:40–9.
24. Kurre A, Bastiaenen CH, van Gool CJ, *et al.* Exploratory factor analysis of the Dizziness Handicap Inventory (German version). *BMC Ear Nose Throat Disord* 2010;10:3.
25. Alghadir AH, Al-Momani M, Marchetti GF, *et al.* Cross-cultural adaptation and measurement properties of the Arabic version of the fall efficacy scale international. *Neurosciences* 2015;20:230–5.
26. Glasgow RE, Wagner EH, Schaefer J, *et al.* Development and validation of the Patient Assessment of Chronic Illness Care (PACIC). *Med Care* 2005;43:436–44.
27. Seidl H, Bowles D, Bock JO, *et al.* [FIMA—questionnaire for health-related resource use in an elderly population: development and pilot study]. *Gesundheitswesen* 2015;77:46–52.
28. Bock JO, Brettschneider C, Seidl H, *et al.* [Calculation of standardised unit costs from a societal perspective for health economic evaluation]. *Gesundheitswesen* 2015;77:53–61.
29. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–13.
30. Grill E, Penger M, Kentala E. Health care utilization, prognosis and outcomes of vestibular disease in primary care settings: systematic review. *J Neurol* 2016;263(Suppl 1):36–44.
31. Vega S, Benito-León J, Bermejo-Pareja F, *et al.* Several factors influenced attrition in a population-based elderly cohort: neurological disorders in Central Spain Study. *J Clin Epidemiol* 2010;63:215–22.
32. Littenberg B, MacLean CD. Intra-cluster correlation coefficients in adults with diabetes in primary care practices: the Vermont Diabetes Information System field survey. *BMC Med Res Methodol* 2006;6:20.
33. Bonevski B, Magin P, Horton G, *et al.* Response rates in GP surveys—trialling two recruitment strategies. *Aust Fam Physician* 2011;40:427–30.
34. Singer JD, Willett JB. *Applied longitudinal data analysis: Modeling change and event occurrence*. Oxford: Oxford university press, 2003.
35. Graf von der Schulenburg JM, Greiner W, Jost F, *et al.* German recommendations on health economic evaluation: third and updated version of the Hanover Consensus. *Value Health* 2008;11:539–44.
36. Husereau D, Drummond M, Petrou S, *et al.* CHEERS Task Force. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *BMJ* 2013;346:f1049.
37. Caro JJ, Briggs AH, Siebert U, *et al.* Modeling good research practices—overview: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force-1. *Med Decis Making* 2012;32:667–77.
38. Andersen R, Newman JF. Societal and individual determinants of medical care utilization in the United States. *Milbank Mem Fund Q Health Soc* 1973;51:95–124.
39. Stephan AJ, Kovacs E, Phillips A, *et al.* Barriers and facilitators for the management of vertigo: a qualitative study with primary care providers. *Implement Sci* 2018;13:25.
40. Schneider AM, Klases B, Jennerwein C, *et al.* [Intersectoral pain treatment. The Algesiologikum group]. *Schmerz* 2013;27:166–73.