

Zygote 23 (April), pp. 307–311. © Cambridge University Press 2013
doi:10.1017/S0967199413000427 First Published Online 28 August 2013

Short communication

Expression pattern of *STAT5A* gene during early bovine embryogenesis

Krzysztof Flisikowski¹, Marc Boulhaeve², Fabiola P. Lopes², Eckhard Wolf² and Lech Zwierzchowski³
Technische Universität München, Freising, Germany; Ludwig-Maximilians-Universität München, Oberschleissheim, Germany; and Institute of Genetics and Animal Breeding, Polish Academy of Sciences, Magdalenka, Poland

Date submitted: 29.04.2013. Date revised: 28.06.2013. Date accepted: 04.07.2013

Summary

Growth hormone (*GH*) plays an important role in early embryo development. It has been shown to activate multiple pathways, the most comprehensively studied being the STAT/JAK (Signal transducers and activators of transcription/Janus kinase) pathway. The objective of the present study was to investigate *STAT5A* gene expression during early bovine embryogenesis. Real-time polymerase chain reaction (RT-PCR) was used to measure the abundance of *STAT5A* transcripts. The mRNA was present at all stages of preimplantation bovine embryos investigated. The most abundant *STAT5A* expression occurred at the 2-cell stage. Expression was markedly reduced between the 4-cell and 8-cell stages, coinciding with the known time of embryo genome activation and loss of maternal mRNAs. This finding suggests that the embryonic *STAT5A* gene is primarily activated by maternal gene products.

Introduction

Signal transducers and activators of transcription (STATs) are transcription factors implicated in cellular proliferation, differentiation and metabolism. After ligand binding-induced activation of cognate receptors, STAT proteins become phosphorylated, hetero- or homodimerize, and enter the nucleus (Wakao *et al.*, 1994). *STAT5A* mediates, through the STAT/JAK pathway, signals of hormones with important functions in bovine reproduction – growth hormone (*GH*), leptin and prolactin (Darnell *et al.*, 1994; Duncan *et al.*, 1997). This signalling cascade is extraordinarily versatile and leads both to general phenomena, such as cell proliferation, and to highly specialized signalling, such as tissue-specific expression of individual genes. It is also known that *GH* exerts distinct effects on the development, differentiation and metabolism of preimplantation embryos. The development of

bovine embryos cultured *in vitro* can be improved by supplementing the medium with *GH* (Kolle *et al.*, 2001). *GH* and its receptor (*GHR*) mRNA are detectable from the second stage (2-cell stage) of bovine embryogenesis onwards (Kolle *et al.*, 1998, 2002; Joudrey *et al.*, 2003). Leptin, another possible ligand for the *STAT5* gene family, is known to regulate diverse reproductive functions, and recent studies suggest its involvement in early embryo development (Levy *et al.*, 2001; Hansis & Edwards, 2003; Madeja *et al.*, 2009). In the study by Madeja *et al.* (2009), the expression of leptin and its functional receptor was shown in mouse oocytes and preimplantation embryos.

The aim of the present study was to evaluate the level of *STAT5A* mRNA expression by quantitative real-time reverse transcription-polymerase chain reaction (RT-PCR) in bovine preimplantation embryos at different stages. The results obtained enhance understanding of the functioning of the *GH/STAT5A* pathways during embryo development.

Materials and methods

In vitro production of bovine embryos

Bovine ovaries were collected at a local slaughterhouse (University of Munich) and washed in pre-heated

¹All correspondence to: Krzysztof Flisikowski. Technische Universität München, Liesel-Beckmann-Str. 1, 85354 Freising, Germany. Tel. +49 8161 712036. Fax. +49 8161 712108. e-mail: flisikowski@wzw.tum.de

²Ludwig-Maximilians-Universität München, Hackerstrasse 27, 85764 Oberschleissheim, Germany.

³Institute of Genetics and Animal Breeding, Polish Academy of Sciences, Jastrzebiec, 05–552 Magdalenka, Poland.

Table 1 Primers used for RT-qPCR determination of mRNA abundance and their localization in respective genes

Gene	Primer sequence 5'→3'	Amplified fragment length (bp)	Accession number GenBank
STAT5A	ACGGTACCTTCTTGTTGCGC CAGGTTACGGTCAGAGAGTCAAAC	85	Z72482
H2A	TTCGAAATGGCTGGCGG GGAAGTCAAACCGGCTCT	91	NM174809

phosphate-buffered saline (PBS) (30°C). Cumulus-oocyte complexes (COCs) were obtained by aspirating follicles (2–8 mm diameter) with a 20-gauge needle and a vacuum of approximately 13 kPa. Only COCs with a complete dense cumulus and dark, evenly granulated cytoplasm were selected for *in vitro* maturation. COCs were collected in a 50 ml centrifuge tube and washed twice with pre-incubated (39°C, 5% CO₂ in air) tissue culture media 199 (Invitrogen) supplemented with 10% heat-inactivated serum from cows at estrus (OCS). They were washed again in medium 199 supplemented with 10% OCS and 10 µg/ml bovine follicle stimulating hormone (FSH) and leutenizing hormone (LH) and matured in this medium for 22–24 h at 39 °C in an atmosphere of 5% CO₂ in air and maximum humidity. After maturation, COCs were maintained in Tyrode's–albumin–lactate–pyruvate (TALP) medium that contains 6 mg/ml BSA, 100 µg/ml heparin (Sigma). Motile spermatozoa were obtained by the swim-up procedure and by centrifugation of semen from the upper phase for 10 min at 700 g at room temperature. Viable spermatozoa (final concentration 10⁶ spermatozoa/ml) were added to matured COCs for 22 h under the same conditions as used for *in vitro* maturation. Semen from the same bulls was used for all experiments.

For *in vitro* culture, cumulus cells were removed from presumptive zygotes by vortexing for 3 min. Groups of 30–40 presumptive zygotes were washed three times and cultured for 8 days in 400 µl synthetic oviduct fluid (SOF) supplemented with 10% OCS and covered with pre-equilibrated mineral oil (Sigma). Embryos were cultured in incubator chambers at 38.5 °C in an atmosphere of 5% CO₂, 10% O₂, 85% N₂. The cleavage rate was evaluated 72 h post-insemination (hpi) and the number of blastocysts scored at 186 hpi. Three replicates of each experiment were performed. Overall, high rates of maturation (82 ± 4%), fertilization (68 ± 2%), cleavage (52 ± 3%) were obtained, and a moderate rate of blastocyst formation (32 ± 3%), in the *in vitro* produced (IVP) experiments described.

Oocyte and embryo collection

Oocytes were collected 22–24 h after maturation and cumulus cells removed by vortexing. Zygotes were

collected 18–20 hpi, 2-cells 26–30 hpi, 4-cells 37–39 hpi, 8-cells at day 3 (d3), 16-cells d4, morula d5, expanded d7 and hatched blastocysts d8. Embryos were stored in groups of four in 10 µl RNeasy Lysis Buffer (Qiagen) at –80 °C until extraction of RNA. Three replicates of pools of five *in vitro* matured oocytes and pools of five *in vitro* embryos were used for each developmental stage.

RNA isolation and reverse transcription

Total RNA was isolated with TriZol reagents (Invitrogen), essentially according to the method of Chomczynski & Sacchi (1987). RNA preparations were treated with 0.25 µl DNase (10 IU/µl; Invitrogen) by incubation at 25 °C for 15 min. The enzyme was inactivated by adding 1 µl 25 mM EDTA and heating at 65 °C for 10 min. Samples were chilled rapidly on ice and reverse transcription directly followed. Reverse transcription was carried out in a total volume of 40 µl according to the Superscript II reverse transcriptase protocol (Invitrogen). Samples were stored at –80 °C until quantitative PCR (qPCR).

Gene cloning

Amplification primers (Table 1) were designed according to the sequences found in GenBank and using the software Primer Express 3 (Applied Biosystems).

For qPCR, plasmid standards were established for each gene of interest with the TOPO cloning kit (Invitrogen). Prior to the cloning step PCR was performed in a total volume of 20 µl with 0.25 µl HotStarTaq (Qiagen), 1 µl of each primer listed in Table 1 (final concentration 300 nM), 1 µl cDNA, 2 µl 1 mM dNTPs (Fermentas), 1.25 µl MgCl₂ (0.25 mM) and 11.5 µl water. Thermal cycling conditions were as follows: 15 min at 95 °C, then 35 cycles of 95 °C for 15 s, 60 °C for 30 s and 72 °C for 30 s, terminating with 10 min at 72 °C. Samples comprising 10% of the PCR reaction were analysed by 2.5% agarose gel electrophoresis and ethidium bromide fluorescence.

For the cloning step, a PCR aliquot of 2 µl was mixed with 1 µl salt solution, 1 µl TOPO vector and 2 µl water (Invitrogen). Reactions were incubated for 10 min at room temperature and 2 µl ligation mix added to the bacteria. Plasmids were isolated using the QIAprep® spin Mini kit (Qiagen) following the




Figure 1 Relative expression levels of the *STAT5A* gene during early embryogenesis. Expression of *STAT5A* was normalized to the *H2A* gene. Mean \pm standard error (SE) for nine determinations, five embryos each. Abbreviations: 2-C, 2-cell stage; 4-C, 4-cell stage; 8-C, 8-cell stage; 16-C, 16-cell stage; EB, expanded blastocyst; HB, hatched blastocyst; M, morula; MO, matured oocyte; Z, zygote. *Differences statistically significant at $P < 0.05$. **Differences statistically significant at $P < 0.01$.

manufacturer's instructions. Sequence integration was tested by digestion of 2 μ l plasmid isolate with *EcoRI* (Fermentas) and gel electrophoresis.

Quantitative PCR (qPCR)

Quantification of mRNA abundance was performed by SybrGreen-based real-time PCR detection using an ABI PRISM 7000 apparatus (Applied Biosystems). Amplification mixes (25 μ l) contained 2 μ l cDNA, 12.5 μ l SybrGreen PCR Mix, 0.25 IU AmpErase uracil *N*-glycosylase (Applied Biosystems), 300 nM each primer and 7.25 μ l water.

Standard curves were obtained using plasmids prepared as described above, and seven serial dilutions (1:5) used for all selected genes, providing a range of 1 million to 64 copies. The abundance of *STAT5A* mRNA was calculated using the standard curve method, with determination of PCR amplification efficiency and normalization for histone 2A mRNA as the internal reference. qPCR was started with 2 min at 50°C for AmpErase activation and 10 min at 95 °C for denaturation. The programme continued with 45 cycles of 15 s at 95°C, 30 s at 60°C and 30 s at 72°C. Each assay included triplicates of cDNA primed separately for the two genes of interest. The significance of the differences in the expression levels of *STAT5A* was estimated using Duncan's new multiple range test using data from different embryo stages (mean \pm standard deviation (SD)).

Results and Discussion

Real-time RT-PCR was performed to detect and quantify *STAT5A* mRNA expression. Real-time RT-PCR with an external calibration curve is a fully quantitative method, and thus allows an absolute comparison of the abundance of individual transcripts

in RNA preparations at each embryonic stage. RT-PCR products showed single bands of the lengths predicted: 85 bp for *STAT5A* and 91 bp for *H2A* (data not shown). The abundance of mRNA encoding *STAT5A* was normalized relative to *H2A* mRNA. The C_T values obtained for *H2A* mRNA were similar in all embryonic stages (data not shown), as reported earlier (Robert *et al.*, 2002).

Figure 1 shows relative expression measurements for *STAT5A*. The expression level at the mature oocyte stage was significantly lower than at zygote ($P < 0.05$) or 2-cell stage ($P < 0.01$). *STAT5A* expression reached a peak at the 2-cell stage and showed a sudden drop at the 8-cell stage ($P < 0.01$). *STAT5A* expression in hatched blastocysts appeared somewhat higher than in 8-cell, morula and expanded blastocyst stages ($P < 0.05$), but was still approximately 100-fold lower than at the 4-cell stage. Quantification of *STAT5A* expression in bovine embryos revealed that the abundance of the transcript decreased markedly at the 8-cell stage.

Low fertilization rate and embryonic mortality are significant factors contributing to failed pregnancies in cattle (Diskin & Morris, 2008). Among the genetic factors, the genes of the pituitary-specific positive transcription factor 1 (*POU1F1*) pathway play a crucial role in embryonic survival rate (Khatib *et al.*, 2009a). Growth hormone, prolactin and their main mediator, *STAT5A*, belong to this pathway. Growth hormone supports mouse (Wright & Bondioli, 1981) and bovine (Kolle *et al.*, 2001) preimplantation embryo development *in vitro*. Also, in both species *GH* transcripts were detected at different developmental stages of early embryos. In bovine embryos the *GH* transcript has been detected in oocytes and all pre-attachment stages including blastocysts (Kolle *et al.*, 1998; Joudrey *et al.*, 2003). It has also been shown that the quality of bovine blastocysts derived from non-early cleaving zygotes is improved by *GH* supplementation (Pers-Kamczyc *et al.*, 2010). However,

little information is known about the major mediator of the GH signal, embryonic *STAT5A*. *Stat5a*^{-/-} double knockout mice exhibit female infertility (Teglund *et al.*, 1998), thus *STAT5A* is considered to be involved in oogenesis and early embryo development. Two studies showed upregulation of *STAT5A* expression in bovine degenerate embryos compared with blastocysts (Khatib *et al.*, 2009b; Laporta *et al.*, 2011).

In mice, *STAT5A* transcription decreases between MII and 4-cell stages, and then increases from the morula stage (Nakasato *et al.*, 2006). The present paper demonstrates developmental changes in *STAT5A* gene expression during early bovine preimplantation embryogenesis. *STAT5A* mRNAs are present throughout the early developmental stages but fall to low levels after embryonic genome activation, which is different to mice. The bovine embryonic genome undergoes a major transcriptional activation at the 8-cell stage (66–72 hpi) and *in vitro* cultured bovine embryos can struggle to pass the so-called ‘8–16-cell developmental block’ (Wolf *et al.*, 2003). The marked drop in *STAT5A* expression after the 4-cell stage suggests that abundant *STAT5A* expression observed in the early stage embryos is a consequence of activation by maternal gene products, and the subsequent low level of *STAT5A* expression represents regulation by the embryonic genome. The different expression pattern before and after developmental block suggests different activation mechanism for *STAT5A*. It is known that mammary gland *STAT5A* expression is induced by GH treatment (Boutinaud & Jammes, 2004). Therefore, maternal GH or other cytokines may also stimulate expression of *STAT5A* before the 8-cell stage in cattle. Indeed, mice experiments showed that dynamic changes in the expression of cytokine receptors activate the STAT5 signalling pathway in preimplantation embryos (Nakasato *et al.*, 2006). The constantly low level of *STAT5A* expression suggests that autocrine cytokine stimulation of *STAT5A* pathway may be activated at later stages in bovine embryos. It should also be mentioned that the presence of GH in unknown concentrations during maturation and culture could alter *STAT5A* expression. In contrast, expression of the related factor *STAT3* declines steadily from the oocyte to the 16-cell stage and then increases until hatched blastocyst (Leidenfrost *et al.*, 2011). A similar expression pattern to *STAT3* has also been observed for the related *LEP* gene in *in vitro* produced bovine embryos (Madeja *et al.*, 2009). In the same experiment *LEPR* mRNA expression was detected in all but 4-cell bovine embryos (Madeja *et al.*, 2009). These data illustrate the complexity and dynamic nature of mRNA expression changes in preimplantation embryos.

These results provide new data regarding the embryonic *STAT5A* expression and new insight for the

GH pathways regulation in bovine early embryonic development.

References

- Boutinaud, M. & Jammes, H. (2004). Growth hormone increases Stat5 and Stat1 expression in lactating goat mammary gland: a specific effect compared to milking frequency. *Domest. Anim. Endocrinol.* **27**, 363–78.
- Chomczynski, P. & Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**, 156–9.
- Darnell, J.E., Jr., Kerr, I.M. & Stark, G.R. (1994). Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* **264**, 1415–21.
- Diskin, M.G. & Morris, D.G. (2008). Embryonic and early foetal losses in cattle and other ruminants. *Reprod. Domest. Anim.* **43**(Suppl 2), 260–7.
- Duncan, S.A., Zhong, Z., Wen, Z. & Darnell, J.E., Jr. (1997). STAT signaling is active during early mammalian development. *Dev. Dyn.* **208**, 190–8.
- Hansis, C. & Edwards, R.G. (2003). Cell differentiation in the preimplantation human embryo. *Reprod. Biomed. Online* **6**, 215–20.
- Joudrey, E.M., Lechniak, D., Petrik, J. & King, W.A. (2003). Expression of growth hormone and its transcription factor, Pit-1, in early bovine development. *Mol. Reprod. Dev.* **64**, 275–83.
- Khatib, H., Huang, W., Wang, X., Tran, A.H., Bindrim, A.B., Schutzkus, V., Monson, R.L. & Yandell, B.S. (2009a). Single gene and gene interaction effects on fertilization and embryonic survival rates in cattle. *J. Dairy Sci.* **92**, 2238–47.
- Khatib, H., Maltecca, C., Monson, R.L., Schutzkus, V. & Rutledge, J.J. (2009b). Monoallelic maternal expression of *STAT5A* affects embryonic survival in cattle. *BMC Genet.* **10**, 13.
- Kolle, S., Sinowatz, F., Boie, G., Lincoln, D., Palma, G., Stojkovic, M. & Wolf, E. (1998). Topography of growth hormone receptor expression in the bovine embryo. *Histochem. Cell Biol.* **109**, 417–9.
- Kolle, S., Stojkovic, M., Prella, K., Waters, M., Wolf, E. & Sinowatz, F. (2001). Growth hormone (GH)/GH receptor expression and GH-mediated effects during early bovine embryogenesis. *Biol. Reprod.* **64**, 1826–34.
- Kolle, S., Stojkovic, M., Boie, G., Wolf, E. & Sinowatz, F. (2002). Growth hormone inhibits apoptosis in *in vitro* produced bovine embryos. *Mol. Reprod. Dev.* **61**, 180–6.
- Laporta, J., Driver, A. & Khatib, H. (2011). Short communication: expression and alternative splicing of POU1F1 pathway genes in preimplantation bovine embryos. *J. Dairy Sci.* **94**, 4220–3.
- Leidenfrost, S., Boelhauve, M., Reichenbach, M., Gungor, T., Reichenbach, H.D., Sinowatz, F., Wolf, E. & Habermann, F.A. (2011). Cell arrest and cell death in mammalian preimplantation development: lessons from the bovine model. *PLoS One* **6**, e22121.
- Levy, R.R., Cordonier, H., Czyba, J.C. & Guerin, J.F. (2001). Apoptosis in preimplantation mammalian embryo and genetics. *Ital. J. Anat. Embryol.* **106**(2 Suppl 2), 101–8.

- Madeja, Z.E., Warzych, E., Peippo, J., Lechniak, D. & Switonski, M. (2009). Gene expression and protein distribution of leptin and its receptor in bovine oocytes and preattachment embryos produced *in vitro*. *Animal* **3**, 568–78.
- Nakasato, M., Shirakura, Y., Ooga, M., Iwatsuki, M., Ito, M., Kageyama, S., Sakai, S., Nagata, M. & Aoki, F. (2006). Involvement of the STAT5 signaling pathway in the regulation of mouse preimplantation development. *Biol. Reprod.* **75**, 508–17.
- Pers-Kamczyc, E., Warzych, E., Peippo, J. & Lechniak, D. (2010). Growth hormone exerts no effect on the timing of the first zygotic cleavage in cattle. *Theriogenology* **74**, 581–95.
- Robert, C., McGraw, S., Massicotte, L., Pravetoni, M., Gandolfi, F. & Sirard, M.A. (2002). Quantification of housekeeping transcript levels during the development of bovine preimplantation embryos. *Biol. Reprod.* **67**, 1465–72.
- Teglund, S., McKay, C., Schuetz, E., van Deursen, J.M., Stravopodis, D., Wang, D., Brown, M., Bodner, S., Grosveld, G. & Ihle, J.N. (1998). Stat5a and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell* **93**, 841–50.
- Wakao, H., Gouilleux, F. & Groner, B. (1994). Mammary gland factor (MGF) is a novel member of the cytokine regulated transcription factor gene family and confers the prolactin response. *EMBO J.* **13**, 2182–91.
- Wolf, E., Arnold, G.J., Bauersachs, S., Beier, H.M., Blum, H., Einspanier, R., Frohlich, T., Herrler, A., Hiendleder, S., Kolle, S., Prella, K., Reichenbach, H.D., Stojkovic, M., Wenigerkind, H. & Sinowatz, F. (2003). Embryo-maternal communication in bovine - strategies for deciphering a complex cross-talk. *Reprod. Domest. Anim.* **38**, 276–89.
- Wright, R.W., Jr. & Bondioli, K.R. (1981). Aspects of *in vitro* fertilization and embryo culture in domestic animals. *J. Anim. Sci.* **53**, 702–29.