



Archived at the Flinders Academic Commons:

<http://dspace.flinders.edu.au/dspace/>

'This is the peer reviewed version of the following article:

FULL CITATION,

which has been published in final form at

<http://dx.doi.org/>

PUBLISHER SET STATEMENT/MADE AVAILABLE WITH xxx PERMISSION (IF APPLIC.) This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving'. //// "This is an Accepted Manuscript of an article published by Taylor & Francis in [JOURNAL TITLE] on [date of publication], available online: [http://www.tandfonline.com/\[Article DOI\]](http://www.tandfonline.com/[Article DOI])." //// © <year> Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

COPYRIGHT YEAR/HOLDER Copyright (201?) John Wiley & Sons, Inc. All rights reserved.



Archived at the Flinders Academic Commons:

<http://dspace.flinders.edu.au/dspace/>

'This is the peer reviewed version of the following article:
Weinstein, S.A., Everest, E., Purdell-Lewis, J., Harrison, M.,
Tavender, F., Alfred, S., Marrack, L., Davenport-Klunder,
C., Wearn, N., White, J., Neurotoxicity with persistent
unilateral ophthalmoplegia from envenoming by a wild
inland taipan (*Oxyuranus microlepidotus*, Elapidae) in
remote outback South Australia, *Toxicon* (2017),
doi:10.1016/j.toxicon.2017.07.006

which has been published in final form at

<http://dx.doi.org/10.1016/j.toxicon.2017.07.006>

© Crown. This manuscript version is made available under
the CC-BY-NC-ND 4.0 license [http://creativecommons.org/
licenses/by-nc-nd/4.0/](http://creativecommons.org/licenses/by-nc-nd/4.0/)

Accepted Manuscript

Neurotoxicity with persistent unilateral ophthalmoplegia from envenoming by a wild inland taipan (*Oxyuranus microlepidotus*, Elapidae) in remote outback South Australia

Scott A. Weinstein, Evan Everest, Jeremy Purdell-Lewis, Michael Harrison, Fiona Tavender, Sam Alfred, Liz Marrack, Chris Davenport-Klunder, Neralie Wearn, Julian White

PII: S0041-0101(17)30211-8

DOI: [10.1016/j.toxicon.2017.07.006](https://doi.org/10.1016/j.toxicon.2017.07.006)

Reference: TOXCON 5665

To appear in: *Toxicon*

Received Date: 11 June 2017

Revised Date: 3 July 2017

Accepted Date: 5 July 2017

Please cite this article as: Weinstein, S.A., Everest, E., Purdell-Lewis, J., Harrison, M., Tavender, F., Alfred, S., Marrack, L., Davenport-Klunder, C., Wearn, N., White, J., Neurotoxicity with persistent unilateral ophthalmoplegia from envenoming by a wild inland taipan (*Oxyuranus microlepidotus*, Elapidae) in remote outback South Australia, *Toxicon* (2017), doi: 10.1016/j.toxicon.2017.07.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Neurotoxicity with persistent unilateral ophthalmoplegia from envenoming by a wild inland taipan (*Oxyuranus microlepidotus*, Elapidae) in remote outback South Australia.

Scott A Weinstein^{1,2*}, Evan Everest^{3,4}, Jeremy Purdell-Lewis⁵, Michael Harrison⁵, Fiona Tavender⁴, Sam Alfred⁶, Liz Marrack⁵, Chris Davenport-Klunder⁵, Neralie Wearn⁴, Julian White^{1,2}

¹Toxinology Department, Women's and Children's Hospital, 72 King William Street, North Adelaide, South Australia 5006; ²Department of Paediatrics and Reproductive Medicine, University of Adelaide School of Medicine, 30 Frome Street, Adelaide, South Australia 5005; ³Intensive and Critical Care Unit, Flinders Medical Centre, Bedford Park, South Australia 5042; ⁴MedSTAR Retrieval Service, South Australian Ambulance Service; ⁵Royal Flying Doctor Service, Central Operations, 1 Tower Road, Adelaide Airport, South Australia 5950; ⁶Emergency Department, Royal Adelaide Hospital, North Terrace, Adelaide 5000

Abstract

Introduction

A case of life threatening envenoming by a wild specimen of the inland taipan, *Oxyuranus microlepidotus*, is described. There have been 11 previously well-documented envenomings by *O. microlepidotus*, but only 2 were inflicted by wild snakes. Envenomed patients have presented predominantly with defibrinating coagulopathy and neurotoxicity.

Case Report

The victim was seeking to observe members of an isolated population of this species and was envenomed while attempting to photograph an approximately 1.5 m specimen. He reported feeling "drowsiness" and blurred vision that

*Corresponding author: P 61-8-8-161-8044; e-mail herptoxmed@msn.com

progressed to ptosis; he later developed dysphagia and dysarthria. The patient was treated with 1 vial of polyvalent antivenom, which was later followed with an additional two vials of taipan monovalent. He was intubated during retrieval, and recovered after 3 days of intensive care. He had a right ophthalmoplegia that persisted for approximately 1 week post-envenoming. Despite a positive 20-minute whole blood clotting test, defibrination coagulopathy was absent, and there was no myotoxicity, or acute kidney injury.

Discussion

Physicians presented with a patient envenomed by *O. microlepidotus* should remain cognizant of the possible variability of medically important venom toxins in some populations of this species. Some patients seriously envenomed by this species may develop persistent cranial nerve palsies. When clinically indicated, prompt provision of adequate antivenom is the cornerstone of managing *O. microlepidotus* envenoming. Rapid application of pressure-bandage immobilization and efficient retrieval of victims envenomed in remote locales, preferably by medically well-equipped aircraft, probably improves the likelihood of a positive outcome.

Introduction

The genus *Oxyuranus* contains 3 species of highly venomous elapid snakes. Of these, the coastal taipan (*O. scutellatus*) has been most medically important, although in comparison to the early Twentieth Century, is now a less common cause of serious envenoming in Australia. However, the Papuan taipan, *O. s. canni*, has major medical importance in Papua New Guinea. It is noteworthy that some authors consider *O. s. canni* synonymous with *O. scutellatus*, but this

impression is probably based on a single study (Wüster et al. 2005) that investigated several mitochondrial DNA sequences derived from a single specimen of *O. scutellatus*, and from two specimens of *O. s. canni*. Wüster et al. (2005) appropriately indicated that for several reasons (e.g. the limited sample size), further investigation would be necessary in order to confirm synonymy of *O. s. canni* with *O. scutellatus*. We agree, and although their synonymy may soon be confirmed, we prefer to reference the subspecies until further supporting evidence is published.

There have not yet been any documented envenomings by the newly discovered western desert taipan, *O. temporalis*. The inland taipan, *Oxyuranus microlepidotus*, attains an average adult length of 1.5 m, and occurs in remote inland regions of channel country (south-western Queensland with extension into sections of north-eastern South Australia and north-western New South Wales) and surrounding areas of dunes, as well as gibber plains (or, 'stony downs', flat, pebble/stone-covered plains) near the junction of Queensland, New South Wales and South Australia (Mirtschin et al., 2017). *Oxyuranus microlepidotus* venom is well known for having the so far highest experimental lethal potency (subcutaneous LD₅₀ 0.025 mg/kg; Broad et al., 1979) of any tested snake venom. However, experimental venom lethal potency tested in mice doesn't reliably predict medical risk, and there have been no fatalities among the handful of patients seriously envenomed by *O. microlepidotus* (Mirtschin et al., 2017). Most patients envenomed by *O. microlepidotus* have presented with defibrinating coagulopathy and neurotoxicity.

We report here a case of an *O. microlepidotus* envenoming that caused progressive and persistent neurotoxicity without defibrinating coagulopathy.

The case also highlights some of the difficulties that may be encountered when assessing a victim bitten by a highly venomous snake in a remote location.

Clinical Record

A 28 yr old male professional snake catcher sought to photograph a specimen of inland taipan (*O. microlepidotus*) in a location (north of Coober Pedy, South Australia; 526 miles north of Adelaide; general regional coordinates: 29.0139° South, 134.7533° East) known to have a possibly isolated population of this species. An approximately 1.5 m specimen of *O. microlepidotus* was found (photographs of the specimen have been reviewed and the taxon verified by the senior author) in late afternoon (at about 1730 hours), and carefully positioned for photography by the victim. The snake moved away from the selected site, and the victim gently pulled the snake back by the tail. The snake 'looped' backwards, and the victim re-positioned it by turning it while maintaining the grip on its tail. As the victim's companion prepared to take the photo, the victim noted that he was bleeding from the right forearm, and understood that he had been grazed or 'raked' by a glancing bite or single fang scratch. Within a few minutes he felt nauseated, light-headed and stated that he felt "drowsy"; he and his companion promptly applied a pressure-bandage immobilization (PBI) then sought help at a nearby hospital. It should be noted that the area is serviced by community and family physicians, and has no intensive care facilities. Critically ill patients are moved by a statewide specialist medical retrieval service to a tertiary referral hospital in the state's capital city (a 2hr flight to the south).

The victim was evaluated approximately 45 minutes after the bite, and the treating physician consulted with the South Australian retrieval service

(MedSTAR) who then contacted the on-call toxicologist (SAW). At approximately one hour post-envenoming, the patient began vomiting, and reported increasingly blurred vision; early bilateral ptosis and slurred speech were noted, but there was no evidence of respiratory compromise. Given the local unavailability of formal laboratory services, a 20-minute whole blood clotting test (20WBCT) was performed in a glass tube and remained non-clotted after 25+ minutes, a 'positive' result suggesting the presence of coagulopathy. Because of the limited clinical personnel in the Coober Pedy local hospital, urgency of the patient's condition and process of arranging retrieval, a blood sample from a healthy volunteer was not tested as a 20WBCT control sample. Two hours after the envenoming, one vial of Polyvalent snake antivenom (Seqirus®, Melbourne, VIC, Australia; contains 12000 *in vitro* neutralizing units for *O. scutellatus* venom; this is equal to the neutralizing capacity of monospecific taipan antivenom) was administered IV to the patient, and a two-stage air retrieval from Coober Pedy was arranged. The Polyvalent antivenom infusion was completed, and the patient departed with the first aircraft 3.5 hours after the 'bite'. In order to slow the uptake of any possible sequestered venom, the PBI was intentionally left in place during the entire retrieval process. As the patient was transferred from the first to the second aircraft mid-retrieval (5 hours after the envenoming), his vital signs remained within normal limits, but he was unable to significantly crease his forehead, exhibited progressive dysarthria, dysphagia with increasing drooling, and severe ptosis. In light of this progression, the patient was given rocuronium, intubated, ventilated and provided with an additional two vials of Taipan antivenom (Seqirus®, Melbourne, VIC, Australia; as noted previously, each vial also contains 12000 *in*

vitro neutralizing units for *O. scutellatus* venom), 5.5 hours post-bite. The retrieval was further complicated by an aircraft malfunction necessitating transfer to a third aircraft.

After arrival at the tertiary medical center, 10.3 hours post-bite, formal laboratory investigations revealed no evidence of coagulopathy including an absence of elevated D-dimer. Aside from a transiently elevated C-reactive protein (60 mg/L, peaked approximately 40-42 hr post-envenoming; likely related to moderate local inflammation at the wound site), all laboratory investigations remained within normal limits, as did the patient's vital signs. He remained intubated for 48 hours, and exhibited ptosis, as well as ophthalmoplegia post-extubation. Deep tendon reflexes were reduced (1+) in upper and lower extremities. Both pupils reacted sluggishly, but his right pupil appeared qualitatively more so; the patient was unable to tolerate comprehensive testing of his right eye, and exhibited right-sided photophobia. He complained of nausea and growing ocular fatigue with further attempted examination/testing. He was discharged five days after admission with a persistent right ophthalmoplegia; ophthalmological and neurological examination prior to discharge was otherwise unremarkable. The patient reported a full recovery 7 days after the envenoming.

Discussion

Approximately 11 bites and/or envenomings by *O. microlepidotus* have been reported (only 2/11 were inflicted by wild specimens) (Trinca, 1969; Mirtschin et al., 1984; Covacevitch et al., 1995; White *et al.*, 1992; Smith and Ambikapathy, 1992; Johnston et al., 2017), and a number of formally undocumented bites by

this species have also occurred. Almost all of the previously documented patients presented with defibrination coagulopathy, most also had paralysis, and at least one had myotoxicity. Our patient only developed neurotoxicity, that manifested as delayed descending flaccid paralysis commencing with cranial nerve involvement about 1 hour post-bite, then progressing approximately 4-5 hours post-bite.

The results of a 20WBCT suggested coagulopathy, but instrumented testing about 9 hours later did not find any fibrin-split products, and the aPTT and INR were normal. Relevantly, Smith and Ambikapathy (1992) reported an *O. microlepidotus* envenoming that featured persistent neurotoxicity with a prolonged aPTT, thrombocytopenia and giant platelets. Their patient received one vial of Polyvalent antivenom, and the aPTT returned to normal 4 hours later; the fibrin split products remained within normal limits, but the thrombocytopenia remained abnormal for 10 days. They did not specifically mention the INR, or fibrinogen, although omission of the INR may have been due to the mixed use of the term at that time because 'INR' was standardized after re-assessment of the International Sensitivity Index (ISI) in 1983 (WHO, 1983), and for several years thereafter its standardization was discussed or reviewed by several authors (e.g. see Hirsh et al., 1989). Thus, the report by Smith and Ambikapathy (1992) is difficult to interpret because it could suggest a possible transient anticoagulant coagulopathy with thrombocytopenia and altered platelet morphology. In a recent study of 33 patients systemically envenomed by *Oxyuranus* spp., 31 (93%) developed defibrinating coagulopathy and the median time for the INR to decrease to less than 2.0 was 12.3 hours (IQR: 12.3–18.7 hours; range 8.5–42 hours; Johnston et al., 2017). Recent investigation suggests

that D-dimer has a half-life of approximately 15 hours (Rühl et al., 2015). Smith and Ambikapathy (1992) speculated that the venom had a “direct marrow effect”, but their report contains no further information to support their hypothesis. Therefore, the possibility that our patient developed a transient anticoagulant coagulopathy cannot be excluded. However, given the clinical evidence in our case and other well-documented *O. microlepidotus* envenoming, in our opinion a false positive 20WBCT is more likely. The 20WBCT is commonly used globally in the rural tropics where medical laboratories may be difficult to access, and has certainly been a useful test in a variety of clinical settings (e.g. see Sano-Martins et al., 1994; Gaus et al. 2013; Iliyasa et al., 2015). However, the assay has variable sensitivity and specificity and thus can yield false negatives or positives under some clinical circumstances (Isbister et al., 2013; Ratnayake et al., 2017). In Australian rural settings without laboratory instrumentation it may be the only option, but in our case the available evidence suggests that it probably yielded a false positive, as has been occasionally noted after coagulopathic envenoming from some Australian elapids (White, 2013). It should be stressed that failure of the 20WBCT may often be due to “operator error” (e.g. extent of training of personnel performing the test, assay temperature, use of incorrect test vessels, misinterpretation of results, etc.; Isbister et al., 2013; Auerbach et al., 2016; Ratnayake et al., 2017).

It is also noteworthy that although our patient had correctly applied a PBI within a few minutes after contact with the taipan, he developed signs and symptoms of systemic envenoming in less than 30 minutes. While it is possible that early provision of antivenom prevented a more rapidly progressive paralysis, respiratory support was still deemed clinically necessary for two days. In the

aforementioned series of *Oxyuranus* spp. envenoming, Johnston et al. (2017) reported 7 bites by captive *O. microlepidotus*; 6 patients were envenomed, and these all received antivenom. All developed consumption coagulopathy, 57% had neurotoxicity, and 43% had an acute kidney injury (AKI) and/or microangiopathic hemolytic anemia (MAHA); none developed myotoxicity (Johnston et al., 2017).

Studied samples of *O. microlepidotus* venom contain paradoxin, which is closely similar to taipoxin, the heteropentameric presynaptic neurotoxin described from coastal taipan (*O. scutellatus*) venom, several postsynaptic neurotoxins, procoagulants that function as Va and Xa homologues, myotoxic phospholipases A₂, natriuretic peptides, and numerous other components (Fohlman, 1979; Hamilton et al., 1980; Bell et al., 1988; Crachi et al., 1999; Fry et al., 2005; Clarke et al., 2006; Vink et al., 2010; Earl et al., 2012). Venom composition variability has been thoroughly documented in several medically important viperids and elapids (e.g. Minton and Weinstein, 1986; Chippaux et al., 1991). It is probable that the specimen that envenomed our patient secreted venom that either lacked, or contained much lower concentrations, of several medically important toxins (procoagulants) commonly found in most tested venoms of this taxon.

Johnston et al. (2017) asserted that one vial of antivenom “should be sufficient to treat most taipan envenoming because all of the detectable venom antigens were bound by antibody present in one vial of antivenom”. This view has been previously criticised (Weinstein et al., 2016) because of problems and pitfalls of the methods (ELISA assays) used to support their recommendations. Treatment must be provided per individual patient with consideration for their co-morbidities and individual risks; guidelines provide a starting reference, but,

ultimately, one “size” does not necessarily “fit all”. Coastal taipan (*O. scutellatus*) envenoming can rapidly (in several minutes) clinically progress after removal of the PBI (Southern et al., 1996; SAW, personal observations), and thus it is essential to initially provide adequate antivenom, as clinically indicated.

The persistence of a unilateral ophthalmoplegia in this case is uncommon and not readily explained, as venom paralytic effects are typically bilateral because of the systemic distribution of venom, a non-systemic exception being localised effects from paralysis ticks attached close to a cranial nerve (White, 1995). Delayed resolution of bilateral ptosis and partial ophthalmoplegia does occur in some patients who develop paralytic features after Australian elapid envenoming and longer term residual hyposmia or anosmia also may occur, and not always in association with neurotoxic envenoming (Sutherland and Tibballs, 2001; White, 2010; 2013; Sethi et al., 2017). A patient seriously envenomed by a common tiger snake (*Notechis scutatus*, Elapidae) experienced a similarly persistent unilateral (left) ophthalmoplegia that lasted for at least ten months and did not fully resolve (lost to follow-up after one year; SAW personal observations). Although it is speculative to consider possible causes of these patients’ persistent unilateral ophthalmoplegia, it is noteworthy that it may present in patients with internuclear ophthalmoplegia (INO) that arises from an interruption of fibres connecting the oculomotor nuclei, and can be unilateral or bilateral (Karski et al., 2013). Unilateral INOs are most commonly associated with ischaemia (Kaski et al., 2013). Unilateral ophthalmoplegia also has been rarely reported in other disease states such as acute myeloid leukemia (Mott and Carlson, 2012), ophthalmoplegic migraine (Patel et al., 2015), and occasionally with various central nervous system infarcts.

In conclusion, we have described a serious envenoming inflicted by a wild *O. microlepidotus* that featured progressive neurotoxicity and without defibrinating coagulopathy. Management of our patient emphasized several important observations: the 20WBCT may give false positive results and must be considered together with clinical observations of the patient; prompt and correct application of PBI may delay onset of venom effects; early administration of antivenom may provide a clinically important window for successful patient retrieval from remote locations; prompt intubation and airway protection should be performed whenever possible, as clinically needed, especially during retrievals from remote locales; some populations of *O. microlepidotus* may produce venom without clinically significant levels of defibrinating procoagulants, and possibly with anticoagulant components; clinically indicated antivenom should be administered as early as possible when medically feasible, and the dose should be based on the severity of the individual patient's envenoming, as well as the patient's therapeutic response.

Acknowledgements

We gratefully thank our patient who generously provided informed consent for our reporting this case. We also gratefully acknowledge the important contributions of Dr John Woodall, Chief Medical Officer, RFDS, Central Operations, and Holly Carroll, the on-call pharmacist at the Royal Adelaide Hospital who facilitated prompt procurement of taipan antivenom.

Declaration of Interest Statement

One of the authors reports the following: Julian White does provide advice to the antivenom manufacturer (Seqirus, formerly CSL, Ltd), as part of a contract between his employing hospital and Seqirus. He is not paid by Seqirus and Seqirus has no input or influence on any of his reports, clinical practice or comments.

All of the remaining authors declare no conflicts of interest.

References

- Auerbach PS, Cushing TA, Stuart Harris N. 2016. Auerbach's Wilderness Medicine E-Book, 7th Edition, Elsevier, Waltham, pp. 2280.
- Bell KL, Sutherland SK, Hodgson WC. 1988. Some pharmacological studies of venom from the inland taipan (*Oxyuranus microlepidotus*). *Toxicon* 36, 63-74.
- Broad AJ, Sutherland SK, Coulter AR. 1979. The lethality in mice of dangerous Australian and other snake venoms. *Toxicon* 17, 661-664.
- Chippaux JP, Williams V, White J. 1991. Snake venom variability: methods of study, results and interpretation. *Toxicon* 29, 1279-1303.
- Clarke C, Kuruppu S, Reeve S, Ian Smith A, Hodgson WC. 2006. Oxylepitoxin-1, a reversible neurotoxin from the venom of the inland taipan (*Oxyuranus microlepidotus*). *Peptides* 27, 2655-2660.
- Covacevich J, Pearn J, White J. 1995. The World's most dangerous snake. In: Pearn J, Covacevich J, Eds., *Venoms and Victims* Brisbane, The Queensland Museum and Amphion Press, pp. 111-120.

Crachi MT, Hammer LW, Hodgson WC. 1999. The effects of antivenom on the *in vitro* neurotoxicity of venoms from the taipans *Oxyuranus scutellatus*, *Oxyuranus microlepidotus* and *Oxyuranus scutellatus canni*. *Toxicon* 37, 1771-1778.

Earl ST, Richards R, Johnson LA, Flight S, Anderson S, Liao A, de Jersey J, Masci PP, Lavin MF. 2012. Identification and characterisation of Kunitz-type plasma kallikrein inhibitors unique to *Oxyuranus* sp. snake venoms. *Biochimie* 94, 365-373.

Fohlman, J. 1979. Comparison of two highly toxic Australian venoms: the taipan (*Oxyuranus s. scutellatus*) and the fierce snake (*Parademansia microlepidotus*). *Toxicon* 17, 170-172.

Fry BG, Wickmaratana JC, Lemme S, Beuve A, Garbers D, Hodgson WC, Alewood P. 2005. Novel natriuretic peptides from the venom of the inland taipan (*Oxyuranus microlepidotus*): isolation, chemical and biological characterization. *Biochem. Biophys. Res. Commun.* 327, 1011-1015.

Gaus DP, Herrera DF, Troya CJ, Guevara AH. 2013. Management of snakebite and systemic envenomation in rural Ecuador using the 20-minute whole blood clotting test. *Wild. Environ. Med.* 24, 345-350.

Hamilton RC, Broad AJ, Sutherland SK. 1980. Ultrastructural effects of the venom of the small-scaled snake (*Parademansia microlepidotus*) on the nerve terminals of the rat diaphragm. *Aust. J. Exp. Biol. Med. Sci.* 58, 377-380.

Hirsh J, Poller L, Deykin D, Levine M, Dalen JE. 1989. Optimal therapeutic range for oral anticoagulants. *Chest* 95 (2_Supplement), 12S-18S, doi:10.1378/chest.95.2_Supplement.12S.

- Iliyasu G, Tihamiyu AB, Daiyab FM, Tambuwal SH, Habib ZG, Habib AG. 2015. Effect of distance and delay in access to care on outcome of snakebite in rural north-eastern Nigeria. *Rural Remote Health* 15, 3496.
- Isbister GK, Maduwage K, Shahmy S, Mohamed F, Abeysinghe C, Karunathilake CA, Buckley NA. 2013. Diagnostic 20-min whole blood clotting test in Russell's viper envenoming delays antivenom administration. *QJM* 106, 925-932.
- Johnston CI, Ryan NM, O'Leary MA, Brown SGA, Isbister GK. 2017. Australian taipan (*Oxyuranus* spp.) envenoming: clinical effects and potential benefits of early antivenom therapy – Australian Snakebite Project (ASP-25). *Clin. Toxicol. (Philadelphia)* 55, 115-122.
- Kaski D, Domínguez RO, Bronstein AM. 2013. Motion diplopia in an isolated unilateral internuclear ophthalmoplegia: A new neurological symptom? *Clin. Neurol. Neurosurg.* 115, 636-637.
- Minton SA, Weinstein SA. 1986. Geographic and ontogenetic variation in venom of the Western diamondback rattlesnake (*Crotalus atrox*). *Toxicon* 24, 71-80.
- Mirtschin PJ, Rasmussen AR, Weinstein SA. 2017. Australia's Dangerous Snakes: Identification, Biology, Envenoming. Melbourne, CSIRO (in press).
- Mirtschin PJ, Crowe GR, Thomas MW. 1984. Envenomation by the inland taipan, *Oxyuranus microlepidotus*. *Med J. Aust.* 22, 850-851.
- Mott J, Carlson MD. 2012. Acute unilateral ophthalmoplegia as the presenting sign of acute myeloid leukemia in a 15-month-old girl. *Pediatr. Neurol.* 47, 366-368.
- Patel R, Harper-Shankie M, Patel E, Sivaswamy L. 2015. Unilateral ptosis. *J. Pediatr.* 167, 1160.

Ratnayake I, Shihana F, Dissanayake DM, Buckley NA, Maduwage K, Isbister GK. 2017. Performance of the 20-minute whole blood clotting test in detecting venom induced consumption coagulopathy from Russell's viper (*Daboia russelii*) bites. *Thromb Haemost.* 117, 500-507.

Rühl H, Berens C, Winterhagen A, Müller J, Oldenburg J, Pötzsch B. 2015. Label-free kinetic studies of haemostasis-related biomarkers including D-dimer using autologous serum transfusion. *PLoS One* 10:e0145012. doi: 10.1371/journal.pone.0145012.

Sano-Martins IS, Fan HW, Castro SC, Tomy SC, Franca FO, Jorge MT, Kamiguti AS, Warrell DA, Theakston RD. 1994. Reliability of the simple 20 minute whole blood clotting test (WBCT20) as an indicator of low plasma fibrinogen concentration in patients envenomed by *Bothrops* snakes. Butantan Institute Antivenom Study Group. *Toxicon* 32, 1045-1050.

Sethi M, Cook M, Winkel KD. 2017. Persistent anosmia and olfactory bulb atrophy after mulga (*Pseudechis australis*) snakebite. *J. Clin. Neurosci.* (in press; <http://dx.doi.org/10.1016/j.jocn.2015.12.019>).

Smith KGC, Ambikapathy A. 1992. Envenomation by the small-scaled snake: the world's most venomous snake. *Med. J. Aust.* 157, 140.

Southern DA, Callanan VI, Gordon GS. 1996. Severe envenomation by the taipan (*Oxyuranus scutellatus*). *Med. J. Aust.* 165, 662-664.

Sutherland SK, Tibballs J (2001) *Australian Animal Toxins*. 2nd Edition, Oxford University Press, Melbourne, pp. 856.

Trinca JC. 1969. Report of Recovery from taipan Bite. *The Med. J. Aust.* 1, 514-516.

- Vink S, Jin AH, Poth KJ, Head GA, Alewood PF. 2010. Natriuretic peptide drug leads from snake venom. *Toxicon* 59, 434-445.
- Weinstein SA, White J, Ou J, Haiart S, Galluccio S. 2016. Reply to Isbister and Page: Further discussion of an illuminated case of presumed brown snake (*Pseudonaja* spp.) envenoming. *Clin. Toxicol.* 53, 926-927.
- White J. 1995. Clinical toxicology of tick bites. In: Meier J, White J, Eds., *Handbook of Clinical Toxicology of Animal Venoms and Poisons*. CRC Press: Boca Raton, pp. 191-204.
- White J. 2010. Envenomation; prevention and treatment in Australia. In: Mackessy SP, Ed., *Handbook of Venoms and Toxins of Reptiles*. CRC Press: Boca Raton, pp. 423-451.
- White J. 2013. *A Clinician's Guide to Australian Venomous Bites and Stings*. bioCSL Pty Ltd, Parkville, pp. 70-71.
- White J, Duncan B, Wilson C, Williams V, Lloyd J. 1992. Coagulopathy following Australian elapid snakebite; a review of 20 cases. In: Gopalakrishnakone P, Tan CK, Eds., *Recent Advances in Toxinology Research (volume 1)*. National University of Singapore, Singapore, pp. 337-344.
- World Health Organization (WHO). 1983. WHO Expert Committee on Biological Standardization. 33rd Report. *Wld Hlth Org. Techn. Rep. Ser.* 687, 81.
- Wüster W, Dumbrella AJ, Hay C, Pook CE, Williams DJ, Fry BG. 2005. Snakes across the Strait: trans-Torresian phylogeographic relationships in three genera of Australasian snakes (Serpentes: Elapidae: *Acanthophis*, *Oxyuranus*, and *Pseudechis*). *Mol. Phylogen. Evol.* 34, 1-14.

Neurotoxicity with persistent unilateral ophthalmoplegia from envenoming by a wild inland taipan (*Oxyuranus microlepidotus*, Elapidae) in remote outback South Australia

Highlights

- Documentation of a serious envenoming by a wild inland taipan (*Oxyuranus microlepidotus*) in the remote Australian outback
- Defibrinating coagulopathy was absent; a 20 minute whole blood clotting test may have been a false positive, or possibly indicated a transient anti-coagulant coagulopathy
- The patient developed progressive neurotoxicity; had a complicated retrieval from the outback; was ultimately treated with Polyvalent and Taipan monospecific antivenom
- An initial bilateral ophthalmoplegia was notably more pronounced on the right side, and persisted as a unilateral right sided ophthalmoplegia
 - Full recovery achieved after 1 week