

# 1 Next generation sequencing to investigate genomic diversity in Caryophyllales

2 Boas Pucker<sup>1,2\*</sup>, Tao Feng<sup>1,3</sup>, Samuel F. Brockington<sup>1,2</sup>

3 1 Evolution and Diversity, Plant Sciences, University of Cambridge, Cambridge, United Kingdom

4 2 Genetics and Genomics of Plants, CeBiTec & Faculty of Biology, Bielefeld University, Bielefeld,  
5 Germany

6 3 Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan, China

7 \* corresponding author: Boas Pucker, [bpucker@cebitec.uni-bielefeld.de](mailto:bpucker@cebitec.uni-bielefeld.de)

8

9 BP: [bpucker@cebitec.uni-bielefeld.de](mailto:bpucker@cebitec.uni-bielefeld.de), ORCID: 0000-0002-3321-7471

10 TF: [fengtao@wbgcas.cn](mailto:fengtao@wbgcas.cn), ORCID: 0000-0002-0489-2021

11 SFB: [sb771@cam.ac.uk](mailto:sb771@cam.ac.uk), ORCID: 0000-0003-1216-219X

12

13 Key words: whole genome sequencing, genome assembly, anthocyanin, betalain, *Kewa caespitosa*,  
14 *Macarthuria australis*, *Pharnaceum exiguum*, *Caryophyllales*

15

## 16 Abstract

17 Caryophyllales are a highly diverse and large order of plants with a global distribution. While some  
18 species are important crops like *Beta vulgaris*, many others can survive under extreme conditions.  
19 This order is well known for the complex pigment evolution, because the red pigments anthocyanin  
20 and betalain occur with mutual exclusion in species of the Caryophyllales. Here we report about  
21 genome assemblies of *Kewa caespitosa* (Kewaceae), *Macarthuria australis* (Macarthuraceae), and  
22 *Pharnaceum exiguum* (Molluginaceae) which are representing different taxonomic groups in the  
23 Caryophyllales. The availability of these assemblies enhances molecular investigation of these species  
24 e.g. with respect to certain genes of interest.

25

26

27

## 28 **Introduction**

29 Caryophyllales form the largest flowering plant order and are recognized for their outstanding ability  
30 to colonise extreme environments. Examples are the evolution of Cactaceae in deserts, extremely  
31 fast radiation [1–3] e.g. in arid-adapted Aizoaceae and in carnivorous species in nitrogen-poor  
32 conditions. Caryophyllales harbor the greatest concentration of halophytic plant species and display  
33 repeated shifts to alpine and arctic habitats in Caryophyllaceae and Montiaceae. Due to these  
34 extreme environments, species exhibit many adaptations [2–4] such as specialized betalain pigments  
35 to protect photosystems in high salt and high light conditions [5]. There are several examples for  
36 repeated evolution in the Caryophyllales e.g. leaf and stem succulence for water storage, various  
37 mechanisms for salt tolerance, arid-adapted C<sub>4</sub> and CAM photosynthesis [4], and insect trapping  
38 mechanisms to acquire nitrogen [6].

39 In addition, to their fascinating trait evolution, the Caryophyllales are well known for important crops  
40 and horticultural species like sugar beet, quinoa and spinach. Most prominent is the genome  
41 sequence of *Beta vulgaris* [7] which was often used as a reference for studies within Caryophyllales  
42 [7–10]. In addition, genomes of *Spinacia oleracea* [7,11], *Dianthus caryophyllus* [12], *Amaranthus*  
43 *hypochondriacus* [13], *Chenopodium quinoa* [14] were sequenced. Besides *Carnegiea gigantea* and  
44 several other cacti [15], recent genome sequencing projects were focused on crops due to their  
45 economical relevance. However, genome sequences of other species within the Caryophyllales, are  
46 needed to provide insights into unusual patterns of trait evolution.

47 The evolution of pigmentation is known to be complex within the Caryophyllales [8] with a single  
48 origin of betalain and at least three reversals to anthocyanin pigmentation. The biosynthetic  
49 pathways for betalain and anthocyanin pigmentation are both well characterized. While previous  
50 studies have demonstrated that the genes essential for anthocyanin synthesis persists in betalain  
51 pigmented taxa, the fate of the betalain pathway in the multiple reversals to anthocyanin  
52 pigmentations is unknown. Here, we sequenced three species from different families to contribute to  
53 the genomic knowledge about Caryophyllales: *Kewia caespitosa* (Kewaceae), *Macarthuria australis*  
54 (Macarthuraceae), and *Pharnaceum exiguum* (Molluginaceae) were selected as representatives of  
55 anthocyanic lineages within the predominantly betalain pigmented Caryophyllales. *K. caespitosa* and  
56 *P. exiguum* are examples of putative reversals from betalain pigmentation to anthocyanic  
57 pigmentation, while *Macarthuria* is a lineage that diverged before the inferred origin of betalain  
58 pigmentation [8].

59 Several transcript sequences of the three plants investigated here were assembled as part of the 1KP  
60 project [16]. Since the sampling for this transcriptome project was restricted to leaf tissue, available  
61 sequences are limited to genes expressed there. Here we report three draft genome sequences to

62 complement the available gene set and to enable analysis of untranscribed sequences like  
63 promoters, regulatory elements, pseudogenes, and transposable elements.

64

65

## 66 **Material & Methods**

### 67 **Plant material**

68 The seeds of *Kewa caespitosa* (Friedrich) Christenh., *Marcarthuria australis* Hügel ex Endl., and  
69 *Pharnaceum exiguum* Adamson were obtained from Millennium Seed Bank (London, UK) and were  
70 germinated at the Cambridge University Botanic Garden. The plants were grown in controlled  
71 glasshouse under conditions: long-day (16 h light and 8 h dark), 20 °C, 60% humidity. About 100 mg  
72 fresh young shoots were collected and immediately frozen in liquid nitrogen. Tissue was ground in  
73 liquid nitrogen using a mortar and pestle. DNA was extracted using the QIAGEN DNeasy Plant Mini Kit  
74 (Hilden, Germany) and RNA was removed by the QIAGEN DNase-Free RNase Set. DNA quantity and  
75 quality were assessed by Nanodrop (ThermoFisher Scientific, Waltham, MA, USA) and agarose gel  
76 electrophoresis. DNA samples were sent to BGI Technology (Hongkong) for library construction and  
77 Illumina sequencing.

78

### 79 **Sequencing**

80 Libraries of *K. caespitosa*, *M. australis*, and *P. exiguum* were sequenced on an Illumina HiSeq X-Ten  
81 generating 2x150nt reads (AdditionalFile 1). Trimmomatic v0.36 [17] was applied for adapter removal  
82 and quality trimming as described previously [18]. Due to remaining adapter sequences, the last 10  
83 bases of each read were clipped. FastQC [19] was applied to check the quality of the reads.

84

### 85 **Genome size estimation**

86 The size of all three investigated genomes was estimated based on k-mer frequencies in the  
87 sequencing reads. Jellyfish v2 [20] was applied for the construction of a k-mer table with parameters  
88 described by [21]. The derived histogram was further analyzed by GenomeScope [21] to predict a  
89 genome size. This process was repeated for all odd k-mer sizes between 17 and 25 (AdditionalFile 2).  
90 Finally, an average value was selected from all successful analyses.

91

## 92 **Genome assembly**

93 The performance of different assemblers on the data sets was tested (AdditionalFile 3, AdditionalFile  
94 4, AdditionalFile 5). While CLC Genomics Workbench performed best for the *M. australis* assembly,  
95 SOAPdenovo2 [22] showed the best results for *K. caespitosa* and *P. exiguum* and was therefore  
96 selected for the final assemblies. To optimize the assemblies, different k-mer sizes were tested as  
97 this parameter can best be adjusted empirically [23]. First, k-mer sizes from 67 to 127 in steps of 10  
98 were evaluated, while most parameters remained on default values (AdditionalFile 6). Second,  
99 assemblies with k-mer sizes around the best value of the first round were tested. In addition,  
100 different insert sizes were evaluated without substantial effect on the assembly quality. In  
101 accordance with good practice, assembled sequences shorter than 500 bp were discarded prior to  
102 downstream analyses. Custom Python scripts [18,24] were deployed for assembly evaluation based  
103 on simple statistics (e.g. N50, N90, assembly size, number of contigs), number of genes predicted by  
104 AUGUSTUS v3.2 [25] *ab initio*, average size of predicted genes, and number of complete BUSCOs  
105 [26]. Scripts are available on github: <https://github.com/bpucker/GenomeAssemblies2018>.

106 BWA-MEM v0.7 [27] was used with the `-m` flag to map all sequencing reads back against the  
107 assembly. REAPR v1.0.18 [28] was applied on the selected assemblies to identify putative assembly  
108 errors through inspection of paired-end mappings and to break sequences at those points.

109 The resulting assemblies were further polished by removal of non-plant sequences. First, all  
110 assembled sequences were subjected to a BLASTn [29] against the sugar beet reference genome  
111 sequence RefBeet v1.5 [7,30] and the genome sequences of *Chenopodium quinoa* [14], *Carnegieia*  
112 *gigantea* [15], *Amaranthus hypochondriacus* [13], and *Dianthus caryophyllus* [12]. Hits below the e-  
113 value threshold of  $10^{-10}$  were considered to be of plant origin. Second, all sequences without hits in  
114 this first round were subjected to a BLASTn search against the non-redundant nucleotide database  
115 nt. Sequences with strong hits against bacterial and fungal sequences were removed as previously  
116 described [18,24]. BLASTn against the *B. vulgaris* plastome (KR230391.1, [31]) and chondrome  
117 (BA000009.3, [32]) sequences was performed to identify and remove sequences from these  
118 organelle subgenomes.

119

## 120 **Assembly quality assessment**

121 Mapping of sequencing reads against the assembly and processing with REAPR [28] was the first  
122 quality control step. RNA-Seq reads (AdditionalFile 7) were mapped against the assemblies to assess  
123 completeness of the gene space and to validate the assembly with an independent data set. STAR  
124 v2.5.1b [33] was used for the RNA-Seq read mapping as previously described [24].

125

## 126 **Genome annotation**

127 RepeatMasker [34] was applied using crossmatch [35] to identify and mask repetitive regions prior to  
128 gene prediction. Masking was performed in sensitive mode (-s) without screening for bacterial IS  
129 elements (-no\_is) and skipping interspersed repeats (-noint). Repeat sequences of the Caryophyllales  
130 (-species caryophyllales) were used and the GC content was calculated per sequence (-gccalc).  
131 Protein coding sequences of transcriptome assemblies (AdditionalFile 7) were mapped to the  
132 respective genome assembly via BLAT [36] to generate hints for the gene prediction process as  
133 previously described [37]. BUSCO v3 [26] was deployed to optimize species-specific parameter sets  
134 for all three species based on the sugar beet parameter set [38]. AUGUSTUS v.3.2.2 [25] was applied  
135 to incorporate all available hints with previously described parameter settings to optimize the  
136 prediction of non-canonical splice sites [37]. Different combinations of hints and parameters were  
137 evaluated to achieve an optimal annotation of all three assemblies. A customized Python script was  
138 deployed to remove all genes with premature termination codons in their CDS or spanning positions  
139 with ambiguous bases. Representative transcripts and peptides per locus were identified based on  
140 maximization of the encoded peptide length. INFERNAL (cmscan) [39] was used for the prediction of  
141 non-coding RNAs based on models from Rfam13 [40].

142 Functional annotation was transferred from *Arabidopsis thaliana* (Araport11) [41] via reciprocal best  
143 BLAST hits as previously described [24]. In addition, GO terms were assigned to protein coding genes  
144 through an InterProScan5 [42]-based pipeline [24].

145

## 146 **Comparison between transcriptome and genome assembly**

147 The assembled genome sequences were compared against previously published transcriptome  
148 assemblies (AdditionalFile 7) in a reciprocal way to assess completeness and differences. BLAT [36]  
149 was used to align protein coding sequences against each other. This comparison was limited to the  
150 protein coding sequences to avoid biases due to UTR sequences, which are in general less reliably  
151 predicted or assembled, respectively [37]. The initial alignments were filtered via filterPSL.pl [43]  
152 based on recommended criteria for gene prediction hint generation to remove spurious hits and to  
153 reduce the set to the best hit per locus e.g. caused by multiple splice variants.

154

155

156

## 157 **Results**

### 158 **Genome size estimation and genome sequence assembly**

159 Prior to the *de novo* genome assembly, the genome sizes of *Kewa caespitosa*, *Macarthuria australis*,  
160 and *Pharnaceum exiguum* were estimated from the sequencing reads (Table 1, AdditionalFile 1). The  
161 estimated genome sizes range from 265 Mbp (*P. exiguum*) to 623 Mbp (*M. caespitosa*). Based on  
162 these genome sizes, the sequencing coverage ranges from 111x (*K. caespitosa*) to 251x (*M. australis*).

163 Different assembly tools and parameters were evaluated to optimize the assembly process  
164 (AdditionalFile 3, AdditionalFile 4, AdditionalFile 5). Sizes of the final assemblies ranged from  
165 254.5 Mbp (*P. exiguum*) to 531 Mbp (*K. caespitosa*) (Table 1, AdditionalFile 8). The best continuity  
166 was achieved for *P. exiguum* with an N50 of 515 Mbp.

167 **Table 1: Genome size estimation and *de novo* assembly statistics.**

	<i>Kewa caespitosa</i>	<i>Macarthuria australis</i>	<i>Pharnaceum exiguum</i>
Accession	GCA_900322205	GCA_900322265	GCA_900322385
Estimated genome size [Mbp]	623	497.5	265
Sequencing coverage	111x	251x	206x
Assembly size (-N)	531,205,354	525,292,167	254,526,612
Number of sequence	55,159	271,872	16,641
N50	28,527	2,804	56,812
Max. sequence length	340,297	211,626	514,701
GC content	38.1%	36.6%	37.4%
Complete BUSCOs	83.6%	44.4%	84.3%
Assembler	SOAPdenovo2	CLC Genomics Workbench v9	SOAPdenovo2
k-mer size	79	Automatic	117

168

169

### 170 **Assembly validation**

171 The mapping of sequencing reads against the assembled sequences resulted in mating rates of 99.5%  
172 (*K. caespitosa*), 98% (*M. australis*), and 94.8% (*P. exiguum*). REAPR identified between 1390 (*P.*  
173 *exiguum*) and 16181 (*M. australis*) FCD errors which were corrected by breaking assembled  
174 sequences. The mapping of RNA-Seq reads to the polished assembly resulted in mapping rates of

175 53.9% (*K. caespitosa*) and 43.1% (*M. australis*), respectively, when only considering uniquely mapped  
176 reads. Quality assessment via BUSCO revealed 83.6% (*K. caespitosa*), 44.4% (*M. australis*), and 84.3%  
177 (*P. exiguum*) complete benchmarking universal single copy ortholog genes (n=1440). In addition,  
178 6.5% (*K. caespitosa*), 21.7% (*M. australis*), and 4.0% (*P. exiguum*) fragmented BUSCOs as well as 9.9%  
179 (*K. caespitosa*), 33.9% (*M. australis*), and 11.7% (*P. exiguum*) missing BUSCOs were identified. The  
180 proportion of duplicated BUSCOs ranges from 1.5% (*K. caespitosa*) to 2.1% (*P. exiguum*). The number  
181 of duplicated BUSCOs was high in *M. australis* (11.8%) compared to both other genome assemblies  
182 (1.5% and 2.1%, respectively).

183

#### 184 **Genome annotation**

185 After intensive optimization (AdditionalFile 9), the polished structural annotation contains between  
186 26,155 (*P. exiguum*) and 80,236 (*M. australis*) protein encoding genes per genome (Table 2). The  
187 average number of exons per genes ranged from 2.9 (*M. australis*) to 6.6 (*K. caespitosa*). Predicted  
188 peptide sequence lengths vary between 241 (*M. australis*) and 447 (*K. caespitosa*) amino acids. High  
189 numbers of recovered BUSCO genes support the assembly quality (Fig. 1). Functional annotations  
190 were assigned to between 50% (*K. caespitosa*) and 70% (*P. exiguum*) of the predicted genes per  
191 species. These assemblies revealed between 598 (*P. exiguum*) and 1604 (*M. australis*) putative rRNA,  
192 821 (*K. caespitosa*) to 1492 (*M. australis*) tRNA genes, and additional non-protein-coding RNA genes  
193 (Table 2).

194

#### 195 **Fig. 1. Assembly completeness.**

196 Assembly completeness was assessed based on the proportion of complete, fragmented, and missing BUSCOs.

197

#### 198 **Table 2: Assembly annotation statistics.**

	<i>Kewa caespitosa</i>	<i>Macarthuria australis</i>	<i>Pharnaceum exiguum</i>
Final gene number	50661	80236	26,155
Functional annotation assigned	25,058 (49.46%)	50,536 (62.98%)	18,372 (70.24%)
Average gene lengths [bp]	5494	1936	5090
Average mRNA length [bp]	2143	1018	2154
Average peptide length [aa]	447	241	435
RBHs vs. BeetSet2	9,968	10,568	10,045
Average number of exons per	6.6	2.9	6

---

gene

Number of predicted tRNAs	821	1491	1260
---------------------------	-----	------	------

Number of predicted rRNAs	720	1604	598
---------------------------	-----	------	-----

Link to data set	<a href="https://docs.cebitec.uni-bielefeld.de/s/pZ4kGpPEDtTPgiW">https://docs.cebitec.uni-bielefeld.de/s/pZ4kGpPEDtTPgiW</a>		
------------------	---	--	--

(TEMPORARY LINK FOR PEER-REVIEW)

---

199

200

## 201 **Comparison between transcriptome and genome assemblies**

202 Previously released transcriptome assemblies were compared to the genome assemblies to assess  
203 completeness and to identify differences. In total 44,169 of 65,062 (67.9%) coding sequences of *the*  
204 *K. caespitosa* transcriptome assembly were recovered in the corresponding genome assembly. This  
205 recovery rate is lower for both *M. australis* assemblies, where only 27,894 of 58,953 (47.3%) coding  
206 sequences were detected in the genome assembly. The highest rate was observed for *P. exiguum*,  
207 where 37,318 of 42,850 (87.1%) coding sequences were found in the genome assembly. When  
208 screening the transcriptome assemblies for transcript sequences predicted based on the genome  
209 sequences, the recovery rate was lower (Fig. 2). The number of predicted representative coding  
210 sequences with best hits against the transcriptome assembly ranged from 16.3% in *K. caespitosa* to  
211 29.7% in *P. exiguum* thus leaving most predicted coding sequences without a good full length hit in  
212 the transcriptome assemblies.

213

### 214 **Fig. 2. Recovery of sequences between transcriptome and genome assemblies.**

215 The figure displays the percentage of sequences present in one assembly that are recovered or missing in the  
216 other assembly type.

217

218

## 219 **Discussion**

220 An almost perfect match between the predicted genome size and the final assembly size was  
221 observed for *P. exiguum*. When taking gaps within scaffolds into account the *K. caespitosa* assembly  
222 size reached the estimated genome size. High heterozygosity could be one explanation for the  
223 assembly size exceeding the estimated haploid genome size of *M. australis*. The two independent  
224 genome size estimations for *M. australis* based on different read data sets indicate almost perfect  
225 reproducibility of this method. Although centromeric regions and other low complexity regions were



226 probably underestimated in the genome size estimation as well as in the assembly process, this  
227 agreement between estimated genome size and final assembly size indicates a high assembly quality.  
228 The continuity of the *P. exiguum* assembly is similar to the assembly continuity of *Dianthus*  
229 *caryophyllus* [12] with a scaffold N50 of 60.7 kb. Additional quality indicators are the high proportion  
230 of detected BUSCOs in the final assemblies as well as the high mapping rate of reads against the  
231 assemblies. The percentage of complete BUSCOs is in the same range as the value of the  
232 *D. caryophyllales* genome assembly which revealed 88.9% complete BUSCOs based on our BUSCO  
233 settings. We demonstrate a cost-effective generation of draft genome assemblies of three different  
234 plant species. Investing into more paired-end sequencing based on Illumina technology would not  
235 substantially increase the continuity of the presented assemblies. This was revealed by initial  
236 assemblies for *M. australis* performed with less than half of all generated sequencing reads. Although  
237 the total assembly size increased when doubling the amount of incorporated sequencing reads, the  
238 continuity is still relatively low. No direct correlation between the sequencing depth and the  
239 assembly quality was observed in this study. Genome properties seem to be more influential than  
240 the amount of sequencing data. Even very deep sequencing with short reads in previous studies  
241 [12,18] was unable to compete with the potential of long reads in genome assembly projects [13,14].  
242 No major breakthroughs were achieved in the development of publicly available assemblers during  
243 the last years partly due to the availability of long reads which made it less interesting.

244 The number of predicted genes in *P. exiguum* is in the range expected for most plants [44,45]. While  
245 the predicted gene numbers for *K. caespitosa* and *M. australis* are much higher, they are only slightly  
246 exceeding the number of genes predicted for other plants [44,45]. Nevertheless, the assembly  
247 continuity and the heterozygosity of *M. australis* are probably the most important factors for the  
248 artificially high number of predicted genes. The high percentage of duplicated BUSCOs (11.8%)  
249 indicates the presence of both alleles for several genes. As the average gene length in *M. australis* is  
250 shorter than in both other assemblies, some gene model predictions might be too short. This gene  
251 prediction could be improved by an increase in assembly continuity.

252 There is a substantial difference between the transcriptome sequences and the predicted transcripts  
253 of the genome assembly. The presence of alternative transcripts and fragmented transcripts in the  
254 transcriptome assemblies are one explanation why not all transcripts were assigned to a genomic  
255 locus. Some transcripts probably represent genes which are not properly resolved in the genome  
256 assemblies. This is especially the case for *M. australis*. The high percentage of complete BUSCOs of  
257 the *K. caespitosa* and *P. exiguum* genome assemblies indicate that missing sequences in the genome  
258 assemblies account only for a minority of the differences. The complete BUSCO percentage of the  
259 *P. exiguum* genome assembly even exceeds the value assigned to the corresponding transcriptome  
260 assembly. Although BUSCOs are selected in a robust way, it is likely that some of these genes are not

261 present in the genomes investigated here, since *B. vulgaris* is the closest relative with an almost  
262 completely sequenced genome [7]. Our genome assemblies provide additional sequences of genes  
263 which are not expressed in the tissues sampled for the generation of the transcriptome assembly. In  
264 addition, coding sequences might be complete in the genome assemblies, while low expression  
265 caused a fragmented assembly based on RNA-Seq reads. This explains why only a small fraction of  
266 the predicted coding sequences of the genome assemblies was mapped to the coding sequences  
267 derived from the corresponding transcriptome assembly.

268 The availability of assembled sequences as well as large sequencing read data sets enables the  
269 investigation of repeats e.g. transposable elements across a large phylogenetic distance within the  
270 Caryophyllales. It also allows the extension of genome-wide analysis like gene family investigations  
271 from *B. vulgaris* across Caryophyllales. As all three species produce anthocyanins, we provide the  
272 basis to study the underlying biosynthetic genes. Due to the huge evolutionary distance to other  
273 anthocyanin producing species, the availability of these sequences could facilitate the identification  
274 of common and unique features of the involved enzymes.

275

#### 276 **Author contribution**

277 TF isolated DNA. BP and TF performed data processing, assembly, and annotation. BP, TF, and SFB  
278 interpreted the results. BP wrote the initial draft. All authors read and approved the final version of  
279 this manuscript.

280

#### 281 **Acknowledgements**

282 We thank the CeBiTec Bioinformatic Resource Facility team for great technical support.

283

#### 284 **References**

- 285 1. Brockington SF, Walker RH, Glover BJ, Soltis PS, Soltis DE. Complex pigment evolution in the  
286 Caryophyllales. *New Phytol.* 2011;190: 854–864. doi:10.1111/j.1469-8137.2011.03687.x
- 287 2. Yang Y, Moore MJ, Brockington SF, Soltis DE, Wong GK-S, Carpenter EJ, et al. Dissecting  
288 Molecular Evolution in the Highly Diverse Plant Clade Caryophyllales Using Transcriptome  
289 Sequencing. *Mol Biol Evol.* 2015;32: 2001–2014. doi:10.1093/molbev/msv081
- 290 3. Smith SA, Brown JW, Yang Y, Bruenn R, Drummond CP, Brockington SF, et al. Disparity,  
291 diversity, and duplications in the Caryophyllales. *New Phytol.* 2018;217: 836–854.  
292 doi:10.1111/nph.14772

- 293 4. Kadereit G, Ackerly D, Pirie MD. A broader model for C4 photosynthesis evolution in plants  
294 inferred from the goosefoot family (Chenopodiaceae s.s.). *Proc R Soc B Biol Sci.* 2012;279:  
295 3304–3311. doi:10.1098/rspb.2012.0440
  
- 296 5. Jain G, Schwinn KE, Gould KS. Betalain induction by l-DOPA application confers photoprotection  
297 to saline-exposed leaves of *Disphyma australe*. *New Phytol.* 2015;207: 1075–1083.  
298 doi:10.1111/nph.13409
  
- 299 6. Thorogood CJ, Bauer U, Hiscock SJ. Convergent and divergent evolution in carnivorous pitcher  
300 plant traps. *New Phytol.* 2018;217: 1035–1041. doi:10.1111/nph.14879
  
- 301 7. Dohm JC, Minoche AE, Holtgräwe D, Capella-Gutiérrez S, Zakrzewski F, Tafer H, et al. The  
302 genome of the recently domesticated crop plant sugar beet (*Beta vulgaris*). *Nature.* 2014;505:  
303 546–549. doi:10.1038/nature12817
  
- 304 8. Brockington SF, Yang Y, Gandia-Herrero F, Covshoff S, Hibberd JM, Sage RF, et al. Lineage-  
305 specific gene radiations underlie the evolution of novel betalain pigmentation in Caryophyllales.  
306 *New Phytol.* 2015;207: 1170–1180. doi:10.1111/nph.13441
  
- 307 9. Stevanato P, Trebbi D, Saccomani M. Single nucleotide polymorphism markers linked to root  
308 elongation rate in sugar beet. *Biol Plant.* 2017;61: 48–54. doi:10.1007/s10535-016-0643-1
  
- 309 10. Kong W, Yang S, Wang Y, Bendahmane M, Fu X. Genome-wide identification and  
310 characterization of aquaporin gene family in *Beta vulgaris*. *PeerJ.* 2017;5.  
311 doi:10.7717/peerj.3747
  
- 312 11. Xu C, Jiao C, Sun H, Cai X, Wang X, Ge C, et al. Draft genome of spinach and transcriptome  
313 diversity of 120 *Spinacia* accessions. *Nat Commun.* 2017;8. doi:10.1038/ncomms15275
  
- 314 12. Yagi M, Kosugi S, Hirakawa H, Ohmiya A, Tanase K, Harada T, et al. Sequence Analysis of the  
315 Genome of Carnation (*Dianthus caryophyllus* L.). *DNA Res Int J Rapid Publ Rep Genes Genomes.*  
316 2014;21: 231–241. doi:10.1093/dnares/dst053
  
- 317 13. Lightfoot DJ, Jarvis DE, Ramaraj T, Lee R, Jellen EN, Maughan PJ. Single-molecule sequencing  
318 and Hi-C-based proximity-guided assembly of amaranth (*Amaranthus hypochondriacus*)  
319 chromosomes provide insights into genome evolution. *BMC Biol.* 2017;15: 74.  
320 doi:10.1186/s12915-017-0412-4
  
- 321 14. Jarvis DE, Ho YS, Lightfoot DJ, Schmöckel SM, Li B, Borm TJA, et al. The genome of *Chenopodium*  
322 *quinoa*. *Nature.* 2017;542: 307–312. doi:10.1038/nature21370
  
- 323 15. Copetti D, Búrquez A, Bustamante E, Charboneau JLM, Childs KL, Eguiarte LE, et al. Extensive  
324 gene tree discordance and hemiplasy shaped the genomes of North American columnar cacti.  
325 *Proc Natl Acad Sci.* 2017;114: 12003–12008. doi:10.1073/pnas.1706367114
  
- 326 16. Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, et al. Phylotranscriptomic  
327 analysis of the origin and early diversification of land plants. *Proc Natl Acad Sci U S A.* 2014;111:  
328 E4859–E4868. doi:10.1073/pnas.1323926111
  
- 329 17. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.  
330 *Bioinforma Oxf Engl.* 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170
  
- 331 18. Pucker B, Holtgräwe D, Sørensen TR, Stracke R, Viehöver P, Weisshaar B. A *De Novo* Genome  
332 Sequence Assembly of the *Arabidopsis thaliana* Accession Niederzenz-1 Displays

- 333 Presence/Absence Variation and Strong Synteny. PLOS ONE. 2016;11: e0164321.  
334 doi:10.1371/journal.pone.0164321
- 335 19. Andrews S. FastQC A Quality Control tool for High Throughput Sequence Data. 2010. Available:  
336 <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- 337 20. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of occurrences  
338 of k-mers. Bioinformatics. 2011;27: 764–770. doi:10.1093/bioinformatics/btr011
- 339 21. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al.  
340 GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics. 2017;33:  
341 2202–2204. doi:10.1093/bioinformatics/btx153
- 342 22. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved memory-  
343 efficient short-read *de novo* assembler. GigaScience. 2012;1: 18. doi:10.1186/2047-217X-1-18
- 344 23. Cha S, Bird DM. Optimizing k-mer size using a variant grid search to enhance *de novo* genome  
345 assembly. Bioinformation. 2016;12: 36–40. doi:10.6026/97320630012036
- 346 24. Haak M, Vinke S, Keller W, Droste J, Rückert C, Kalinowski J, et al. High Quality *de Novo*  
347 Transcriptome Assembly of *Croton tiglium*. Front Mol Biosci. 2018;5.  
348 doi:<https://doi.org/10.3389/fmolb.2018.00062>
- 349 25. Keller O, Kollmar M, Stanke M, Waack S. A novel hybrid gene prediction method employing  
350 protein multiple sequence alignments. Bioinforma Oxf Engl. 2011;27: 757–763.  
351 doi:10.1093/bioinformatics/btr010
- 352 26. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing  
353 genome assembly and annotation completeness with single-copy orthologs. Bioinforma Oxf  
354 Engl. 2015;31: 3210–3212. doi:10.1093/bioinformatics/btv351
- 355 27. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.  
356 ArXiv13033997 Q-Bio. 2013; Available: <http://arxiv.org/abs/1303.3997>
- 357 28. Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD. REAPR: a universal tool for  
358 genome assembly evaluation. Genome Biol. 2013;14: R47. doi:10.1186/gb-2013-14-5-r47
- 359 29. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol  
360 Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2
- 361 30. Holtgräwe D, Rosleff Sörensen T, Parol-Kryger R, Pucker B, Kleinbölting N, Viehöver P, et al. Low  
362 coverage re-sequencing in sugar beet for anchoring assembly sequences to genomic positions  
363 [Internet]. 2017. Available: <https://jbrowse.cebitec.uni-bielefeld.de/RefBeet1.5/>
- 364 31. Stadermann KB, Weisshaar B, Holtgräwe D. SMRT sequencing only *de novo* assembly of the  
365 sugar beet (*Beta vulgaris*) chloroplast genome. BMC Bioinformatics. 2015;16.  
366 doi:10.1186/s12859-015-0726-6
- 367 32. Kubo T, Nishizawa S, Sugawara A, Itchoda N, Estiati A, Mikami T. The complete nucleotide  
368 sequence of the mitochondrial genome of sugar beet (*Beta vulgaris* L.) reveals a novel gene for  
369 tRNACys(GCA). Nucleic Acids Res. 2000;28: 2571–2576.
- 370 33. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal  
371 RNA-seq aligner. Bioinforma Oxf Engl. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

- 372 34. Smit A, Hubley R, Green P. RepeatMasker Frequently Open-4.0 [Internet]. 2015. Available:  
373 <http://www.repeatmasker.org/>
- 374 35. Green P. Consed--A Finishing Package [Internet]. Available:  
375 <http://www.phrap.org/consed/consed.html#howToGet>
- 376 36. Kent WJ. BLAT—The BLAST-Like Alignment Tool. *Genome Res.* 2002;12: 656–664.  
377 doi:10.1101/gr.229202
- 378 37. Pucker B, Holtgräwe D, Weisshaar B. Consideration of non-canonical splice sites improves gene  
379 prediction on the *Arabidopsis thaliana* Niederzenz-1 genome sequence. *BMC Res Notes.*  
380 2017;10. doi:<https://doi.org/10.1186/s13104-017-2985-y>
- 381 38. Minoche AE, Dohm JC, Schneider J, Holtgräwe D, Viehöver P, Montfort M, et al. Exploiting  
382 single-molecule transcript sequencing for eukaryotic gene prediction. *Genome Biol.* 2015;16:  
383 184. doi:10.1186/s13059-015-0729-7
- 384 39. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics.*  
385 2013;29: 2933–2935. doi:10.1093/bioinformatics/btt509
- 386 40. Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, et al. Rfam 13.0:  
387 shifting to a genome-centric resource for non-coding RNA families. *Nucleic Acids Res.* 2018;46:  
388 D335–D342. doi:10.1093/nar/gkx1038
- 389 41. Cheng C-Y, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD. Araport11: a  
390 complete reannotation of the *Arabidopsis thaliana* reference genome. *Plant J.* 2017;89: 789–  
391 804. doi:10.1111/tpj.13415
- 392 42. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale  
393 protein function classification. *Bioinformatics.* 2014;30: 1236–1240.  
394 doi:10.1093/bioinformatics/btu031
- 395 43. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: *ab initio*  
396 prediction of alternative transcripts. *Nucleic Acids Res.* 2006;34: W435–W439.  
397 doi:10.1093/nar/gkl200
- 398 44. Sterck L, Rombauts S, Vandepoele K, Rouzé P, Van de Peer Y. How many genes are there in  
399 plants (... and why are they there)? *Curr Opin Plant Biol.* 2007;10: 199–203.  
400 doi:10.1016/j.pbi.2007.01.004
- 401 45. Pucker B, Brockington SF. Genome-wide analyses supported by RNA-Seq reveal non-canonical  
402 splice sites in plant genomes. *BMC Genomics.* 2018;19: 980. doi:10.1186/s12864-018-5360-z

403

404

405 **Supporting Information**

406 **AdditionalFile 1. Sequencing result overview.**

407 **AdditionalFile 2. Genome size estimation results.** Genome size estimations with GenomeScope [21]  
408 are listed for various k-mer sizes. Two different read sets of *M. australis* were used for the genome  
409 size estimation (1=ERR2401802, 2=ERR2401614) to check the reproducibility.

410 **AdditionalFile 3. Evaluation of assembly attempts of *K. caespitosa*.**

411 **AdditionalFile 4. Evaluation of assembly attempts of *M. australis*.**

412 **AdditionalFile 5. Evaluation of assembly attempts of *P. exiguum*.**

413 **AdditionalFile 6. Detailed list of assembly parameters.**

414 **AdditionalFile 7. Gene prediction hint sources.** These RNA-Seq read data sets and transcriptome  
415 assemblies were incorporated in the gene annotation process as hints.

416 **AdditionalFile 8. Assembly attempt evaluation results.** Statistics of raw assemblies were calculated  
417 to identify the best parameter settings. Since k-mer size was previously reported as the most  
418 important parameter, extensive optimization was performed. In addition, different settings for insert  
419 sizes were evaluated for *P. exiguum* (phe001-phe006). Parameter optimization for *M. australis* was  
420 performed on a subset of all reads due to availability.

421 **AdditionalFile 9. Gene prediction statistics.** Different gene prediction approaches were performed  
422 during the optimization process. Results of these predictions include *ab initio* gene prediction and  
423 hint-based approaches. RNA-Seq reads and coding sequences derived from previous transcriptome  
424 assemblies are two incorporated hint types. In addition, we assessed the impact of repeat masking  
425 prior to gene prediction.

426

percentage of detected BUSCOs



