

Draft Genome Sequence of “*Candidatus Phytoplasma oryzae*” Strain Mbita1, the Causative Agent of Napier Grass Stunt Disease in Kenya

Anne Fischer,^{a,b} Ivette Santana-Cruz,^c Lillian Wambua,^{a,d} Cassandra Olds,^{b,e} Charles Midega,^a Matthew Dickinson,^f Praphat Kawicha,^g Zeyaur Khan,^a Daniel Masiga,^a Joerg Jores,^{b,h} Bernd Schneiderⁱ

International Centre of Insect Physiology and Ecology, Nairobi, Kenya^a; International Livestock Research Institute, Nairobi, Kenya^b; Institute for Genome Sciences, Baltimore, Maryland, USA^c; School of Biological Sciences, University of Nairobi, Nairobi, Kenya^d; Washington State University, Pullman, Washington, USA^e; Faculty of Science, The University of Nottingham, Sutton Bonington Campus, Sutton Bonington, Leicestershire, United Kingdom^f; Faculty of Natural Resources and Agro-industry, Chalermphrakiat Sakon Nakhon Province Campus, Kasetsart University, Sakon Nakhon, Thailand^g; Institute of Veterinary Bacteriology, University of Bern, Bern, Switzerland^h; Julius Kühn-Institut, Institut für Pflanzenschutz in Obst- und Weinbau, Dossenheim, Germanyⁱ

Phytoplasmas are bacterial plant pathogens with devastating impact on agricultural production worldwide. In eastern Africa, Napier grass stunt disease causes serious economic losses in the smallholder dairy industry. This draft genome sequence of “*Candidatus Phytoplasma oryzae*” strain Mbita1 provides insight into its genomic organization and the molecular basis of pathogenicity.

Received 29 February 2016 Accepted 7 March 2016 Published 21 April 2016

Citation Fischer A, Santana-Cruz I, Wambua L, Olds C, Midega C, Dickinson M, Kawicha P, Khan Z, Masiga D, Jores J, Schneider B. 2016. Draft genome sequence of “*Candidatus Phytoplasma oryzae*” strain Mbita1, the causative agent of Napier grass stunt disease in Kenya. *Genome Announc* 4(2):e00297-16. doi:10.1128/genomeA.00297-16.

Copyright © 2016 Fischer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](http://creativecommons.org/licenses/by/4.0/).

Address correspondence to Joerg Jores, jores.anne@gmail.com.

Phytoplasmas are bacteria that parasitize the plant phloem tissue causing a wide range of symptoms, like yellowing, stunting, little leaves, witches’ brooms, virescence, and phyllody (1). They are transmitted by insect vectors and more than 1,000 plant species have been described as host plants, many of agricultural importance. Current phytoplasma classification is based on restriction fragment length polymorphism (RFLP) patterns of their 16SrDNA sequence (2). Because phytoplasmas are unculturable in axenic media, their molecular characterization has been hampered and only five phytoplasma genomes have been deposited in public databases to date (3–7). Napier grass, or elephant grass (*Pennisetum purpureum*), is the major fodder grass for the dairy industry in eastern Africa. Napier grass stunt disease (NSD) severely impairs the growth of Napier grass, with losses of up to 70% biomass in infected plants. In Kenya and Uganda, NSD is caused by a phytoplasma belonging to group 16SrXI (“*Candidatus Phytoplasma oryzae*” or rice yellow dwarf) (8, 9). We report here the first draft genome of Napier grass phytoplasma strain Mbita1.

Infected Napier grass plants were collected at the International Centre of Insect Physiology and Ecology (ICIPE) Mbita Research Station, western Kenya. A pulse-field-electrophoresis gel was done with a pilot DNA extract, which allowed the genome size of the Napier grass phytoplasma to be estimated to approximately 550 kbp. Afterward, bacterial DNA was further extracted from plant leaves using a Cs-Cl gradient, purified, and the genome was sequenced using the Illumina HiSeq sequencing platform. Illumina raw reads were split into 31 data sets. Each set was assembled independently using the Celera Assembler v8.2 at 150× coverage. Resulting contigs (approximately 30 contigs per assembly) were overlapped using Minimus (10) to find potential gaps in coverage from the independent assemblies. The final assembly resulted in 28 contigs and a genome size of 533,195 total bp (16 contigs > 10 kbp adding up to 455,349 bp). The 16S rRNA operon was

absent from the assembly, due to low coverage in the region. We therefore amplified the 16S rRNA operon using PCR. Sanger sequencing of the PCR products confirmed that Napier grass phytoplasma belonged to the group 16SrXI. The average G+C content of the reported genome was 19.3%, which is the lowest value reported for a phytoplasma species to date, as the G+C content for other species ranges between 21.4% and 27.3%. Open reading frames (ORFs) were predicted using Prodigal 2.50. A total of 465 ORFs and 10 tRNAs were obtained. Functional annotation was produced by the Institute for Genome Sciences Annotation Engine (11), <http://ae.igs.umaryland.edu/cgi/index.cgi>. The average gene length was 888 bp. Among the ORFs, we identified the immunodominant encoding genes *imp*, *secA*, and *secY*, which are essential to the Sec protein translocation system. We also found one *hflB* gene and three of its truncated copies.

This annotated genome sequence is the first from the “*Candidatus Phytoplasma oryzae*” species. It will give insight into the genetic diversity of phytoplasmas and further our understanding of molecular pathogenicity mechanisms.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [LTBM000000000](https://www.ncbi.nlm.nih.gov/nuccore/LTBM000000000). The version described in this paper is version [LTBM010000000](https://www.ncbi.nlm.nih.gov/nuccore/LTBM010000000). The raw reads are available under accession number [SRP069757](https://www.ncbi.nlm.nih.gov/nuccore/SRP069757).

ACKNOWLEDGMENTS

This work was funded by the German Federal Ministry for Economic Cooperation and Development (contract 81156063, project 11.7860.7-001.00) and received additional support from the Consultative Group for International Agricultural Research (CGIAR) Program on Livestock and Fish. The Centrum of International Migration and Development (CIM) supported A.F. L.W. was supported by a postdoctoral fellowship from the Department for International Development (DFID).

REFERENCES

- Hogenhout SA, Oshima K, Ammar D, Kakizawa S, Kingdom HN, Namba S. 2008. Phytoplasmas: bacteria that manipulate plants and insects. *Mol Plant Pathol* 9:403–423. <http://dx.doi.org/10.1111/j.1364-3703.2008.00472.x>.
- Lee IM, Gundersen-Rindal DE, Bertaccini A. 1998. Phytoplasma: ecology and genomic diversity. *Phytopathology* 88:1359–1366. <http://dx.doi.org/10.1094/PHYTO.1998.88.12.1359>.
- Kakizawa S, Makino A, Ishii Y, Tamaki H, Kamagata Y. 2014. Draft genome sequence of “*Candidatus* Phytoplasma asteris” strain OY-V, an unculturable plant-pathogenic bacterium. *Genome Announc* 2(5): e00944-14. <http://dx.doi.org/10.1128/genomeA.00944-14>.
- Kube M, Schneider B, Kuhl H, Dandekar T, Heitmann K, Migdoll AM, Reinhardt R, Seemüller E. 2008. The linear chromosome of the plant-pathogenic mycoplasma “*Candidatus* Phytoplasma mali”. *BMC Genomics* 9:306. <http://dx.doi.org/10.1186/1471-2164-9-306>.
- Oshima K, Kakizawa S, Nishigawa H, Jung HY, Wei W, Suzuki S, Arashida R, Nakata D, Miyata S, Ugaki M, Namba S. 2004. Reductive evolution suggested from the complete genome sequence of a plant-pathogenic phytoplasma. *Nat Genet* 36:27–29. <http://dx.doi.org/10.1038/ng1277>.
- Tran-Nguyen LT, Kube M, Schneider B, Reinhardt R, Gibb KS. 2008. Comparative genome analysis of “*Candidatus* Phytoplasma australiense” (subgroup *tuf*-Australia I; *rp*-A) and “*Ca.* Phytoplasma asteris” strains OY-M and AY-WB. *J Bacteriol* 190:3979–3991. <http://dx.doi.org/10.1128/JB.01301-07>.
- Andersen MT, Liefing LW, Havukkala I, Beever RE. 2013. Comparison of the complete genome sequence of two closely related isolates of “*Candidatus* Phytoplasma australiense” reveals genome plasticity. *BMC Genomics* 14:529. <http://dx.doi.org/10.1186/1471-2164-14-529>.
- Jones P, Devonshire BJ, Holman TJ, Ajanga S. 2004. Napier grass stunt: a new disease associated with a 16SrXI group phytoplasma in Kenya. *Plant Pathol* 53:519. <http://dx.doi.org/10.1111/j.1365-3059.2004.01038.x>.
- Nielsen SL, Ebong C, Kabirizi J, Nicolaisen M. 2007. First report of a 16SrXI group phytoplasma (*Candidatus* Phytoplasma oryzae) associated with Napier grass stunt disease in Uganda. *Plant Pathol* 56:1039. <http://dx.doi.org/10.1111/j.1365-3059.2007.01638.x>.
- Sommer DD, Delcher AL, Salzberg SL, Pop M. 2007. Minimus: a fast, lightweight genome assembler. *BMC Bioinformatics* 8:64. <http://dx.doi.org/10.1186/1471-2105-8-64>.
- Galens K, Orvis J, Daugherty S, Creasy HH, Angiuoli S, White O, Wortman J, Mahurkar A, Giglio MG. 2011. The IGS standard operating procedure for automated prokaryotic annotation. *Stand Genomic Sci* 4:244–251. <http://dx.doi.org/10.4056/sigs.1223234>.