

1 **Assessing Phosphatidylethanol (PEth) Levels Reflecting Different Drinking**  
2 **Habits in Comparison to the Alcohol Use Disorders Identification Test – C**  
3 **(AUDIT-C)**

4

5 **Authors:**

6 MSc. Alexandra Schröck<sup>1,5</sup>, PD Dr. Friedrich M. Wurst<sup>2,3</sup>, MSc. Natasha Thon<sup>4</sup>, Prof. Dr.  
7 Wolfgang Weinmann<sup>1</sup>

8

9 **Affiliations:**

10 <sup>1</sup>Institute of Forensic Medicine, Forensic Toxicology and Chemistry, University of Bern,  
11 Switzerland

12 <sup>2</sup>Center for Interdisciplinary Addiction Research Hamburg, Germany

13 <sup>3</sup> University of Basel, Basel, Switzerland

14 <sup>4</sup>Paracelsus Medical University, Salzburg, Austria

15 <sup>5</sup>Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland

16

17 **Corresponding author:**

18 Wolfgang Weinmann

19 Institute of Forensic Medicine, University of Bern

20 Bühlstrasse 20

21 3012 Bern

22 Switzerland

23 Tel. +41 31 631 5668

24 Fax. +41 31 631 8580

25 Email: Wolfgang.Weinmann@irm.unibe.ch

1 **Acknowledgement**

2 We would like to thank the employees of the “Interregionale Blutspende SRK AG”  
3 (Interregional Blood Donation Center Swiss Red Cross) for the collection of 300 blood  
4 samples from volunteers for generation of reference levels, and especially Ina Krebber for her  
5 help in organizing this project.

6

7 **Funding**

8 This study was supported by the Swiss Foundation of Alcohol Research (Grant 254/2014:  
9 Studies on phosphatidylethanol – a promising biomarker for the detection of harmful ethanol  
10 consumption – and its possible use for abstinence monitoring).

## 1 **Abstract**

2 In addition to monitoring problematic or harmful alcohol consumption, drinking experiments  
3 indicated the potential of phosphatidylethanol (PEth) in abstinence monitoring. To date no  
4 profound evaluation of thresholds for the differentiation of abstinence from moderate drinking  
5 and for detection of excessive consumption based on PEth homologues exists. Investigations  
6 with a large group of healthy volunteers (n=300) were performed to establish PEth reference  
7 values reflecting different drinking habits.

8 Blood samples were analyzed for PEth 16:0/18:1 and 16:0/18:2 by online-SPE-LC-MS/MS  
9 method. Results were compared to AUDIT-C questionnaires, to the amounts of alcohol  
10 consumed during the two-weeks prior to blood sampling, and were statistically evaluated.

11 PEth concentrations were significantly correlated with self-reported alcohol consumption  
12 ( $r>0.69$ ) and with AUDIT-C scores ( $r>0.65$ ). 4.0% of 300 volunteers reported abstinence  
13 (AUDIT-C score: 0), no PEth was detectable in their blood. PEth 16:0/18:1 concentrations  
14 below the limit of detection of 10.0ng/mL match with abstinence and light drinking habits  
15 ( $\leq 10$ g pure alcohol/day). However, some volunteers classified as “excessive alcohol  
16 consumers” had negative PEth results.

17 In the group of volunteers classified as “moderate drinkers” (AUDIT-C score: 1–3 (women)  
18 and 1–4 (men)), 95% of the test persons had PEth 16:0/18:1 ranging from not detected to  
19 112ng/mL, and PEth 16:0/18:2 ranging from not detected to 67.0ng/mL.

20 Combination of self-reported alcohol consumption and AUDIT-C score showed that negative  
21 PEth results match with abstinence or light drinking. Moderate alcohol consumption resulted  
22 in PEth 16:0/18:1 from 0 to 112ng/mL and for PEth 16:0/18:2 ranged from 0 to 67.0ng/mL.  
23 Higher PEth concentrations indicated excessive alcohol consumption.

24 **Keywords:** Phosphatidylethanol (PEth), Alcohol Use Disorders Identification Test – C  
25 (AUDIT-C), Online-SPE-LC-MS/MS, Abstinence, Moderate drinking

# 1 **Introduction**

2

3 Alcohol, as one of the most commonly used recreational drugs, is an important cause of  
4 morbidity and mortality (Kechagias et al., 2015). To gain information about a person's  
5 alcohol consumption habits, clinicians often rely on questionnaires such as Alcohol Use  
6 Disorders Identification Test (AUDIT) or its short form, the AUDIT-C (SAMHSA, WHO,  
7 2001). As these may be influenced by memory bias and under-reporting by the desirability to  
8 report less alcohol ingestion (Bajunirwe et al., 2014), also biomarkers of alcohol intake are  
9 highly useful to identify drinking patterns and alcohol-related disorders, even at an early stage  
10 (Schröck et al., 2014, Wurst et al., 2015, Mann K et al., 2016).

11 Very recent alcohol ingestion can be verified by breath alcohol testing, determination of the  
12 blood alcohol concentration (BAC) or by the highly specific direct markers ethyl glucuronide  
13 (EtG) and ethyl sulphate (EtS) in blood and/or urine (Halter et al., 2008, Dresen et al., 2004).

14 Besides hair analysis for EtG, which is used in forensic applications (Kharbouche et al., 2012,  
15 SOHT, 2014, Boscolo-Berto et al., 2013), for the identification of excessive alcohol  
16 consumption the traditional indirect markers carbohydrate deficient transferrin (CDT) and  $\gamma$ -  
17 glutamyltranspeptidase (GGT) are in clinical use. However, these markers lack in specificity  
18 as they are not only influenced by alcohol intake, but also by age, gender, smoking, and  
19 various diseases (Hannuksela et al., 2007, Thierauf et al., 2011). Additionally, these markers  
20 are suboptimal with regard to sensitivity for the detection of moderate alcohol consumption.

21 For persons with moderate drinking habits or binge drinkers more sensitive markers are  
22 needed, such as phosphatidylethanol (PEth), which is a more specific and sensitive alcohol  
23 biomarker concerning this matter (Kechagias et al., 2015). PEth represents a group of  
24 phospholipids present in cell membranes, which are formed directly after alcohol intake  
25 (Gnann et al., 2009, Gnann et al., 2012) via the enzyme phospholipase D (PL D) from  
26 phosphatidylcholine (PC), as long as ethanol is present (Kobayashi and Kanfer, 1987). The

1 homologues PEth 16:0/18:1 and PEth 16:0/18:2 are the most abundant homologues in human  
2 blood (Helander and Zheng, 2009, Zheng et al., 2011, Nalesso et al., 2011).

3 Due to an elimination half-life of approximately four days (Varga et al., 2000), which  
4 however can vary strongly between individuals (Gnann et al., 2012), PEth accumulates in the  
5 body after repeated drinking, and is therefore suitable for the differentiation of problematic  
6 excessive alcohol consumption and moderate drinking (Aradottir et al., 2006, Viel et al.,  
7 2012, Winkler et al., 2013, Gnann et al., 2009).

8 Recently, drinking studies with moderate alcohol consumption have shown that PEth  
9 concentrations correlate with the consumed amounts of alcohol, and that PEth has a potential  
10 in abstinence monitoring as it is formed directly after alcohol intake (Gnann, 2011, Gnann et  
11 al., 2012, Schröck et al., 2016c). Another pilot study showed that combining EtG and EtS  
12 testing in urine with PEth analysis in blood seems to be effective in providing additional  
13 information on recent drinking (Skipper et al., 2013).

14 PEth also has a potential in driving aptitude assessment (DAA) as additional marker to EtG in  
15 hair, which is the most frequently routinely used direct marker for the detection of excessive  
16 alcohol consumption and also for abstinence monitoring in this context (Kharbouche et al.,  
17 2012). In a recent study, the classification into drinking groups (abstainers, moderate drinkers  
18 and excessive drinkers) according to EtG concentrations in hair corresponded to the detected  
19 PEth values in blood in 68 % of the tested samples (Schröck et al., 2016b).

20 So far, several PEth thresholds, such as 210 ng/mL (Helander and Hansson, 2013), 700 ng/mL  
21 (Schröck et al., 2016a) , and 800 ng/mL (Gnann, 2011) were proposed for the differentiation  
22 of moderate and excessive alcohol consumption, which were obtained either from alcoholics  
23 starting withdrawal or which were obtained during drinking studies.

24 By now, there are no reference levels established for PEth homologues allowing the  
25 differentiation of abstinence, moderate alcohol consumption and excessive consumption in  
26 addition to AUDIT-C or self-reports. On this background, our study was performed with 300

1 volunteers to investigate the potential of PEth in reflecting different drinking habits, such as  
2 abstinence, moderate and excessive alcohol consumption, and to assess reference values for  
3 setting-up a database of PEth 16:0/18:1 and PEth 16:0/18:2 concentrations, which can be  
4 expected after consumption of alcohol in a harmless way ( $\leq 20$  g/day females,  $\leq 40$  g/day  
5 males, (WHO, 2000)). The PEth concentrations were statistically evaluated in comparison  
6 with AUDIT-C questionnaires and with self-reported amounts of ingested alcohol of the last  
7 two weeks prior to blood sampling.

8

## 9 **Material and methods**

10

### 11 **Chemicals and materials**

12 PEth 16:0/18:1, PEth 16:0/18:2 were purchased from Avanti Polar Lipids (Alabaster, USA).  
13 Acetonitrile was supplied by Agros Organics (New Jersey, USA), 2-propanol was obtained  
14 from Fisher Scientific (Loughborough, UK), and formic acid (HCOOH) was from Sigma  
15 Aldrich (Buchs, Switzerland). Ammonium acetate was provided by Merck (Darmstadt,  
16 Germany). Deionized H<sub>2</sub>O was produced in-house with a Milli-Q water system from  
17 Millipore (Billerica, USA).

18 Deuterated internal standards were synthesized in our laboratory from phosphatidylcholine  
19 16:0/18:1 and phosphatidylcholine 16:0/18:2 and D<sub>6</sub>-ethanol catalyzed by phospholipase D  
20 (Schröck et al., 2016a). Heparin Vacuettes (volume: 9 mL) were obtained from Greiner Bio-  
21 one (Kremsmünster, Austria). Lithium-heparinized whole blood, which was used as blank  
22 blood, was obtained from volunteers who were abstinent from alcohol for at least 4 weeks.

23

### 24 **Study design**

25 To establish thresholds for PEth as diagnostic marker for the differentiation of abstinence and  
26 moderate drinking habits, as well as excessive drinking habits, reference values were

1 determined from 300 volunteers. Blood samples were collected after blood donation at the  
2 blood donation centers in Burgdorf, Bern and Thun (Switzerland). To prevent contamination  
3 with ethanol during veno-puncture, Ethanol-free swabs were used for disinfection, and the  
4 blood samples were stored at 4 °C directly after sampling and were analyzed for PEth within  
5 three days to prevent in-vitro formation of PEth. The volunteers additionally answered the  
6 AUDIT-C questionnaire and retrospectively documented their alcohol consumption of the last  
7 two weeks prior to blood sampling. The concentrations of the PEth homologues 16:0/18:1 and  
8 16:0/18:2 in blood were compared to the AUDIT-C scores, to the self-reported amounts of  
9 consumed alcohol, and these results were statistically evaluated. This study has been  
10 approved by the Cantonal Ethics Commission Bern (064/13) on December 10, 2015.

11

## 12 **AUDIT-C**

13 The 10-item Alcohol Use Disorders Identification Test (AUDIT) is a screening questionnaire  
14 developed by the World Health Organization (WHO) to identify harmful or hazardous alcohol  
15 consumption. Response options for each item range from 0 to 4, resulting in a total possible  
16 score of 40.

17 The AUDIT-C is a shortened version of the 10-items AUDIT, which was developed as an  
18 even briefer and easier screening method of the alcohol consumption habits of the last year  
19 (SAMHSA, WHO, 2001). It consists of the first three questions concerning alcohol  
20 consumption habits such as quantity, frequency and binge drinking (Bush et al., 1998). An  
21 AUDIT-C score equal to zero indicates abstinence. For women an AUDIT-C score  $\geq 4$ , and  
22 for men a score  $\geq 5$  indicates excessive alcohol consumption in a German speaking population  
23 according to the recent evidence- and consensus-based guidelines of the grade S3 (Mann K et  
24 al., 2016).

25

26

## 1 **Self-reported alcohol consumption**

2

3 A modified form of the time line follow back questionnaire, originally developed by Sobell  
4 and Sobell restricted to the last two weeks prior to the study was used to obtain retrospective  
5 estimates of drinking (Sobell and Sobell 1992). This questionnaire contained a choice of  
6 different alcoholic beverages with typical percent alcohol by volume (e.g. beer 5 %, wine 12  
7 %, liquors 40 %). The volunteers were asked to retrospectively fill in the type of consumed  
8 alcoholic beverages and consumed amounts (volumes/sizes of units, and numbers of units per  
9 week)- for each of the two weeks prior to blood sampling. Participants with an alcohol  
10 consumption below  $\leq 10$  g/day were classified as light drinkers.

11

## 12 **Determination of PEth**

13 PEth 16:0/18:1 and PEth 16:0/18:2 were analyzed in 200  $\mu$ L of whole blood with D<sub>5</sub>-PEth  
14 16:0/18:1 and D<sub>5</sub>-PEth 16:0/18:2 as internal standards. The analysis was performed by online-  
15 SPE-LC-MS/MS with a QTrap 3200 mass spectrometer (Sciex, Toronto, Canada). The two  
16 homologues were chromatographically separated on a Luna RP-C5 column, 50 mm x 2 mm, 5  
17  $\mu$ m particle size (Phenomenex, Brechbühler, Schlieren, Switzerland). The limit of detection  
18 (LOD) was 10.0 ng/mL and the limit of quantification (LOQ) was 20.0 ng/mL for both PEth  
19 homologues. The precise details about sample preparation and the validated online-SPE-LC-  
20 MS/MS method are described in (Schröck et al., 2016c).

21

## 22 **Statistic evaluation**

23 Statistical analysis was performed with GraphPad Prism 6 (GraphPad Software Inc., La Jolla,  
24 California, USA). For the Kruskal-Wallis-Test, a non-parametric test used to detect  
25 differences between three or more groups, the results of this study were classified into three  
26 groups (group A: AUDIT-C score 0: abstainers; group B: AUDIT-C score 1 – 3 (women) and



1 1 – 4 (men): moderate alcohol consumption; group C: AUDIT-C score  $\geq 4$  (women) and  $\geq 5$   
2 (men): excessive alcohol consumption) depending on the achieved points of the AUDIT-C  
3 (Mann K et al., 2016).

4 For receiver operating characteristics (ROC) analysis, the results were classified into two  
5 groups (group A and B: abstainers/moderate consumers versus group C: excessive  
6 consumers). The AUDIT-C scores served as primary reference standard for the evaluation of  
7 cut-offs for PEth 16:0/18:1 and 16:0/18:2 for differentiation of these populations with high  
8 sensitivity and specificity. Furthermore, the daily average of self-reported amount of alcohol  
9 consumed over the last 2 weeks prior to blood sampling was compared to PEth 16:0/18:1 and  
10 16:0/18:2 concentrations using Spearman's rank correlation coefficient.

11

## 12 **Results**

13

### 14 **Comparison of PEth concentrations to AUDIT-C scores**

15 The 300 volunteers (203 male, 94 female, 3 not specified; 18 – 78 years, mean:  $47 \pm 13$  years)  
16 were divided into the three groups A, B and C (see above: statistic evaluation; figures 1 and 2,  
17 table 1). Overall, male participants tended to have slightly higher alcohol consumption.

18 Group A: Four percent (6 female and 6 male) of the 300 participating volunteers reported  
19 abstinence (AUDIT-C score 0) and PEth was not detectable in any of these samples (with a  
20 limit of detection (LOD) of 10 ng/mL for PEth 16:0/18:1 and 16:0/18:2).

21 Group B: 71 percent (214 volunteers: 143 male, 68 female, 3 not specified) showed moderate  
22 alcohol consumption (AUDIT-C scores 1 – 3 for women, 1 – 4 for men). PEth 16:0/18:1  
23 values ranged from not detected to 619 ng/mL (mean 31.8 ng/mL). PEth 16:0/18:2 values  
24 ranged from not detected to 286 ng/mL (mean 18.4 ng/mL). Among those “moderate  
25 drinkers” 61.7 % (n = 132) had positive PEth results (PEth  $\geq 10.0$  ng/mL (LOD)).

1 Twelve persons (8 female and 4 male, 31 – 67 years, mean  $50 \pm 12$  years) reported “moderate  
2 drinking habits” (AUDIT-C scores representing moderate alcohol consumption), but did not  
3 report having consumed alcohol in the two weeks prior to blood sampling for this study. Only  
4 one of these 12 persons had PEth 16:0/18:1 ( $> 10.0$  ng/mL) above the LOD (10.0 ng/mL) of  
5 the online-SPE-LC-MS/MS method. PEth 16:0/18:2 was not detectable in any of these  
6 persons. Another 9 persons (all with AUDIT-C scores representing moderate alcohol  
7 consumption) reported no alcohol ingestion in the week prior to blood sampling. Among  
8 them, four persons had PEth 16:0/18:1 concentrations above the LOD, but no PEth 16:0/18:2  
9 was detectable. Another person had 28.3 ng/mL PEth16:0/18:1 and 17.0 ng/mL PEth  
10 16:0/18:2. This means that 44.4 % (4 of 9) of the test persons being abstinent for one week  
11 prior to blood sampling were PEth negative (PEth  $< 10.0$  ng/mL).

12 Group C: 25 percent (74 volunteers: 54 male and 20 female) showed excessive alcohol  
13 consumption (AUDIT scores  $\geq 4$  for women,  $\geq 5$  for men). PEth 16:0/18:1 values ranged from  
14 not detected to 937 ng/mL (mean 110 ng/mL). PEth 16:0/18:2 values ranged from not  
15 detected to 472 ng/mL (mean 63.2 ng/mL). Among those persons with excessive alcohol  
16 consumption 95.9 % (n = 71) had positive PEth results (PEth  $\geq 10.0$  ng/mL (LOD)). Detailed  
17 information can be found in table 1 and figure 2.

18 Five data points of figure 2 (PEth 16:0/18:1 and PEth 16:0/18:2, respectively) were outside  
19 the axis limit in the column of moderate and excessive alcohol consumption (moderate: 1.  
20 PEth 16:0/18:1: 414 ng/mL, PEth 16:0/18:2: 224 ng/mL; 2. PEth 16:0/18:1: 619 ng/mL, PEth  
21 16:0/18:2: 286 ng/mL; excessive: 3. PEth 16:0/18:1: 506 ng/mL, whereas PEth 16:0/18:2: 168  
22 ng/mL is still included; 4. PEth 16:0/18:1: 524 ng/mL, PEth 16:0/18:2: 376 ng/mL; 5. PEth  
23 16:0/18:1: 937 ng/mL, PEth 16:0/18:2: 472 ng/mL).

24

25

26

1 **Comparison of PEth concentrations to self-reported alcohol consumption two weeks**  
2 **prior to blood sampling**

3 12 persons (6 female and 6 male) of the 300 participating volunteers had not-detectable PEth  
4 concentrations (< 10 ng/mL for both PEth 16:0/18:1 and 16:0/18:2) and reported abstinence.

5 A total of 85 test persons had (PEth not detected) but reported drinking in the last two weeks.

6 These persons were mainly light drinkers (35 female, 49 male, 1 not specified) with AUDIT-

7 C scores were between 1 - 4, and averaged amounts of consumed alcohol in the past two

8 weeks ranged from 0 – 10 g/day. Concerning PEth 16:0/18:2 concentrations, 121 test persons

9 (47 female, 71 male, 3 not specified) had PEth 16:0/18:2 < 10.0 ng/mL, with AUDIT-C

10 scores between 1 - 5, and averaged amounts of consumed alcohol in the past two weeks

11 ranged from 0 – 10 g/day.

12 161 test persons had PEth 16:0/18:1 concentrations between 10.0 and 100 ng/mL (44 female,

13 115 male, 2 not specified). AUDIT-C scores were between 1-7, and averaged amounts of

14 consumed alcohol in the past two weeks ranged from 0 – 50 g/day. Concerning

15 concentrations, 150 test persons had PEth 16:0/18:2 between 10.0 and 100 ng/mL (38 female,

16 112 male). AUDIT-C scores were between 1-8, and averaged amounts of consumed alcohol in

17 the past two weeks ranged from 0 – 50 g/day. According to WHO (WHO, 2000), these ranges

18 are still in the region of medium risk consumption (41 to 60 g) for men, but for women this

19 amount lies in the range of high risk consumption (41 to 60 g).

20 42 test persons had PEth 16:0/18:1 concentrations > 100 ng/mL (9 female, 33 male), whereas

21 only 17 test persons had PEth 16:0/18:2 concentrations > 100 ng/mL (3 female, 14 male).

22 AUDIT-C scores were between 2-8, and averaged amounts of consumed alcohol in the past

23 two weeks ranged from 2.0 – 33 g/day.

24

25

26

## 1 **Statistic evaluation**

2 We found statistically significant correlations of PEth 16:0/18:1, as well as of PEth 16:0/18:2  
3 concentrations, with the measures of alcohol consumption (AUDIT-C and self-reported  
4 amount of consumed alcohol in the last two weeks prior to blood sampling, respectively)  
5 (table 2). The correlation of PEth 16:0/18:1 with the measures of alcohol consumption led to  
6 Spearman correlation coefficients of  $r > 0.68$  (AUDIT-C) and  $r > 0.70$  (self-reported amount  
7 of consumed alcohol)); the respective correlation of PEth 16:0/18:2 led to Spearman  
8 correlation coefficients of  $r > 0.65$  (AUDIT-C) and  $r > 0.69$  (self-reported amount of  
9 consumed alcohol)). There was a slightly higher significance of PEth to self-reported amounts  
10 of consumed alcohol than to the AUDIT-C score.

11 The Kruskal-Wallis-Test showed significant differences between the three alcohol  
12 consumption groups (abstinence, moderate and excessive alcohol consumption) with  
13 significantly varying P values (PEth 16:0/18:1: P value  $< 0.0001$ ; PEth 16:0/18:2: P value  $<$   
14  $0.0001$ ).

15 For the extrapolation of reference values of PEth 16:0/18:1 and PEth 16:0/18:2 for moderate  
16 alcohol consumption, the group of 214 volunteers (71 % of 300 participants) was evaluated,  
17 as these participants reported AUDIT-C scores corresponding to moderate alcohol  
18 consumption. 95 % of the samples ranged between “not detected” and 112 ng/mL for PEth  
19 16:0/18:1, and between “not detected” and 67.0 ng/mL for PEth 16:0/18:2 (figure 2).

20 Applying the suggested PEth 16:0/18:1 range of “not detected” to 112 ng/mL, which indicates  
21 abstinence or moderate alcohol consumption, to all participants of this study ( $n = 300$ ), 87.7  
22 % of the test persons ( $n = 263$ ) were within this range, with average amounts of consumed  
23 alcohol of 0 – 50 g/day and a mean amount of consumed alcohol of 5.7 g/day.

24 However, only 36.5 % of those persons with AUDIT-C scores  $\geq 4$  (women) and  $\geq 5$  (men),  
25 respectively, had PEth 16:0/18:1 concentrations above 112 ng/mL and PEth 16:0/18:2  
26 concentrations above 67.0 ng/mL.

## 1 **Discussion**

2

3 The aim of this study was to establish reference levels of PEth 16:0/18:1 and PEth 16:0/18:2  
4 in comparison to AUDIT-C and self-reports reflecting different drinking habits, such as  
5 abstinence, moderate and excessive alcohol consumption in 300 volunteers. As this study was  
6 anonymous (only information about age and sex was available) and without consequences for  
7 the volunteers whatever resulted from the AUDIT-C, self-reports or PEth analysis, the  
8 answers to the questionnaires were supposed to be correct with no significant over- or under-  
9 reporting. However, there were two cases of clear under-reporting of alcohol consumption  
10 habits. For one female participant, AUDIT-C score was 2 (moderate alcohol consumption)  
11 and self-reported mean alcohol consumption was 2.1 g/day, whereas high PEth concentrations  
12 were found (PEth 16:0/18:1: 414 ng/mL, PEth 16:0/18:2: 224 ng/mL). For one male  
13 participant with an AUDIT-C score of 3 (moderate alcohol consumption) and self-reported  
14 mean alcohol consumption of 7.8 g/day, PEth concentrations were even higher (PEth  
15 16:0/18:1: 619 ng/mL, PEth 16:0/18:2: 286 ng/mL). In both cases, PEth concentrations  
16 clearly exceeded the above mentioned thresholds for both PEth homologues for differentiation  
17 of moderate and excessive consumption.

18 All tested persons who declared to be abstainers were PEth negative, resulting in an  
19 accordance of 100 % between AUDIT-C score and PEth regarding abstinence, meaning that  
20 no “endogenous” PEth levels have been detected in the blood of abstainers. 91.7 % (n = 11)  
21 of the test persons (n = 12), who did not consume alcohol in the two weeks prior to blood  
22 sampling for this study, but were classified as moderate drinkers according to AUDIT-C  
23 score, were PEth negative (PEth < 10.0 ng/mL). In addition, 85 test persons, who declared  
24 drinking in the last two weeks (mean alcohol amount ranging from 0 – 10 g/day), had  
25 negative PEth 16:0/18:1 results. As a consequence, persons who are negative for PEth do not  
26 necessarily have to be abstainers, but might belong to the group of moderate consumers.

1 However, there were also some PEth negative volunteers in the group classified as excessive  
2 consumers (PEth 16:0:18:1 < 10.0 ng/mL: n = 9; PEth 16:0:18:2 < 10.0 ng/mL: n = 30).  
3 PEth levels reflecting moderate alcohol consumption (AUDIT-C score 1 – 3 (women) and 1 –  
4 4 (men)) ranged from not detected to 112 ng/mL for PEth 16:0/18:1, and from not detected to  
5 67.0 ng/mL for PEth 16:0/18:2, corresponding to 95 % of the tested persons in the group with  
6 AUDIT-C scores representing moderate alcohol consumption.

7 This cut-off concentration for PEth 16:0/18:1 (112 ng/mL) for the differentiation of moderate  
8 and excessive alcohol consumption was higher than the findings from a recently published  
9 study among a group of persons with hepatic diseases, in which a threshold of 80.0 ng/mL  
10 PEth 16:0/18:1 for averaging at least 4 drinks/day was identified, which would impair or  
11 cause liver disease (Stewart et al., 2014).

12 After evaluating the suggested PEth 16:0/18:1 cut-off level of 112 ng/mL by ROC analysis  
13 (figure 3) (comparing PEth to AUDIT-C scores), 95 % of the group of moderate consumers  
14 and abstainers had PEth 16:0/18:1 concentrations ranging between not detected and 112  
15 ng/mL, but only 36.5 % of the participants with AUDIT-C scores representing excessive  
16 alcohol consumption had PEth 16:0/18:1 concentrations above 112 ng/mL.

17 Comparing PEth 16:0/18:2 to AUDIT-C scores by ROC analysis (figure 4) to evaluate the  
18 suggested PEth 16:0/18:2 cut-off level of 67.0 ng/mL, 95 % of the group of “social  
19 consumers” and abstainers had PEth 16:0/18:2 concentrations ranging between not detected  
20 and 67.0 ng/mL, and 36.5 % of the participants with AUDIT-C scores representing excessive  
21 alcohol consumption had PEth 16:0/18:2 concentrations above 67.0 ng/mL.

22 Using the PEth 16:0/18:1 cut-off of 80.0 ng/mL from the literature (Stewart et al., 2014) for  
23 our test persons, only 93.3 % of the participants would be identified as “social consumers”  
24 and abstainers, but 46.0 % of the participants with AUDIT-C scores representing excessive  
25 alcohol consumption would be classified correctly for excessive drinking habits.

1 A limitation of our study is that excessive alcohol consumption is underrepresented in the  
2 participants of this study (only 74 of 300 test persons). Therefore, it was not our aim to  
3 evaluate a threshold for the detection of excessive alcohol consumption with high sensitivity.  
4 We found overlapping ranges of PEth concentrations between the groups with moderate and  
5 excessive alcohol consumption, respectively. This is mainly due to the fact, that AUDIT-C  
6 covers a rather long time period (1 year), whereas PEth can only be detected for some days to  
7 weeks after excessive drinking behavior.

8 The correlations of PEth 16:0/18:1 and PEth 16:0/18:2 to AUDIT-C score and to self-reports  
9 (amount of consumed alcohol in the last two weeks prior to blood sampling) were statistically  
10 significant – with slightly higher correlations of PEth to the amount of consumed alcohol than  
11 to AUDIT-C scores, as PEth can only cover the alcohol consumption of the last days to  
12 weeks. This is in agreement with other studies among moderate drinkers (Stewart et al., 2014,  
13 Bajunirwe et al., 2014, Jain et al., 2014, Kechagias et al., 2015).

14 Under-reporting of alcohol consumption might lead to unexpectedly (high) PEth  
15 concentrations – however, “false-positive” PEth results might be another source of  
16 discrepancies. To the best of our knowledge, with the exception of post-sampling in-vitro  
17 formation of PEth in the presence of ethanol in blood, no other sources for false-positive PEth  
18 results have been detected yet, but relevant amounts of hidden alcohol in nutrition or  
19 medication (which might not be reported in a questionnaire) as well as inter-individual  
20 variations of enzyme activities for the formation of PEth might cause unexpected elevated  
21 concentration levels of PEth.

22

23 We conclude, that the alcohol biomarker PEth can be applied to confirm or to disprove self-  
24 reported alcohol consumption. In the present study a high correlation between PEth values  
25 and self-reports was found. PEth 16:0/18:1 values below the LOD of 10.0 ng/mL are strongly  
26 correlated with light drinking habits ( $\leq 10$  g pure alcohol/day) or abstinence.

1 As PEth 16:0/18:1 concentrations ranging from not detected to 112 ng/mL (for PEth 16:0/18:2  
2 values from not detected to 67.0 ng/mL) correspond to 95 % of the tested persons in the group  
3 with AUDIT-C scores representing moderate alcohol consumption, we suggest to employ  
4 these reference ranges as optimal indicators for alcohol consumption in a harmless way ( $\leq 20$   
5 g/day females,  $\leq 40$  g/day males, (WHO, 2000)). PEth values above this range point to recent  
6 excessive alcohol consumption (AUDIT-C score  $\geq 4$  females, AUDIT-C score  $\geq 5$  males).

7

8

9

10

11

12

13

#### 14 **Tables**

15

16 **Table 1.** Descriptive statistics for the volunteers regarding drinking types according to  
17 AUDIT-C

18

19 **Table 2.** Correlations between the measures of alcohol consumption and the PEth 16:0/18:1  
20 and PEth 16:0/18:2 values

21

22



1 **Figure Legends**

2

3 **Figure 1.** Distribution of the AUDIT-C (AUDIT: Alcohol Use Disorders Identification Test)  
4 scores.

5

6 **Figure 2.** Classification of PEth 16:0/18:1 and 16:0/18:2 values in comparison to  
7 consumption according to AUDIT-C score. The crosses represent the mean, the boxes show  
8 the interquartile range, the line within the boxes is the median, and the tails represent the 5 –  
9 95 percentiles with the dots as outliers.

10 Five data points (PEth 16:0/18:1 and PEth 16:0/18:2, respectively) are outside the axis limit in  
11 the column of moderate and excessive alcohol consumption (moderate: 1. PEth 16:0/18:1: 414  
12 ng/mL, PEth 16:0/18:2: 224 ng/mL; 2. PEth 16:0/18:1: 619 ng/mL, PEth 16:0/18:2: 286  
13 ng/mL; excessive: 3. PEth 16:0/18:1: 506 ng/mL, whereas PEth 16:0/18:2: 168 ng/mL is still  
14 included; 4. PEth 16:0/18:1: 524 ng/mL, PEth 16:0/18:2: 376 ng/mL; 5. PEth 16:0/18:1: 937  
15 ng/mL, PEth 16:0/18:2: 472 ng/mL).

16

17 **Figure 3.** ROC analysis for PEth 16:0/18:1: For the threshold of 112 ng/mL, a specificity of  
18 96.6 % and a sensitivity of 22.9 % were determined

19

20

21 **Figure 4.** ROC analysis for PEth 16:0/18:2: For the threshold of 67 ng/mL, a specificity of  
22 96.6 % and a sensitivity of 22.1 % were determined

23

24

## 1 **References**

- 2 Aradottir S, Asanovska G, Gjerss S, Hansson P & Alling C (2006). Phosphatidylethanol  
3 (PEth) concentrations in blood are correlated to reported alcohol intake in alcohol-  
4 dependent patients. *Alcohol Alcohol*, 41, 431-7.
- 5 Bajunirwe F, Haberer JE, Boum Y, 2nd, Hunt P, Mocello R, Martin JN, Bangsberg DR &  
6 Hahn JA (2014). Comparison of self-reported alcohol consumption to  
7 phosphatidylethanol measurement among HIV-infected patients initiating antiretroviral  
8 treatment in southwestern Uganda. *PLoS One*, 9, e113152.
- 9 Boscolo-Berto, R., Viel, G., Montisci, M., Terranova, C., Favretto, D. & Ferrara, S. D. 2013.  
10 Ethyl glucuronide concentration in hair for detecting heavy drinking and/or abstinence: a  
11 meta-analysis. *Int J Legal Med*, 127, 611-9.
- 12 Bush K, Kivlahan DR, Mcdonell MB, Fihn SD & Bradley KA (1998). The AUDIT alcohol  
13 consumption questions (AUDIT-C): an effective brief screening test for problem  
14 drinking. *Arch Intern Med*, 158, 1789-95.
- 15 Dresen S, Weinmann W & Wurst FM (2004). Forensic confirmatory analysis of ethyl sulfate  
16 – a new marker for alcohol consumption – by liquid-chromatography/electrospray  
17 ionization/tandem mass spectrometry. *J Am Soc Mass Spectrom*, 15, 1644-8.
- 18 Gnann H (2011). Phosphatidylethanol - Ein Alkoholkonsummarker auf dem Weg in die  
19 klinisch-forensische Routinediagnostik. Dissertation, University of Freiburg, Germany.  
20 [http://www.freidok.uni-freiburg.de/volltexte/8428/pdf/Diss\\_Heike\\_Gnann.pdf](http://www.freidok.uni-freiburg.de/volltexte/8428/pdf/Diss_Heike_Gnann.pdf). Accessed  
21 15 April 2014.
- 22 Gnann H, Weinmann W, Engelmann C, Wurst FM, Skopp G, Winkler M, Thierauf A,  
23 Auwarter V, Dresen S & Ferreiros Bouzas N (2009). Selective detection of  
24 phosphatidylethanol homologues in blood as biomarkers for alcohol consumption by LC-  
25 ESI-MS/MS. *J Mass Spectrom*, 44, 1293-9.
- 26 Gnann H, Weinmann W & Thierauf A (2012). Formation of phosphatidylethanol and its  
27 subsequent elimination during an extensive drinking experiment over 5 days. *Alcohol*  
28 *Clin Exp Res*, 36, 1507-11.
- 29 Halter CC, Dresen S, Auwaerter V, Wurst FM & Weinmann W (2008). Kinetics in serum and  
30 urinary excretion of ethyl sulfate and ethyl glucuronide after medium dose ethanol intake.  
31 *Int J Legal Med*, 122, 123-8.
- 32 Hannuksela ML, Liisanantti MK, Nissinen AE & Savolainen MJ (2007). Biochemical  
33 markers of alcoholism. *Clin Chem Lab Med*, 45, 953-61.



1 Schröck A, Thierauf-Emberger A, Schürch S & Weinmann W (2016c). Phosphatidylethanol  
2 (PEth) detected in blood for 3 to 12 days after single consumption of alcohol – a drinking  
3 study with 16 volunteers. *Int J Legal Med*, 131, 153-160.

4 Schröck A, Thierauf A, Wurst FM, Thon N & Weinmann W (2014). Progress in monitoring  
5 alcohol consumption and alcohol abuse by phosphatidylethanol. *Bioanalysis*, 6, 2285-94.

6 Skipper GE, Thon N, Dupont RL, Baxter L & Wurst FM (2013). Phosphatidylethanol: the  
7 potential role in further evaluating low positive urinary ethyl glucuronide and ethyl  
8 sulfate results. *Alcohol Clin Exp Res*, 37, 1582-6.

9 Sobell LC & Sobell MB (1992). Timeline Follow-back: A technique for assessing self-  
10 reported ethanol consumption. In J. Allen & R. Z. Litten (Eds.), *Measuring Alcohol*  
11 *Consumption: Psychosocial and Biological Methods* (pp. 41-72). Totowa, NJ: Humana  
12 Press.

13 Society of Hair Testing (SOHT) (2014). 2014 Consensus for the Use of Alcohol Markers in  
14 Hair for Assessment of both Abstinence and Chronic Excessive Alcohol Consumption.  
15 [http://www.soht.org/images/pdf/2014%20Alcohol%20markers%20revision%2013JUN14](http://www.soht.org/images/pdf/2014%20Alcohol%20markers%20revision%2013JUN14%20FINAL.pdf)  
16 [%20FINAL.pdf](http://www.soht.org/images/pdf/2014%20Alcohol%20markers%20revision%2013JUN14%20FINAL.pdf). Accessed 05 April 2016.

17 Stewart SH, Koch DG, Willner IR, Anton RF & Reuben A (2014). Validation of blood  
18 phosphatidylethanol as an alcohol consumption biomarker in patients with chronic liver  
19 disease. *Alcohol Clin Exp Res*, 38, 1706-11.

20 Thierauf A, Grosse-Perdekamp M, Weinmann W & Auwärter V (2011).  
21 Alkoholkonsummarker. *Rechtsmedizin*. DOI 10.1007/s00194-010-0729-6.

22 Varga A, Hansson P, Johnson G & Alling C (2000). Normalization rate and cellular  
23 localization of phosphatidylethanol in whole blood from chronic alcoholics. *Clin Chim*  
24 *Acta*, 299, 141-50.

25 Viel G, Boscolo-Berto R, Cecchetto G, Fais P, Nalesso A & Ferrara SD (2012).  
26 Phosphatidylethanol in blood as a marker of chronic alcohol use: a systematic review and  
27 meta-analysis. *Int J Mol Sci*, 13, 14788-812.

28 World Health Organization (WHO) (2000). International Guide for Monitoring Alcohol  
29 Consumption And Related Harm.  
30 [http://apps.who.int/iris/bitstream/10665/66529/1/WHO\\_MSD\\_MSB\\_00.4.pdf](http://apps.who.int/iris/bitstream/10665/66529/1/WHO_MSD_MSB_00.4.pdf). Accessed  
31 13 April 2016, (pp. 50 – 54).

32 World Health Organization (WHO) (2001). The Alcohol Use Disorders Identification Test.  
33 [http://www.talkingalcohol.com/files/pdfs/WHO\\_audit.pdf](http://www.talkingalcohol.com/files/pdfs/WHO_audit.pdf). Accessed 21 January.2016.

- 1 Winkler M, Skopp G, Alt A, Miltner E, Jochum T, Daenhardt C, Sporkert F, Gnann H,  
2 Weinmann W & Thierauf A (2013). Comparison of direct and indirect alcohol markers  
3 with PEth in blood and urine in alcohol dependent inpatients during detoxication. *Int J*  
4 *Legal Med*, 127, 761-8.
- 5 Wurst FM, Thon N, Yegles M, Schröck A, Preuss UW & Weinmann W (2015). Ethanol  
6 Metabolites: Their Role in the Assessment of Alcohol Intake. *Alcohol Clin Exp Res*, 39,  
7 2060-72.
- 8 Zheng Y, Beck O & Helander A (2011). Method development for routine liquid  
9 chromatography–mass spectrometry measurement of the alcohol biomarker  
10 phosphatidylethanol (PEth) in blood. *Clin Chim Acta*, 412, 1428-1435.
- 11