

Tumour Review

Palliative treatment of germ cell cancer

Christoph Oing^{a,1}, Patrizia Giannatempo^{b,1}, Friedemann Honecker^c, Karin Oechsle^a,
Carsten Bokemeyer^a, Jörg Beyer^{d,*}

^a Department of Oncology, Hematology and Bone Marrow Transplantation with Department of Pneumology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

^b Department of Medical Oncology, Fondazione IRCCS, Istituto Nazionale dei Tumori, Milan, Italy

^c ZeTUP Breast and Tumor Center, St. Gallen, Switzerland

^d Department of Medical Oncology, Inselspital, Bern University Hospital, University of Bern, Freiburgstrasse, CH-3031 Bern, Switzerland



ARTICLE INFO

Keywords:

Germ cell cancer
Germ cell tumor
Platinum-refractory
Palliative treatment
Drug therapy

ABSTRACT

Most patients with metastatic germ-cell cancer (GCC) can be cured by cisplatin-based combination chemotherapy. Yet, about 10–15% of metastatic GCC patients will eventually die of their disease. This narrative review focuses on treatment options when cure is no longer realistic.

Introduction

Germ-cell cancers (GCC) are a unique example of a curable malignant disease [1]. Therefore, palliative treatment is rarely being discussed. Yet, 10–15% of metastatic patients will fail first as well as subsequent lines of platinum-based treatments and eventually die of their disease [2]. Palliative treatment of patients with GCC remains challenging with little high-level evidence to support decision making in this situation.

Who is “palliative” among GCC patients?

There is good evidence that subsets of patients with multiple relapses, and, to a lesser extent, patients with primary cisplatin-refractory disease may still be cured by conventional-dose (CDCT), high-dose chemotherapy (HDCT) and/or desperation surgery delivered as first, second or subsequent salvage treatment [2–8]. Therefore, patients will usually have had multiple lines of chemotherapy including HDCT before being considered palliative. The same applies to patients unfit for HDCT failing three or more lines of CDCT [2,9]. The majority of patients will have received modern type cisplatin- and ifosfamide-based salvage regimens such as cisplatin, ifosfamide and vinblastine (VeIP), etoposide (VIP) or paclitaxel (TIP) [2,9]. Desperation surgery might still salvage individual patients of these two cohorts [5–7]. However, if desperation surgery cannot successfully be performed, patients failing

HDCT or three or more lines of CDCT face little or no realistic hope for cure. Therefore, a palliative treatment population can be defined as patients failing salvage HDCT or patients unfit for HDCT who fail three or more lines of CDCT in whom desperation surgery cannot be successfully be performed. Rare exceptions to this definition may occur. With respect to the many early clinical trials investigating different systemic treatment approaches for refractory GCC patients, life-extension remains the primary goal for the majority of patients. As reported in the following sections, responses are generally short-lived, which is why alleviation of pain and other symptoms to enhance the patients' quality-of-life is also desired. However, quality-of-life was not routinely assessed as a trial end point in the majority of available studies and has to be considered individually as a main goal, whichever treatment is being chosen. To provide a better understanding of the efficacy and toxicities associated with palliative treatments, common adverse events for the different treatment options are mentioned in the tables.

Single agent chemotherapy

Single agent activity has been reported for paclitaxel [10–14], gemcitabine [15,16], oxaliplatin [17,18], oral etoposide [19], ifosfamide [20], and oral temozolomide [21] mainly in small phase I/II trials in patients with cisplatin-refractory disease. Responses were mainly partial remissions or disease stabilizations. However, sequential single agent treatment is an appropriate option in addition to best-supportive-

* Corresponding author at: Dept. of Medical Oncology, Inselspital, Bern University Hospital, University of Bern, Freiburgstrasse, CH-3031 Bern, Switzerland.

E-mail address: joerg.beyer@insel.ch (J. Beyer).

¹ Equal contribution.

Table 1
Single agent regimens.

Drug	Schedule		N Patients	Previous HDCT	ORR	Duration of benefit for responders (measure, median, range) [months]	Relevant toxicities ≥ grade 3	Ref	
Gemcitabine	1000 mg/m ² iv.	d1,8,15	q3w	31	71%	19%	TTF, 4+ (2–9+)	Granulocytopenia 13%	15
	1200 mg/m ² iv.	d1,8,15	q3w	20	55%	15%	PFS, n.r. (2–6)	Thrombocytopenia 22%	16
Oxaliplatin	60 mg/m ² iv.	d1,8,15	q3w	16	78%	6%	PFS, n.r. (2–8)	Neutropenia 10%	17
	130 mg/m ² iv.	d1	q2w	16		19%		Thrombocytopenia 30%	
Paclitaxel	130 mg/m ² iv.	d1	q3w	8	NR	25%	PFS, n.r.	Nausea 5%	18
	135–310 mg/m ² iv.	d1	q3w	10	NR	20%	DOR, n.r. (3–5)	Granulocytopenia 9%	14
	170 mg/m ² iv.	d1	q3w	18	16%	11%	PFS, n.r. (1.5–2)	Thrombocytopenia 16%	13
								Nausea/vomiting 6%	
								Neurotoxicity 3%	11
	225 mg/m ² iv.	d1	q3w	24	50%	25%	DOR, 8+ (3–16+)	n.r.	
250 mg/m ² iv.	d1	q3w	15	13%	13%	DOR, n.r. (9–10 weeks)	Granulocytopenia 90%	12	
250 mg/m ² iv.	d1	q3w	31	16%	26%	n.r.	Infection 10%	10	
Ifosfamide	2000 mg/m ² iv.	d1-5	q3w	30	NR	23%	DOR, 3.5 (2–5.5)	Granulocytopenia 67%	20
							Anemia 7%		
Oral Etoposide	50 mg/m ² po.	d1-14	q3w	21	29%	14%	n.r.	Neutropenic fever 17%	19
							Neurotoxicity 13%		
Oral Temozolomide	150–200 mg/m ² po.	d1-5	q4w	20	40%	10%	DOR, 1.5 (3.5–9)	Granulocytopenia 50%	21
							Severe infections 12%		
							Neurotoxicity ≥ 'II 27%		
							Mucositis 'II/III 16%		
							Neutropenia 87%		
							Thrombocytopenia 73%		
							Peripheral neuropathy 'II/III 53%		
							Mainly hematologic		
							Neurotoxicity 'III 19%		
							Mucositis 'II/III 16%		
							Liver function test > 5xULN 10%		
							Neutropenia 52%		
							Thrombocytopenia 20%		
							Nausea/vomiting 43%		
							Granulocytopenia 36%		
							Febrile granulocytopenia 23%		
							Thrombocytopenia 10%		
							Anemia 5%		

DOR, duration of response; PFS, progression-free survival; TTF, time to treatment failure; n.r., not reported.

care in the palliative setting of patients with a poor performance status asking for further active treatment in order to avoid the more frequent side-effects of combination chemotherapy. Table 1 provides an overview of single-agent chemotherapy regimens.

Combination chemotherapy

Combinations of drugs with single-agent activity result in higher response rates compared to using single-agent treatment alone. The triple combination of gemcitabine, oxaliplatin and paclitaxel (GOP) reported the highest response rates so far with more than 50% objective responses and even few long-term remissions (Table 2) [22–25]. Doublet combinations of these drugs may also provide favorable responses ranging from 4% to 46% (Table 3) [26–31]. Other doublets include cisplatin, plus epirubicin [32] or irinotecan [33], or oxaliplatin together with the VEGF-targeted antibody bevacizumab [34] or again irinotecan [35]. Other triple combination regimens comprise of paclitaxel, cisplatin and gemcitabine (TPG) (ORR 49%) [36], or oxaliplatin, irinotecan and paclitaxel (IPO) followed by one cycle of topotecan-based HDCT (ORR 49%) [37]. Recently, clinical activity and good tolerability of the combination of etoposide, methotrexate and actinomycin D, alternating with cyclophosphamide and vincristine (EMA/CO), as established for female gestational trophoblastic disease, has been reported in patients with refractory GCC and elevated levels of human chorionic gonadotrophin [38] (Table 2).

Taken together, combination chemotherapy seems to be the most promising palliative approach in medically fit patients with a good performance status in whom the toxicities of a doublet or a triple combination seem acceptable. Direct comparisons of combination

treatments have not been published, and none of the regimens have proven unequivocal superiority.

Targeted treatments and immunotherapy

Investigational agents include tyrosine kinase inhibitors, i.e. sunitinib [39,40], sorafenib [41], imatinib [42], tivantinib [43] and pazopanib [44] as well as anti-angiogenic agents, i.e. thalidomide [45] and lenalidomide [46], and the mTOR inhibitor everolimus [47] amongst many others (Table 4). However, apart from single case reports, all these agents failed to induce clinically meaningful responses in unselected patient cohorts [48]. Similarly, antibody-drug conjugate brentuximab-vedotin was investigated in CD30-positive, refractory GCC patients. Some responses were observed but tended to be short-lived [49,50]. Thus, the results of investigational agents are conflicting, and at present these drugs have no established role in the palliative treatment setting of GCC.

Immunomodulatory treatment with immune checkpoint inhibitors active against the programmed- cell death receptor (PD-1) or its ligand (PD-L1) are currently being investigated also in refractory GCC as many tumors were shown to express PD-L1 [51]. However, application of the PD-1-directed monoclonal antibody pembrolizumab failed to induce clinically meaningful responses in a phase II trial enrolling 12 patients, which was terminated prematurely [52]. In another phase II trial, the anti-PD-L1 antibody Durvalumab alone yielded no responses among 9 GCC patients, whereas the combination with the CTLA-4 antibody tremelimumab achieved clinical responses in 2 out of 9 heavily pretreated GCC patients [53]. Other immune checkpoint inhibitors are currently under investigation, i.e. single agent treatment with the PD-L1 antibody

Table 2
Triple-drug regimens.

Drug	Schedule		Patients	Previous HDCT	ORR	Duration of benefit for responders (measure, median, range) [months]	Relevant toxicities \geq grade 3	Ref	
Gemcitabine Oxaliplatin Paclitaxel	800 mg/m ^b iv. 130 mg/m ^b iv. 80 mg/m ^b iv.	d1&8 d1 d1&8	q3w q3w q3w	41	78%	51%	PFS, 3 (1–17+)	Granulocytopenia 15% Anemia 7% Thrombocytopenia 49% Neurotoxicity 2% Nausea/Emesis 2% Diarrhea 2%	24
Gemcitabine Oxaliplatin Paclitaxel	800 mg/m ^b iv. 100–125 mg/m ^b iv. 170 mg/m ^b iv.	d1&8 d1 d1	q2w q2w q2w	30	20%	31%	PFS, 6.5 (95%CI, 3.2–27.5)	Neutropenia 17% Febrile Neutropenia 7% Gastrointestinal ^{II/III} 12% Neuropathy 5%	25
Cisplatin	50 mg/m ^b iv.	d1&8	q3w	75	13%	DCR 67%	PFS, 5 (95%CI, 4–8)	Anemia 28% Neutropenia 29%	36
Gemcitabine Paclitaxel	800 mg/m ^b iv. 80 mg/m ^b iv.	d1&8 d1&8	q3w q3w					Febrile Neutropenia 5% Thrombocytopenia 40% Ototoxicity 4%	
Oxaliplatin	100 mg/m ^b iv.	d1	q3w	43	0% (74% subsequent HDCT)	49%	2-year PFS 21%, (median n.r.)	Neutropenia 33%	37
Irinotecan Paclitaxel	200 mg/m ^b iv. 80 mg/m ^b iv.	d1 d1,8,15	q3w q3w					Thrombocytopenia 16% Neuropathy 2% Nause/Emesis 4% Diarrhea 12%	
Etoposide Methotrexate ^a Folinic acid ^b Actinomycin D Vincristine Cyclophosphamide	100 mg/m ^b iv. 300 mg/m ^b iv. 15 mg \times 4 iv./po. 0.5 mg/m ^b iv. 1 mg/m ^b iv. 600 mg/m ^b iv.	d1&2 d1 d2&3 d1&2 d8 d8	q2w q2w q2w q2w q2w q2w	41	39%	29%	PFS, 3 (95%CI, 2–4)	Hematologic 39%	38

DCR, disease control rate (complete + partial remissions + stable disease); DOR, duration of response; PFS, progression-free survival; TTF, time to treatment failure, HDCT, high-dose chemotherapy.

^a Methotrexate given as continuous infusion over 12 h.

^b Folinic acid started 24 h after methotrexate for 4 doses every 12 h.

Atezolizumab (NCT02458638), or the PD-1 antibody Nivolumab (NCT02832167) in phase II basket trials. Moreover, combined treatment approaches, such as the anti-PD-1 antibody Nivolumab plus anti-CTLA-4 antibody Ipilimumab, are also currently being evaluated (NCT02834013).

Choosing the right treatment for the right patient

Finding the right treatment for the right patient at the right time is obviously key to any palliative strategy in GCC patients. A formal comparison of available palliative regimens was not attempted in this review, as cohorts of published series are notoriously small and pre-treatments too heterogeneous to allow a meaningful approach. However, the large number of small studies indicates the lack of unequivocal superiority of any given regimen. Many palliative patients will receive sequential lines of treatments often switching from a combination to a single-agent regimen until available options are exhausted. An algorithm to support treatment selection is given in Fig. 1.

Challenging scenarios in palliative GCC patients

Brain metastases

Palliative patients with GCC and brain metastases should still receive radiotherapy for symptom control, delivered either as stereotactic or total brain irradiation depending on the number and location of the metastases as well as the local availability of the respective techniques. Patients who are “fit” for chemotherapy may be treated with multimodality strategies of chemotherapy and irradiation. In the palliative setting neurosurgery is rarely indicated [54].

Bone metastases

Patients suffering metastatic bone disease may undergo multimodal treatment including chemotherapy and radiotherapy to alleviate skeletal symptoms, if a limited number of symptomatic bone metastases are present. The use of anti-resorptive agents, e.g. bisphosphonates or RANKL-inhibitors can be considered, although data on their use in GCC patients is lacking [55].

Peripheral neuropathy

The majority of palliative GCC patients experience peripheral neuropathy as a result of exposure to cisplatin, to taxanes or to HDCT. Treatment options and high-level evidence are limited, but the use of co-analgesic medication, e.g. the antidepressants duloxetine, amitriptyline or venlafaxine and/or the anticonvulsants gabapentin or pregabalin may alleviate symptoms [56]. Often peripheral neuropathy will limit or prohibit the further use of cisplatin, oxaliplatin or the taxanes.

Early integration of palliative care and psychosocial support

Similar to patients suffering from other malignancies, palliative patients with GCC may benefit from early integration of a specialized palliative care team in order to professionally identify and treat many of the additional symptoms and needs common to palliative cancer patients [57]. Particularly, the young age of patients with GCC increases their vulnerability for psychological distress. Studies have emphasized a strong correlation between young age and higher levels of anxiety and depression among patients and their relatives [57]. Therefore, psychological support must be an integral part of any palliative strategy in GCC patients.

Table 3
Doublet regimens.

Drug	Schedule			Patients	Previous HDCT	ORR	Duration of benefit for responders (measure, median, range) [months]	Relevant toxicities ≥ grade 3	Ref
Gemcitabine Oxaliplatin	1000 mg/m ² iv. 130 mg/m ² iv.	d1&8 d1	q3w q3w	35	89%	46%	DFS, n.r. (2–16 +)	Leukocytopenia 54% Thrombocytopenia 48% Anemia 11% Febrile Neutropenia 9% Treatment-related death 3% Neurotoxicity 9% Nausea/Emesis 16% Diarrhea 6%	26
Gemcitabine Oxaliplatin	1000 mg/m ² iv. 130 mg/m ² iv.	d1&8 d1	q3w q3w	28	14%	32%	DFS, n.r. (2–28 +)	Neutropenia 62% Febrile Neutropenia 10% Thrombocytopenia 41% Anemia 21% Neurotoxicity 10% Nausea/Emesis 27% Fatigue 17%	27
Gemcitabine Oxaliplatin	1250 mg/m ² iv. 130 mg/m ² iv.	d1&8 d1	q3w q3w	18	22%	17%	DFS, n.r. (18+ - 44 +)	Neutropenia 39% Thrombocytopenia 22% Anemia 11% Peripheral Neuropathy 17% Asthenia 11%	28
Paclitaxel Oxaliplatin	175 mg/m ² iv. 130 mg/m ² iv.	d1 d1	q3w q3w	26	19%	4%	TTP, 1.4 (95%CI 0–14.8)	Nausea/Vomiting 5% Neutropenia 30% Thrombocytopenia 4% Anemia 12% Neuropathy 1% Nausea/Emesis 8% Fatigue 4%	29
Bevacizumab Oxaliplatin	10 mg/kg iv. 85 mg/m ² iv.	d1&8 d1	q3w q2w	24	54%	29%	DOR, 6 (4–22)	Peripheral neuropathy Hypertension Seizures Deep vein thrombosi	34
Oxaliplatin Irinotecan	85 mg/m ² iv. 80 mg/m ² iv.	d1&15 d1,8,15	q4w q4w	18	0	39%	DFS, n.r. (2–19 +)	Neutropenia 17% Thrombocytopenia 17% Anemia 12% Neurotoxicity 11% Nausea/Emesis 28% Diarrhea 22% Mucositis 11%	36
Cisplatin Epirubicin	20 mg/m ² iv. 90 mg/m ² iv.	d1-5 d1	q3w q3w	30	13%	57%	NED, n.r. (n.r. - 48 +)	Neutropenia 13% Febrile Neutropenia 3% Thrombocytopenia 7% Anemia 17% Neuropathy 7% Nausea/Vomiting 27% Diarrhea 3% Nephrotoxicity 7%	32
Cisplatin Irinotecan	20 mg/m ² iv. 100–150 mg/m ² iv.	d1-5 d1&15	q4w q4w	11	27%	45%	DOR, n.r. (12+ -140 +)	Neutropenia 18% Thrombocytopenia 17% Anemia 14% Nausea/Vomiting 8% Diarrhea 5%	33
Paclitaxel Gemcitabine	110 mg/m ² iv. 1000 mg/m ² iv.	d1,8,15 d1,8,15	q4w q4w	28	36%	21%	DOR, n.r. (2–25 +)	Neutropenia 54% Febrile Neutropenia 4% Thrombocytopenia 32% Anemia 11% Neuropathy 4% Nausea/Emesis 8% Fatigue 8%	30
Paclitaxel Gemcitabine	100 mg/m ² iv. 1000 mg/m ² iv.	d1,8,15 d1,8,15	q4w q4w	32	100%	31%	NED, n.r. (22+ -54 +)	n.r.	31

DFS, disease-free survival; DOR, duration of response; NED, no evidence of disease; PFS, progression-free survival; TTF, time to treatment failure; n.r., not reported.

Conclusion

Guidance of treatment and care of refractory GCC patients in the palliative setting is challenging and emotionally demanding. Although some cytotoxic agents have shown efficacy either as single agents or as part of drug combinations, survival is limited and durable remissions are rare. The most difficult decision is when to stop cancer-directed

treatments altogether and switch to best-supportive care alone. On the technical side, we suggest as one cornerstone of care close contact with an expert GCC center that cares for such patients regularly. Locally, early integration of specialized palliative care as outlined by the American Society of Oncology [58] is the second cornerstone that supports symptom control and alleviates suffering for all parties involved.

Table 4
Targeted treatment approaches and immunotherapy.

Drug	Molecular target	Patients (n)	ORR (%)	Med. PFS (months)	Ref
Sunitinib	VEGFR, KIT, PDGFR, RET	33	13	2.0	39
		10	0	n.r.	40
Sorafenib	VEGFR, KIT, PDGFR, RAF	18	0	n.r.	41
Pazopanib	VEGFR, PDGFR	9	0	n.r.	44
Imatinib mesylate	KIT	6	0	n.r.	42
Tivantinib	MET	21	0	1.0	43
Everolimus	mTOR	15	0	1.7	47
Thalidomide	Anti-angiogenic	15	0	3.0	45
Lenalidomide	Anti-angiogenic	4	0	n.r.	46
Brentuximab- vedotin	CD30	5	40	n.r.	49
		9	22	n.r.	50
Pembrolizumab	PD-1	12	0	n.r.	52
Durvalumab	PD-L1	9	0	n.r.	53
Durvalumab + Tremelimumab	PD-L1	9	22	n.r.	

CTLA-4, cytotoxic T-lymphocyte-associated protein 4; KIT, stem cell growth factor receptor; MET, hepatocyte growth factor receptor; ORR, objective response rate (complete and/or partial remissions); PD-1, programmed death 1; PD-L1, programmed death ligand 1; PDGFR, platelet-derived growth factor receptor; PFS, progression-free survival; RAF, rapidly accelerated fibrosarcome oncogene family; RET, rearranged during transfection proto-oncogene; VEGFR, vascular endothelial growth factor receptor; n.r., not reported.




Fig. 1. Treatment algorithm for selection of palliative treatments.

References

- [1] Einhorn LH. Treatment of testicular cancer: a new and improved model. *J Clin Oncol* 1990;8:1777–81.
- [2] International Prognostic Factors Study G, Lorch A, Beyer J, et al. Prognostic factors in patients with metastatic germ cell tumors who experienced treatment failure with cisplatin-based first-line chemotherapy. *J Clin Oncol* 2010;28:4906–11.
- [3] Lorch A, Bascoul-Mollevi C, Kramar A, et al. Conventional-dose versus high-dose chemotherapy as first salvage treatment in male patients with metastatic germ cell tumors: evidence from a large international database. *J Clin Oncol* 2011;29:2178–84.
- [4] Lorch A, Neubauer A, Hackenthal M, et al. High-dose chemotherapy (HDCT) as second-salvage treatment in patients with multiple relapsed or refractory germ-cell tumors. *Ann Oncol* 2010;21:820–5.
- [5] Murphy BR, Breeden ES, Donohue JP, et al. Surgical salvage of chemorefractory germ cell tumors. *J Clin Oncol* 1993;11:324–9.
- [6] Albers P, Ganz A, Hannig E, et al. Salvage surgery of chemorefractory germ cell tumors with elevated tumor markers. *J Urol* 2000;164:381–4.
- [7] Beck SD, Foster RS, Bihle R, et al. Pathologic findings and therapeutic outcome of desperation post-chemotherapy retroperitoneal lymph node dissection in advanced germ cell cancer. *Urol Oncol* 2005;23:423–30.
- [8] Oechsle K, Kollmannsberger C, Honecker F, et al. Long-term survival after treatment with gemcitabine and oxaliplatin with and without paclitaxel plus secondary surgery in patients with cisplatin-refractory and/or multiply relapsed germ cell tumors.

- Eur Urol 2011;60:850–5.
- [9] Lorch A, Kleinhans A, Kramar A, et al. Sequential versus single high-dose chemotherapy in patients with relapsed or refractory germ cell tumors: long-term results of a prospective randomized trial. *J Clin Oncol* 2012;30:800–5.
- [10] Motzer RJ, Bajorin DF, Schwartz LH, et al. Phase II trial of paclitaxel shows anti-tumor activity in patients with previously treated germ cell tumors. *J Clin Oncol* 1994;12:2277–83.
- [11] Bokemeyer C, Beyer J, Metzner B, et al. Phase II study of paclitaxel in patients with relapsed or cisplatin-refractory testicular cancer. *Ann Oncol* 1996;7:31–4.
- [12] Nazario A, Amato RJ, Hutchinson L, et al. Paclitaxel in extensively pretreated nonseminomatous germ cell tumors. *Urol Oncol* 1995;1:184–7.
- [13] Sandler AB, Cristou A, Fox S, et al. A phase II trial of paclitaxel in refractory germ cell tumors. *Cancer* 1998;82:1381–6.
- [14] Bokemeyer C, Schmoll HJ, Natt F, et al. Preliminary results of a phase I/II trial of paclitaxel in patients with relapsed or cisplatin-refractory testicular cancer. *J Cancer Res Clin Oncol* 1994;120:754–7.
- [15] Bokemeyer C, Gerl A, Schoffski P, et al. Gemcitabine in patients with relapsed or cisplatin-refractory testicular cancer. *J Clin Oncol* 1999;17:512–6.
- [16] Einhorn LH, Stender MJ, Williams SD. Phase II trial of gemcitabine in refractory germ cell tumors. *J Clin Oncol* 1999;17:509–11.
- [17] Kollmannsberger C, Rick O, Derigs HG, et al. Activity of oxaliplatin in patients with relapsed or cisplatin-refractory germ cell cancer: a study of the German Testicular Cancer Study Group. *J Clin Oncol* 2002;20:2031–7.
- [18] Fizazi K, Culine S, Chen I. Oxaliplatin in non-seminomatous germ-cell tumors. *Ann Oncol* 2004;15:1295.
- [19] Miller JC, Einhorn LH. Phase II study of daily oral etoposide in refractory germ cell tumors. *Semin Oncol* 1990;17:36–9.
- [20] Wheeler BM, Loehrer PJ, Williams SD, Einhorn LH. Ifosfamide in refractory male germ cell tumors. *J Clin Oncol* 1986;4:28–34.
- [21] Maroto P, Huddart R, Garcia del Muro X, et al. Brief report: phase II multicenter study of temozolomide in patients with cisplatin-resistant germ cell tumors. *Oncology* 2011;80:219–22.
- [22] Necchi A, Nicolai N, Piva L, et al. Long-term results of a combination of paclitaxel, oxaliplatin and gemcitabine as far rescue in heavily pre-treated male germ cell tumors (GCT). *J Clin Oncol* 2008;26(suppl:abstr):16036.
- [23] Seidel C, Oechsle K, Lorch A, et al. Efficacy and safety of gemcitabine, oxaliplatin, and paclitaxel in cisplatin-refractory germ cell cancer in routine care-Registry data from an outcomes research project of the German Testicular Cancer Study Group. *Urol Oncol* 2016;34(168):e121–68.
- [24] Bokemeyer C, Oechsle K, Honecker F, et al. Combination chemotherapy with gemcitabine, oxaliplatin, and paclitaxel in patients with cisplatin-refractory or multiply relapsed germ-cell tumors: a study of the German Testicular Cancer Study Group. *Ann Oncol* 2008;19:448–53.
- [25] Sadeghi S, Quinn D, Tsao-Wei D, et al. Phase II study of gemcitabine, oxaliplatin, and paclitaxel (GOT) on a 2-weekly schedule in patients (pts) with refractory germ cell tumor (rGCT): Final results. *J Clin Oncol* 2013;31(suppl: abstr):4531.
- [26] Kollmannsberger C, Beyer J, Liersch R, et al. Combination chemotherapy with gemcitabine plus oxaliplatin in patients with intensively pretreated or refractory germ cell cancer: a study of the German Testicular Cancer Study Group. *J Clin Oncol* 2004;22:108–14.
- [27] Pectasides D, Pectasides M, Farmakis D, et al. Gemcitabine and oxaliplatin (GEMOX) in patients with cisplatin-refractory germ cell tumors: a phase II study. *Ann Oncol* 2004;15:493–7.
- [28] De Giorgi U, Rosti G, Aieta M, et al. Phase II study of oxaliplatin and gemcitabine salvage chemotherapy in patients with cisplatin-refractory nonseminomatous germ cell tumor. *Eur Urol* 2006;50:1032–8. discussion 1038–1039.
- [29] Theodore C, Chevreau C, Yataqhe Y, et al. A phase II multicenter study of oxaliplatin in combination with paclitaxel in poor prognosis patients who failed cisplatin-based chemotherapy for germ-cell tumors. *Ann Oncol* 2008;19:1465–9.
- [30] Hinton S, Catalano P, Einhorn LH, et al. Phase II study of paclitaxel plus gemcitabine in refractory germ cell tumors (E9897): a trial of the Eastern Cooperative Oncology Group. *J Clin Oncol* 2002;20:1859–63.
- [31] Einhorn LH, Brames MJ, Juliar B, Williams SD. Phase II study of paclitaxel plus gemcitabine salvage chemotherapy for germ cell tumors after progression following high-dose chemotherapy with tandem transplant. *J Clin Oncol* 2007;25:513–6.
- [32] Bedano PM, Brames MJ, Williams SD, et al. Phase II study of cisplatin plus epirubicin salvage chemotherapy in refractory germ cell tumors. *J Clin Oncol* 2006;24:5403–7.
- [33] Miki T, Mizutani Y, Nonomura N, et al. Irinotecan plus cisplatin has substantial antitumor effect as salvage chemotherapy against germ cell tumors. *Cancer* 2002;95:1879–85.
- [34] Jain A, Brames MJ, Vaughn D, Einhorn L. Phase II clinical trial of oxaliplatin and bevacizumab in refractory metastatic germ cell tumors (GCT). *J Clin Oncol* 2011;29(suppl:abstr):4579.
- [35] Pectasides D, Pectasides M, Farmakis D, et al. Oxaliplatin and irinotecan plus granulocyte-colony stimulating factor as third-line treatment in relapsed or cisplatin-refractory germ-cell tumor patients: a phase II study. *Eur Urol* 2004;46:216–21.
- [36] Necchi A, Nicolai N, Mariani L, et al. Combination of paclitaxel, cisplatin, and gemcitabine (TPG) for multiple relapses or platinum-resistant germ cell tumors: long-term outcomes. *Clin Genitourin Cancer* 2014;12(63–69):e61.
- [37] Badreldin W, Krell J, Chowdhury S, et al. The efficacy of irinotecan, paclitaxel, and oxaliplatin (IPO) in relapsed germ cell tumours with high-dose chemotherapy as consolidation: a non-cisplatin-based induction approach. *BJU Int* 2016;117:418–23.
- [38] Raggi D, Giannatempo P, Etoposide Miceli R, et al. methotrexate, and dactinomycin alternating with cyclophosphamide and vincristine (EMA/CO) for males with HCG-expressing, chemorefractory germ cell tumors (GCT): Long-term efficacy and safety outcomes. *J Clin Oncol* 2014;32:4561.
- [39] Oechsle K, Honecker F, Cheng T, et al. Preclinical and clinical activity of sunitinib in patients with cisplatin-refractory or multiply relapsed germ cell tumors: a Canadian Urologic Oncology Group/German Testicular Cancer Study Group cooperative study. *Ann Oncol* 2011;22:2654–60.
- [40] Feldman DR, Turkula S, Ginsberg MS, et al. Phase II trial of sunitinib in patients with relapsed or refractory germ cell tumors. *Invest New Drugs* 2010;28:523–8.
- [41] Skoneczna IA, Natarska U, Tacikowska M, et al. Sorafenib monotherapy in patients with inoperable/recurrent germ cell tumors (GCT) refractory to chemotherapy: Phase II study. *J Clin Oncol* 2014;32(suppl4:abstr):367.
- [42] Einhorn LH, Brames MJ, Heinrich MC, et al. Phase II study of imatinib mesylate in chemotherapy refractory germ cell tumors expressing KIT. *Am J Clin Oncol* 2006;29:12–3.
- [43] Feldman DR, Einhorn LH, Quinn DI, et al. A phase 2 multicenter study of tivantinib (ARQ 197) monotherapy in patients with relapsed or refractory germ cell tumors. *Invest New Drugs* 2013;31:1016–22.
- [44] Necchi A, Lo Vullo S, Giannatempo P, et al. Pazopanib in advanced germ cell tumors after chemotherapy failure: results of the open-label, single-arm, phase 2 Pazotest trial. *Ann Oncol* 2017;28:1346–51.
- [45] Rick O, Braun T, Siegert W, Beyer J. Activity of thalidomide in patients with platinum-refractory germ-cell tumours. *Eur J Cancer* 2006;42:1775–9.
- [46] Oechsle K, Bokemeyer C, Honecker F. Lenalidomide in patients with cisplatin-refractory and multiply relapsed germ cell tumors. *J Cancer Res Clin Oncol* 2010;136:165–7.
- [47] Mego M, Svetlovska D, Miskovska V, et al. Phase II study of everolimus in refractory testicular germ cell tumors. *Urol Oncol* 2016;34(122):e117–22.
- [48] Oing C, Kollmannsberger C, Oechsle K, Bokemeyer C. Investigational targeted therapies for the treatment of testicular germ cell tumors. *Expert Opin Investig Drugs* 2016;25:1033–43.
- [49] Necchi A, Magazzu D, Anichini A, et al. An open-label, single-group, phase 2 study of brentuximab vedotin as salvage therapy for males with relapsed germ-cell tumors (GCT): Results at the end of first stage (FM12GCT01). *J Clin Oncol* 2016;34: abstr:480.
- [50] Albany C, Einhorn L, Garbo L, et al. Treatment of CD30-expressing germ cell tumors and sex cord stromal tumors with brentuximab vedotin: identification and report of seven cases. *Oncologist* 2018;23:316–23.
- [51] Fankhauser CD, Curioni-Fontecedro A, Allmann V, et al. Frequent PD-L1 expression in testicular germ cell tumors. *Br J Cancer* 2015;113:411–3.
- [52] Adra N, Einhorn LH, Althouse SK, et al. Phase II trial of pembrolizumab in patients with platinum refractory germ cell tumors: a hoosier cancer research network study GU14-206. *Ann Oncol* 2017.
- [53] Raggi D, Giannatempo P, Mariani L, et al. Apache: An open label, randomized, phase 2 study of durvalumab (Durva), alone or in combination with tremelimumab (Treme), in patients (pts) with advanced germ cell tumors (GCT): Results at the end of first stage. *J Clin Oncol* 2018;36. 4547-4547.
- [54] Feldman DR, Lorch A, Kramar A, et al. Brain metastases in patients with germ cell tumors: prognostic factors and treatment options—an analysis from the global germ cell cancer group. *J Clin Oncol* 2016;34:345–51.
- [55] Coleman R, Body JJ, Aapro M, et al. Bone health in cancer patients: ESMO Clinical Practice Guidelines. *Ann Oncol* 2014;25(Suppl 3). iii124-137.
- [56] Hershman DL, Lacchetti C, Dworkin RH, et al. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol* 2014;32:1941–67.
- [57] Vehling S, Mehnert A, Hartmann M, et al. Anxiety and depression in long-term testicular germ cell tumor survivors. *Gen Hosp Psychiatry* 2016;38:21–5.
- [58] Smith TJ, Temin S, Alesi ER, et al. American Society of Clinical Oncology provisional clinical opinion: the integration of palliative care into standard oncology care. *J Clin Oncol* 2012;30:880–7.