

RELATIVE BRAUER GROUPS OF TORSORS OF PERIOD TWO

BRENDAN CREUTZ

ABSTRACT. We consider the problem of computing the relative Brauer group of a torsor of period two under an elliptic curve. We show how this problem can be reduced to finding a set of generators for the group of rational points on the elliptic curve. This extends work of Haile and Han to the case of torsors with unequal period and index. Our results also apply to torsors under higher dimensional abelian varieties. Several numerical examples are given.

1. INTRODUCTION

For a geometrically integral variety V over a field K , the relative Brauer group $\text{Br}(V/K)$ is the subgroup of the Brauer group $\text{Br}(K)$ consisting of classes which are split by the function field $\mathbf{k}(V)$ of V . The index of V is the greatest common divisor of the degrees of field extensions K'/K for which V possesses a K' -rational point. These two invariants are closely related. For example, the relative Brauer group can only be nontrivial when the index is greater than one. The simplest case in which this occurs is a conic without a rational point, for which the relative Brauer group is generated by the class of the conic in $\text{Br}(K)$.

For curves with nontrivial Jacobian, the relative Brauer group need not be cyclic. Here the simplest examples occur among genus one curves of index 2, which (assuming the characteristic of K is not 2) are curves admitting a model of the form $y^2 = g(x)$ with $g(x) \in K[x]$ a square free quartic polynomial. For such a curve V results of Shick [Shi94], Han [Han03] and Haile and Han [HH07] reduce the problem of determining the relative Brauer group to determining the group of K -rational points on the Jacobian J of V . Their approach can be explained by an old result of Lichtenbaum [Lic69]: corresponding to any torsor V there is a canonically defined class in $\mathcal{A}_V \in \text{Br}(J)$ such that the evaluation map induces a homomorphism $J(K) \rightarrow \text{Br}(V/K)$ which, as noted by Çiperiani and Krashen [ÇK12], will be surjective when the index of V and the order of \mathcal{A}_V (called the period of V) coincide.

In [HH07] Lichtenbaum's result is made explicit by constructing a representative for \mathcal{A}_V as an Azumaya algebra over the coordinate ring of J . Their work was motivated by earlier work of Haile [Hai84], who showed that the Clifford algebra of a binary cubic form $F(x, y)$ plays an analogous role for the genus one curve $z^3 = F(x, y)$. This has been further developed by Kuo [Kuo11] to construct the algebra \mathcal{A}_V corresponding to an arbitrary cubic curve, i.e., a torsor of period and index dividing 3. A method applicable to torsors of cyclic type (necessarily having equal period and index) is given in [ÇK12] and applied in [HHW12].

For a hyperelliptic curve X , Creutz and Viray [CV15] have recently given an explicit means of constructing unramified central simple $\mathbf{k}(X)$ -algebras representing 2-torsion elements in the Brauer group of X . Our primary purpose here is to explore applications of these algebras to the computation of relative Brauer groups. We show that each of these algebras corresponds to a torsor V under the Jacobian of X in such a way that the specializations of the algebra give elements of $\text{Br}(V/K)$. When X is a genus one curve, the corresponding

torsors are intersections of quadric hypersurfaces in \mathbb{P}^3 , and every torsor of index dividing 4 is given this way, including those with unequal period and index. The results also apply to higher genus hyperelliptic curves allowing for computation of relative Brauer groups of higher dimensional torsors under abelian varieties. In Section 6 we give several numerical examples illustrating the practicality of the method.

1.1. Statement of the results. Throughout the paper we consider a smooth and projective hyperelliptic curve X defined over a field K of characteristic different from 2. We fix an equation $y^2 = cf(x)$ defining X , where $c \in K^\times$ and $f(x) \in K[x]$ is a monic square free polynomial. We set $L = K[\theta]/f(\theta)$ and let $J = \text{Jac}(X)$ denote the Jacobian of X . For any $\ell \in L^\times$, let \mathcal{A}_ℓ'' be the $L(x)$ -algebra obtained by adjoining anticommuting square roots of $x - \theta$ and ℓ to $L(x)$. The corestriction $\mathcal{A}_\ell' := \text{Cor}_{L(x)/K(x)}(\mathcal{A}_\ell'')$ is a tensor product of quaternion algebras over $K(x)$ which can be explicitly computed using Rosset-Tate reciprocity (see [CV15, Proposition 3.1]). In [CV15] it is shown that, under suitable conditions on the norm of ℓ , the algebra \mathcal{A}_ℓ' pulls back to an unramified central simple $\mathbf{k}(X)$ -algebra $\mathcal{A}_\ell := \mathcal{A}_\ell' \otimes_{K(x)} \mathbf{k}(X)$.

1.2. Elliptic curves. When $\deg(f) = 3$, $X = J$ is an elliptic curve and every torsor of period 2 under X is given by a classical construction associating to an element $\ell \in L^\times$ of square norm the genus one curve $V_\ell \subset \mathbb{P}_K^3$ defined by

$$(1.1) \quad V_\ell : Q_\ell^{(1)}(u, v, w) + t^2 = Q_\ell^{(2)}(u, v, w) = 0,$$

where, for $0 \leq i \leq 2$, $Q_\ell^{(i)}(u, v, w) \in K[u, v, w]$ is the ternary quadratic form uniquely determined by requiring that $\sum \theta^i Q_\ell^{(i)} = \ell(u + \theta v + \theta^2 w)^2$. See [Cas91, Section 15]. In the appendix we prove that every torsor of period 2 arises in this way.

The following theorem, proven in Section 5.1, reduces computation of the relative Brauer group of V_ℓ to a computation of a basis for $X(K)/2X(K)$.

Theorem 1.1. *Let V be a genus one curve of period 2 and suppose $X : y^2 = f(x)$ is a Weierstrass model for $X = \text{Jac}(V)$ (i.e., $\deg(f) = 3$). Then*

- (1) *There exist $\ell \in L^\times$ such that $V \simeq V_\ell$.*
- (2) *Evaluation of \mathcal{A}_ℓ induces an exact sequence*

$$\frac{X(K)}{2X(K)} \xrightarrow{\mathcal{A}_\ell} \text{Br}(V/K) \longrightarrow \frac{P(V)}{I(V)} \longrightarrow 0,$$

where $P(V)$ and $I(V)$ are the period and index subgroups in Definition 2.4.

- (3) *$\text{Br}(V/K)$ is generated by $\{\mathcal{A}_\ell(P) : P \in X(K)\}$ and the class $[C_\ell] \in \text{Br}(K)$ corresponding to the conic $C_\ell \subset \mathbb{P}_K^2$ defined by $Q_\ell^{(2)}(u, v, w) = 0$.*
- (4) *$P(V) = I(V)$ if and only if $[C_\ell] \in \mathcal{A}_\ell(X(K))$.*

The connection between these torsors and those considered in [HH07] is as follows. Projecting away from the point $u = v = w = 0$ in \mathbb{P}_K^3 induces a degree 2 morphism $V_\ell \rightarrow C_\ell$, where $C_\ell \subset \mathbb{P}_K^2$ is the conic defined by $Q_\ell^{(2)}(u, v, w) = 0$. If C_ℓ has a K -rational point, then $C_\ell \simeq \mathbb{P}_K^1$ and so V_ℓ is a double cover of \mathbb{P}_K^1 with a model of the form $y^2 = \text{quartic}$. However, as first shown by Cassels [Cas63], it is possible that V_ℓ has no rational points over any quadratic extension of K , in which case $P(V_\ell) \neq I(V_\ell)$ and no degree 2 map to \mathbb{P}_K^1 exists. In fact, the torsors V_ℓ with equal period and index are not generic; they correspond to those $\ell \in L^\times$

whose classes in $L^\times/L^{\times 2}$ can be represented by a linear polynomial in $K[\theta] = L$ (This follows easily from (1.1); see also [CF09, Theorem 13]). Theorem 1.1 therefore recovers results of [HH07], and extends them to all torsors of period 2 under an elliptic curve.

1.3. Genus one curves. When $X : y^2 = cf(x)$ is defined by a quartic polynomial $cf(x)$, a construction analogous to (1.1) yields an intersection of quadric surfaces $V_\ell \subset \mathbb{P}^3$. Given $\ell \in L^\times$ such that $N_{L/K}(\ell)$ is a square times the leading coefficient of $cf(x)$, one defines

$$(1.2) \quad V_\ell : Q_\ell^{(2)}(t, u, v, w) = Q_\ell^{(3)}(t, u, v, w) = 0,$$

where, for $0 \leq i \leq 3$, $Q_\ell^{(i)}(t, u, v, w)$ is the quadratic form defined by

$$\sum_{0 \leq i \leq 3} \theta^i Q_\ell^{(i)}(t, u, v, w) = \ell(t + \theta u + \theta^2 v + \theta^3 w)^2.$$

We associate to V_ℓ the central simple $\mathbf{k}(X)$ -algebra \mathcal{A}_ℓ constructed above. In Section 5.2 we prove the following.

Theorem 1.2. *Let V be a genus one curve of period and index 4. Then there exists a quartic polynomial $cf(x) \in K[x]$ and $\ell \in L^\times$, where $L = K[\theta]/f(\theta)$ such that:*

- (1) $\text{Jac}(V) \simeq \text{Jac}(X)$ where $X : y^2 = cf(x)$.
- (2) $2V = X$ in the Weil-Châtelet group of the Jacobian.
- (3) $V \simeq V_\ell$.
- (4) Evaluation of \mathcal{A}_ℓ on degree zero K -rational divisors of X gives rise to an exact sequence

$$0 \rightarrow \text{Pic}^0(V) \xrightarrow{\iota} \text{Pic}^0(X) \xrightarrow{\mathcal{A}_\ell} \text{Br}(V/K) \xrightarrow{2} \text{Br}(X/K) \rightarrow 0,$$

where ι is induced by the inclusion $\text{Pic}^0(V) \subset \text{Pic}^0(\overline{V}) = \text{Pic}^0(\overline{X})$ and the last map is induced by multiplication by 2 in $\text{Br}(K)$.

The first three statements of this theorem are well known. If V has index dividing 4, then the complete linear system associated to any K -rational divisor of degree 4 gives a model for V as an intersection of quadrics in \mathbb{P}_K^3 . The quartic polynomial is $\det(M_1 x - M_2)$, where M_1 and M_2 are the symmetric matrices corresponding to these quadrics, and a formula for ℓ is given in [Fis08, Section 5].

It is worth noting that while Theorem 1.2 may not yield enough to completely determine $\text{Br}(V/K)$, Theorems 1.1 and 1.2 together allow one to determine whether a given K -rational point on the Jacobian is represented by a K -rational divisor of degree 0 on V . See Section 6.4 for an example.

1.4. Hyperelliptic curves. When the degree of $cf(x)$ is arbitrary, work of Cassels [Cas83], Schaefer [Sch95] and Poonen-Schaefer [PS97] recalled in Section 4 similarly allows one to associate a torsor V_ℓ of period 2 under J to any $\ell \in L^\times$ of square norm. Consequently, to any such torsor we may associate an algebra \mathcal{A}_ℓ . The following more general version of 1.1 is proven in Section 4.

Theorem 1.3. *Suppose V_ℓ is the torsor of period 2 under the Jacobian of a hyperelliptic curve X corresponding to an element $\ell \in L^\times$ of square norm. Suppose that $\text{Br}(X/K) = 0$*

and let \mathcal{A}_ℓ be the corresponding $\mathbf{k}(X)$ -algebra. Then evaluation of \mathcal{A}_ℓ on degree zero K -rational divisors of X induces an exact sequence

$$J(K)/2J(K) \xrightarrow{\mathcal{A}_\ell} \mathrm{Br}(V_\ell/K) \longrightarrow \frac{P(V_\ell)}{I(V_\ell)} \longrightarrow 0,$$

where $P(V_\ell)$ and $I(V_\ell)$ are the period and index subgroups defined in Definition 2.4.

As mentioned above, Lichtenbaum's result shows that the relative Brauer group of a torsor with equal period and index under an elliptic curve J can be obtained by specialization of a Brauer class on J . More generally, Çiperiani and Krashen [CK12, Theorem 3.5] have shown this to be true for torsors under any abelian variety. In Section 3.1 we use their result to deduce the following more general version.

Theorem 1.4. *Let Y/K be a smooth projective and geometrically integral variety over K with Picard variety $A = \mathrm{Pic}_Y^0$. There is a class $\mathcal{B}_Y \in \mathrm{Br}(A)$ canonically associated to Y (see Definition 3.1) such that evaluation induces an exact sequence*

$$0 \longrightarrow \frac{A(K)}{\mathrm{Pic}^0(Y)} \xrightarrow{\mathcal{B}_Y} \mathrm{Br}(Y/K) \longrightarrow \frac{P(Y)}{I(Y)} \longrightarrow 0,$$

where $P(Y)$ and $I(Y)$ are the period and index subgroups of Definition 2.4.

In the context of Theorem 1.3, we have a hyperelliptic curve X and a torsor V_ℓ . One may identify $J = \mathrm{Jac}(X) = \mathrm{Pic}_X^0 = \mathrm{Pic}_{V_\ell}^0$. When X has a K -rational point, there is an embedding $X \hookrightarrow J$ sending the point to the identity element of $A(K)$. As we shall see, the class \mathcal{B}_{V_ℓ} featuring in Theorem 1.4 pulls back via this embedding to give a class in $\mathrm{Br}(X)$ represented by the algebra \mathcal{A}_ℓ .

1.5. Notation. The following notation will be used throughout.

Let K be a field of characteristic not equal to 2, with a separable closure \overline{K} and absolute Galois group $G_K := \mathrm{Gal}(\overline{K}/K)$. For a G_K -module M (with the discrete topology) and an integer $i \geq 0$, $H^i(G_K, M)$ denotes the i th Galois cohomology group. For an étale K -algebra L and elements $a, b \in L^\times$ define

$$(a, b)_2 := \frac{L[i, j]}{\langle i^2 = a, j^2 = b, ij = -ij \rangle}.$$

Kummer theory gives an isomorphism $\kappa : L^\times/L^{\times 2} \simeq H^1(G_K, \mu_2(\overline{L}))$. The L -algebra $(a, b)_2$ represents the class of the cup product $\kappa(a) \cup \kappa(b)$ in $H^2(G_K, \mu_2(\overline{L})) = \mathrm{Br}(L)[2]$ (see Definition 2.1). We denote this class by $[a, b]_2$.

Suppose Y is a smooth, projective and geometrically integral variety over K . We use \overline{Y} to denote the base change of Y to \overline{K} . The function field of Y is denoted $\mathbf{k}(Y)$. Let $\mathrm{Pic}(Y)$ be the Picard group of Y . Then $\mathrm{Pic}(Y) = \mathrm{Div}(Y)/\mathrm{Princ}(Y)$, where $\mathrm{Div}(Y)$ (resp. $\mathrm{Princ}(Y)$) is the group of Weil divisors (resp. principal Weil divisors) of Y defined over K . If $D \in \mathrm{Div}(Y)$, then $[D]$ denotes its class in $\mathrm{Pic}(Y)$. Let Pic_Y denote the (reduced) Picard scheme of Y , and let $\mathrm{Pic}_Y^0 \subseteq \mathrm{Pic}_Y$ denote the connected component of the identity. There is a bijective map $\mathrm{Pic}(\overline{Y})^{G_K} \rightarrow \mathrm{Pic}_Y(K)$, but in general the map $\mathrm{Pic}(Y) \rightarrow \mathrm{Pic}(\overline{Y})^{G_K}$ will not be surjective. Let $\mathrm{Pic}^0(Y)$ be the subgroup of $\mathrm{Pic}(Y)$ mapping into $\mathrm{Pic}_Y^0(K)$. Then $\mathrm{NS}(Y) := \mathrm{Pic}(Y)/\mathrm{Pic}^0(Y)$ is the Néron-Severi group of Y . If $\lambda \in \mathrm{NS}(\overline{Y})^{G_K}$, let Pic_Y^λ denote the corresponding component of the Picard scheme and use $\mathrm{Pic}^\lambda(Y)$ and $\mathrm{Div}^\lambda(Y)$ to denote

the subsets of $\text{Pic}(Y)$ and $\text{Div}(Y)$ mapping into $\text{Pic}_Y^\lambda(K)$. We write Alb_Y for the Albanese scheme of Y parameterizing equivalence classes of zero-cycles on Y and, for $i \in \mathbb{Z}$, write Alb_Y^i for the degree i component of Alb_Y . Then Alb_Y^0 is an abelian variety, and its dual abelian variety is Pic_Y^0 . When Y is a curve, $\text{NS}(\overline{Y}) = \mathbb{Z}$, $\text{Pic}_Y^i = \text{Alb}_Y^i$ for all $i \in \mathbb{Z}$ and $\text{Jac}(Y) := \text{Pic}_Y^0 = \text{Alb}_Y^0$ is called the Jacobian of Y .

A **torsor** under an abelian variety A over K is a variety V , together with an algebraic group action of A on V such that $A(\overline{K})$ acts simply transitively on $V(\overline{K})$. For any torsor V under A , there is an \overline{K} -isomorphism of torsors $\overline{V} \simeq \overline{A}$, determined up to translation by an element in $A(\overline{K})$. Since the action of a translation on $\text{Pic}^0(\overline{A})$ is trivial, this determines an isomorphism of G_K -modules $\text{Pic}^0(\overline{V}) \simeq \text{Pic}^0(\overline{A})$ which is independent of the choice of translation. For any $i \in \mathbb{Z}$, Alb_Y^i is a torsor under Alb_Y^0 . There is a canonical map $Y \rightarrow \text{Alb}_Y^1$ induced by sending $y \in Y(\overline{K})$ to the class of the 0-cycle y . This gives a canonical isomorphism of G_K -modules, $\text{Pic}_Y^0(\overline{K}) \simeq \text{Pic}_{\text{Alb}_Y^1}^0(\overline{K})$.

The isomorphism classes of torsors under A are parametrized by the Weil-Châtelet group, $H^1(K, A(\overline{K}))$. We define the **period** of a torsor to be its order in the Weil-Châtelet group, and define the period of Y to be the period of Alb_Y^1 .

2. THE RELATIVE BRAUER GROUP OF A VARIETY

2.1. The Brauer group of a variety.

Definition 2.1. *Let Y be a smooth, projective and geometrically integral variety over a field K . The algebraic Brauer group of Y is*

$$\text{Br}(Y) := \ker(\text{div}_* : H^2(G_K, \mathbf{k}(\overline{Y})^\times) \rightarrow H^2(G_K, \text{Div}(\overline{Y})^\times))$$

where div_* is induced by the map sending a rational function on \overline{Y} to its divisor.

When $Y = \text{Spec}(K)$ we abbreviate $\text{Br}(\text{Spec}(K))$ to $\text{Br}(K)$. In this case div_* is trivial and so $\text{Br}(K) = H^2(G_K, \overline{K}^\times)$ is the usual Brauer group of the field K . In general, the inflation-restriction sequence identifies $H^2(G_K, \mathbf{k}(\overline{Y})^\times)$ with the relative Brauer group, $\text{Br}(\mathbf{k}(\overline{Y})/\mathbf{k}(Y))$. Thus, $\text{Br}(Y)$ can (and tacitly will) be viewed as the subgroup of $\text{Br}(\mathbf{k}(Y))$ consisting of classes that are unramified at all divisors and split by extension of scalars to \overline{K} . Although we will not need it, let us also mention the étale-cohomological definition of the algebraic Brauer group as

$$\text{Br}(Y) := \ker(H_{\text{ét}}^2(Y, \mathbb{G}_m) \rightarrow H_{\text{ét}}^2(\overline{Y}, \mathbb{G}_m)) .$$

It is shown in [Lic69, Appendix] when Y is a curve, and in [CTS77, Lemme 14] in general, that these definitions are equivalent.

2.2. The basic exact sequence. The Hochschild-Serre spectral sequence gives rise to a well known exact sequence,

$$(2.1) \quad 0 \rightarrow \text{Pic}(Y) \rightarrow \text{Pic}(\overline{Y})^{G_K} \xrightarrow{a_Y} \text{Br}(K) \rightarrow \text{Br}(Y) \xrightarrow{h_Y} H^1(G_K, \text{Pic}(\overline{Y})) \rightarrow H^3(G_K, \overline{K}^\times) .$$

Following [Lic69, Section 2] (in the case that Y is a curve), we will give a more direct derivation of (2.1) using Galois cohomology. The exact sequences,

$$(2.2) \quad 1 \rightarrow \mathbf{k}(\overline{Y})^\times / \overline{K}^\times \xrightarrow{\text{div}} \text{Div}(\overline{Y}) \rightarrow \text{Pic}(\overline{Y}) \rightarrow 0 ,$$

and

$$1 \rightarrow \overline{K}^\times \rightarrow \mathbf{k}(\overline{Y})^\times \rightarrow \mathbf{k}(\overline{Y})^\times / \overline{K}^\times \rightarrow 1,$$

induce a commutative diagram with exact rows and columns,

(2.3)

$$\begin{array}{ccccccc}
& & \text{Br}(Y) & & \text{H}^1(G_K, \text{Pic}(\overline{Y})) & & \\
& & \downarrow & & \downarrow \rho & & \\
\text{H}^1\left(G_K, \frac{\mathbf{k}(\overline{Y})^\times}{\overline{K}^\times}\right) & \xrightarrow{\delta} & \text{Br}(K) & \longrightarrow & \text{H}^2(G_K, \mathbf{k}(\overline{Y})^\times) & \xrightarrow{\epsilon} & \text{H}^1\left(G_K, \frac{\mathbf{k}(\overline{Y})^\times}{\overline{K}^\times}\right) \longrightarrow \text{H}^3(G_K, \overline{K}^\times) \\
& & \downarrow \text{div}_* & & \downarrow & & \\
& & \text{H}^2(G_K, \text{Div}(\overline{Y})) & \xlongequal{\quad} & \text{H}^2(G_K, \text{Div}(\overline{Y})) & &
\end{array}$$

The image of $\text{Br}(K)$ is contained in the kernel of div_* (a consequence of the fact that the divisor of a constant function is 0). Also, ρ is injective, since $\text{H}^1(G_K, \text{Div}(\overline{Y})) = 0$ (a consequence of Shapiro's Lemma). From the diagram we see that ϵ induces a unique map h_Y making (2.1) exact at all terms to the right of $\text{Br}(K)$. We define \mathbf{a}_Y to be the composition of δ and the connecting homomorphism arising from (2.2). The connecting homomorphism is surjective (for the same reason that ρ is injective) and δ is injective (by Hilbert's Theorem 90). This shows that (2.1) is exact at $\text{Br}(K)$. The map $\text{Pic}(Y) \rightarrow \text{Pic}(\overline{Y})^{G_K}$ is injective by Hilbert's Theorem 90 and its image is equal to $\text{image}(\text{Div}(\overline{Y})^{G_K} \rightarrow \text{Pic}(\overline{Y})^{G_K})$, which is equal to $\ker(\mathbf{a}_X)$ by definition. Therefore (2.1) is exact.

Lemma 2.2. *Suppose $[D] \in \text{Pic}(\overline{Y})^{G_K}$ is represented by a base point free divisor $D \in \text{Div}(\overline{Y})$. Then the complete linear system of effective divisors linearly equivalent to D forms a Brauer-Severi variety S over K , and $\mathbf{a}_Y([D])$ is the class of S in $\text{Br}(K)$.*

Proof. Since $[D]$ is fixed by Galois, for each $\sigma \in G_K$ there exists $f_\sigma \in \mathbf{k}(\overline{Y})^\times$ such that $\text{div}(f_\sigma) = \sigma(D) - (D)$. For any $\sigma, \tau \in G_K$, the divisor of $a_{\sigma, \tau} := \sigma(f_\tau) f_\sigma / f_{\sigma\tau}$ is 0, which implies $a_{\sigma, \tau} \in \overline{K}^\times$. The definition of \mathbf{a}_Y shows that $\mathbf{a}_Y([D])$ is represented by the 2-cocycle, $(\sigma, \tau) \mapsto a_{\sigma, \tau}$.

The set $\mathcal{L}(D) = \{f \in \mathbf{k}(\overline{Y})^\times : \text{div}(f) + D \geq 0\} \cup \{0\}$ is a \overline{K} -vector space. For each $\sigma \in G_K$, the map $\phi_\sigma : f \mapsto \sigma(f) f_\sigma$ is an automorphism of $\mathcal{L}(D)$. For $\sigma, \tau \in G_K$, these maps satisfy $a_{\sigma, \tau} \phi_{\sigma\tau} = \phi_\sigma \circ \phi_\tau$, and using this one easily verifies that the maps $\tilde{\phi}_\sigma : \mathbb{P}(\mathcal{L}(D)) \rightarrow \mathbb{P}(\mathcal{L}(D))$ induced by the ϕ_σ satisfy Weil's criteria for Galois descent (see [Wei56, Theorem 1]). We conclude that there is a K -variety S and a \overline{K} -isomorphism $\psi : S_{\overline{K}} \simeq \mathbb{P}(\mathcal{L}(D))$ such that $\tilde{\phi}_\sigma = \psi^\sigma \circ \psi^{-1}$, for every $\sigma \in G_K$. In particular, S is a Brauer-Severi variety, and its class in $\text{H}^1(G_K, \text{Aut}(\mathbb{P}(\mathcal{L}(D)))) \simeq \text{H}^1(G_K, \text{PGL}(\mathcal{L}(D)))$ is represented by the 1-cocycle $\sigma \mapsto \tilde{\phi}_\sigma$. The image of S in $\text{Br}(K) = \text{H}^2(G_K, \overline{K}^\times)$ is then given by the 2-cocycle $(\sigma, \tau) \mapsto \sigma(\phi_\tau) \circ \phi_\sigma \circ \phi_{\sigma\tau}^{-1} = a_{\sigma, \tau}$. As shown above, this also represents $\mathbf{a}_Y([D])$. \square

2.3. Specialization of Brauer classes. There is a natural *evaluation pairing*,

$$\text{ev} : Y(K) \times \text{Br}(Y) \longrightarrow \text{Br}(K).$$

For $P \in Y(K)$, define $\text{Br}(Y, P) := \ker(\text{ev}(P, \cdot) : \text{Br}(Y) \rightarrow \text{Br}(K))$.

Lemma 2.3. *For any K -rational point $P \in Y(K)$ the map h_Y of (2.1) restricts to an isomorphism $\text{Br}(Y, P) \simeq H^1(G_K, \text{Pic}(\overline{Y}))$.*

Proof. The evaluation map gives a retraction of the exact sequence

$$\text{Br}(K) \rightarrow \text{Br}(Y) \xrightarrow{h_Y} H^1(G_K, \text{Pic}(\overline{Y})).$$

□

2.4. Period and index subgroups. Following [CK12] we make the following definitions

Definition 2.4. *Let Y and \mathfrak{a}_Y be as in (2.1).*

- (i) $\text{Br}^0(Y/K)$ is the image of \mathfrak{a}_Y restricted to $\text{Pic}^0(\overline{Y})^{G_K}$.
- (ii) $P(Y)$ is the image of $\text{Pic}(\overline{Y})^{G_K}$ in $\text{NS}(\overline{Y})$.
- (iii) $I(Y)$ is the image of $\text{Pic}(Y)$ in $\text{NS}(\overline{Y})$.

Remark 2.5. *If Y is a curve of period p and index i , then $\text{NS}(\overline{Y}) \simeq \mathbb{Z}$ and under this isomorphism $P(Y) = p\mathbb{Z}$ and $I(Y) = i\mathbb{Z}$. The integer p (resp. i) is the least positive degree of a K -rational divisor class (resp. K -rational divisor) on Y .*

By definition of $\text{Br}^0(Y/K)$ we have a commutative diagram with exact rows

$$(2.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}^0(Y) & \longrightarrow & \text{Pic}^0(\overline{Y})^{G_K} & \xrightarrow{\mathfrak{a}_Y} & \text{Br}^0(Y/K) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Pic}(Y) & \longrightarrow & \text{Pic}(\overline{Y})^{G_K} & \xrightarrow{\mathfrak{a}_Y} & \text{Br}(Y/K) \longrightarrow 0 \end{array}$$

Applying the snake lemma immediately yields an exact sequence

$$(2.5) \quad 0 \rightarrow \text{Br}^0(Y/K) \rightarrow \text{Br}(Y/K) \rightarrow \frac{P(Y)}{I(Y)} \rightarrow 0.$$

If $\phi : Y \rightarrow Z$ is a morphism of smooth projective and geometrically integral varieties over K . Then the identity map on $\text{Br}(K)$ induces an injection $\text{Br}(Z/K) \hookrightarrow \text{Br}(Y/K)$ and ϕ induces a morphism of short exact sequences,

$$(2.6) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}^0(Z) & \longrightarrow & \text{Pic}^0(\overline{Z})^{G_K} & \xrightarrow{\mathfrak{a}_Z} & \text{Br}^0(Z/K) \longrightarrow 0 \\ & & \downarrow \phi^* & & \downarrow \phi^* & & \downarrow \\ 0 & \longrightarrow & \text{Pic}^0(Y) & \longrightarrow & \text{Pic}^0(\overline{Y})^{G_K} & \xrightarrow{\mathfrak{a}_Y} & \text{Br}^0(Y/K) \longrightarrow 0. \end{array}$$

This has the following consequences.

Lemma 2.6. *Let $\phi : Y \rightarrow Z$ be as above.*

- (1) *If ϕ is the canonical map $Y \rightarrow \text{Alb}_Y^1$, then (2.6) is an isomorphism of exact sequences.*
- (2) *If $D \in \text{Div}(\overline{Z})$ represents a class in $\text{Pic}^0(\overline{Z})^{G_K}$ and ϕ^*D is principal, then D is linearly equivalent to a K -rational divisor.*

Proof. Both statements follow by applying the snake lemma to (2.6). For the first statement, one also uses the fact that $\phi^* : \text{Pic}^0(\text{Alb}_Y^1)^{G_K} \rightarrow \text{Pic}^0(\overline{Y})^{G_K}$ is an isomorphism. □

3. RELATIVE BRAUER GROUPS VIA SPECIALIZATION

3.1. Brauer classes on the Picard variety. Suppose A is an abelian variety with dual $B = \text{Pic}_A^0$. There is a canonical choice of rational point on A , namely the identity 0_A . So by Lemma 2.3, h_A restricts to an isomorphism $\text{Br}(A, 0_A) \simeq H^1(G_K, \text{Pic}(\overline{A}))$. Therefore, the inclusion $\text{Pic}^0(\overline{A}) \subset \text{Pic}(\overline{A})$ induces a map

$$(3.1) \quad H^1(G_K, B(\overline{K})) = H^1(G_K, \text{Pic}^0(\overline{A})) \rightarrow H^1(G_K, \text{Pic}(\overline{A})) \simeq \text{Br}(A, 0_A)$$

Definition 3.1. For a torsor V under an abelian variety B with dual abelian variety A , let $\mathcal{B}_V \in \text{Br}(A, 0_A)$ be the image of the class of V under the map in (3.1). For a smooth projective and geometrically integral variety Y with Picard variety $A = \text{Pic}_Y^0$, define $\mathcal{B}_Y = \mathcal{B}_{\text{Alb}_Y^1} \in \text{Br}(A, 0_A)$ where Alb_Y^1 is the torsor under $B = \text{Alb}_Y^0$ parameterizing classes of 0-cycles of degree 1 on Y .

Remark 3.2. When Y is a torsor under B these two definitions of \mathcal{B}_Y coincide, since in this case $B = \text{Alb}_Y^0$ and $Y = \text{Alb}_Y^1$.

Proof of Theorem 1.4. Let Y be a smooth projective and geometrically integral variety, let $\mathcal{B}_Y \in \text{Br}(A, 0_A)$ as in Definition 3.1 and let $V = \text{Alb}_Y^1$. Evaluation of \mathcal{B}_Y gives a homomorphism $A(K) \rightarrow \text{Br}(K)$, which by [CK12, Theorem 3.5] coincides with $\mathfrak{a}_V : \text{Pic}^0(\overline{V})^{G_K} = A(K) \rightarrow \text{Br}(K)$. Hence evaluation of \mathcal{B}_Y induces a short exact sequence,

$$(3.2) \quad 0 \rightarrow \text{Pic}^0(V) \longrightarrow A(K) \xrightarrow{\mathcal{B}_Y} \text{Br}^0(V/K) \rightarrow 0.$$

Using Lemma 2.6 we may replace V by Y in (3.2) without affecting the exactness. The theorem then follows from the exactness of (2.5). \square

3.2. Brauer classes on a curve.

Definition 3.3. Let X be a smooth projective and geometrically integral curve. A torsor V under the Jacobian of X is *Brauer-like* if its image in $H^1(G_K, \text{Pic}(\overline{X}))$ lies in the image of the map h_X in (2.1). For a Brauer-like torsor V , let $\mathcal{A}_V \in \text{Br}(X)$ be any class such that $h_X(\mathcal{A}_V)$ is the image of V in $H^1(G_K, \text{Pic}(\overline{X}))$.

We note that \mathcal{A}_V is only defined up to the image of $\text{Br}(K)$ in $\text{Br}(X)$. The relationship between \mathcal{A}_V and the class \mathcal{B}_V of Definition 3.1 is given by the following.

Lemma 3.4. Suppose X has a rational point $P \in X(K)$ and let V be a torsor under the Jacobian J of X . Then

- (1) V is Brauer-like,
- (2) there is a unique choice for \mathcal{A}_V which lies in $\text{Br}(X, P)$ and
- (3) $\mathcal{A}_V = i^*(\mathcal{B}_V)$, where $i^* : X \rightarrow J$ is the map induced by sending $Q \in X(\overline{K})$ to $[Q - P]$.

Proof. By Lemma 2.3 h_X gives an isomorphism $\text{Br}(X, P) \simeq H^1(G_K, \text{Pic}(\overline{X}))$. So it is clear that V is Brauer-like and that the choice for \mathcal{A}_V in $\text{Br}(X, P)$ is unique. The cokernel of the map $H^1(G_K, \text{Pic}^0(\overline{X})) \rightarrow H^1(G_K, \text{Pic}(\overline{X}))$ induced by the inclusion is $H^1(G_K, \text{NS}(\overline{X})) =$

$H^1(G_K, \mathbb{Z}) = 0$. So the map i^* induces a commutative diagram

$$\begin{array}{ccccc} \mathrm{Br}(X, P) & \xrightarrow{h_X} & H^1(G_K, \mathrm{Pic}(\overline{X})) & \xlongequal{\quad} & H^1(G_K, \mathrm{Pic}^0(\overline{X})) \\ \uparrow i^* & & & & \uparrow i^* \\ \mathrm{Br}(J, 0_J) & \xrightarrow{h_J} & H^1(G_K, \mathrm{Pic}(\overline{J})) & \longleftarrow & H^1(G_K, \mathrm{Pic}^0(\overline{J})) \end{array}$$

Both h_X and h_J are isomorphisms (again by Lemma 2.3). The vertical map on the right of the diagram is also an isomorphism, since $i^* : \mathrm{Pic}^0(\overline{J}) \rightarrow \mathrm{Pic}^0(\overline{X})$ is an isomorphism. The lemma then follows from a diagram chase. \square

3.3. Lichtenbaum's pairings. For a proper, smooth, geometrically integral curve X over K , Lichtenbaum defined in [Lic69] three pairings and showed that they are compatible in the sense that there is a commutative diagram:

$$(3.3) \quad \begin{array}{ccccc} \langle \cdot, \cdot \rangle_1 : & \mathrm{Pic}^0(\overline{X})^{G_K} & \times & H^1(G_K, \mathrm{Pic}^0(\overline{X})) & \rightarrow & \mathrm{Br}(K) \\ & \uparrow & & \downarrow & & \parallel \\ \langle \cdot, \cdot \rangle_2 : & \mathrm{Pic}^0(X) & \times & H^1(G_K, \mathrm{Pic}(\overline{X})) & \rightarrow & \mathrm{Br}(K) \\ & \downarrow & & \uparrow & & \parallel \\ \langle \cdot, \cdot \rangle_3 : & \mathrm{Pic}(X) & \times & \mathrm{Br}(X) & \rightarrow & \mathrm{Br}(K) \end{array}$$

The pairing $\langle \cdot, \cdot \rangle_1$ was first defined by Tate; its definition may be found in [CK12, Section 3]. The pairing $\langle \cdot, \cdot \rangle_2$ is induced by $\langle \cdot, \cdot \rangle_1$; that it is well defined is verified in [Lic69, Section 4]. The third pairing is induced by the evaluation map. Specifically, for $D \in \mathrm{Pic}(X)$ and $\mathcal{A} \in \mathrm{Br}(X)$ one has $\langle D, \mathcal{A} \rangle_3 = \bigotimes_P \mathcal{A}(P)^{\otimes n_P}$, where $\sum_P n_P P$ is an integral linear sum of closed points of X representing the divisor class D (see [Lic69, Section 4]). The vertical maps of the left column of (3.3) are the obvious ones. The upper-middle map is induced by the inclusion $\mathrm{Pic}^0(X) \rightarrow \mathrm{Pic}^0(\overline{X})$, and the lower-middle map is h_X of (2.1).

The following lemma relates the map \mathfrak{a}_V with Tate's pairing. It was first proven by Lichtenbaum in the case of elliptic curves; for general abelian varieties see [CK12, Theorem 3.5].

Lemma 3.5. *For $V \in H^1(G_K, \mathrm{Pic}^0(\overline{X}))$ and $P \in \mathrm{Pic}^0(\overline{V})^{G_K} = \mathrm{Pic}^0(\overline{X})^{G_K}$ one has $\langle P, V \rangle_1 = \mathfrak{a}_V(P)$.*

Proposition 3.6. *Let X be a curve and let V be a Brauer-like torsor under the Jacobian of X . Evaluation of \mathcal{A}_V induces an exact sequence*

$$0 \rightarrow (\mathrm{Pic}^0(V) \cap \mathrm{Pic}^0(X)) \rightarrow \mathrm{Pic}^0(X) \xrightarrow{\mathcal{A}_V} \mathrm{Br}^0(V/K) \longrightarrow \frac{\mathrm{Br}^0(X/K)}{\mathfrak{a}_X(\mathrm{Pic}^0(V))} \rightarrow 0,$$

where $\mathrm{Pic}^0(V)$ is identified with a subgroup of $\mathrm{Pic}^0(\overline{X})^{G_K}$ via the canonical identification of $\mathrm{Pic}^0(\overline{X})$ and $\mathrm{Pic}^0(\overline{V})$.

The proposition is proved below. First we state a corollary.

Corollary 3.7. *If X is a curve such that $\mathrm{Br}^0(X/K) = 0$ and V is a Brauer-like torsor under its Jacobian, then evaluation of \mathcal{A}_V induces an exact sequence,*

$$J(K) \xrightarrow{\mathcal{A}_V} \mathrm{Br}(V/K) \rightarrow \frac{P(V)}{I(V)} \rightarrow 0,$$

where $J = \text{Jac}(X)$ is the Jacobian of X .

Proof of Corollary 3.7. If $\text{Br}^0(X/K) = 0$, then $\text{Pic}^0(X) = \text{Pic}^0(\overline{X})^{G_K} = J(K)$. By Proposition 3.6, evaluation of \mathcal{A}_V gives a surjective map $J(K) \rightarrow \text{Br}^0(V/K)$. The exactness of (2.5) then implies that the sequence in the corollary is exact. \square

Proof of Proposition 3.6. Let $P \in \text{Pic}^0(X) \subset \text{Pic}^0(\overline{X})^{G_K}$. First note that $\langle P, \mathcal{A}_V \rangle_3$ does not depend on the choice for \mathcal{A}_V , since P is a divisor class of degree 0. By the commutativity in (3.3) and Lemma 3.5 we have $\langle P, \mathcal{A}_V \rangle_3 = \langle P, V \rangle_1 = \mathfrak{a}_V(P)$. Hence evaluation of \mathcal{A}_V induces a commutative and exact diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}^0(X) & \longrightarrow & \text{Pic}^0(\overline{X})^{G_K} & \xrightarrow{\mathfrak{a}_X} & \text{Br}^0(X/K) \longrightarrow 0 \\ & & \downarrow \langle \cdot, \mathcal{A}_V \rangle_3 & & \downarrow \mathfrak{a}_V & & \downarrow \\ 0 & \longrightarrow & \text{Br}^0(V/K) & \xlongequal{\quad} & \text{Br}^0(V/K) & \longrightarrow & 0. \end{array}$$

Every element $a \in \text{Br}^0(V/K)$ may be written as $a = \mathfrak{a}_V([D])$ for some $[D] \in \text{Pic}^0(\overline{X})^{G_K}$, with $[D]$ determined uniquely modulo $\text{Pic}^0(V)$. Therefore the map

$$r : \text{Br}^0(V/K) \rightarrow \text{Br}^0(X/K)/\mathfrak{a}_X(\text{Pic}^0(V))$$

sending a to $\mathfrak{a}_X([D])$ is well defined. Exactness in the statement in the proposition now follows by applying the snake lemma to the diagram above. \square

4. HYPERELLIPTIC CURVES

Consider a hyperelliptic curve $X : y^2 = cf(x)$, where $f(x) \in K[x]$ is a square free monic polynomial over a field of characteristic not equal to 2. The x -coordinate defines a degree 2 map $\pi : X \rightarrow \mathbb{P}_K^1$, embedding $K(x)$ into $\mathbf{k}(X)$ as an index 2 subfield. We set $L = K[\theta]/f(\theta)$, and for $\ell \in L^\times$ set

$$\mathcal{A}_\ell := (\text{Cor}_{L(x)/K(x)}(\ell, x - \theta)_2) \otimes_{K(x)} \mathbf{k}(X).$$

This is a central simple algebra over $\mathbf{k}(X)$ which can be written explicitly as a tensor product of quaternion algebras (see [CV15, Proposition 2.4]). In this section we show that for suitable torsors V the class $\mathcal{A}_V \in \text{Br}(X)$ can be represented by an algebra of the form \mathcal{A}_ℓ .

We say that X is **odd** if $\deg(f)$ is odd. Otherwise we say that X is **even**. If X is odd, we can perform a change of coordinates to arrange that $c = 1$. Let $\Omega \subseteq X$ be the set of ramification points of π , and let $L_\circ = \text{Map}_K(\Omega, \overline{K})$ denote the étale K -algebra corresponding to Ω . When X is even we may identify L and L_\circ . When X is odd, L can be identified with the subalgebra of L_\circ consisting of elements in $\text{Map}_K(\Omega, \overline{K})$ that take the value 1 at the ramification point above $\infty \in \mathbb{P}^1(K)$. In the odd case this gives an isomorphism $L_\circ \cong L \times K$.

Set

$$\mathfrak{L} = \frac{L_\circ^\times}{K^\times L_\circ^{\times 2}}$$

where $K^\times \subset L_\circ^\times = \text{Map}_K(\Omega, \overline{K}^\times)$ here denotes the multiplicative group of nonzero constant maps. For $a \in K^\times$ and $\ell \in L^\times$, we use \bar{a} and $\bar{\ell}$ to denote the corresponding classes in $K^\times/K^{\times 2}$ and \mathfrak{L} , and set

$$\mathfrak{L}_a = \{ \bar{\ell} \in \mathfrak{L} : \text{Norm}_{L/K}(\bar{\ell}) \in \langle \bar{a} \rangle \},$$

where $\text{Norm}_{L/K}$ denotes the map $\mathfrak{L} \rightarrow K^\times/K^{\times 2}$ induced by the norm on L . Note that when X is odd we have an isomorphism $\mathfrak{L} \simeq L^\times/L^{\times 2}$ under which $\text{Norm}_{L/K}$ coincides with the map induced by the norm on L .

4.1. Odd hyperelliptic curves. Suppose that X is odd (i.e., that $f(x)$ has odd degree). Schaefer [Sch95], building on work of Cassels [Cas83] defined an isomorphism of G_K -modules between $J[2]$ and $\ker(\text{Norm}_{L/K} : \mu_2(\overline{L}) \rightarrow \mu_2(\overline{K}))$. From this one obtains an isomorphism $e : \mathfrak{L}_1 \simeq H^1(G_K, J[2])$. The Kummer sequence associated to J gives a surjective map $H^1(G_K, J[2]) \rightarrow H^1(G_K, J(\overline{K}))[2] = H^1(G_K, \text{Pic}^0(\overline{X}))[2]$. Furthermore, the K -rational point $0_X \in X(K)$ lying above $\infty \in \mathbb{P}_K^1(K)$ gives rise to isomorphisms,

$$\text{Br}(X, 0_X) \xrightarrow{h_X} H^1(G_K, \text{Pic}(\overline{X})) \simeq H^1(G_K, \text{Pic}^0(\overline{X})).$$

Hence there exists a homomorphism γ , uniquely determined by the requirement that the following diagram commute.

$$(4.1) \quad \begin{array}{ccc} H^1(G_K, \text{Pic}^0(\overline{X})[2]) & \twoheadrightarrow & H^1(G_K, \text{Pic}^0(\overline{X}))[2] \\ e \uparrow \wr & & h_X \uparrow \wr \\ \mathfrak{L}_1 & \xrightarrow{\gamma} & \text{Br}(X, 0_X)[2] \end{array}$$

Given $\ell \in L^\times$ of square norm, let V_ℓ be a torsor under J representing the image of ℓ in the top right corner of the diagram. When $f(x)$ has degree 3 (so that $X = J$ is an elliptic curve in Weierstrass form) the class of this torsor coincides with the torsor constructed in (1.1) (see Theorem A.1).

In [CV15] it is shown that γ is induced by the map $\ell \mapsto \mathcal{A}_\ell$. The commutativity of the diagram therefore implies that \mathcal{A}_ℓ represents the class \mathcal{A}_{V_ℓ} of Definition 3.3. Applying Corollary 3.7 we obtain the following.

Theorem 4.1. *Suppose X is an odd hyperelliptic curve and V is a torsor of period 2 under J . Then $V \simeq V_\ell$ for some $\ell \in L^\times$ of square norm. The algebra \mathcal{A}_ℓ represents the class $\mathcal{A}_V \in \text{Br}(X, 0_X)$ and is split if and only if $V(K) \neq \emptyset$. Moreover, evaluation of \mathcal{A}_ℓ on K -rational divisors of degree 0 induces an exact sequence.*

$$\frac{J(K)}{2J(K)} \xrightarrow{\mathcal{A}_\ell} \text{Br}(V/K) \rightarrow \frac{P(V)}{I(V)} \rightarrow 0.$$

4.2. Even hyperelliptic curves. The case of even hyperelliptic curves is somewhat more involved. Suppose $\ell \in L^\times$ represents a class in \mathfrak{L}_c and let $n \in \{0, 1\}$ be such that $c^n N_{L/K}(\ell) \in K^{\times 2}$. Then ℓ , together with a choice of square root for $c^n N_{L/K}(\ell)$, can be used to construct a torsor V_ℓ under J . For $n = 0$ this follows from work of Poonen and Schaefer [PS97]. When X has genus 2 Flynn, Testa and Van Luijk [FTvL12] give an explicit construction of V_ℓ as an intersection of 72 quadrics in \mathbb{P}^{15} . For $n = 1$ this is developed in [Cre13]. See also [BGW, Theorems 23 and 24] for an alternative description of this torsor. When X has genus 1, the proof of [Cre13, Proposition 5.4] shows that V_ℓ is given by the construction in (1.2). In this case the V_ℓ are the torsors produced by a 4-descent as described in [MSS96].

The class of V_ℓ in $H^1(G_K, J(\overline{K}))$ may depend up to inverse on the choice of square root for $c^n N_{L/K}(\ell)$, but the image of this class in $H^1(G_K, J(\overline{K}))/\langle \text{Pic}_X^1 \rangle$ does not. These torsors also

have the property that $2V_\ell = n \text{Pic}_X^1$ in $H^1(G_K, J(\overline{K}))$. This gives rise to a homomorphism

$$(4.2) \quad \mathfrak{L}_c \longrightarrow \left(\frac{H^1(G_K, J(\overline{K}))}{\langle \text{Pic}_X^1 \rangle} \right) [2] \simeq H^1(G_K, \text{Pic}(\overline{X})) [2].$$

It follows from [CV15] that this homomorphism coincides with the composition

$$\mathfrak{L}_c \xrightarrow{\gamma} \text{Br}(X) / \text{Br}_0(X) \xrightarrow{h_X} H^1(G_K, \text{Pic}(\overline{X})),$$

where γ is induced by the map $\ell \mapsto \mathcal{A}_\ell$ and $\text{Br}_0(X)$ denotes the image of $\text{Br}(K)$ in $\text{Br}(X)$. This implies that V_ℓ is Brauer-like and that the algebra \mathcal{A}_ℓ is a representative for \mathcal{A}_{V_ℓ} . Combining this with Corollary 3.7 we obtain the following.

Theorem 4.2. *Suppose $\text{Br}^0(X/K) = 0$ and that the norm of $\ell \in L^\times$ lies in $c^n K^{\times 2}$. Let V_ℓ be a torsor under J corresponding to a choice of square root for $c^n N_{L/K}(\ell)$, and let \mathcal{A}_ℓ be the $\mathbf{k}(X)$ -algebra defined above. Then evaluation of \mathcal{A}_ℓ on K -rational degree 0 divisors induces an exact sequence*

$$\frac{J(K)}{2J(K)} \xrightarrow{\mathcal{A}_\ell} \text{Br}(V_\ell/K) \longrightarrow \frac{P(V_\ell)}{I(V_\ell)} \longrightarrow 0.$$

It is worth mentioning that the homomorphism in (4.2) need not be surjective. In general there exist Brauer-like torsors of period 2 under J which are not of the form V_ℓ . For example, see [CV15, Proposition 5.1 and Remark 5.4]. We also note that the hypothesis $\text{Br}^0(X/K) = 0$ is satisfied quite frequently, as suggested by the following lemma.

Lemma 4.3. *Suppose X is a hyperelliptic curve of even genus defined over a local or global field. Then $\text{Br}^0(X/K) = 0$.*

Proof. The hypothesis implies that $\text{Pic}^0(X) \rightarrow \text{Pic}^0(\overline{X})^{G_K}$ is an isomorphism; see [PS97, Propositions 3.3 and 3.4]. \square

5. GENUS ONE CURVES

We now specialize to the case that X is a genus 1 hyperelliptic curve.

5.1. Elliptic curves. If X is odd, then it is an elliptic curve in Weierstrass form. In this section we prove Theorem 1.1 which allows us to compute the relative Brauer group of any torsor under $X = \text{Jac}(X)$ of period 2 (assuming we are able to compute the group $X(K)$ of rational points on X). Recall from Section 1.2 that for $\ell \in L^\times$ of square norm, the corresponding torsor under $X = \text{Jac}(X)$ is the curve $V_\ell \subset \mathbb{P}_K^3$ defined by

$$(5.1) \quad V_\ell : Q_\ell^{(1)}(u, v, w) + t^2 = Q_\ell^{(2)}(u, v, w) = 0,$$

where, for $0 \leq i \leq 2$, $Q_\ell^{(i)}(u, v, w) \in K[u, v, w]$ are the ternary quadratic forms uniquely determined by requiring that $\sum \theta^i Q_\ell^{(i)} = \ell(u + \theta v + \theta^2 w)^2$. Let $C_\ell \subset \mathbb{P}_K^2$ be the conic defined by $Q_\ell^{(2)}(u, v, w) = 0$.

Lemma 5.1. *Let $P_0 = (u_0 : v_0 : w_0) \in C_\ell(\overline{K})$ be any point, and let*

$$D = \left(u_0 : v_0 : w_0 : \sqrt{-Q_\ell^{(1)}(u_0, v_0, w_0)} \right) + \left(u_0 : v_0 : w_0 : -\sqrt{-Q_\ell^{(1)}(u_0, v_0, w_0)} \right) \in \text{Div}(\overline{V}_\ell).$$

Then

- (1) The class $[D]$ of D in $\text{Pic}(\overline{V}_\ell)$ is G_K -invariant, and
- (2) $\mathfrak{a}_{V_\ell}([D])$ is represented by the class of the conic C_ℓ in $\text{Br}(K)$

Proof. Let $\pi : V_\ell \rightarrow C_\ell$ be the degree 2 map obtained by projecting away from the point $(0 : 0 : 0 : 1) \in \mathbb{P}_K^3$. Then $D = \pi^*P_0$ and for $\sigma \in G_K$ we have $\sigma(D) = \pi^*(\sigma(P_0))$ which shows that D and $\sigma(D)$ are linearly equivalent. This proves the first statement. The second follows from Lemma 2.2. \square

We are now in a position to prove Theorem 1.1.

Proof of Theorem 1.1. If $V_\ell(K) \neq \emptyset$ the statement holds trivially. Hence we may assume that V_ℓ has no K -rational point. This implies (using Riemann-Roch) that V_ℓ has no K -rational divisor class of degree 1. It follows that $P(X)$ is generated by the image of any K -rational divisor class of degree 2. By Lemma 5.1, the class of C_ℓ is equal to $\mathfrak{a}_{V_\ell}([D])$, where $[D]$ is a K -rational divisor class of degree 2 on V_ℓ . So the class of C_ℓ in $\text{Br}(K)$ must generate $\text{Br}(V_\ell/K)/\text{Br}^0(V_\ell/K)$. Since, by Theorem 1.3, evaluation of \mathcal{A}_ℓ induces a surjective map $X(K) \rightarrow \text{Br}^0(V_\ell/K)$, we have shown that $\text{Br}(V_\ell/K)$ is generated by the set $\{\mathcal{A}_\ell(P) : P \in X(K)\} \cup \{[C_\ell]\}$.

To prove the second statement of the theorem, note that when $V_\ell(K) = \emptyset$ we have $P(V_\ell) = I(V_\ell)$ if and only if $\text{Pic}^2(V_\ell) \neq \emptyset$, which is the case if and only if there is some $M \in \text{Pic}^2(\overline{V}_\ell)^{G_K}$ such that $\mathfrak{a}_{V_\ell}(M) = 0$. Since $\text{Pic}^2(\overline{V}_\ell)^{G_K} = [D] + \text{Pic}^0(\overline{V}_\ell)^{G_K}$ and \mathfrak{a}_{V_ℓ} is a homomorphism, we conclude that $P(V_\ell) = I(V_\ell)$ if and only if $[C_\ell]$ lies in $\{\mathcal{A}_\ell(P) : P \in X(K)\}$. \square

5.2. Curves of index 4. Now suppose X is even, that is $X : y^2 = cf(x)$ where $f(x)$ is a square free monic quartic polynomial. As described in Section 4.2 an element $\ell \in L^\times$ such that $cN_{L/K}(\ell) \in K^{\times 2}$ gives rise to a torsor V_ℓ under J such that $2V_\ell = X$ in the Weil-Châtelet group of the Jacobian. This may be given explicitly as an intersection of quadric surfaces in \mathbb{P}^3 . Namely, $V_\ell : Q_\ell^{(2)}(t, u, v, w) = Q_\ell^{(3)}(t, u, v, w) = 0$, where for $0 \leq i \leq 3$, $Q_\ell^{(i)}(t, u, v, w)$ are the quadratic forms defined by

$$\sum_{0 \leq i \leq 3} \theta^i Q_\ell^{(i)}(t, u, v, w) = \ell(t + \theta u + \theta^2 v + \theta^3 w)^2.$$

This defines a genus one curve of index dividing 4.

Conversely, given a genus one curve V of index 4, the complete linear system corresponding to any K -rational divisor of degree 4 defines an embedding of V in \mathbb{P}^3 as an intersection of quadric surfaces. If M_1 and M_2 denote the symmetric matrices corresponding to these quadrics, then $g(x) := \det(M_1 x + M_2)$ is a quartic polynomial, V is a torsor under the Jacobian of the hyperelliptic curve $X : y^2 = g(x)$, and $2V = X$ in the Weil-Châtelet group of the Jacobian (see [MSS96]). Moreover, there exists some $\ell \in L^\times$ of norm a square times the leading coefficient of $g(x)$ such that $V \simeq V_\ell$ (an explicit formula for ℓ is given in [Fis08, Section 5]).

Proof of Theorem 1.2. Statements (1), (2) and (3) in the theorem follow from the discussion above. It remains to prove (4).

By assumption V has period 4, so X has period 2. By construction, the index of X divides 2, so it must in fact be 2. Then both V and X have equal period and index, so $\text{Br}(V/K) = \text{Br}^0(V/K)$ and $\text{Br}(X/K) = \text{Br}^0(X/K)$.

Proposition 3.6 gives an exact sequence

$$(5.2) \quad 0 \rightarrow \text{Pic}^0(V) \cap \text{Pic}^0(X) \rightarrow \text{Pic}^0(X) \xrightarrow{\mathcal{A}_\ell} \text{Br}^0(V_\ell/K) \xrightarrow{r} \frac{\text{Br}^0(X/K)}{\mathfrak{a}_X(\text{Pic}^0(V))} \rightarrow 0,$$

where the map r is induced by sending $\mathfrak{a}_V([D]) \in \text{Br}^0(V/K)$ to the class of $\mathfrak{a}_X([D]) \in \text{Br}^0(X/K)$. Let J denote the Jacobian of V and X . Since the pairing $\langle \cdot, \cdot \rangle_1$ is bilinear, Lemma 3.5 implies that $\mathfrak{a}_X(P) = 2\mathfrak{a}_V(P)$, for any $P \in J(K) \cong \text{Pic}^0(\overline{V})^{G_K} \cong \text{Pic}^0(\overline{X})^{G_K}$. This shows that $\ker(\mathfrak{a}_V) \subset \ker(\mathfrak{a}_X)$. Hence we have $\text{Pic}^0(V) = \ker(\mathfrak{a}_V) \cap J(K) \subset \ker(\mathfrak{a}_X) \cap J(K) = \text{Pic}^0(X)$ and $\mathfrak{a}_X(\text{Pic}^0(V)) = 2\mathfrak{a}_V(\text{Pic}^0(V)) = 0$. The fact that $\mathfrak{a}_X(P) = 2\mathfrak{a}_V(P) = \mathfrak{a}_V(2P)$ also shows that r is induced by multiplication by 2. From this we see that (5.2) can be rewritten as

$$(5.3) \quad 0 \rightarrow \text{Pic}^0(V) \rightarrow \text{Pic}^0(X) \xrightarrow{\mathcal{A}_\ell} \text{Br}(V/K) \xrightarrow{2} \text{Br}(X/K) \rightarrow 0.$$

□

6. EXAMPLES

All computations in the examples below were performed using Magma (described in [BCP97]). The code used to perform these computations may be found in the source of the arXiv distribution of this article.

6.1. A torsor of index 2 under an elliptic curve. We begin by illustrating how the results of this paper can be used to reproduce calculations of [HH07]. Namely, we show how to compute the relative Brauer group of a torsor of index 2 under an elliptic curve (provided generators for the Mordell-Weil group are known). We will compute the relative Brauer group of

$$V/\mathbb{Q} : y^2 = -2x^4 + 6x^2 + 4x - 14,$$

considered in [HH07, Example 4].

The Jacobian of V is the elliptic curve $X/\mathbb{Q} : y^2 = f(x)$, where $f(x) = x^3 - 27\mathcal{I}x - 27\mathcal{J}$ with $\mathcal{I} = 372$ and $\mathcal{J} = 12528$ the classical rational invariants associated to the quartic $g(x) = -2x^4 + 6x^2 + 4x - 14$. Let $L = \mathbb{Q}[\theta]/f(\theta)$. We can find $\ell \in L^\times$ such that $V \simeq V_\ell$ using [Cre01, Proposition 3.1] (see also [CF09, Section 2]). The *cubic invariant* of $g(x)$ is $-\frac{8}{3}\varphi + 32$, where φ is a root of the resolvent cubic $h(x) = x^3 - 3\mathcal{I}x + \mathcal{J}$ of $g(x)$. Noting that $\theta = -3\varphi$, it then follows from [CF09, Theorem 11] that $V \simeq V_\ell$ for

$$\ell = \frac{8}{9}\theta + 32 \equiv 2\theta + 72 \pmod{L^{\times 2}}$$

The corresponding algebra \mathcal{A}_ℓ is obtained by extension of scalars of the $\mathbb{Q}(x)$ -algebra,

$$\mathcal{A}'_\ell = \text{Cor}_{L(x)/\mathbb{Q}(x)}((2\theta + 72, x - \theta)_2).$$

Let $r(x) = 2x + 72$ and $a = -2^5 3^6 = f(x) \pmod{r(x)}$. By [CV15, Proposition 2.4],

$$[\mathcal{A}'_\ell] = [f(x), r(x)]_2 + [r(x), a]_2 + [1, 2]_2 + [2, a]_2.$$

Since $(1, 2)_2$, $(2, -2)_2$ and $(2, a)_2$ are split and $f(x)$ is a square in $\mathbf{k}(X)$, it follows that

$$[\mathcal{A}_\ell] = [2x + 72, -2]_2 = [x + 36, -2]_2.$$

(This should be compared with the algebra A_g defined in [HH07, Section 2].)

The group $E(\mathbb{Q})$ is generated by the points $P_1 = (-72, 108)$ and $P_2 = (450, 9288)$. Evaluating \mathcal{A}_ℓ at these points gives

$$\begin{aligned} [\mathcal{A}_\ell(P_1)] &= [-36, -2]_2 = [-1, -1]_2, \quad \text{and} \\ [\mathcal{A}_\ell(P_2)] &= [486, -2]_2 = [6, -2]_2 = 0. \end{aligned}$$

So, by Theorem 1.1, $\text{Br}(V/\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z}$ is generated by $[-1, -1]_2$.

Remark 6.1. In [HH07, Example 4] the Jacobian is given as $E : y^2 = x^3 - 12x^2 - 76x - 32$. The map $(x, y) \mapsto (9x - 36, 27y)$ defines an isomorphism between E and X under which P_1 and P_2 are the images of the points $(-4, 4)$ and $(54, 344)$, respectively.

6.2. Torsors of period 2 under elliptic curves with full rational 2-torsion. Let us consider an elliptic curve over K with full rational 2-torsion, say

$$J : y^2 = x(x - a)(x - b), \quad \text{with } a, b \in K^\times$$

Then every torsor of period 2 under J can be written in the form

$$V_\ell : \left\{ \begin{array}{l} au^2 - mv^2 + nt^2 = 0 \\ bu^2 - mv^2 + mnw^2 = 0 \end{array} \right\} \subset \mathbb{P}_{(u:v:w:t)}^3,$$

where $\ell = (m, n, mn) \in K^\times \times K^\times \times K^\times = L^\times$. The algebra \mathcal{A}_ℓ corresponding to ℓ is

$$\mathcal{A}_\ell = (x, m)_2 \otimes (x - a, n)_2 \otimes (x - b, mn)_2$$

The class of \mathcal{A}_ℓ in $\text{Br}(\mathbf{k}(X))$ can be written in any of the following equivalent ways:

$$\begin{aligned} [\mathcal{A}_\ell] &= [x, n]_2 + [x - a, m]_2, \\ &= [x, mn]_2 + [x - b, m]_2, \\ &= [x - a, mn]_2 + [x - b, n]_2. \end{aligned}$$

Evaluating this class at the nontrivial points in $J[2] \subset J(K)$ we obtain the following elements of $\text{Br}^0(V_\ell/K)$:

$$\begin{aligned} [\mathcal{A}_\ell(0, 0)] &= [-a, mn]_2 + [-b, n]_2, \\ [\mathcal{A}_\ell(a, 0)] &= [a, mn]_2 + [a - b, m]_2, \\ [\mathcal{A}_\ell(b, 0)] &= [b, n]_2 + [b - a, m]_2. \end{aligned}$$

If $J(K) = J[2]$, then these generate $\text{Br}^0(V_\ell/K)$ and $\text{Br}(V_\ell/K)/\text{Br}^0(V_\ell/K)$ is generated by the class of the conic $C_\ell : bu^2 - mv^2 + mnw^2$, which is $[C_\ell] = [bmn, -n]_2$.

For a concrete example we may take $K = \mathbb{Q}$, $a = 16$ and $b = -16$. Then $J(\mathbb{Q}) = J[2]$, and for $\ell = (-11, 3, -33)$ we find

$$\text{Br}^0(V_\ell/\mathbb{Q}) = \{[-1, -33]_2, [2, 3]_2, [-2, -11]_2, [1, 1]_2\}$$

and $[C_\ell] = [33, -1]_2 \notin \text{Br}^0(V_\ell/\mathbb{Q})$. In particular, we recover the result of Cassels [Cas63] that $P(V_\ell) \neq I(V_\ell)$.

Over an algebraically closed field, 2-torsion points on the Jacobian of a curve give rise to unramified double covers of the curve. In the presence of a nontrivial relative Brauer group this can fail, as shown in the following lemma.

Proposition 6.2. *Let V_ℓ/\mathbb{Q} be the curve corresponding to $\ell = (-11, 3, -33)$ and let E be its Jacobian. There does not exist an unramified double cover of V_ℓ defined over \mathbb{Q} (despite the fact that $E(\mathbb{Q})[2] \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$).*

Proof. Suppose $\pi : Y \rightarrow V_\ell$ is an unramified double cover. The kernel of $\pi^* : J \rightarrow \text{Jac}(Y)$ is generated by some $P \in J[2](\mathbb{Q})$ with $P \neq 0$. Lemma 2.6(2) implies that P is represented by a \mathbb{Q} -rational divisor on V_ℓ . However, as shown above, none of the nontrivial elements of $J[2]$ is represented by a \mathbb{Q} -rational divisor. \square

6.3. A generic torsor of period 2 and index 4. The presence of rational 2-torsion in the preceding example simplifies the computations, but it is by no means necessary. We now compute the relative Brauer group of a torsor of period 2 and index 4 in an example where there is no rational 2-torsion, but the Mordell-Weil group of the Jacobian is nontrivial (which has the potential to make the relative Brauer group larger).

Consider the elliptic curve $X : y^2 = f(x) = x^3 - 1296x + 11664$ labeled 37a in Cremona's Database. Set $L = \mathbb{Q}[\theta]/f(\theta)$ and $\ell = \frac{-11\theta^2 + 322\theta - 1584}{2} \in L^\times$. One can check that ℓ has norm $2^4 \times 3^6 \times 569^2$ and, hence, that ℓ gives rise (as described in Section 1.2) to a torsor V_ℓ of period 2 under X .

The Mordell-Weil group of X is generated by the point $(0, 108)$ of infinite order. Carrying out the computations as described in the example of Section 6.1 we obtain $[\mathcal{A}_\ell(P)] = [-1, -1]_2$, which shows that $\text{Br}^0(V_\ell/\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z}$. On the other hand, the conic C_ℓ is defined by the vanishing of

$$Q_\ell^{(2)} = 11u^2 - 644uv + 31680uw + 15840v^2 - 1091232vw + 24284448w^2,$$

whose class in $\text{Br}(\mathbb{Q})$ is $[-3, 569]_2$, which ramifies at 2 and 569. In particular it is not equal to $[-1, -1]_2$. Thus $\text{Br}(V_\ell/\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is strictly larger than $\text{Br}^0(V_\ell/\mathbb{Q})$, which shows that V_ℓ has index 4.

6.4. A torsor with period and index 4. Consider the genus one hyperelliptic curve X/\mathbb{Q} defined by $y^2 = f(x)$, where $f(x) = 5x^4 + 3x^2 + x + 3$. Set $L = \mathbb{Q}[\theta]/f(x)$. The element $\ell = 10\theta^3 - \theta^2 + 8\theta - 1 \in L^\times$ has norm $N_{L/\mathbb{Q}}(\ell) = 3920 \in 5\mathbb{Q}^{\times 2}$. So, as described in Section 1.3, ℓ corresponds to a torsor V_ℓ of index dividing 4 under the Jacobian J of X , which is the elliptic curve $J : y^2 = x^3 - 63x - 113$ with $J(\mathbb{Q})$ free of rank 2 generated by $P_1 = (-3, 7)$ and $P_2 = (-6, 7)$.

The curve X has no points over \mathbb{Q}_5 , and so X has period 2. Since $2V_\ell = X$ in the Weil-Châtelet group, V_ℓ has period and index 4. We will use Theorem 1.2 to compute the relative Brauer group of V_ℓ and determine which \mathbb{Q} -rational points on the Jacobian are represented by \mathbb{Q} -rational divisors on V_ℓ . Specifically we will show:

Example 6.3. $\text{Br}(V_\ell/\mathbb{Q})$ is generated by $[5, 7]_2 \in \text{Br}(\mathbb{Q})$ and a point $P \in J(\mathbb{Q})$ is represented by a \mathbb{Q} -rational divisor on V_ℓ if and only if its image in $J(\mathbb{Q})/2J(\mathbb{Q})$ lies in the subgroup generated by P_1 .

While X has no \mathbb{Q}_5 -points, it is locally solvable at all primes not equal to 5 and over the reals. It follows that $\text{Br}(X/\mathbb{Q}) = 0$. Hence the points P_1 and P_2 must be represented by \mathbb{Q} -rational divisors on X . The group of such divisors are generated by differences of traces of K -rational points on X as K ranges over quadratic extensions of \mathbb{Q} . A naive search over

small quadratic fields finds the $\mathbb{Q}(\sqrt{3})$ -points

$$Q_0 = (0, \sqrt{3}), \quad Q_1 = \left(\frac{1 + \sqrt{3}}{2}, -\frac{7 + 2\sqrt{3}}{2} \right), \text{ and}$$

$$Q_2 = \left(\frac{34 + 19\sqrt{3}}{2}, \frac{5011 + 2888\sqrt{3}}{4} \right).$$

Let σ denote the generator of $\text{Gal}(\mathbb{Q}(\sqrt{3})/\mathbb{Q})$. Then

$$D_1 = (Q_1 + \sigma(Q_1)) - (Q_0 + \sigma(Q_0)), \text{ and}$$

$$D_2 = (Q_2 + \sigma(Q_2)) - (Q_0 + \sigma(Q_0))$$

are \mathbb{Q} -rational divisors of degree 0 that represent $P_1, P_2 \in J(\mathbb{Q}) = \text{Pic}^0(\overline{X})^{\text{Gal}(\mathbb{Q})}$.

By Theorem 1.2 we have an exact sequence

$$0 \rightarrow \text{Pic}^0(V_\ell) \rightarrow \text{Pic}^0(X) \xrightarrow{\mathcal{A}_\ell} \text{Br}(V_\ell/\mathbb{Q}) \rightarrow 0,$$

where $\mathcal{A}_\ell = \text{Cor}_{\mathbf{k}(X_\ell)/\mathbf{k}(X)}(x - \alpha, \ell)_2$. Thus $\text{Br}(V_\ell/\mathbb{Q})$ is generated by $\mathcal{A}_\ell([D_i])$ for $i = 1, 2$. Since the x -coordinates of Q_0 and $\sigma(Q_0)$ are the same, $(x - \alpha)([D_i])$ is obtained by evaluating the minimal polynomial over \mathbb{Q} of the x -coordinate of Q_i at θ . Thus we obtain

$$(x - \alpha)([D_1]) = \theta^2 - \theta - 1/2,$$

$$(x - \alpha)([D_2]) = \theta^2 - 34\theta + 73/4.$$

Using Rosset-Tate reciprocity (as described in [GS06, Corollary 7.4.10 and Remark 7.4.12]) we find that $\mathcal{A}_\ell([D_1]) \equiv \sum_{i=1}^4 \mathcal{B}_i$ and $\mathcal{A}_\ell([D_2]) \equiv \sum_{i=1}^4 \mathcal{C}_i$, where

$$\begin{aligned} \mathcal{B}_1 &= [-483, -129]_2 \\ \mathcal{B}_2 &= [-3, -31]_2 \\ \mathcal{B}_3 &= [-401730, -645]_2 \\ \mathcal{B}_4 &= [-72345, -155]_2 \\ \mathcal{C}_1 &= [14717515766140493089374, 54870144093800907126]_2 \\ \mathcal{C}_2 &= [-89459567484222, -94665799875986]_2 \\ \mathcal{C}_3 &= [1051000391530337870055, 274350720469004535630]_2 \\ \mathcal{C}_4 &= [-6948230469054914235, -473328999379930]_2. \end{aligned}$$

Computing Hilbert symbols we find that $\mathcal{A}_\ell([D_1])$ is split, while $\mathcal{A}_\ell([D_2])$ is Brauer equivalent to $[5, 7]_2$, which ramifies only at 5 and 7.

Remark 6.4. *The torsor V_ℓ has points over \mathbb{Q}_p for all primes p other than 5 and 7. So in fact, we knew a priori that the relative Brauer group would be contained in the subgroup generated by $[5, 7]_2$. We take this as strong evidence of the correctness of our computations.*

6.5. Relative Brauer groups of higher dimensional torsors. As a final example we compute $\text{Br}^0(V_\ell/\mathbb{Q})$ for a torsor of period 2 under the Jacobian of a hyperelliptic curve of genus 2. Let X/\mathbb{Q} be the hyperelliptic curve defined by

$$X : y^2 = f(x) = x^6 + x^5 + x^4 + x^3 + x^2 + x + 1,$$

and let J be its Jacobian. Using Magma [BCP97] we compute that $J(\mathbb{Q}) \simeq \mathbb{Z}^2$ generated by divisors $P_1 = D_1 - \pi^*\infty$ and $P_2 = D_2 - \pi^*\infty$, where D_1 and D_2 are divisors on X whose x -coordinates are the roots of $t^2 + t + 1$ and $t^2 - t + 1$, and $\pi^*\infty$ is the pull-back of $\infty \in \mathbb{P}^1$ under the map $\pi : X \rightarrow \mathbb{P}^1$ sending a point on X to its x -coordinate.

For any $\ell \in L = \mathbb{Q}[\theta]/f(\theta)$ of square norm we can compute \mathcal{A}_ℓ , and as described in the previous example, $\mathcal{A}_\ell(P_1) = \text{Cor}_{L/\mathbb{Q}}(\theta^2 + \theta + 1, \ell)$ and $\mathcal{A}_\ell(P_2) = \text{Cor}_{L/\mathbb{Q}}(\theta^2 - \theta + 1, \ell)$. Moreover, the classes of these algebras in $\text{Br}(\mathbb{Q})$ generate $\text{Br}^0(V_\ell/\mathbb{Q})$, by Theorem 4.2.

For a concrete example, take $\ell = 3\theta^4 + 2\theta^2 + 2\theta + 3$ which has norm 43^2 . Then $\mathcal{A}_\ell(P_1)$ ramifies at 2 and 43, while $\mathcal{A}_\ell(P_2)$ is split. In particular, $\text{Br}^0(V_\ell/\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z}$ and a rational point on J is represented by a \mathbb{Q} -rational divisor on V_ℓ if and only if its image in $J(\mathbb{Q})/2J(\mathbb{Q})$ lies in the subgroup generated by P_2 .

A. APPENDIX

Let $E : y^2 = f(x)$ be an elliptic curve defined over a field K of characteristic not equal to 2, where $f(x) \in K[x]$ is a monic polynomial of degree 3 and set $L = K[x]/f(x)$. The Weil pairing gives an isomorphism $E[2] \simeq \text{Hom}(E[2], \mu_2) = \ker(\text{Norm}_{L/K} : \mu_2(\overline{L}) \rightarrow \mu_2(\overline{K}))$, thus inducing an isomorphism $e : H^1(K, E[2]) \simeq \mathfrak{L}_1 := \ker(\text{Norm}_{L/K} : L^\times/L^{\times 2} \rightarrow K^\times/K^{\times 2})$. Composing e with the connecting homomorphism δ yields Cassel's $x - \theta$ map (see [Sch95, Theorems 1.1 and 1.2]). This gives the following commutative and exact diagram.

$$\begin{array}{ccccccc}
0 & \longrightarrow & \frac{E(K)}{2E(K)} & \xrightarrow{\delta} & H^1(K, E[2]) & \longrightarrow & H^1(K, E)[2] \longrightarrow 0 \\
& & & & \parallel & & \nearrow v \\
& & & \searrow x-\theta & \mathfrak{L}_1 & &
\end{array}$$

Theorem A.1. *The map v in the diagram above sends the class of $\ell \in L^\times$ to the class of the torsor V_ℓ defined by (1.1).*

Proof. We will avail ourselves of the notation introduced at the beginning of Section 4. Denote the elements of Ω by $0_E, \omega_1, \omega_2, \omega_3$. Given $\ell \in L^\times \subset L_\circ^\times = \text{Map}_K(\Omega, \overline{K}^\times)$ such that $N_{L/K}(\ell) = a^2$, define $D_\ell \subset \mathbb{P}^3 \times E$ by declaring that $((u_0 : u_1 : u_2 : u_3), (x, y)) \in D_\ell$ if and only if

$$u_0^2(x - x(\omega_i)) = \ell_{\omega_i} u_i^2, \text{ for } i = 1, 2, 3, \text{ and } y = au_1 u_2 u_3.$$

(Here we are using ℓ_{ω_i} to denote the value of $\ell \in \text{Map}(\Omega, \overline{K}^\times)$ at $\omega_i \in \Omega$.) This condition is invariant under the action of G_K , so that D_ℓ is defined over K . Projection onto the first factor gives an isomorphism of D_ℓ and V_ℓ . Since V_ℓ is a complete intersection, D_ℓ is geometrically connected. Projection onto the second factor defines a morphism $\pi_\ell : D_\ell \rightarrow E$ which one checks is étale of degree 4. This implies that D_ℓ is a smooth curve of genus 1.

Any $\alpha \in \overline{L}^\times$ gives an automorphism $(u_0 : u_1 : u_2 : u_3) \mapsto (\alpha_{0_E} u_0 : \alpha_{\omega_1} u_1 : \alpha_{\omega_2} u_2 : \alpha_{\omega_3} u_3)$ of \mathbb{P}^3 . If $\alpha^2 = \ell$, then this induces an isomorphism of E -schemes $\overline{D}_\ell \simeq \overline{D}_1$. The projection D of D_1 onto \mathbb{P}^3 contains the K -rational point $0_D := (0 : 1 : 1 : 1)$ which lies in the fiber above 0_E under the composition $\pi : D \simeq D_1 \rightarrow E$. Thus $\pi : (D, 0_D) \rightarrow (E, 0_E)$ is a degree 4 isogeny of elliptic curves. The points in the kernel of this isogeny all lie on the hyperplane $u_0 = 0$. Since $4 \cdot 0_D$ is also a hyperplane section, the points in the kernel sum to 0_D . This

shows that the kernel of π is not cyclic. Thus the only possibility is that $D \simeq E$ and π is multiplication by 2.

We have shown that $\pi_\ell : D_\ell \rightarrow E$ is the twist of $[2] : E \rightarrow E$ by the cocycle $\xi_\ell : \sigma \mapsto \sigma(\alpha)/\alpha$, where $\alpha^2 = \ell$. The cocycle ξ_ℓ takes values in $\ker(\text{Norm}_{L/K} : \mu_2(\overline{L}) \rightarrow \mu_2(\overline{K})) \simeq E[2]$ and the image of its class under e is the class of ℓ in \mathfrak{L}_1 . On the other hand the isomorphism $\overline{D}_\ell \simeq \overline{D}_1 \simeq \overline{E}$ endows D_ℓ with the structure of an E -torsor over K and the image of ξ_ℓ in $H^1(K, E)[2]$ is represented by $D_\ell \simeq V_\ell$. \square

REFERENCES

- [BGW] Manjul Bhargava, Benedict Gross, and Xiaoheng Wang, *Pencils of quadrics and the arithmetic of hyperelliptic curves*, available at [arXiv:1310.7692](https://arxiv.org/abs/1310.7692). [↑11](#)
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). [↑14, 18](#)
- [Cas63] J. W. S. Cassels, *Arithmetic on curves of genus 1. V. Two counterexamples*, J. London Math. Soc. **38** (1963), 244–248. [↑2, 15](#)
- [Cas83] ———, *The Mordell-Weil group of curves of genus 2*, Arithmetic and geometry, Vol. I, Progr. Math., vol. 35, Birkhäuser, Boston, Mass., 1983, pp. 27–60. [↑3, 11](#)
- [Cas91] ———, *Lectures on elliptic curves*, London Mathematical Society Student Texts, vol. 24, Cambridge University Press, Cambridge, 1991. [↑2](#)
- [ÇK12] Mirela Çiperiani and Daniel Krashen, *Relative Brauer groups of genus 1 curves*, Israel J. Math. **192** (2012), no. 2, 921–949. [↑1, 4, 7, 8, 9](#)
- [CTS77] Jean-Louis Colliot-Thélène and Jean-Jacques Sansuc, *La R-équivalence sur les tores*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 2, 175–229 (French). [↑5](#)
- [Cre01] J. E. Cremona, *Classical invariants and 2-descent on elliptic curves*, J. Symbolic Comput. **31** (2001), no. 1-2, 71–87. Computational algebra and number theory (Milwaukee, WI, 1996). [↑14](#)
- [CF09] J. E. Cremona and T. A. Fisher, *On the equivalence of binary quartics*, J. Symbolic Comput. **44** (2009), no. 6, 673–682. [↑3, 14](#)
- [Cre13] Brendan Creutz, *Explicit descent in the Picard group of a cyclic cover of the projective line*, ANTS X—Proceedings of the Tenth Algorithmic Number Theory Symposium, Open Book Ser., vol. 1, Math. Sci. Publ., Berkeley, CA, 2013, pp. 295–315. [↑11](#)
- [CV15] Brendan Creutz and Bianca Viray, *Two torsion in the Brauer group of a hyperelliptic curve*, Manuscripta Math. **147** (2015), no. 1-2, 139–167. [↑1, 2, 10, 11, 12, 14](#)
- [Fis08] Tom Fisher, *Some improvements to 4-descent on an elliptic curve*, Algorithmic number theory, Lecture Notes in Comput. Sci., vol. 5011, Springer, Berlin, 2008, pp. 125–138. [↑3, 13](#)
- [FTvL12] E. Victor Flynn, Damiano Testa, and Ronald van Luijk, *Two-coverings of Jacobians of curves of genus 2*, Proc. Lond. Math. Soc. (3) **104** (2012), no. 2, 387–429. [↑11](#)
- [GS06] Philippe Gille and Tamás Szamuely, *Central simple algebras and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol. 101, Cambridge University Press, Cambridge, 2006. [↑17](#)
- [Hai84] Darrell E. Haile, *On the Clifford algebra of a binary cubic form*, Amer. J. Math. **106** (1984), no. 6, 1269–1280. [↑1](#)
- [HH07] Darrell Haile and Ilseop Han, *On an algebra determined by a quartic curve of genus one*, J. Algebra **313** (2007), no. 2, 811–823. [↑1, 2, 3, 14, 15](#)
- [HHW12] Darrell E. Haile, Ilseop Han, and Adrian R. Wadsworth, *Curves C that are cyclic twists of $Y^2 = X^3 + c$ and the relative Brauer groups $Br(k(C)/k)$* , Trans. Amer. Math. Soc. **364** (2012), no. 9, 4875–4908. [↑1](#)
- [Han03] Ilseop Han, *Relative Brauer groups of function fields of curves of genus one*, Comm. Algebra **31** (2003), no. 9, 4301–4328. [↑1](#)
- [Kuo11] Jung-Miao Kuo, *On an algebra associated to a ternary cubic curve*, J. Algebra **330** (2011), 86–102. [↑1](#)

- [Lic69] Stephen Lichtenbaum, *Duality theorems for curves over p -adic fields*, Invent. Math. **7** (1969), 120–136. [↑1](#), [5](#), [9](#)
- [MSS96] J. R. Merriman, S. Siksek, and N. P. Smart, *Explicit 4-descents on an elliptic curve*, Acta Arith. **77** (1996), no. 4, 385–404. [↑11](#), [13](#)
- [PS97] Bjorn Poonen and Edward F. Schaefer, *Explicit descent for Jacobians of cyclic covers of the projective line*, J. Reine Angew. Math. **488** (1997), 141–188. [↑3](#), [11](#), [12](#)
- [Sch95] Edward F. Schaefer, *2-descent on the Jacobians of hyperelliptic curves*, J. Number Theory **51** (1995), no. 2, 219–232. [↑3](#), [11](#), [18](#)
- [Shi94] Jonathan Shick, *On Witt-kernels of function fields of curves*, Recent advances in real algebraic geometry and quadratic forms (Berkeley, CA, 1990/1991; San Francisco, CA, 1991), Contemp. Math., vol. 155, Amer. Math. Soc., Providence, RI, 1994, pp. 389–398. [↑1](#)
- [Wei56] André Weil, *The field of definition of a variety*, Amer. J. Math. **78** (1956), 509–524. [↑6](#)

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CANTERBURY, PRIVATE BAG 4800,
CHRISTCHURCH 8140, NEW ZEALAND

E-mail address: `brendan.creutz@canterbury.ac.nz`

URL: `http://www.math.canterbury.ac.nz/~bcreutz`