

An Affine Regular Icosahedron Inscribed in an Affine Regular Octahedron in a GS-Quasigroup

An Affine Regular Icosahedron Inscribed in an Affine Regular Octahedron in a GS-Quasigroup

ABSTRACT

A golden section quasigroup or shortly a GS–quasigroup is an idempotent quasigroup which satisfies the identities $a(ab \cdot c) \cdot c = b$, $a \cdot (a \cdot bc)c = b$. The concept of a GS–quasigroup was introduced by VOLENEC. A number of geometric concepts can be introduced in a general GS–quasigroup by means of the binary quasigroup operation. In this paper, it is proved that for any affine regular octahedron there is an affine regular icosahedron which is inscribed in the given affine regular octahedron. This is proved by means of the identities and relations which are valid in a general GS–quasigroup. The geometrical presentation in the GS–quasigroup $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ suggests how a geometrical consequence may be derived from the statements proven in a purely algebraic manner.

Key words: GS–quasigroup, GS–trapezoid, affine regular icosahedron, affine regular octahedron

MSC2010: 20N05

Afino pravilan ikozaedar upisan u afino pravilan oktaedar u GS–kvazigrupi

SAŽETAK

Kvazigrupa zlatnog reza ili kraće GS–kvazigrupa idempotentna je kvazigrupa u kojoj vrijede identiteti $a(ab \cdot c) \cdot c = b$, $a \cdot (a \cdot bc)c = b$. Pojam GS–kvazigrupe uveo je VOLENEC. Razni geometrijski pojmovi mogu biti uvedeni u GS–kvazigrupi pomoću binarne operacije te kvazigrupe. Korištenjem relacija i identiteta u općoj GS–kvazigrupi u ovom je radu pokazano da se svakom afino pravilnom oktaedru može upisati afino pravilan ikozaedar. Geometrijski prikaz u kvazigrupi $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ pokazuje kako geometrijske tvrdnje mogu biti posljedica potpuno algebarskih razmatranja.

Ključne riječi: GS–kvazigrupa, GS–trapezoid, afino pravilan ikozaedar, afino pravilan oktaedar

1 Introduction

The concept of a GS–quasigroup was introduced by Volenec in [1].

A quasigroup (Q, \cdot) , which satisfies the identity of idempotency

$$aa = a \quad (1)$$

and the (mutually equivalent) identities

$$a(ab \cdot c) \cdot c = b, \quad (2)$$

$$a \cdot (a \cdot bc)c = b, \quad (2)'$$

is called a golden section quasigroup or shortly a GS–quasigroup.

It can be proved that the considered GS–quasigroup (Q, \cdot) satisfies the identities of mediality, elasticity, left and right distributivity, i.e., the identities

$$ab \cdot cd = ac \cdot bd, \quad (3)$$

$$a \cdot ba = ab \cdot a, \quad (4)$$

$$a \cdot bc = ab \cdot ac, \quad (5)$$

$$ab \cdot c = ac \cdot bc. \quad (5)'$$

are valid. Some other identities, e.g.

$$a(ab \cdot b) = b, \quad (6)$$

$$(b \cdot ba)a = b, \quad (6)'$$

are also valid in a general GS–quasigroup.

Let us mention the best known example of a GS–quasigroup. Let \mathbb{C} be the set of points of the Euclidean plane. For any two different points a, b we define $ab = c$ if the point b divides the pair a, c in the ratio of the golden section. In [1], it is proved that (\mathbb{C}, \cdot) is a GS–quasigroup. This quasigroup is denoted by $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$ because we have $c = \frac{1}{2}(1 + \sqrt{5})a$ if $a = 0$ and $b = 1$. As this quasigroup has a nice geometrical interpretation, we shall use this quasigroup for the purpose of illustrating the identities and relationships holding in a general GS–quasigroup.

From now on, let (Q, \cdot) be any GS–quasigroup. The elements of the set Q are said to be points.

Some geometric concepts, such as parallelograms, GS–trapezoids, affine regular pentagons, an affine regular icosahedron and an affine regular octahedron, can be introduced by means of the given binary quasigroup operation (see [1], [2], [3], [4]).

The points a, b, c, d are said to be the vertices of a parallelogram and we write $Par(a, b, c, d)$ if the identity

$$a \cdot b(ca \cdot a) = d \tag{7}$$

holds. In [1], some properties of the quaternary relation Par on the set Q are proved. It is proved that the structure (Q, Par) is a parallelogram space, i.e., that the following three properties hold:

- (P1) For any three points a, b, c , there exists one and only one point d such that there holds $Par(a, b, c, d)$.
- (P2) From $Par(a, b, c, d)$ there follows $Par(e, f, g, h)$, where (e, f, g, h) is any cyclic permutation of (a, b, c, d) or (d, c, b, a) .
- (P3) From $Par(a, b, c, d)$ and $Par(c, d, e, f)$ there follows $Par(a, b, f, e)$.

In [1], the following statement is proved.

Lemma 1 From $Par(a, b, d, e)$ and $Par(b, c, e, f)$ there follows $Par(c, d, f, a)$.

We shall say that b is the midpoint of the pair of points a, c and write $M(a, b, c)$ if $Par(a, b, c, b)$.

In [2], the concept of a GS–trapezoid is defined. The points a, b, c, d successively are said to be the vertices of the golden section trapezoid and it is denoted by $GST(a, b, c, d)$ if the identity

$$a \cdot ab = d \cdot dc \tag{8}$$

holds.

In [2], it is proved that any two of the five statements

$$GST(a, b, c, d), GST(b, c, d, e), GST(c, d, e, a), GST(d, e, a, b), GST(e, a, b, c) \tag{9}$$

imply the remaining statement.

In [3], the concept of an affine regular pentagon is defined. The points a, b, c, d, e successively are said to be the vertices of the affine regular pentagon and it is denoted by $ARP(a, b, c, d, e)$ if any two (and then all five) of the five statements (9) are valid.




Figure 1: Affine regular icosahedron in $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$

In [4], the concepts of an affine regular icosahedron and an affine regular octahedron in a general GS–quasigroup are introduced.

We shall say that the points $a, b, c, d, e, f, a', b', c', d', e', f'$ are the vertices of an affine regular icosahedron (Figure 1) and write $ARI(a, b, c, d, e, f, a', b', c', d', e', f')$ if the following twelve statements are valid

$$\begin{aligned} &ARP(b, c, f, a', e), \quad ARP(c, a, d, b', f), \\ &ARP(a, b, e, c', d), \quad ARP(b', c', f', a, e'), \\ &ARP(c', d', a', b, f'), \quad ARP(d', b', e', c, d'), \\ &ARP(b, c, e', d, f'), \quad ARP(c, a, f', e, d'), \\ &ARP(a, b, d', f, e'), \quad ARP(b', c', e, d', f), \\ &ARP(c', a', f, e', d), \quad ARP(d', b', d, f', e). \end{aligned}$$

The points a, b, c, a', b', c' are the vertices of an affine regular octahedron with the center o if the statements $M(a, o, a'), M(b, o, b'), M(c, o, c')$ are valid (Figure 2).




Figure 2: Affine regular octahedron in $\mathbb{C}(\frac{1}{2}(1 + \sqrt{5}))$

The vertices of an affine regular octahedron are the vertices of three parallelograms. We have the statement.

Lemma 2 From $M(c, o, c')$ and $M(b, o, b')$ there follows $Par(b', c', b, c)$.

Proof. The statements $M(c, o, c')$ and $M(b, o, b')$ are equivalent to $Par(c, o, c', o)$ and $Par(b, o, b', o)$, from where, according to (P2), we get $Par(o, b', o, b)$, whence by Lemma 1 we obtain $Par(b', c', b, c)$. \square

In [4], it is proved that for any regular octahedron there is a regular icosahedron which is circumscribed to the given affine regular octahedron. Now, we shall prove that for any affine regular octahedron there is a regular icosahedron which is inscribed in the given affine regular octahedron. So, we have the following statement.

Theorem 1 If a, b, c, a', b', c' are the vertices of an affine regular octahedron, then $ARI(a_b, b_c, c_a, a'_b, b'_c, c'_a, a_b, b_c, c_a, a'_b, b'_c, c'_a)$, where $a_b = ab \cdot b, b_c = bc \cdot c, \dots, c'_a = c'a' \cdot a'$, (Figure 3).




Figure 3.

Proof. It is enough to prove the statement $ARP(a_b, c_a, a'_b, b'_c, b'_c)$. According to (9), it is sufficient to prove that the statements $GST(a_b, c_a, a'_b, b'_c)$ and $GST(a'_b, c_a, a_b, b'_c)$ are valid. In fact, it is enough to prove only the first one because the second statement follows from the first one by substitution $b \leftrightarrow b'$. According to the definition of a GS-trapezoid, because of (8), for the proof of the first statement we have to prove

$$a_b \cdot a_b c_a = b'_c \cdot b'_c a'_b.$$

We obtain

$$\begin{aligned} a_b \cdot a_b c_a &= (ab \cdot b) \cdot (ab \cdot b)(ca \cdot a) \\ &\stackrel{(3)}{=} (ab)(ab \cdot b) \cdot b(ca \cdot a) \\ &\stackrel{(5)}{=} (a \cdot ab)b \cdot b(ca \cdot a) \stackrel{(6')}{=} a \cdot b(ca \cdot a), \end{aligned}$$

from where, owing to (7), we get that the point $a_b \cdot a_b c_a$ is the fourth vertex of the parallelogram with the vertices a, b, c , i.e., $Par(a, b, c, a_b \cdot a_b c_a)$. We can get $Par(b', c', a, b'_c \cdot b'_c a'_b)$ in the same way. It is necessary to prove that the fourth vertices of these two parallelograms are coincident, i.e., the statement $Par(a, b, c, x)$ implies the statement $Par(b', c', a, x)$. However, because the points b', c', b, c are the vertices of the octahedron, according to Lemma 2, $Par(b', c', b, c)$ holds, which, together with $Par(b, c, x, a)$ and by P3, gives $Par(b', c', a, x)$. \square

In this case, we shall say that an affine regular icosahedron is *inscribed* in the given affine regular octahedron (Figure 4).




Figure 4.

References

- [1] V. VOLENEC, GS-quasigroups, *Čas. pěst. mat.* **115** (1990), 307–318.
- [2] V. VOLENEC, Z. KOLAR-BEGOVIĆ, GS-trapezoids in GS-quasigroups, *Math. Commun.* **7** (2002), 143–158.
- [3] V. VOLENEC, Z. KOLAR-BEGOVIĆ, Affine regular pentagons in GS-quasigroups, *Quasigroups Related Systems* **12** (2004), 103–112.
- [4] V. VOLENEC, Z. KOLAR-BEGOVIĆ, R. KOLAR-ŠUPER, Affine regular icosahedron circumscribed around the affine regular octahedron in GS-quasigroup, *Comment. Math. Univ. Carolin.* **53** (2012), 501–507.

Zdenka Kolar-Begović

orcid.org/0000-0001-8710-8628

e-mail: zkolar@mathos.hr

Department of Mathematics, University of Osijek
Gajev trg 6, HR-31 000 Osijek, Croatia