

Influence of Gibberellin on Increasing of Sodium Chloride Tolerance via Some Morpho-Physiological Changes in Two Olive Cultivars

Akhtar SHEKAFANDEH (✉)
Shahrnaz SIROOEENEJAD
Elham ALSMOUSHTAGHI

Summary

Cultivars of Olive (*Olea europaea* L.) have different ability for adaption to saline environments. Salinity is one of the major factors limiting plant growth and development. The aim of this research was to study the salinity status tolerance of two olive cultivars: 'Zard' and 'Shiraz' and also the possibility of using gibberellin to increase plant salt tolerance. Treatments were different concentrations of sodium chloride: 0, 50, 100 and 200 mg L⁻¹, which were added to the pots via irrigation water, and foliage application of gibberellin (GA₃) at 0, 10 and 100 mg L⁻¹, which were applied one month after salinity treatments on plants. The results showed that increasing salinity declined leaf area; shoots and roots length; leaf, shoots and roots fresh and dry weights, and increased the amount of leaf proline, and leaf and roots Na⁺ and Cl⁻. Application of gibberellin reduced sodium and chloride concentrations in plants, but increased the amount of potassium and chlorophyll. The rate of Na⁺ accumulation in leaves and roots was lower in 'Shiraz' than in 'Zard'. In different levels of sodium chloride, gibberellin increased the synthesis and accumulation of proline in two cultivars, but this increase was more pronounced in 'Shiraz'. 'Shiraz' showed more vegetative growth than 'Zard'.

Key words

growth, ions accumulation, *Olea europaea*, salinity stress, tolerance mechanisms

¹ Department of Horticultural Science, College of Agriculture, Shiraz University, Shiraz Iran
✉ e-mail: shekafan@shirazu.ac.ir

Received: April 22, 2017 · Accepted: September 2, 2017

ACKNOWLEDGEMENTS

The authors wish to thank Mrs. Fatemeh Esfandiyari for technical assistance in the laboratory.

Introduction

Salinity is an important factor affecting plant productivity and constitutes a problem concerning a significant portion of the planet, especially in regions with hot, dry climates (Chartzoulakis et al., 2006; Bazakos et al., 2015). The availability of fresh water is one of the major limitations for crop production. Therefore, the use of non-conventional water resources, such as saline water and reclaimed sewage effluent, has increased in recent years. The utilization of such water resources accelerates the salinization of the upper layer of the soil, where most root activity takes place, and generally decreases crop production (Demiral et al., 2011).

Plants face two basic problems in saline environments. First, excess salt in soil lowers the osmotic potential of soil solution and leads to decreased water uptake and consequently water deficit in plants. This in turn leads to perturbations in cell division and/or extension and influences the integrity of metabolic reactions in plants. Second, increased uptake and accumulation of Na^+ and Cl^- ions decreases the absorption of essential minerals and imposes toxicity to plants (Munns, 1993; Tester and Davenport, 2003). The significance of salinity for the agronomical and physiological aspects of plants is enormous.

All salts can affect plant growth, but not all inhibit growth. Among the most common effects of salinity is growth inhibition by NaCl . For some plants, particularly fruit trees such as citrus and grapevines, accumulation of both Na^+ and Cl^- in the roots and aerial parts is the most damaging to the plants often by inhibiting photosynthesis (Munns, 1993; Flower and Yeo, 1988; White and Broadley, 2001). Na^+ is the primary cause of ion-specific damage (such as reduction in K^+ activity) (Tabatabaei, 2006). Improving plant resistance to salinity may provide yield stability in subsistence agriculture (Flowers and Yeo, 1995; Flowers, 1999; Asghari, 2008).

Usage of plant growth regulators as an easy and rapid tool can help to ameliorate the adverse effects of abiotic stresses such as salinity. Gibberellins (GA_3) play a vital role in the detoxification of heavy metals and in tolerance to salt stress by improving plants growth, chlorophyll synthesis and activities of antioxidant enzymes, and by preventing lipid peroxidation (Saeidi-Sar et al., 2007; Maggio et al., 2010). Some researchers have used plant growth regulators (PGRs) for reducing or eradicating the negative effects of salinity (Angrish et al., 2001; Chakrabarti and Mukherji, 2002). For example, the exogenous application of PGRs [auxins (Khan et al., 2004), gibberellins (Shaheena et al., 2005), cytokinins (Gul et al., 2000)] produces some benefit in alleviating the adverse effects of salt stress and also improves germination, growth, development, seed yield, and yield quality (Egamberdieva, 2009). Abd-El-Samad (1998), Abd-El-Samad and El-Komy (1998) and Azooz et al. (2004) reported that indole acetic acid (IAA) and gibberellic acid stimulate growth in sorghum under stress conditions.

Olive (*Olea europaea* L.) tree cultivation is widespread throughout the Mediterranean, where it has been adapted to the dry climate of the region. However, irrigation at primary stages of plant establishment and during certain growth phases is essential for satisfactory fruit production. Nevertheless, the availability of good-quality irrigation water is limited, and in some areas the use of saline water is unavoidable. Although the olive trees are moderately tolerant to salinity (Mass and Hoffman, 1977), significant differences in salt tolerance have been reported among cultivars (Therios and Misopolinos, 1988; Tattini, 1994; Chartzoulakis et al., 2002).

The present study was therefore designed as an attempt to characterize the influence of GA_3 on the adverse effects of salt stress in two olive cultivars.

Materials and methods

Experiment was conducted in the greenhouse of Shiraz University of Agricultural Sciences during 2011-2012. One year-old own rooted plants of two olive (*Olea europaea* L.) cultivars ('Zard' and 'Shiraz') were transplanted into 8-liter pots filled with sand and perlite (1/1, v/v) and irrigated with 200 ml Hoagland nutrient solution. The pots were kept in the glasshouse with natural sunlight and at temperatures 30 ± 3 during the day and $20 \pm 3^\circ\text{C}$ during the night. Treatments were four salinity concentrations as NaCl (0, 50, 100, 200 mg L^{-1}) combined with three gibberellin (GA_3) concentrations (0, 10, 100 mg L^{-1}). Treatments were established in four replications. After four weeks of transplanting, salinity treatments were started. For avoiding osmotic shock, they were added into the pots in three steps with irrigation water. After 30 days of salt application, the plants were subjected to GA_3 (Sigma Chemical Co., St. Louis, U.S.A.) sprays. Plants were harvested 16 weeks later for measuring growth parameters, ionic contents, proline, and chlorophyll. The shoots and roots length were measured using ruler. Leaf area surface was determined by leaf area meter (LI-300- USA). The harvested shoots and roots were weighed, dried in an oven for two days at 85°C , re-weighed and grounded for determination of ion compositions. Sodium and potassium contents were measured by flame photometry (model JENWAY, PEP-7) after digesting about 100 mg of plant material in a mixture of concentrated nitric and perchloric acid (3:1) at 175°C . Chloride content was measured at 450 nm according to the method of Diatloff and Rengel (2001) using $\text{Hg}(\text{SCN})_2$ and FeNO_3 . To measure the chlorophyll content, after weighing 0.2 g fresh tissue of leaves, they were grounded with a pestle and mortar in 5 ml of distilled water in cool temperature of about 10°C and dim light condition to change into unified mass and then its volume was made up to 25 ml with distilled water. The amount of 4.5 ml of 80% acetone was added to 0.5 ml of the mixture and centrifuged at 3500 ($\times g$) for 15 minutes. Then, the absorbance of the supernatant solution was measured at 645 and 663 nm in a BioQuest CE 2502 spectrophotometer (Arnon, 1949). To measure the proline content in the leaves, 0.1 g fresh leaves grounded well with pestle and mortar in 10 ml of 3.3% sulphosalicylic acid. After filtering, the filtrate was collected in a test tube and kept in an ice and water mixture. In the next step, 2 ml of ninhydrin solution (1.25 ninhydrin + 20 ml 6M phosphoric acid + 30 ml pure acetic acid) and 2 ml of acetic acid were added to every test tube containing the extract. The tubes were placed in a boiling water bath maintained at 100°C for one hour and then the mixtures were cooled. Under the hood, 6 ml Toluene was added to each tube and shook well for 15-20 seconds. In this condition, two different phases were made. The absorbance was measured at 520 nm by spectrophotometer for 1 ml of the upper phase containing proline (Bates et al., 1973).

The experiment was carried out in a completely randomized factorial design. The factors were four levels of salinity and three levels of GA_3 with four replications. Data were subjected to analysis of variance (ANOVA) using the SPSS version 15.0 statistical package (SPSS, Inc., Chicago, USA). For comparison of means the Tukey's test was applied ($P < 0.05$).

Results

The results of analysis of variance showed a high interaction between cultivar, salinity and gibberellin for all measured growth traits except for shoot and root dry weight (DW) (Table 1).

Table 1. ANOVA results of the significance of the effect of cultivar (Cul), salinity (Sal) and gibberellin (GA) interactions (Cul×Sal, Cul×GA, Sal×GA and Cul×Sal×GA) on growth traits of two olive cultivars

Source of Variation	Leaf area	Shoot FW	Root FW	Shoot DW	Root DW	Shoot L	Root L
Cul	**	**	**	**	**	**	**
Sal	**	**	**	**	**	**	**
GA	**	**	**	**	**	**	**
Cul×Sal	**	**	ns	**	ns	**	**
Cul×GA	**	**	**	ns	ns	**	**
Sal×GA	**	**	*	**	*	**	**
Cul×Sal×GA	**	**	**	ns	ns	**	**

Note: * significant at $P = 0.05$, ** significant at $P = 0.01$, ns - not significant

Effects of GA₃ on plant growth, biomass and leaf area

Growth of plants (shoot and root length indexes) was reduced when NaCl concentrations was increased to 200 mg L⁻¹ in the pots. The application of GA₃ at 100 mg L⁻¹ improved the growth of both olive cultivars. At 200 mg L⁻¹ NaCl, GA₃ spraying with 100 mg L⁻¹ significantly increased the length of roots in both cultivars and the shoot length in ‘Shiraz’ cultivar. Generally, gibberellin was more effective in increasing the length of shoots and roots in ‘Shiraz’ than in ‘Zard’ (Fig. 1 a and b).

In control plants (no treatments), the largest leaf area surface was obtained in ‘Shiraz’ cultivar, compared with the corresponding values in ‘Zard’ cultivar. High concentration of salinity (200 mg L⁻¹) decreased the leaf area by 43% and 40% in ‘Zard’ and ‘Shiraz’, respectively, compared to their controls. Interactive effect of salinity and GA₃ concentration on leaf area was significant in both cultivars. It was interesting that in 200 mg L⁻¹ of NaCl, 100 mg L⁻¹ GA₃ promoted the leaf growth to the same surface as in the control plants (Fig. 1c).

The results of analysis of variance showed a high interaction between cultivar, salinity and gibberellin on all measured physiological traits except for chlorophyll content (Table 2).

Table 2. ANOVA results of significance of the effect of cultivar (Cul), salinity (Sal), and gibberellin (GA) interactions (Cul×Sal, Cul×GA, Sal×GA and Cul×Sal×GA) on physiological traits of two olive cultivars

Source of Variation	Proline	Chlorophyll	Leaf Na+	Root Na+	Leaf K+	Root K+	Leaf Cl-	Root Cl-
Cul	**	**	**	**	**	**	**	**
Sal	**	**	**	**	**	**	**	**
GA	**	**	**	**	**	**	**	**
Cul×Sal	**	ns	**	**	**	**	**	**
Cul×GA	**	ns	**	**	**	**	**	**
Sal×GA	**	*	**	**	**	**	**	**
Cul×Sal×GA	**	ns	**	**	**	**	**	**

Note: * significant at $P = 0.05$, ** significant at $P = 0.01$, ns - not significant




Figure 1. Combined effects of NaCl and GA₃ on shoot and root length (a and b) and leaf area (c) of two olive cultivars (Note: means with the same letters are not significantly different at $P < 0.05$ using Tukey’s test).




Figure 2. Combined effects of NaCl and GA₃ on shoot fresh and dry weight (a and b), root fresh and dry weight (c and d) of two olive cultivars (Note: means with the same letters are not significantly different at $P < 0.05$ using Tukey's test).

Effects of GA₃ on proline and chlorophyll content

Proline concentration of both cultivars significantly increased with increasing salinity. The proline accumulation was greater in the 'Shiraz' cultivar compared to the 'Zard' cultivar. In both cultivars, the highest proline concentrations were obtained in plants treated with 200 mg L⁻¹ NaCl. In each level of salinity, foliar application of GA₃, especially at 100 mg L⁻¹, also significantly increased proline concentrations (Fig. 3a).

The results showed that leaf chlorophyll content was significantly reduced in the salt-treated plants. The maximum of chlorophyll reduction was observed in plants under 200 mg L⁻¹ of NaCl. In all levels of salinity, application of GA₃ at 100 mg L⁻¹ increased the chlorophyll content of olive plants in both cultivars. However, chlorophyll content in 'Zard' was higher than in 'Shiraz' (Fig. 3 b).




Figure 3. Combined effects of NaCl and GA₃ on proline (a) and chlorophyll (b) content of two olive cultivars (Note: means with the same letters are not significantly different at $P < 0.05$ using Tukey's test).

Effects of GA₃ on leaf and root Na⁺, K⁺ and Cl⁻ contents

In both cultivars, the leaf Na⁺ content was below 5 mg g⁻¹ DW in control plants, while the amount of Na⁺ in roots of 'Shiraz' was considerably lower than 'Zard'. The Na⁺ content in the leaves and roots increased as salinity concentration increased in root zone. Foliar application of GA₃ significantly decreased the Na⁺ concentration in both organs (Fig. 4 a, b).

Salinity in the root zone led to a significant decrease in K⁺ concentration in the plant tissue in both cultivars regardless of the GA₃ levels. The lowest concentrations of K⁺ occurred in the treatments




Figure 4. Combined effects of NaCl and GA₃ on leaf and root Na⁺ (a and b), K⁺ (c and d) and Cl⁻ (e and f) contents of two olive cultivars (Note: means with the same letters are not significantly different at $P < 0.05$ using Tukey's test).

with the highest NaCl levels (200 mg L⁻¹). In all levels of salinity, spraying the plants with GA₃ increased K content of leaves and roots (Fig. 4 c, d).

Leaf and root Cl⁻ contents were also affected by increasing salinity levels. The application of GA₃ under such conditions was found to decrease Cl⁻ concentration in both cultivars (Fig. 4 e, f).

Discussion

Exposure to high salinity induced a general modification in all parameters studied. In addition, higher concentration of NaCl (200 mg L⁻¹) provoked stronger deleterious effects. Meanwhile, treatment with GA₃ clearly mitigated the adverse effects of salt stress based on the improved parameters studied as compared to the untreated plants. The drastic increase in the concentration of Na⁺ and Cl⁻ in

tissues that following plant exposure to salinity led to toxicity as it was evidenced by reduced plant growth. It is well documented that salt tolerance in glycophytes is associated with the ability to limit the uptake and/or transport of salt ions (mainly Na^+ and Cl^-) from root to shoot (Greenway and Munns, 1980; Tester and Davenport, 2003; Ben-Ahmed et al., 2006; Raza et al., 2007). The exposure of plants to salinity commonly results in a water deficit in plant cells, and maintaining osmotic homeostasis requires an adjustment in osmosis, either by the uptake of soil solutes or by the synthesis of metabolically compatible compounds (Tester and Devenport, 2003). It is known that olive cultivars have an effective salt-exclusion mechanism operating in their thin roots. The control mechanism includes limiting salt translocation to the leaves rather than salt absorption by the roots (Demiral, 2005). Additionally, the process is also supported by some other mechanisms such as stomata closure (Fernandez et al., 1997) and leaf tolerance to dehydration (Dichio et al., 2002). Leaf abscission at high salinity (200 mg L^{-1}) occurred in 'Shiraz' and 'Zard' and led to further reduction in total leaf area. This may be caused by ions accumulation in the leaves, particularly old leaves (Greenway and Munns, 1980). The reduction in plant growth was due to the reduced leaf growth, which agrees with finding of Cramer (2002) and Tabatabaei (2006). The reduction observed in leaf area and dry matter of the salt-treated plants (Fig 1 c, Fig 2 b) can be attributed to the changes in plant water relations under salt stress, which cause a reduction in meristem activity as well as cell elongation (Shah, 2007) thereby inhibiting leaf expansion (Bernstein, 1993). Furthermore, high salinity is known to induce ionic stress, which causes premature abscission and senescence of adult leaves, thus reducing the available photosynthetic area (Munns, 2002). Thus, the observed decrease in dry matter of the salt-stressed plants can be traced to the scanty recovery of leaves following limited photosynthesis production. Aldesuquy and Ibrahim (2001) proposed that hormones used during salt stress may reduce water loss rates and cause a concomitant increase in leaf water potential and carbon gain rates. In the present study, foliar application of GA_3 might have de-repressed the leaf area expansion and caused increased dry weight production in the salt-treated plants. The observed chlorophyll depletion may be considered to be a result of the inhibition of chlorophyll biosynthesis following an increase in ethylene production brought about by the elevated NaCl content (Khan, 2003). Further, chlorophyllase activity increases during stress conditions (Singh and Jain, 1981), suggesting that the observed low chlorophyll content could be a result of both decreased synthesis and increased degradation under salt stress. However, treatment of the salt-stressed plants with GA_3 restored normal chlorophyll levels. This may well be attributed to the GA_3 -generated enhancement of ultra-structural morphogenesis of plastids coupled with retention of chlorophyll and delay of senescence caused by the hormone treatment (Khan et al., 2010). Similarly, the enhanced accumulation of proline in the plants raised from GA_3 may represent: a major biochemical adaptation in plants, osmotic adjustment (Siddiqui et al., 2008; Khan et al., 2010), membrane stabilizer (Bandurska, 2001), and reactive oxygen species (ROS) scavenger (Matysik et al., 2002). The reduction of K^+ in olive leaves found in this study (Fig 4 c, d) confirmed previous results by Tattini et al. (1994) and Loupassaki et al. (2002). Usually, K^+ concentration is reduced by increased salinity, although there are some tree species, like citrus, where salinity does not significantly affect leaf

K^+ content, depending on the rootstock used (Storey and Walker, 1999). Therefore, plant metabolism is directly affected by the alteration of K^+/Na^+ ratio under saline conditions. It has been suggested that K^+ is one of the osmolytes and that its accumulation in plant cells might facilitate osmotic adjustment, lowers the internal osmotic potential, and contributes to salt tolerance (Hasegawa et al., 2000). An important mechanism to avoid the deleterious effects of salinity in olive trees is the ability to limit uptake and/or transport of saline ions (sodium and chloride) from the root zone to aerial parts; salt tolerance in olive cultivars is associated with effective mechanisms of ion exclusion and retention of sodium and chloride in the root (Kchaou et al., 2010). GA_3 induces Ca^{2+} and other nutrients uptake that may be involved in plant tolerance to stress by regulating antioxidant metabolism and reduction of the lipid peroxidation of cell membrane (Siddiqui et al., 2008; Khan et al., 2010; Badr-uz-Zaman et al., 2010). Furthermore, these may also be involved in signal transduction (McAinsh et al., 1996) and gene expression (Braam, 1992). Tattini (1994) reported that the resistance mechanism of salt tolerant olive cultivars is probably related to the ability to maintain an appropriate K^+/Na^+ ratio in actively growing tissue. Therefore, it may be speculated that higher K^+/Na^+ and $(\text{K}^+ + \text{Ca}^{2+} + \text{Mg}^{2+})/\text{Na}$ ratios of the plant leaves can be accepted as key indicators reflecting the levels of adaptation of the cultivar to salt stress (Kasirga and Demiral, 2016).

Conclusion

Our results indicate that high concentrations of salt (NaCl) provoke a general modification in all morphological and physiological parameters studied. Application of GA_3 ameliorates the deleterious effect of Na^+ and Cl^- ions. Cultivar 'Shiraz' has the ability to maintain higher K^+ and K^+/Na^+ ratio than 'Zard' and better growth performance. Therefore, it can be concluded that 'Shiraz' is possibly less sensitive to salinity.

References

- Abd-El-Samad H.M., (1998). The counteraction effect of GA_3 or IAA with endogenous ethylene of wheat plants under salt stress condition. *J Bot* 1: 79-110.
- Abd-El-Samad H.M., El-Komy H.M., (1998). Effect of salinity, gibberellic acid and azospirillum inoculation on growth and nitrogen uptake of *Zea mays*. *Biol Planta* 40:109-120.
- Aldesuquy H.S., Ibrahim A.H., (2001). Interactive effect of seawater and growth bio-regulators on water relations, abscisic acid concentration, and yield of wheat plants. *J Agron Crop Sci* 187: 185-193.
- Angrish A., Kumar B., Datta K.S., (2001). Effect of gibberellic acid and kinetin on nitrogen content and nitrate reductase activity in wheat under saline conditions. *Ind J Plant Physiol* 6: 172-177.
- Arnon D.I., (1949). Copper enzymes in isolated chloroplast. Polyphenoloxidases in *Beta vulgaris*. *Plant Physiol* 24:1-15.
- Asghari H.R., (2008). Vesicular-arbuscular (VA) mycorrhizae improve salinity tolerance in pre-inoculation subterranean clover (*Trifolium subterraneum*) seedlings. *Int J Plant Production* 2: 243-256.
- Azooz M.M., Shaddad M.A., Abdel-Latef A.A., (2004). Leaf growth and K^+ , Na^+ ratio as an indication of the salt tolerance of three sorghum cultivars grown under salinity stress and IAA treatment. *Acta Agron Hung* 52:287-296.
- Badr-uz-Zaman S.M., Asghar R., (2010). Role of Ca^{2+} on growth of *Brassica campestris* L. and *B. juncea* (L.) Czern & Coss under Na^+ Stress. *J Int Plant Biol* 52: 549-555.

- Bandurska H., (2001). Proline accumulation during hardening and its involvement in reducing membrane injuries in leaves subjected to severe osmotic stress. *Acta Physiol Plant* 23:483–490.
- Bates L.S., Waldren R.P., Teare I.D., (1973). Rapid determination of free proline for water-stress studies. *Plant Soil* 39:205–207.
- Bernstein N., Silk W.K., Lauchli A., (1993). Growth and development of sorghum leaves under conditions of NaCl stress. *Planta* 191: 433–439.
- Braam J., (1992). Regulated expression of the calmodulin-related TCH genes in cultured arabidopsis cells: induction by calcium and heat shock. *Proc Natl Acad Sci* 89: 3213–3216.
- Bazakos C., Manioudaki M. E., Sarropoulou E., Spano T., Kalaitzis P., (2015). 454 pyrosequencing of olive (*Olea europaea* L.) transcriptome in response to salinity. *PLoS one* 10(11): 1–22.
- Ben-ahmed C., Ben-rouina B., Habib-ur-rehman A., Boukhriss M., (2006). Olive tree (*Olea europaea* L. cv. "chemlali") under salt stress: water relations and ions content. *Pak J Bot* 38(5): 1477–1484.
- Chakrabarti N., Mukherji S., (2002). Effect of phytohormones pretreatment on metabolic changes in *Vigna radiata* under salt stress. *J Env Biol* 23:295–300.
- Chartzoulakis K., Loupassaki M., Bertaki M., Androulakis I., (2002). Effects of NaCl salinity on growth, ion content and CO₂ assimilation rate of six olive cultivars. *Scientia Hort* 96: 235–247.
- Chartzoulakis K., Psarras G., Vemmos S., Loupassaki M., Bertaki M., (2006). Response of two olive cultivars to salt stress and potassium supplement. *J Plant Nut* 29:2063–2078.
- Cramer G.R., (2002). Differential effects of salinity on leaf elongation kinetics of three grass species. *Plant Soil* 253: 233–244.
- Diatloff E., Rengel Z., (2001). Compilation of simple spectrophotometric techniques for the determination of elements in nutrition solutions. *J Plant Nutr* 24:75–85.
- Demiral M.A., (2005). Comparative response of two olive (*Olea europaea* L.) cultivars to salinity. *Turk J Agric For* 29:267–274.
- Demiral M.A., Aktařuygun D., Uygun M., Kasiręa E., Karagozler A.A., (2011). Biochemical response of *Olea europaea* cv. Gemlik to short-term salt stress. *Turk J Biol* 35:433–442.
- Dichio B., Romano M., Nuzzo V., Xiloyannis C., (2002). Soil water availability and relationship between canopy and roots in young olive trees (cv coratina). *Acta Hort* 586: 255–258.
- Egamberdieva D., (2009). Alleviation of salt stress by plant growth regulators and IAA producing bacteria in wheat. *Acta Physiol Plant* 31: 861–864.
- Flowers T.J., Yeo A.R., (1988). Ion relations of salt tolerance. In: Baker, D., Halls, J. (Eds.), *Solute Transport in Plant Cells and Tissue*. Longman, Harlow, pp 392–414.
- Flowers T.J., Yeo A.R., (1995). Breeding for salinity resistance in crop plants: Where next? *Aust J Plant Physiol* 22:857–884.
- Flowers T. J., (1999). Salinization and horticultural production. *Sci Hort* 78:1–4.
- Fernandez J.E., Moreno F., Giron I.F., and Blazquez O.M., (1997). Stomata control of water use in olive tree leaves. *Plant and Soil* 190: 179–192.
- Greenway H., Munns R., (1980). Mechanism of salt tolerance in nonhalophyte. *Ann Rev Plant Physiol* 31: 149–190.
- Gul B., Khan M.A., Weber D.J., (2000). Alleviation salinity and dark enforced dormancy in *Allenrolfea occidentalis* seeds under various thermo periods. *Aust J Bot* 48:745–752.
- Hasegawa P. M., Bressan R. A., Zhu J. K., and Bohnert H. J., (2000). Plant cellular and molecular responses to high salinity. *Ann Rev Plant Physiol Plant Mol Biol* 51:463–475.
- Kasiręa E., Demiral M. A., (2016). Salt stress-mineral nutrient relations in olive (*Olea europaea* L.) plant. *Eur J Soil Sci* 5(4): 307.
- Kchaou H., Larbi A., Gargouri K., Chaieb M., Morales F., Msallem M., (2010). Assessment of tolerance to NaCl salinity of five olive cultivars based on growth characteristics and Na⁺ and Cl⁻ exclusion mechanisms. *Scientia Hort* 124(3): 306–315.
- Khan N.A., (2003). NaCl inhibited chlorophyll synthesis and associated changes in ethylene evolution and antioxidative enzyme activities in wheat. *Biol Plant* 47: 437–440.
- Khan M.A., Gul B., Weber D.J., (2004). Action of plant growth regulators and salinity on seed germination of *Ceratoides lanata*. *Can J Bot* 82: 37–42.
- Khan M.N., Siddiqui M.H., Mohammad F., Naeem M., Khan M.M.A., (2010). Calcium chloride and gibberellic acid protect linseed (*Linum usitatissimum* L.) from NaCl stress by inducing antioxidative defense system and osmo protectant accumulation. *Acta Physiol Plant* 32:121–132.
- Loupassaki M.H., Chartzoulakis K. S., Dagalaki N. B., Androulakis I. I., (2002). Effects of salt stress on concentration of nitrogen, phosphorus, calcium, magnesium and sodium in leaves, shoots and roots of six olive cultivars. *J Plant Nut* 25:2457–2482.
- McAinsh M.R., Clayton H., Mansfield T.A., Hetherington A.M., (1996). Changes in stomatal behavior and guard cell cytosolic free calcium in response to oxidative stress. *Plant Physiol* 111: 1031–1042.
- Maggio A., Barbieri G., Raimondi G., Pascale S.D., (2010). Contrasting effects of GA₃ treatments on tomato plants exposed to increasing salinity. *J Plant Growth Regul* 29: 63–72.
- Mass E.V., Hoffman G.J., (1977). Crop salt tolerance—Current assessment. *J Irrigation Drainage* 103:115–134.
- Munns R., (1993). Physiological processes limiting plant growth in saline soils: some dogmas and hypotheses. *Plant Cell Environ* 16: 15–24.
- Munns R., (2002). Comparative physiology of salt and water stress. *Plant cell Environ* 25: 239–250.
- Matysik J., Alia B.B., Mohanty P., (2002). Molecular mechanisms of quenching of reactive oxygen species by proline under stress in plants. *Curr Sci* 82:525–532.
- Raza S.H., Athar H. R., Ashraf M., Hameed A., (2007). GB-induced modulation of antioxidant enzymes activities and ion accumulation in two wheat cultivars differing in salt tolerance. *Env Exp Bot* 60(3): 368–376.
- Saeidi-Sar S., Khavari-Nejad R., Fahimi H., Ghorbanli M., Majd A., (2007). Interactive effects of gibberellin GA₃ and ascorbic acid on lipid peroxidation and antioxidant enzyme activities in *Glycine max* seedlings under nickel stress. *Russ J Plant Physiol* 54:74–79.
- Shah S.H., (2007). Effects of salt stress on mustard as affected by gibberellic acid application. *Gen Appl Plant Physiol* 33: 97–106.
- Shaheena A., Firoz M., Shamsul H., Manzer H., (2005). Exogenous Application of gibberellic Acid counteracts the III effect of sodium chloride in mustard. *Turk J Biol* 29: 233–236.
- Singh G., Jain S., (1981). Effect of some growth regulators on certain biochemical parameters during seed development in chickpea under salinity. *Indian J Plant Physiol* 20: 167–179.
- Siddiqui M.H., Khan M.N., Mohammad F., Khan M.M.A., (2008). Role of nitrogen and gibberellins (GA₃) in the regulation of enzyme activities and in osmo protectant accumulation in *Brassica juncea* L. under salt stress. *J Agron Crop Sci* 194:214–224.
- Storey R., Walker R. R., (1999). Citrus and salinity. *Scientia Hort* 78:39–81.
- Tester M., Devenport R., (2003). Na⁺ tolerance Na⁺ transport in higher plants. *Ann Bot* 91: 503–527.
- Tattini M. (1994). Ionic relations of aeroponically-grown olive genotypes during salt stress. *Plant Soil* 161: 251–256.
- Therios I. N., Misopolinos N. D., (1988). Genotypic response to sodium chloride salinity of four major olive cultivars (*Olea europea* L.). *Plant Soil* 106: 105–111.
- Tabatabaei S.J., (2006). Effects of salinity and N on the growth, photosynthesis and N status of olive (*Olea europaea* L.) trees. *Scientia Hort* 108:432–438.
- White P.J., Broadley M.R., (2001). Chloride in soils and its uptake and movement within the plant: A review. *Ann Bot* 88:967–988.