

# Carbon Dynamic after Conversion of Permanent Grassland into Arable Soil

---

Lubica POSPÍŠILOVÁ <sup>1</sup>✉

Magdalena HÁBOVÁ <sup>1</sup>

Vítězslav VLČEK <sup>1</sup>

Jiří JANDÁK <sup>1</sup>

Ladislav MENŠÍK <sup>2</sup>

Gabriela BARANČÍKOVÁ <sup>3</sup>

---

## Summary

Carbon input and balance in soils is regarded as the main criterion of agricultural sustainability. Generally, carbon dynamic depends not only on the carbon input and its decomposition rate, but it is also influenced by various agronomic practices. Therefore, changes in organic carbon stock and humic substances quality were evaluated in two different agricultural management systems (permanent grassland and intensive crop sequences). Haplic Cambisol (Czech-Moravian Highland, locality Vatin, Czech Republic) was sampled twice a year (spring and autumn) in the depth 0-20 cm during the period 2010-2016. Soil was sandy-loam textured, with middle organic carbon content and very low humic substances quality. Results showed that crop management practices directly influenced soil cumulative potential, quality of humic substances, soil reaction and amount of nutrients. Statistically significant differences were found.

---

## Key words

carbon forms, arable soil, permanent grassland

<sup>1</sup> Faculty of AgriScience Mendel University in Brno, Department of Agro chemistry, Soil Science, Microbiology and Plant Nutrition, Zemědělská 1, 613 00 Brno, Czech Republic

✉ e-mail: [lubica.pospisilova@mendelu.cz](mailto:lubica.pospisilova@mendelu.cz)

<sup>2</sup> Crop Research Institute, Division of Crop Management Systems, Drnovská 507/73, 161 06 Praha 6 - Ruzyně, Czech Republic

<sup>3</sup> National Agriculture and Food Centre, SSCRI Bratislava, External working place Prešov, Raymannova 1, 080 01 Prešov, Slovakia

Received: June 23, 2017 · Accepted: September 20, 2017

## ACKNOWLEDGEMENTS

Financial support from the National Agricultural Agency of the project QJ1210263, QJ1510179, MZE RO0417, OP VaVpl CZ.1.05/4.1.00/04.0135 project Mendel University in Brno, Czech Republic and the Slovak Research and Development Agency (APVV-0243-11 and APVV-14-0087) is highly acknowledged.

## Introduction

Soil carbon, specifically in the form of soil organic matter, plays a key role in the functioning of soils to produce a wide range of vital environmental functions. Loss of soil carbon degrades these services, resulting in decreased crop yields, environmental and market value of soil. Land management can also enhance soil carbon by optimal crop rotation and fallow cover crops, organic matter addition, and optimal fertilization systems tillage systems (Banwart et al., 2015). Soil organic carbon forms could be divided into stable and labile. Stable carbon forms are represented by humic acids (HA), fulvic acids (FA) and humins. Because of relatively long period of time that stable carbon spends with the soil and is thereby withheld from the atmosphere, it is often referred to as being sequestered. Carbon sequestration depends on soil capacity to store resistant plant residues and protects and accumulates the stable humic substances (HS). The degree of resistance of HA and FA is attributed to their chemical composition, especially the absence of labile (hydrolysable) linkages, and to the lack of regularly repeating units. Humin is generally considered to be a component of HS although some of its units may not have undergone the biological transformation (Swift, 1999 and 2001). Labile carbon forms are mainly introduced by dissolved organic carbon (DOC) and by carbon of microbial biomass ( $C_{mic}$ ) as quoted Leinweber et al. (1995) and Degens and Sparling (1996). DOC is supposed to be very active in carbon pool and its chemical composition and mobility (the smaller and more polar an organic molecule) directly affect biological soil properties. Relationship between DOC and soil was studied by Zsolnay (2003) and Pospíšilová et al. (2011). They concluded that DOC content is controlled by its content in soil. Instead of complicated determination of DOC determination of water extractable organic carbon (WEOC) is frequently used as quoted Šestauberová and Novák (2011). Stability or lability of any organic carbon fraction could be due to either chemical composition or protection within the soil aggregates. It is supposed that chemical properties of HS are integrally linked to the soil's behavior in nutrient cycling and supply, water retention, soil structure and soil biological parameters (Filep and Rekasí, 2011).

Many different mathematical models were proposed to characterize soil organic carbon stock (Coleman and Jenkinson, 2005; Smith et al., 2005; 2007; De Li Liu et al., 2016). They showed that the amount of soil organic carbon that is attained under agriculture largely depends upon the carbon input and its decomposition rate under various agronomic practices. Whether soils are a sink or source of carbon depends on the actual organic carbon stock, agricultural practices over time, soil properties (e.g. clay content, soil depth, content and quality of plant input, organic fertilizer input etc.), and climatic conditions.

The aim of our research was to evaluate dynamic of stable and labile carbon forms and humic substances quality in two different agricultural management systems (permanent grassland and intensive crop sequences) during the period 2010-2016.

## Material and methods

Haplic Cambisol (locality Vatín, Czech-Moravian Upland, 530 m a.s.l.) belongs to the potatoes growing area. Average annual temperature is 6.9 °C. Average annual precipitation 621 mm. Soil was sampled twice a year (spring and autumn) in 0-20 cm

depth. Native grassland (*Sanguisorba-Festucetum comutatae*) was ploughed and intensive crop sequences involved 50 % of cereals, 16.6 % of root crops, and 33.4 % of technical crops. Nutrients were applied at ratios (N-P-K, kg/ha/year): 130-40-80 (winter wheat) and 60-35-80 (spring barley). Soil reaction was determined by potentiometric method in distilled water and in 1M KCl solution (1:2.5). Particle size analysis was determined by pipette method (Zbiral et al., 2010). Total organic carbon content (TOC) was determined by oxidimetric titration method (Nelson and Sommers, 1982). Fractional composition of HS was determined according to Kononova and Beltchikova method (1963) as follows: 5g of air dried soil sample, sieved at mesh size of 1mm and extracted by a mixture (1:1; 0.1M NaOH + 0.1M  $Na_4P_2O_7$ ) for 24h. The sediment was separated by centrifugation at 2800g for 10min, washed with mixture and centrifuge again. Two individual washings were unified with original supernatant. Sum of HS and HA was determined by oxidimetric titration method in aliquot volumes. FA content was calculated as a difference between HS and HA. WEOC was determined as follows: 1 g of soil sample was suspended into 100 ml demineralized distilled water and shaking for 30 min. Then the samples were centrifuged (15 min at 3000 rpm) and filtrated through membranes filter S-PAK™ Millipore (0.45 μm). Content of WEOC was determined according to ČSN EN 1484 and ČSN EN 12260 using high temperature (720 °C) catalytic combustion with chemo-luminescent detection of flue gases (carbon and oxygen) and spectrometric detection of CO<sub>2</sub> in infrared spectral region using analyser Shimadzu TOC-VCSH with auto sampler ASI-5000 (Shimadzu, Kyoto, Japan).  $C_{mic}$  carbon was determined by fumigation-extraction method in fresh soil samples, sieved through 5 mm according to Vance et al. (1987). Statistical analysis, including graphical outputs, was carried out using STATISTICA 12.0 (Stat-Soft Inc., Tulsa USA, StatSoft ČR s.r.o., 2014). Statistical data processing Exploratory Data Analysis /EDA/, ANOVA - Tukey test /HSD-test/ a Fisher test /LSD test/, Principal Component Analysis (PCA) for data evaluation was applied (Meloun and Militký, 2011). PCA was used for interpreting the parameters of soil organic matter (TOC, HS, HA, FA etc.) and physico-chemical properties of soil (pH, content of phosphorus and potassium, etc.). Selected measured characteristics were used as predictors (factors); they were chosen on the basis of an eigenvalue graph. Variables with impaired assumption of normality were converted using logarithmic transformation. As part of step 1, PCA was carried out with all the variables to compute the most important variables. Step 2 involved selecting active and supplementary variables for better interpretation. This stepwise analysis significantly improves the outcome of the PCA analysis in case of a smaller number of samples. PCA was used for calculating a component weight for the investigated variables (Meloun and Militký, 2011). Based on correlations and contributions in convincing factors each of the characteristics was subsequently judged for relevance to explain the multidimensional dependencies (correlations) in the factorial plane. Statistical significance was assessed at a significance level of  $P = 0.05$ .

## Results and discussion

Studied soil was sandy-loam textured, with acid soil reaction, and low cation exchange capacity. Humus content was middle but its quality was very low. Prevalence of FA was determined and HA/FA ratio was less than 1. Humification degree was middle (30-40

Table 1. Statistical evaluation of studied soil parameters in Arable land and Permanent grassland

Characteristic	Site	HSD Test*		LSD Test**	
		Arable land	Permanent grassland	Arable land	Permanent grassland
TOC	Arable land	x	0.000177	x	0.000010
	Permanent grassland	0.000177	x	0.000010	x
HS	Arable land	x	0.000172	x	0.000000
	Permanent grassland	0.000172	x	0.000000	x
HA	Arable land	x	0.004722	x	0.004559
	Permanent grassland	0.004722	x	0.004559	x
FA	Arable land	x	0.000218	x	0.000060
	Permanent grassland	0.000218	x	0.000060	x
HA/FA ratio	Arable land	x	0.323781	x	0.323656
	Permanent grassland	0.323781	x	0.323656	x
DH I	Arable land	x	0.602599	x	0.602452
	Permanent grassland	0.602599	x	0.602452	x
DH II	Arable land	x	0.261271	x	0.261137
	Permanent grassland	0.261271	x	0.261137	x
pH	Arable land	x	0.000262	x	0.000108
	Permanent grassland	0.000262	x	0.000108	x
C <sub>we</sub>	Arable land	x	0.003794	x	0.003638
	Permanent grassland	0.003794	x	0.003638	x
C <sub>mic</sub>	Arable land	x	0.002175	x	0.002034
	Permanent grassland	0.002175	x	0.002034	x
K	Arable land	x	0.000172	x	0.000000
	Permanent grassland	0.000172	x	0.000000	x
P	Arable land	x	0.000172	x	0.000001
	Permanent grassland	0.000172	x	0.000001	x
Ca	Arable land	x	0.051929	x	0.051790
	Permanent grassland	0.051929	x	0.051790	x
Mg	Arable land	x	0.002702	x	0.002546
	Permanent grassland	0.002702	x	0.002546	x

Note: n = 7; \* Tukey test; \*\* Fisher test; P = 0,05

%). Statistical evaluation of studied soil parameters is showed in Table 1. Comparison of different carbon forms in arable soil and permanent grassland showed higher content of stable carbon form (TOC, HS, HA, FA) in grassland (Figure 1). Also labile carbon forms (WEOC, C<sub>mic</sub>) content was higher in grassland to compare with arable soil (Figure 1). On the other hand, amount of nutrients and more favourable soil reaction was reached in arable soil. Permanent grassland was more acid and had lower nutrients content. Our observation also showed that in spite of lower content of HS, HA, and FA in arable soil we reached more favourable HA/FA ratio here soil to compare with grassland. We can say that in spite of lower content of HS their quality is higher than quality of HS in permanent grassland. Similar results were published by Tesařová et al. (2006) and Petrářová et al. (2009). They stressed that decreasing of total carbon content is evident mainly the first two-three years after ploughing and then the new, and lower equilibrium of total carbon content in arable soil took place. This effect is generally called priming effect. Further the ordination diagram (Figure 2) is showing the results of PCA analysis. Two different categories (permanent grassland and arable soil) are evident. Statistically significant separation into two categories according to balanced TOC, potassium and phosphorus content and the soil pH is evident. Total variability (PC1, PC2 and PC3) of studied parameters is higher than 94 %. On axis PC1 potassium content is in strong

positive relationship with phosphorus content ( $r=0.90$ ). Medium positive relationship ( $r=0.80-0.85$ ) between soil reaction and potassium and phosphorus content was determined. We can also conclude that on the axis PC1, TOC is in positive relationship with FA ( $r=0.74$ ) and HA ( $r=0.68$ ) – see Figure 2. No significant correlation on PC2 axis was found. The PC2 axis lacks considerable correlation; rather, directions are characterised, depending on HA/FA ratio (Figure 2). In conclude it should be stress that differences in quantity and quality of soil organic matter are directly influence by crop sequence and plant species, fertilizing system (mineral and organic), tillage system and by intensity of agricultural exploitation as well. Because of depending of labile carbon forms onto total organic carbon amount regulation of carbon balance in sustainable agronomical practices is possible due to appropriate agricultural management system.

### Conclusion

Land use and management practices are directly influence stabile and labile carbon forms in soil. Very high carbon accumulation potential was confirmed under permanent grassland. Accumulation potential of arable soil is depending on intensity of agriculture and management system with respect to soil type and climatic conditions.




Figure 1. Content of different carbon forms and degree of humification in Arable land and Permanent grassland




Figure 2. Multivariate statistical PCA of studied soil parameters in Arable land and Permanent grassland

## References

- Banwart A. S., Noellemeyer E., Milne E. (2015): The Global Challenge for Soil Carbon. Science, Management and Policy for Multiple benefits SCOPE Series volume 71: 1-10.
- Coleman K., Jenkinson D.S. (2005): ROTHC-26.3. A model for the turnover of carbon in soil. Model description and windows users' guide, November 1999 issue (modified April, 2005), p. 45. <http://www.rothamsted.bbsrc.ac.uk/aen/carbon.pdf>
- ČSN ISO 10694 - 83 6410 (1998): Kvalita půdy - Stanovení organického a celkového uhlíku po termickém rozkladu. Soil quality - Determination of organic carbon and total carbon after thermal decomposition. (In Czech).
- ČSN EN 12260 - 757524 (2004): Jakost vod - Stanovení dusíku - Stanovení vázaného dusíku (TNb) po oxidaci na oxidy dusíku. Water quality - Determination of nitrogen - Determination of bound nitrogen (TNb) after oxidation to oxide. (In Czech).
- ČSN EN 1484 (757515). 1998. Jakost vod - Stanovení celkového organického uhlíku (TOC) a rozpuštěného organického uhlíku (DOC). Water quality - Determination of total organic carbon (TOC) and dissolved organic carbon (DOC). (In Czech).
- Degens B., Sparling G. (1996): Changes in aggregation do not correspond with changes in labile organic „C” fractions in soil amended with 14C glucose. *Soil Biology and Biochemistry* 28: 453-462.
- De Liu L., O'Leary G.J., Ma Y., Cowie A., Li F.Y., Mccaskill M., Conyer M., Dalal R., Robertson F., Dougherty, W. (2016): Modelling soil organic carbon 2. Changes under a range of cropping and grazing farming systems in eastern Australia. *Geoderma* 265: 164-175.
- Filep T., Rekas M. (2011): Factors controlling dissolved organic carbon (DOC), dissolved organic nitrogen (DON), and DOC/DON ratio in arable soils based on a dataset from Hungary. *Geoderma* 162 (1-4): 312-318.
- Kononova M. M., Belchikova N.P. (1963): Organiceskoje vescestvo povcy (Soil organic matter). Moscow, AN SSSR, 228-234. (In Russian).
- LECO TruSpec CN. TruSpec CN Carbon/Nitrogen Determinator. Instruction manual. LECO Corporation Michigan (2006).
- Leinweber P., Schulten H.R., Korschens M. (1995): Hot water extracted organic matter: Chemical composition and temporal variations in a long-term field experiment. *Biology and Fertility of Soils* 20: 17-23.
- Meloun M., Militký J. (2011): Statistical Data Analysis. A Practical Guide with 1250 Exercises and Answer key on CD, Woodhead Publishing India, 1600 pages, ISBN: 978-93-80308-11-1.
- Petrášová V., Martinec J., Pospíšilová L. (2009): Total carbon content and humic substances quality in selected subtypes of Cambisols. *Acta universitatis agriculturae et silviculturae Mendelianae Brunensis* vol. 4: 73-81.
- Pospíšilová L., Škarpa P., Konečná M. (2011): Different carbon fractions in soils and their relationship with trace elements content. *Journal of Life Sciences* 5 (4): 316-321.
- Smith J., Smith P., Wattenbach M., Zaehle S., Hiederer R., Jones R.J.A., Montanarella L., Rounsevell M., Reginster I., Ewert, F. (2005): Projected changes in mineral soil carbon of European croplands and grasslands, 1990-2080. *Global Change Biolog.* Online: 21-Nov-2005. DOI: 10.1111/j.1365-2486.2005.001075x.
- Smith J., Smith P., Wattenbach M., Gottschalk P., Romanenkov V.A., Sevcova L.K., Sirotenko O.D., Rukhovic D.I., Korolova P.V., Romanenko I.A., Lisovoj, N.V. (2007): Projected changes in the organic carbon stocks of cropland mineral soils of European Russia and the Ukraine 1990-2070. *Global Change Biol.* 13 (2): 342-354.
- Swift R. S. (1999): Macromolecular properties of soil humic substances: fact, fiction, and opinion. *Soil Sci.* 164: 790-802.
- Swift R. S. (2001): Sequestration of carbon by soil. *Soil Sci.* 166 (11): 858-871.
- Šestauberová M., Novák F. (2011): Comparison of extractable soil carbon and dissolved organic carbon by their molecular characteristics. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* vol. LIX, 41(6): 337-342.
- Tesařová M., Kudlička P., Pospíšilová L., Kalhotka L., Hrabě F. (2006): Comparison of mineralisation and humification of postharvest residues of cereals in conventional and organic cropping practices. *Acta universitatis agriculturae et silviculturae Mendelianae Brunensis* vol. LIV (1):121-126.
- Zbíral J., Honsa I., Malý S. (2010): Jednotné pracovní postupy: Analýza půd I. Brno, ÚKZÚZ. Unified analytical procedures: Soil analysis I. ISBN: 978-80-7401-031-6. (In Czech).
- Zsolnay A. (2003): Dissolved organic matter: artefacts, definitions, and functions. *Geoderma* 113: 187-209.

acs83\_o4