

Original article

DOI: 10.2478/aiht-2018-69-3160

Removing aflatoxin M₁ from milk with native lactic acid bacteria, centrifugation, and filtration

Željka Kuharić¹, Željko Jakopović², Iva Čanak², Jadranka Frece², Jasna Bošnjir¹, Željka Pavlek¹,
Martina Ivešić¹, and Ksenija Markov²

Andrija Štampar Teaching Institute for Public Health¹, University of Zagreb, Faculty of Food Technology and Biotechnology², Zagreb, Croatia

[Received in June 2018; Similarity Check in July 2018; Accepted in November 2018]

In order to minimise human exposure to aflatoxin M₁ (AFM₁) the levels of this highly carcinogenic mycotoxin in milk, heat-treated milk, and other dairy products have been limited to <0.05 µg kg⁻¹. However, its removal from dairy products presents a challenge for dairy producers, as commercial additives change organoleptic properties, and filtration alone yields poor results. The aim of this study was to find a strain of lactic acid bacteria (LAB) from milk or dairy products that most effectively binds AFM₁ and to see whether heat treatment of the selected LAB affects the binding efficiency. We also wanted to investigate whether centrifugation can improve filtering of the obtained AFM₁-LAB complexes from milk. To do that, we isolated and identified 10 native LAB species/strains, incubated their viable or heat-treated cells (10⁸ CFU mL⁻¹) in milk spiked with 0.5 µg L⁻¹ of AFM₁ at 4 °C for 0, 2, 4, and 24 h, and quantified the amount of unbound AFM₁ with HPLC. AFM₁ binding efficiency ranged from 21 to 92 % for viable cells and from 26 to 94 % for the heat-treated ones. Since both viable and heat-treated *Lactobacillus plantarum* KM showed the best results, we used it for the next step in AFM₁ removal from milk. Heat treatment in combination with filtration and centrifugation yielded removal as high as 96 %.

KEY WORDS: HPLC, lactic acid bacteria, *L. plantarum* KM, secondary fungal metabolites, viable and heat-treated cells

Although grain and cereal-based products are the main source of mycotoxins for humans, mycotoxins can enter human body through meat, eggs, milk, and dairy products of animals fed with fodder contaminated with mycotoxins or moulds that synthesise them (*carryover* effect) (1-5). Aflatoxin M₁ (AFM₁), a highly carcinogenic mycotoxin, is a product of aflatoxin B₁ (AFB₁) metabolism in the liver (6). It is excreted in the tissues, biological fluids, and milk of lactating animals, including humans (7, 8). Its content in milk correlates with the AFB₁ level in feed (9). Once it enters milk, AFM₁ cannot be removed by heat treatment such as pasteurisation or sterilisation (10). Instead, it is carried over to dairy products, including milk powder, cheese, yogurt, and butter (11).

For this reason, the EU countries have limited AFM₁ levels in raw milk, heat-treated milk, and milk for the production of dairy products to <0.05 µg kg⁻¹ (12, 13). Milk with AFM₁ levels above this limit should not be used for human consumption or dairy production nor should it be mixed with uncontaminated milk to bring AFM₁ level down to acceptable.

This limit, of course, creates serious problems for dairies who cannot control AFM₁ levels, as their products become

unmarketable. Physical and chemical methods for removing mycotoxins from milk are limited because of safety issues, quality losses in dairy produce, poor efficiency, and high cost (14). One of the ways to remove mycotoxins from milk is to use additives/mycotoxin deactivation products and intestinal adsorbents such as aluminium silicates, clay, or zeolite. Detoxification additives are efficient in bringing contamination levels below the European limit for AFM₁, while their residues in milk present no concern for human health (3). Intestinal adsorbents, in turn, are used with animal feed to reduce AFB₁ absorption in animal body and prevent further toxin metabolism, distribution, and carryover to milk. However, as these adsorbents alter the nutritional properties of food (3, 15-17), alternative solutions have been proposed, such as the use of microorganisms (18), lactic acid bacteria (LAB) in particular. LAB showed success, albeit varying, in binding AFM₁ in phosphate buffer and milk solutions (11, 19-24).

Problems arise when commercial starter cultures that bind AFM₁ compete with native microflora and change traditional organoleptic properties of the fermented product. Another problem is the removal of LAB-AFM₁ complex, since the data about mycotoxin complex removal are still scarce.

The aim of our study was to determine the binding parameters of AFM₁ in milk by native LAB (viable and

Correspondence to: Željko Jakopović, University of Zagreb, Faculty of Food Technology and Biotechnology, Pierottijeva 6, 10000 Zagreb, Croatia, E-mail: zjakopovic@pbf.hr

heat-treated cells) that would not affect the organoleptic properties of a product and also determine the LAB-AFM₁ complex removal efficiency with the membrane filtration method.

MATERIALS AND METHODS

Isolation and identification of LAB

To obtain as many native LAB strains as possible for further analysis, we isolated and identified them from fresh cow milk, cottage cheese, and cream of randomly selected dairy producers. Milk and dairy samples (N=5 of each) were prepared for microbiological testing according to the standard procedure (25) by serially diluting 10 mL samples of milk and cream or 10 g of cottage cheese (n=5) homogenised in 90 mL of sterile buffered solution, until we obtained 0.1 mL samples that were inoculated on MRS agar (Biolife, Milano, Italy) and incubated at 30 °C for 48 h.

Lactic acid bacteria were isolated from the milk, cream, and cheese samples following the ISO 15214:1998 procedure (26). We randomly took 150 LAB colonies cultivated on MRS agar and identified them based on colony and cell morphology and Gram staining. Identification was confirmed with the API 50 CHL biochemical test kit (BioMérieux, Marcy l'Etoile, France) (27, 28) following the manufacturer's instructions. The isolated LAB strains were stored in MRS broth (Biolife) containing 30 % glycerol (in volume) at -70 °C until further analysis.

Standard AFM₁

The working solution of AFM₁ was prepared by diluting the standard 0.5 µg mL⁻¹ solution in acetonitrile (LGC Promochem, Leeds, UK) to 0.05 µg mL⁻¹ and storing it at 4 °C until further analysis.

Determination of AFM₁ to LAB binding in milk

To determine AFM₁ binding to LAB in milk, the LAB were grown in MRS broth at 30 °C for 48 h. For each bacterial strain the cells were harvested by centrifugation (3000×g for 10 min) under aseptic conditions at room temperature, washed three times with 5 mL of sterile deionised water, resuspended in sterile deionised water, and divided in two groups. The first group contained viable cells and the second cells heated in a water bath at 100 °C for 60 min (heat-treated cells). Heat-treated cells have already been reported to bind AFM₁ (11, 19, 20, 24).

Before the experiment began, we verified that the sterile milk samples were not contaminated with AFM₁. Two hundred millilitres of milk were then inoculated with a suspension of either viable or heat-treated LAB cells (10⁸ CFU mL⁻¹) and 0.5 µg AFM₁ per litre of milk. The samples were then incubated at 4 °C to see if the LAB strains could reduce AFM₁ levels in milk at refrigeration temperature. AFM₁ binding was determined with high-

performance liquid chromatography (HPLC, see detailed description below) at 10–15 min after the sample was homogenised (0 h) and then at 2, 4, and 24 h of incubation by analysing the amount of unbound AFM₁ in the supernatant obtained by centrifuging the samples at 3000×g for 20 min.

Determining AFM₁ residues after filtration

To find the most efficient filtration method for the removal of AFM₁ from milk – alone or bound to viable or heat-treated *L. plantarum* KM cells, which showed the best binding results in the previous test (see below) – we tested two methods: the first involved centrifuging and filtering 50 mL milk samples and the second only filtering without centrifugation. The centrifugation parameters were 6000×g for 30 min followed by vacuum filtration on 0.45 µm pore size cellulose-nitrate filters (Sartorius Stedim Biotech GmbH, Goettingen, Germany) at a flow rate of 0.5 mL min⁻¹. Ten to 15 min after centrifugation and/or filtering, 15 mL filtrates were collected and AFM₁ was quantified using HPLC.

HPLC analysis of residual AFM₁

Before HPLC analysis, the samples (25 mL) were purified on immunoaffinity columns (VICAM, Milford, MA, USA) conditioned with 10 mL phosphate buffer saline (PBS). Sample flow rate did not exceed 3 mL min⁻¹. The immunoaffinity columns were then washed with 10 mL of distilled water and dried under vacuum. AFM₁ was eluted with 4 mL of acetonitrile. The eluate was collected in a tube and evaporated to dryness in a stream of nitrogen.

The concentration of residual AFM₁ in the samples was determined on an Agilent 1100 HPLC System (Agilent, Santa Clara, CA, USA) equipped with a CSI-6150 online vacuum column degasser (Cambridge Scientific Instruments, Witchford, Ely, England), an Agilent isoquantine pump (model G 1310 A9), and a fluorescent detector (Agilent 1100, FLD G1321A) with wavelength excitation and emission of 365 nm and 455 nm, respectively. Separation was carried out on a Synergi Polar column (150x3.0 mm, 4 µm) (Agilent). For the mobile phase we used water/acetonitrile (60/40, v/v) at a flow rate of 0.5 mL min⁻¹ (29).

HPLC method validation

For validation we used a standard of AFM₁ in acetonitrile (AFM₁ solution), milk free of AFM₁ (AFM₁-free milk), and milk spiked with AFM₁ (AFM₁-spiked milk). Linearity was tested by injecting seven concentrations of the working standard. Each concentration was injected three times, and the regression line and correlation coefficient determined.

The linear equation was:

$$y=8.788x-1.193$$

where y stands for area and x for concentration.

The correlation coefficient was 0.999. The selectivity of the method was determined by comparing the chromatograms of the standard AFM₁ solution, AFM₁-spiked milk, and AFM₁-free milk. The limit of detection (LOD) of AFM₁ in milk was 0.005 µg kg⁻¹, and the limit of quantitation (LOQ) 0.010 µg kg⁻¹. The recoveries of spiked AFM₁ in milk at different levels show that this method is reliable for the determining AFM₁ in milk (Table 1).

Statistical analysis

All experiments were carried out in triplicate. For statistical analysis we used STATISTICA v. 7.1 for Windows 10.0 (StatSoft, Inc, Tulsa, OK, USA). We compared mean percentages of bound AFM₁ between viable and heat-treated cells for each strain using the two-way analysis of variance (ANOVA) and set the level of significance at p<0.05.

Table 1 HPLC validation of AFM₁ recoveries

Analyte	Spiking (µg kg ⁻¹)	Recovery range (%)	RSD (%)
AFM ₁	0.01	92–96.5	2.2
	0.05	91.5–95.4	
	0.10	91.9–94.3	

RESULTS AND DISCUSSION

AFM₁ binding by LAB

Among the 150 isolated colonies the biochemical API tests identified five LAB species of two genera (*Lactobacillus* 83.3 % and *Lactococcus* 16.7 %) with 99–99.9 % accuracy (Table 2). The predominant species in cow milk was *L. plantarum* (51.9 %), while *Lactobacillus helveticus* and *Lactobacillus paracasei* dominated in the cottage cheese and cream (51.1 % and 45.3 %, respectively).

The results were expressed as the percentage of AFM₁ bound in the LAB+AFM₁ complex to see which LAB species/strain, viable or heat-treated, was the most effective in AFM₁ binding. Every measurement included positive control (only AFM₁ in milk).

Table 2 Native LAB species in cow milk, cottage cheese, and cream

Sample	N	Identified LAB	N isolates/N samples
Fresh cow milk	5	<i>Lactobacillus plantarum</i>	27/5
		<i>Lactobacillus rhamnosus</i>	18/5
		<i>Lactococcus lactis</i>	7/3
Cow cheese	5	<i>Lactobacillus plantarum</i>	5/5
		<i>Lactococcus lactis</i>	7/5
		<i>Lactobacillus helveticus</i>	23/5
Cream	5	<i>Lactobacillus rhamnosus</i>	10/2
		<i>Lactobacillus plantarum</i>	18/5
		<i>Lactobacillus paracasei</i>	24/5
		<i>Lactococcus lactis</i>	11/2

Table 3 shows AFM₁ binding efficiency of the identified native LAB species. The most efficient was *L. plantarum* KM at 4 h of incubation, heat-treated or not. Heat treatment, however, significantly improved its efficiency (from 92.7 % to 94.5 %, p<0.05).

These findings are consistent with the ones reported by Kabak and Var (11), who also observed a rise in AFM₁ binding between viable (7.9–25.9 %) and heat-treated cells (12.9–27.3 %) of *Lactobacillus* and *Bifidobacterium* after 4 h of incubation. Pierides et al. (24) also found significantly improved AFM₁ removal by *Lactobacillus* strains when they were heat-treated.

All these findings, including ours, suggest that LAB do not need to be alive to bind AFM₁. Although the mechanism of action of these microorganisms on aflatoxin has not yet been clarified, some authors suggest that AFM₁ molecules physically attach to bacterial cell components, mainly polysaccharides and peptidoglycans, instead of creating covalent bonds with them or being metabolised by them (30, 31). Furthermore, heat treatment seems to increase the attachment surface for AFM₁ by making available not only the LAB cell wall but also the plasma components, as the cell wall collapses (32).

Our findings have also shown that AFM₁ binding is reversible; all the isolated LAB released some of the bound AFM₁ as incubation continued. In fact, viable *L. plantarum* KM cells released back a great percentage of AFM₁ after 24 h, which underscores the advantages of heat treatment.

AFM₁ removal by filtration

Figure 1 shows the removal efficiency of AFM₁ from milk by filtration with or without centrifugation. Its removal from AFM₁-spiked milk without LAB by filtration alone was 81.2±0.2 % and in combination with centrifugation it reached 83.3±0.2 %. In contrast, the removal of the viable LAB-AFM₁ complex (obtained with the treatment with viable cells) in combination with centrifugation and filtration was less effective: 78.9±0.3 %. However, removal with combined centrifugation and filtration significantly increased to 95.7±0.2 % for the heat-treated LAB-AFM₁ complex. Of course, this increase has to do with the higher




Figure 1 Comparison of AFM₁ removal efficiency between methods. 1 – AFM₁ removal from spiked milk (0.5 µg L⁻¹ of AFM₁) with filtration alone; 2 – AFM₁ removal from spiked milk (0.5 µg L⁻¹ of AFM₁) with filtration and centrifugation; 3 – removal of the AFM₁ + viable *L. plantarum* KM cell complex with filtration and centrifugation; 4 – removal of the AFM₁ + heat-treated *L. plantarum* KM cell complex with filtration and centrifugation. *statistically significant difference from treatments 1, 2, and 3

AFM₁ binding ability of the heat-treated *L. plantarum* KM strain.

We believe that these findings greatly contribute to the current knowledge about AFM₁ removal from milk. Assaf et al. (19) have only recently studied the retention of AFM₁ on filters and proposed that it is owed to AFM₁ adsorption on the filter membrane. In combination with LAB treatment (*L. rhamnosus* GG) and centrifugation, the retention on filter paper was even greater, which the authors explained with greater contact surface between the LAB and AFM₁. Additionally, centrifugation improved the retention, because AFM₁ was bound to milk components such as fat globules or casein separated from milk by centrifugation and subsequent deposition. AFM₁ preference for milk proteins had been demonstrated earlier by Brackett and Marth (33).

However, centrifugation in combination with viable *L. plantarum* KM cells yielded lower AFM₁ removal from milk. We believe that this is because centrifugation breaks up electrostatic bonds between viable cells and AFM₁.

CONCLUSION

Our study has singled out the most efficient AFM₁ removal procedure using the native LAB in order to avoid organoleptic changes in milk. It consists of refrigeration at 4 °C with heat-treated *L. plantarum* KM and then centrifugation and filtering, which in our experiment ensured over 95 % removal of AFM₁ from contaminated milk.

In practical terms, we believe that this procedure can find wide application among dairy producers, as it does not require special skills or equipment, provided that the native *L. plantarum* KM strain is available.

Our future studies shall focus on better understanding the interaction between bacteria and AFM₁. To do that, it

is necessary to examine multiple LAB strains and other cultures used in the food industry.

Conflicts of interest

None to declare.

Acknowledgments

This work was supported by the Croatian Science Foundation (project no. IP-2016-06-4306). The authors wish to thank Dado Čakalo for his edits and help with the language.

REFERENCES

1. Britzi M, Friedman S, Miron J, Solomon R, Cuneah O, Shimshoni JA, Soback S, Ashkenazi R, Armer S, Shlosberg A. Carry-over of aflatoxin B₁ to aflatoxin M₁ in high yielding israeli cows in mid- and late-lactation. *Toxins* 2013;5:173-83. doi: 10.3390/toxins5010173
2. Di Stefano V, Pitonzo R, Cicero N, D'Oca MC. Mycotoxin contamination of animal feedingstuff: detoxification by gamma-irradiation and reduction of aflatoxins and ochratoxin A concentrations. *Food Addit Contam A* 2014;31:2034-9. doi: 10.1080/19440049.2014.968882
3. Giovati L, Magliani W, Ciociola T, Santinoli C, Conti S, Polonelli L. AFM₁ in milk: physical, biological, and prophylactic methods to mitigate contamination. *Toxins* 2015;7:4330-49. doi: 10.3390/toxins7104330
4. Markov K, Pleadin J, Bevardi M, Vahčić N, Sokolić-Mihalak D, Frece J. Natural occurrence of aflatoxin B₁, ochratoxin A and citrinin in Croatian fermented meat products. *Food Control* 2013;34:312-7. doi: 10.1016/j.foodcont.2013.05.002
5. Völkel I, Schröer-Merker E, Czerny CP. The carry-over of mycotoxins in products of animal origin with special regard to its implications for the European Food Safety Legislation. *Food Nutr Sci* 2011;2:852-67. doi: 10.4236/fns.2011.28117

Table 3 AFM₁ binding efficiency by lactic acid bacteria strains (viable and heat-treated) and incubation time

Treatment -- Strain	% of bound AFM ₁ ±SD							
	0 h		2 h		4 h		24 h	
	Viable	Heat-treated	Viable	Heat-treated	Viable	Heat-treated	Viable	Heat-treated
<i>L. plantarum</i> SM1	80.6±0.1	80.2±0.1	24.9±0.1	76.1±0.2*	79.6±0.2	79.2±0.2	35.9±0.2	26.1±0.1
<i>L. plantarum</i> SMB	91.0±0.1	84.6±0.2*	73.1±0.2	75.7±0.1	82.45±0.2	46.94±0.1*	21.4±0.2	64.1±0.2*
<i>L. plantarum</i> MM	80.6±0.2	78.8±0.1	58.2±0.1	55.9±0.2	72.24±0.2	84.3±0.2*	47.6±0.3	48.2±0.2
<i>L. plantarum</i> KM	62.45±0.2	83.3±0.2*	75.5±0.2	62.7±0.1*	92.7±0.1	94.5±0.3	26.1±0.1	85.7±0.1*
<i>L. paracasei</i> KM	43.1±0.1	87.14±0.1*	38.2±0.3	71.84±0.2*	57.4±0.1	86.5±0.2*	68.8±0.2	77.14±0.3
<i>L. rhamnosus</i> KM	41.0±0.1	83.3±0.2*	32.04±0.2	63.3±0.2*	84.7±0.1	77.6±0.1	43.1±0.2	84.1±0.1*
<i>L. plantarum</i> SMA	29.6±0.2	51.44±0.3*	44.14±0.1	43.2±0.2	34.6±0.2	37.6±0.2	35.7±0.3	37.04±0.2
<i>L. plantarum</i> SS1	42.9±0.3	71.2±0.3*	47.0±0.2	55.1±0.3	86.6±0.2	63.2±0.1*	36.5±0.1	46.64±0.2*
<i>L. helveticus</i> S9	56.6±0.2	33.9±0.1*	52.9±0.2	69.4±0.2*	40.5±0.2	47.8±0.3	51.44±0.1	42.4±0.1
<i>L. lactis</i> SMS1	77.9±0.2	71.9±0.2	33.8±0.1	34.2±0.1	38.4±0.3	44.3±0.2	38.6±0.1	37.04±0.2

* statistically significant difference in comparison to viable cells ($p < 0.05$)

- International Agency for Research on Cancer (IARC). Aflatoxins. Monograph on the Evaluation of Carcinogenic Risks to Humans. Vol. 82. Lyon: IARC; 2002.
- Cavaliere C, Foglia P, Pastorini E, Samperi R, Lagana A. Liquid chromatography/tandem mass spectrometric confirmatory method for determining aflatoxin M₁ in cow milk comparison between electrospray and atmospheric pressure photoionization sources. J Chromatogr A 2006;1101:69-78. doi: 10.1016/j.chroma.2005.09.060
- Masoero F, Gallo A, Moschini M, Piva G, Diaz D. Carryover of aflatoxin from feed to milk in dairy cows with low or high somatic cell counts. Animal 2007;1:1344-50. doi: 10.1017/S1751731107000663
- Bakirci I. A study on the occurrence of aflatoxin M₁ in milk and milk products produced in Van province of Turkey. Food Control 2001;12:47-51. doi: 10.1016/S0956-7135(00)00020-7
- Galvano F, Galofaro V, De Angelis A, Galvano M, Bognanno M, Galvano G. Survey of the occurrence of aflatoxin M₁ in dairy products marketed in Italy. J Food Protect 1998;61:738-41. doi: 10.4315/0362-028X-61.6.738
- Kabak B, Var I. Factors affecting the removal of aflatoxin M₁ from food model by *Lactobacillus* and *Bifidobacterium* strains. J Environ Sci Heal B 2008;43:617-24. doi: 10.1080/03601230802234740
- Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs (Text with EEA relevance) [displayed 5 November 2018]. Available at <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32006R1881&from=EN>
- Pravilnik o najvećim dopuštenim količinama određenih kontaminanata u hrani [Ordinance on maximum permissible quantities of certain contaminants in food, in Croatian]. Narodne novine 154/2008.
- Bata A, Lásztity R. Detoxification of mycotoxin-contaminated food and feed by microorganisms. Trends Food Sci Tech 1999;10:223-8. doi: 10.1016/S0924-2244(99)00050-3
- Diaz DE, Hagler WM Jr., Blackwelder JT, Eve JA, Hopkins BA, Anderson KL, Jones FT, Whitlow LW. Aflatoxin binders II: Reduction of aflatoxin M₁ in milk by sequestering agents of cows consuming aflatoxin in feed. Mycopathologia 2004;157:233-41. doi: 10.1023/B:MYCO.0000020587.93872.59
- Di Natale F, Gallo M, Nigro R. Adsorbents selection for aflatoxins removal in bovine milks. J Food Eng 2009;95:186-91. doi: 10.1016/j.jfoodeng.2009.04.023
- Eskandari MH, Zarei M, Shekarforoush SS. Aflatoxin M₁ removal from milk by adsorbent clays. Iranian Vet J 2013;8:13-20.
- Alberts JF, Gelderblomb WCA, Botha A, van Zyl WH. Degradation of aflatoxin B₁ by fungal laccase enzymes. Int J Food Microbiol 2009;135:47-52. doi: 10.1016/j.ijfoodmicro.2009.07.022
- Assaf JC, Atoui A, El Khoury A, Chokr A, Louka N. A comparative study of procedures for binding of aflatoxin M₁ to *Lactobacillus rhamnosus* GG. Braz J Microbiol 2018;49:120-7. doi: 10.1016/j.bjm.2017.05.003
- Bovo F, Corassin CH, Rosim RE, de Oliveira CAF. Efficiency of lactic acid bacteria strains for decontamination of aflatoxin M₁ in phosphate buffer saline solution and in skimmed milk. Food Bioprocess Tech 2013;6:2230-4. doi: 10.1007/s11947-011-0770-9

21. El-Nezami H, Kankaanpää P, Salminen S, Ahokas J. Ability of dairy strains of lactic acid bacteria to bind a common food carcinogen, aflatoxin B₁. *Food Chem Toxicol* 1998;36:321-6. doi: 10.1016/S0278-6915(97)00160-9
22. El-Nezami H, Mykkänen H, Haskard C, Salminen S, Salminen E. Lactic acid bacteria as a tool for enhancing food safety by removal of dietary toxins. In: Alminen S, Von Wright A, Ouwehand A, editors. *Lactic acid bacteria: Microbiological and functional aspects*. 3rd ed. New York (NY): Marcel Dekker, Inc.; 2004. p. 397-406.
23. Markov K, Frece J, Čvek D, Lovrić N, Delaš F. Aflatoksin M₁ u sirovom mlijeku i vezanje aflatoksina pomoću bakterija mliječne kiseline [Aflatoxin M₁ in raw milk and binding of aflatoxin by lactic acid bacteria, in Croatian] *Mljekarstvo* 2010;60:244-51.
24. Pierides M, El-Nezami H, Peltonen K, Salminen S, Ahokas J. Ability of dairy strains of lactic acid bacteria to bind aflatoxin M₁ in a food model. *J Food Protect* 2000;63:645-50. doi: 10.4315/0362-028X-63.5.645
25. ISO 6887-5:2010. Microbiology of food and animal feeding stuffs - Preparation of test samples, initial suspension and decimal dilutions for microbiological examination - Part 5: Specific rules for the preparation of milk and milk products.
26. ISO 15214:1998. Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of mesophilic lactic acid bacteria. Colono-count technique at 30 degrees C.
27. Frece J, Cvrtila J, Topić I, Delaš F, Markov K. *Lactococcus lactis* ssp. *lactis* as potential functional starter culture. *Food Technol Biotech* 2014;52:489-94. doi: 10.17113/ftb.52.04.14.3794
28. Frece J, Vrdoljak M, Filipčić M, Jelić M, Čanak I, Jakopović Ž, Pleadin J, Gobin I, Landeka Dragičević T, Markov K. Microbiological quality and variability of natural microbiota in Croatian cheese maturing in lambskin sacks. *Food Technol Biotech* 2016;54:129-34. doi: 10.17113/ftb.54.02.16.4418
29. ISO 14501:2007. Milk and milk powder. Determination of aflatoxin M₁ content. Clean-up by immunoaffinity chromatography and determination by high-performance liquid chromatography.
30. Lahtinen SJ, Haskard CA, Ouwehand AC, Salminen SJ, Ahokas JT. Binding of aflatoxin B₁ to cell wall components of *Lactobacillus rhamnosus* strain GG. *Food Addit Contam* 2004;21:158-64. doi: 10.1080/02652030310001639521
31. Shetty PH, Jespersen L. *Saccharomyces cerevisiae* and lactic acid bacteria as potential mycotoxin decontaminating agents. *Trends Food Sci Tech* 2006;17:48-55. doi: 10.1016/j.tifs.2005.10.004
32. Haskard CA, El-Nezami HS, Kankaanpää PE, Salminen S, Ahokas JT. Surface binding of aflatoxin B₁ by lactic acid bacteria. *Appl Environ Microb* 2001;67:3086-91. doi: 10.1128/AEM.67.7.3086-3091.2001
33. Brackett RE, Marth EH. Association of aflatoxin M₁ with casein. *Eur Food Res Technol* 1997;174:439-41. doi: 10.1007/BF01042721

Uklanjanje aflatoksina M₁ iz mlijeka s autohtonim bakterijama mliječne kiseline centrifugiranjem i filtriranjem

Kako bi se smanjila izloženost ljudi aflatoksinu M₁ (AFM₁), razina toga visokokancerogenoga mikotoksina u mlijeku, termički obrađeno mlijeku i drugim mliječnim prerađevinama ograničena je na <0,05 μg kg⁻¹. Međutim, njegovo uklanjanje iz mliječnih prerađevina pravi je izazov za proizvođače mlijeka, jer komercijalni aditivi mijenjaju njegova organoleptička svojstva, a sama filtracija daje slabe rezultate. Cilj ovoga rada bio je pronaći soj bakterije mliječne kiseline (BMK) iz mlijeka ili mliječnih prerađevina koji najučinkovitije veže AFM₁, te vidjeti utječe li termičko tretiranje izabrane BMK na učinkovitost vezanja. Također, željeli smo istražiti može li centrifugiranje poboljšati filtriranje dobivenog AFM₁-BMK kompleksa iz mlijeka. Kako bismo to učinili, izolirali smo i identificirali 10 autohtonih vrsta/sojeva BMK, inkubirali žive ili termički tretirane stanice (10⁸ CFU mL⁻¹) u mlijeku s 0,5 μg L⁻¹ AFM₁ na 4 °C tijekom 0, 2, 4 i 24 sata te kvantificirali nevezani AFM₁ koristeći se HPLC-om. Učinkovitost vezanja AFM₁ živim stanicama kretala se od 21 do 92 % te od 26 do 94 % termički tretiranim stanicama. Budući da su i žive i termički tretirane stanice *Lactobacillus plantarum* KM pokazale najbolje rezultate, navedeni se soj koristio u sljedećem koraku uklanjanja AFM₁ iz mlijeka. Tretman toplinom u kombinaciji s filtracijom i centrifugiranjem rezultirao je uklanjanjem do 96 % AFM₁ iz mlijeka.

KLJUČNE RIJEČI: HPLC; *L. plantarum* KM; sekundarni metaboliti plijesni; termički tretirane stanice; žive stanice