

EXPLICIT BOUNDS FOR COMPOSITE LACUNARY POLYNOMIALS

CHRISTINA KAROLUS

University of Salzburg, Austria

ABSTRACT. Let $f, g, h \in \mathbb{C}[x]$ be non-constant complex polynomials satisfying $f(x) = g(h(x))$ and let f be lacunary in the sense that it has at most l non-constant terms. Zannier proved in [9] that there exists a function $B_1(l)$ on \mathbb{N} , depending only on l and with the property that $h(x)$ can be written as the ratio of two polynomials having each at most $B_1(l)$ terms. Here, we give explicit estimates for this function or, more precisely, we prove that one may take for instance

$$B_1(l) = (4l)^{(2l)^{(3l)^{l+1}}}.$$

Moreover, in the case $l = 2$, a better bound is obtained using the same strategy.

1. INTRODUCTION

Let $f, g, h \in \mathbb{C}[x]$ and $f = g \circ h$ be a lacunary polynomial with l non-constant terms, i.e. f is of the form $f(x) = a_0 + a_1x^{n_1} + \dots + a_lx^{n_l}$. Note that only the number of terms is viewed as fixed, while the coefficients and the degrees may vary. In [9], it was shown by Zannier that there exists a function $B_1(l)$ such that $h(x)$ can be written as the ratio of two polynomials in $\mathbb{C}[x]$ both having no more than $B_1(l)$ terms. In order to give explicit estimates for $B_1(l)$, we are following the strategy of [9, Prop. 2]. Therefore, we will recall those parts of Zannier's proof, which are relevant to our arguments, in the very beginning of the paper. We prove

THEOREM 1.1. *Let $f, g, h \in \mathbb{C}[x]$ be non-constant complex polynomials such that $f(x) = g(h(x))$ has at most l non-constant terms. Then $h(x)$ can*

2010 *Mathematics Subject Classification.* 11C08, 11R09.

Key words and phrases. Decomposable polynomials, lacunary polynomials.

The author was supported by Austrian Science Fund (FWF) Grant No. P24574.

be written as the ratio of two polynomials in $\mathbb{C}[x]$ having each at most $B_1(l)$ terms, where

$$B_1(l) = (4l)^{(2l)^{(3l)^{l+1}}} \text{ for } l \geq 1.$$

There are quite different notions of *lacunarity* (lacunary polynomials are also sometimes called *sparse*). Here, we deal with the situation that the number of terms of a given polynomial is fixed. It was conjectured by Erdős that if g is a complex non-constant polynomial with the property that $g(x)^2$ has at most l terms, then $g(x)$ has also boundedly many terms and their number depends only on l . In [7], Schinzel proved a generalized version of Erdős's conjecture, namely the statement not only for $g(x)^2$ but for $g(x)^d$, $d \in \mathbb{N}$. He also extended the conjecture to compositions $f(x) = g(h(x))$, claiming that if f has l terms, then $h(x)$ has at most $B(l)$ terms for some function B on \mathbb{N} . Zannier gave a proof for this (actually in a stronger version), wherein he showed in a first step the existence of a function B_1 such that, under the given assumptions, $h(x)$ can be written as a rational function with at most $B_1(l)$ terms in both the numerator and the denominator ([9]). Using this result, he proved the stated claim for representations as polynomials¹ and moreover, he gave a complete description of general decompositions $f(x) = g(h(x))$ of a given polynomial $f(x)$ with l terms. Also, for an outer composition factor g in $f(x) = g(h(x))$, Zannier gave suitable bounds for the degree of g depending only on the number of terms of f ([8]). Note that in both, the polynomial case and the case of a rational function, the special shapes $h(x) = ax^m + b$ and $h(x) = ax^m + bx^{-m} + c$, respectively, must be taken into account. This follows easily from the following observation. Let $h(x) = ax^m + b$ and $g(x) = g_1(x - b)$. Then $f(x) = g(h(x)) = g_1(ax^m)$ has at most l non-constant terms, whereas the degree of g can be arbitrarily high. Similarly, in the case of rational functions, for instance one can take $h(x) = x + x^{-1}$ and $g(x) = T_n(x)$, the n -th Chebyshev polynomial, to get a contradiction with the given statements.

The results were later extended first to Laurent polynomials ([10]) and then to rational functions ([5]). Recently in [3], Fuchs, Mantova and Zannier achieved a final result for completely general algebraic equations $f(x, g(x)) = 0$. Here, $f(x, y)$ is assumed to be monic and of given degree in y and with boundedly many terms in x . As pointed out in [4], there are also other forms of lacunarity. Here, Fuchs and Pethő considered rational functions having only a given number of zeros and poles and they again studied their decomposability. This can be seen as a multiplicative analogue to the above mentioned problem. Based on their results, by computational experiments Pethő and Tengeley studied the decomposability of rational functions having at most four zeros

¹Examples as $h(x) = (x^n - 1)/(x - 1) = x^{n-1} + x^{n-2} + \dots + x + 1$ show that, written as a polynomial, $h(x)$ can have substantially more terms than in a representation as a rational function.

and poles and they provided parametrizations of all possible solutions and the appropriate varieties in this case ([6]).

The present paper is organized as follows. In the very beginning, we repeat the main results and parts of Zannier's proof, which are crucial for our deductions. Based on this, in Section 3 we prove Theorem 1.1, giving an explicit bound for the function $B_1(l)$ in Zannier's proposition. This bound happens to be triple-exponential. The reason for this is the following. In the proof, an estimate for the ratio n_l/n_1 is used to give an upper bound for an exponent. Such an estimate can be found through a recursive procedure, pointed out by Zannier. The bound we obtain through this method will be double-exponential, which in the end leads to the received order. We do not know whether that can be improved in general or how far away we are from a "good bound". However, our bound surely is far from the truth. At least for the cases $l = 1$ and $l = 2$ we have the smaller estimates 2 and 1377, respectively. For the latter bound, the corresponding statement and its proof is given in the last part of Section 3.

Finally, we mention that, independently of us, Dona in [1] also gave an explicit bound for $B_1(l)$. Namely, he proved that one may as well take

$$B_1(l) = 2^{(2l)^{16(2l)^{l-1}}}.$$

Moreover, he showed that as a polynomial, the number of terms of $h(x)$ is bounded above by $B_1(l)^{5B_1(l)^2}$. His deductions are based on Zannier's proofs too. Therefore it is not surprising that qualitatively his bound is of the same shape (i.e. also triple-exponential) although it is quantitatively better than ours. Nevertheless, and also for the fact that our result is already mentioned in [2], it appears to us that the result is still worth to be found in the literature.

2. ARGUMENTS FROM ZANNIER'S PROOF

In this section we recall the main steps of Zannier's proof ([9, Prop. 2]). For clarity, we keep the original notation. Assuming that $f(x) = g(h(x))$ has at most l non-constant terms, let $\deg f = m = n_l$ and $\deg g = d$, so that we have $\deg h = n_l/d$. From [8, Thm. 1] it follows that $d \leq 2l(l-1)$. Set $y = 1/x$ and $\tilde{h}(y) = x^{-n_l/d}h(x) = y^{n_l/d}h(1/y)$, such that $\tilde{h} \in \mathbb{C}[y]$. Moreover, write $f(x) = ax^m(1 + b_1y^{n_1} + \dots + b_ly^{n_l})$, where $0 =: n_0 < n_1 < \dots < n_l$. We may assume that f has exactly l non-constant terms, i.e. $ab_1 \dots b_l \neq 0$. Instead of $h(x)$, Zannier proves the statement for $\tilde{h}(y)$, which in fact has the same number of terms as $h(x)$. Also, its degree is bounded by $\deg \tilde{h} \leq n_l/d = \deg h$.

Writing $\delta_p(x) = 1 + b_1x^{n_1} + \dots + b_px^{n_p}$ for an integer p with $1 \leq p \leq l-1$, Zannier deduces that, in $\mathbb{C}[[y]]$, \tilde{h} is of the form $\tilde{h}(y) = t_1 + t_2 + \dots + t_L + O(y^{2n_l})$, where L is an integer which may be bounded by a function in l and

n_l/n_{p+1} and the t_1, \dots, t_L are all of the shape

$$(2.1) \quad c\delta_p(y)^{s/d-k}y^{h_1n_{p+1}+\dots+h_{l-p}n_l+(1-s)m/d},$$

for varying $h_1, \dots, h_{l-p} \in \mathbb{N}$, $k = h_1 + \dots + h_{l-p}$, suitable constants $c = c(h_1, \dots, h_{l-p}, s)$ and $s \in \{1, 0, -1, \dots, 1 - 2d\}$, where the exponent of y in such terms is smaller than $2n_l$. An easy argument shows that for L one may take the rough estimate $L \leq (2d + 1)(2n_l/n_{p+1} + 1)^l$.

In the case that $t_1, \dots, t_L, \tilde{h}(y)$ are linearly independent over \mathbb{C} , using a function field variant of Schmidt subspace theorem (see [9, Proposition 1]), the author then proves that

$$(2.2) \quad n_l \leq 16^{l+1}d^3(n_l/n_{p+1})^{2l}(1 + n_p),$$

and furthermore, using $1 + n_p \leq 2n_p$, it follows that

$$(2.3) \quad (n_l/n_p) \leq 2 \cdot 16^{l+1}d^3(n_l/n_{p+1})^{2l}.$$

If on the other hand $t_1, \dots, t_L, \tilde{h}(y)$ are linearly dependent over \mathbb{C} , it inductively follows that $\tilde{h}(y)$ can be written as the ratio of two polynomials in $\mathbb{C}[y]$ having each at most $B_2(l, n_l/n_{p+1})$ terms, where $B_2(l, u)$ is a suitable function which may be estimated in terms of $B_1(l - 1)$ and of $u \geq 0$.

Zannier then distinguishes between those two cases for each $p = l - 1, l - 2, \dots$. Suppose that for $p = l - 1, l - 2, \dots, l - r$ always the first situation occurs. In that case we can recursively determine upper bounds for the quotients n_l/n_p , since for $p = l - 1$ the initial condition $n_l/n_{p+1} = 1$ holds.

The proof then argues via backwards induction on $p = l - 1, l - 2, \dots$. If $t_1, t_2, \dots, t_L, \tilde{h}(y)$ are linearly independent for all $p = l - 1, l - 2, \dots, 1$, one can use (2.3) and (2.2) to get an estimate for n_l , which in fact also gives an estimate for the number of terms of $\tilde{h}(y)$ (and hence of $h(x)$) even written as a polynomial, as the number of terms of \tilde{h} is bounded by its degree $n_l/d \leq n_l$.

On the other hand, if the $t_1, \dots, t_L, \tilde{h}(y)$ are linearly dependent over \mathbb{C} for at least one $p \in \{1, \dots, l - 1\}$, let p_0 denote the last p for which this case occurs, i.e. we may assume that we have linear independency for $p = l - 1, l - 2, \dots, p_0 + 1$ and linear dependency for $p = p_0$. Here, Zannier concludes that $\tilde{h}(y)$ is the sum of at most $L = L(p_0)$ terms of the form (2.1), where $k \leq 2ln_l/n_{p_0+1}$ and $s \in \{1, 0, -1, \dots, 1 - 2d\}$. Now the author uses the linear independency in the cases $p > p_0$ to estimate the quotient n_l/n_{p_0+1} , which occurs in the estimates for L and k to get the desired result.

3. PROOF OF THEOREM 1.1 AND THE CASE $l = 2$

In order to prove Theorem 1.1 we start with the following lemma.

LEMMA 3.1. *Let $f \in \mathbb{C}[x]$ be of the form $f(x) = a_0 + a_1x^{n_1} + \dots + a_lx^{n_l}$, $0 < n_1 < \dots < n_l$, and let for every integer p , $1 \leq p \leq l - 1$, $S = S(p) =$*

$\{t_1, \dots, t_L, \tilde{h}(y)\}$ be the set described in Section 2. If for each $p = l-1, \dots, l-r$ the set S is linearly independent over \mathbb{C} , it holds that

$$\frac{n_l}{n_{l-r}} \leq (16^{l+2}l^6)^{(3l)^{r-1}}.$$

PROOF. We set $\lambda = 2 \cdot 16^{l+1}d^3$. By [8, Thm. 1], we have $d \leq 2l(l-1)$. Applying (2.3), we iteratively get

$$\begin{aligned} \frac{n_l}{n_{l-1}} &\leq 2 \cdot 16^{l+1} \cdot d^3 =: \lambda, \\ \frac{n_l}{n_{l-2}} &\leq 2 \cdot 16^{l+1} \cdot d^3 (2 \cdot 16^{l+1} \cdot d^3)^{2l} = \lambda^{2l+1}, \\ \frac{n_l}{n_{l-3}} &\leq \lambda(\lambda^{2l+1})^{2l} = \lambda^{1+2l(1+2l)}, \\ &\vdots \\ \frac{n_l}{n_{l-r}} &\leq \lambda^{1+2l(1+2l(1+2l(\dots)))} \leq \lambda^{(3l)^{r-1}}. \end{aligned}$$

As $\lambda = 2 \cdot 16^{l+1}d^3 \leq 2 \cdot 16^{l+1}(2l(l-1))^3 \leq 16^{l+2}l^6$, in the case that for $p = l-1, l-2, \dots, l-r$ the set S is always linearly independent over \mathbb{C} , we obtain the claimed result. \square

PROOF OF THEOREM 1.1. Following the proof of [9, Prop. 2], we argue by induction on l . As in the previous section, we keep the notation from Zannier's proof. As pointed out by Zannier, for $l = 1$ we may take $B_1(1) \geq 2$.² Now, for the rest of the proof let us assume that the statement holds for $l-1$, i.e. that B_1 has been suitably defined on $\{1, 2, \dots, l-1\}$.

For $p = l-1, l-2, \dots, 0$ and $\delta_p(y) = 1 + b_1 y^{n_1} + \dots + b_p y^{n_p}$, in $\mathbb{C}[[y]]$ we can write

$$\tilde{h}(y) = t_1 + t_2 + \dots + t_L + O(y^{2n_l}),$$

where the t_i , $1 \leq i \leq L$, are terms of the shape (2.1), which we may assume to be linearly independent over \mathbb{C} , and L is an integer which, as we know from the proof of [9, Prop. 2], can be bounded by $L \leq (2d+1)(2n_l/n_{p+1} + 1)^l$.

We now consider the set $S = \{t_1, \dots, t_L, \tilde{h}(y)\}$ and, following the original proof, we distinguish between two possible cases, namely that $S = S(p)$ is linearly independent over \mathbb{C} for every $p = l-1, l-2, \dots, 1$ or that there is an integer p_0 , $1 \leq p_0 \leq l-1$, such that we have linear dependency, i.e. $S = S(p)$ is linearly independent for $p = l-1, l-2, \dots, p_0+1$ and $S(p_0)$ is linearly dependent over \mathbb{C} .

CASE 1. In the case of linear independency for every $p = l-1, l-2, \dots, 1$, by Lemma 3.1, we get the estimate $n_l/n_1 \leq (16^{l+2}l^6)^{(3l)^{l-2}}$. Now, we may

²If $g(h(x))$ is a monomial ax^m , then g cannot have distinct roots, hence is of the shape $g(x) = b(x-\xi)^n$. Then $h(x) = \xi + \eta x^{m/n}$ (where $b\eta^n = a$) has at most two terms.

apply (2.2) for $p = 0$ (recall that $n_0 = 0$) to get

$$\begin{aligned} n_l &\leq 16^{l+1} d^3 (16^{l+2} l^6)^{(2l)(3l)^{l-2}} \leq 16^{l+1} (2l^2)^3 (16^{l+2} l^6)^{(2l)(3l)^{l-2}} \\ &< (16^{l+2} l^6)^{(2l)(3l)^{l-2}+1} < (16^{l+2} l^6)^{(3l)^{l-1}}. \end{aligned}$$

This clearly gives an estimate for the number of terms of $\tilde{h}(y)$ written as a polynomial (and hence of $h(x)$), bounding its degree by $n_l/d \leq n_l$.

CASE 2. Let us now consider the second case, where we have linear dependency for some $p = p_0$. Since we assume $S = S(p)$ to be linearly independent over \mathbb{C} for each $p > p_0$, we have, again by Lemma 3.1, that n_l/n_{p_0+1} can be bounded by $(16^{l+2} l^6)^{(3l)^{l-1}}$ as well. Let $e = [K : \mathbb{C}(y)]$, where $K = \mathbb{C}(y, \delta_p(y)^{1/d})$, so $e \in \{1, \dots, d\}$ is the least integer such that $\delta_p(y)^e$ is a d -th power in $\mathbb{C}(y)$. We write $\delta_p(y)^e = \eta_p(y)^d$ for a polynomial $\eta_p \in \mathbb{C}[y]$ to express this fact. Also, for the rest of the proof, we simply write p instead of p_0 . As pointed out by Zannier, it then follows that $\tilde{h}(y) = \Lambda_0$, where Λ_0 is the sum of at most L terms of the shape

$$(3.1) \quad c \eta_p(y)^{(s-kd)/e} y^{(1-s)m/d + h_1 n_{p+1} + \dots + h_{l-p} n_l},$$

with $k = h_1 + \dots + h_{l-p} \leq 2ln_l/n_{p+1}$ and where $s \in \{1 - 2d, \dots, 0, 1\}$ is such that $e|s$.

In order to estimate the number of terms in the requested representation of \tilde{h} , we look at $\eta_p(y)^{(s-kd)/e}$, which is the important quantity in (3.1), when it comes to counting terms. Note that $\eta_p(y)^{d/e} = \delta_p(y)$ is a polynomial which has at most $p \leq l - 1$ non-constant terms, so by the induction hypothesis $\eta_p(y)$ can be expressed as a rational function with at most $B_1(l - 1)$ terms in both the numerator and the denominator. That is,

$$\eta_p(y) = \frac{\eta_{p,1}(y)}{\eta_{p,2}(y)},$$

where $\eta_{p,1}, \eta_{p,2}$ are complex polynomials with at most $B_1(l - 1)$ terms.

To find a suitable function for B_1 , we start with estimating the exponent of $\eta_p(y)$ in (3.1). Recall that $s \in \{1 - 2d, \dots, -1, 0, 1\}$, $k = h_1 + \dots + h_{l-p} < 2ln_l/n_{p+1}$ and that $d \leq 2l(l - 1)$. Therefore, if $l \geq 3$, we get the following estimate

$$\begin{aligned} |(s - kd)/e| &\leq |s| + |kd| \leq 2d - 1 + 2l \frac{n_l}{n_{p+1}} d \\ &\leq 2 \cdot 2l(l - 1) \left(1 + l \frac{n_l}{n_{p+1}}\right) - 1 \leq 2^2 l^2 \left(1 + l(2^{4(l+2)} l^6)^{(3l)^{l-1}}\right) - 1 \\ &< 2^2 l^2 \left(2 \cdot 2^{(4l+8)(3l)^{l-1}} l^{(3l)^l}\right) - 1 = 2^{3+(4l+8)(3l)^{l-1}} l^{2+(3l)^l} - 1. \end{aligned}$$

It is easy to check that this is also an upper bound for $|(s - kd)/e|$ in the case $l = 2$. Hence, if we set $M = 2^{3+(4l+8)(3l)^{l-1}}l^{2+(3l)^l}$, we see that

$$(s - kd)/e \in \{-(M - 1), \dots, -1, 0, 1\}.$$

Now, let us again consider $\tilde{h}(y) = \Lambda_0$. After reducing all of the L terms of the shape (3.1) to the common denominator $\eta_{2,p}(y)\eta_{1,p}(y)^{M-1}$, we can make the rough estimate $B_1(l-1)B_1(l-1)^{M-1} = B_1(l-1)^M$ for the number of terms in the denominator and $LB_1(l-1)^M$ for the number of terms in the numerator. Since we are looking for a function that bounds both the number of terms in the numerator and the denominator, it now suffices to define B_1 in such a way that $B_1(l) \geq x_l$, where $(x_l)_{1 \leq l} \subset \mathbb{N}$ is the recurrence sequence defined by $x_1 = 2$ and $x_l = L \cdot x_{l-1}^M$ (recall that for $l = 1$ it already has been shown in [9, Prop. 2] that we may take $B_1(1) \geq 2$). It follows that

$$x_l = L^{1+M+M^2+\dots+M^{l-2}} \cdot 2^{M^{l-1}} < (2L)^{M^{l-1}},$$

hence we may as well define B_1 as any function satisfying $B_1(l) \geq (2L)^{M^{l-1}}$. Similarly as for the exponent, we find an upper bound for $2L$,

$$\begin{aligned} 2L &\leq 2(2d+1)(1+2n_l/n_{p+1})^l \\ &\leq 2(2^{2l}l(l-1)+1)(1+2(2^{4(l+2)}l^6)^{(3l)^{l-1}})^l \\ &< 2^3l^2 \cdot 2^{2l+4l(l+2)(3l)^{l-1}}l^{2(3l)^l} \\ &= 2^{3+2l+4l(l+2)(3l)^{l-1}}l^{2(3l)^l+2} \\ &= 4^{1.5+l+2l(l+2)(3l)^{l-1}}l^{2(3l)^l+2} \\ &< 4^{(3l)^{l-1}(2l^2+4l+1)}l^{2((3l)^l+1)} \\ &< 4^{(3l)^{l-1}(4/3)^l 3l^2}l^{2(4l)^l} \\ &= 4^{2^{2l}l^{l+1}}l^{2^{2l+1}l^l}. \end{aligned}$$

Based on these estimates, we get an upper bound for $(2L)^{M^{l-1}}$:

$$\begin{aligned} (2L)^{M^{l-1}} &\leq (4^{2^{2l}l^{l+1}}l^{2^{2l+1}l^l})^{2^{3+(4l+8)(3l)^{l-1}}(l-1)}l^{(2+(3l)^l)(l-1)} \\ &= 4^{2^{(3l)^{l-1}(4l^2+4l-8)+5l-3}l^{(3l)^l(l-1)+3l-1}}l^{2^{(3l)^{l-1}(4l^2+4l-8)+5l-2}l^{(3l)^l(l-1)+3l-2}}. \end{aligned}$$

Note that for the exponents we have

$$(3l)^{l-1}(4l^2+4l-8+(5l-2)/(3l)^{l-1}) < (3l)^{l-1}(3l)^2 = (3l)^{l+1}$$

and

$$(3l)^l(l-1)+3l-1 < (3l)^l(l-1+1) < (3l)^{l+1}.$$

We therefore get

$$(2L)^{M^{l-1}} \leq 4^{2^{(3l)^{l+1}}l^{(3l)^{l+1}}}l^{2^{(3l)^{l+1}}l^{(3l)^{l+1}}} = (4l)^{(2l)^{(3l)^{l+1}}}.$$

Finally, we have to check that the obtained estimate also holds in CASE 1, i.e. if $(16^{l+2}l^6)^{(3l)^{l-1}} \leq (4l)^{(2l)^{(3l)^{l+1}}}$ for $l \geq 2$. But this follows trivially from the fact that we already used the quantity $(16^{l+2}l^6)^{(3l)^{l-1}}$ in estimating $2L$ in CASE 2, hence the claimed result. \square

Using the same arguments, it is also possible to obtain a better bound for $l = 2$, since in this case we are able to keep the estimates during the proof essentially smaller. Eventually, we get the following.

PROPOSITION 3.2. *Let $f, g, h \in \mathbb{C}[x]$ be non-constant complex polynomials such that $f(x) = g(h(x))$ has at most 2 non-constant terms. Then $h(x)$ may be written as the ratio of two polynomials in $\mathbb{C}[x]$ having each at most 1377 terms.*

PROOF. Assume that f is a polynomial in $\mathbb{C}[x]$ with at most two non-constant terms, i.e. $f(x) = a_0 + a_1x^{n_1} + a_2x^{n_2}$, $0 < n_1 < n_2$. We keep the notations from above. If $\deg g = d = 1$, the number of terms of $g(h(x))$ and of $h(x)$ may only deviate by one (namely the constant term), hence in this case $h(x)$ is a polynomial that has not more than 3 terms. So in the following we assume that $d \geq 2$. Note that we also have $d \leq 4$, for $d \leq 2l(l-1)$.

As before, we start with the observation that $\tilde{h}(y)$ can be written in the shape

$$\tilde{h}(y) = t_1 + t_2 + \dots + t_L + \mathcal{O}(2n_2),$$

where the t_i are terms of the form (2.1), which we may assume to be linearly independent over \mathbb{C} . We again consider the two cases that $S = S(p) = \{t_1, \dots, t_L, \tilde{h}(y)\}$ is linearly dependent or linearly independent over \mathbb{C} , respectively. In general, we had to distinguish between those cases for each $p = l-1, l-2, \dots$, but since $l = 2$, $p = 1$ is the only remaining situation we have to look at. The above approximation was chosen in such a way that in each t_i the exponent of y is smaller than $2n_2$, that is

$$\frac{(1-s)n_2}{d} + h_1n_2 = \left(\frac{1-s}{d} + h_1 \right) n_2 \leq 2n_2,$$

where $s \leq 1$ and $h_1 \geq 0$ are integers. It follows that $1 - 2d \leq s \leq 1$ and $h_1 \in \{0, 1, 2\}$. A rough estimate on the number of such terms t_i would therefore be $3(2d+1) \leq 27$, but checking for each $d \in \{2, 3, 4\}$ separately, we find that under the given conditions there cannot be more than 17 such exponents, hence

$$L \leq 17.$$

CASE 1. If $\{t_1, \dots, t_L, \tilde{h}(y)\}$ is linearly independent over \mathbb{C} , then by [9, Prop.1] we get

$$n_2 \leq L(L+1)d(1+n_1) \leq 17 \cdot 18 \cdot 4 \cdot 2n_1,$$

and consequently $n_2/n_1 \leq 2448$. The starting point in Zannier's proof was to expand $h(x)$ as a Puiseux-series, which led to the observation that, in $\mathbb{C}[[y]]$, for certain $\gamma_{-1}, \gamma_0, \gamma_1, \dots \in \mathbb{C}$ we have

$$\tilde{h}(y) = \gamma_{-1} \tilde{f}(y)^{1/d} + \gamma_0 y^{m/d} + \gamma_1 y^{2m/d} \tilde{f}(y)^{-1/d} + \gamma_2 y^{3m/d} \tilde{f}(y)^{-2/d} + \dots,$$

where the roots $\tilde{f}(y)^{s/d}$ are of the form

$$\tilde{f}(y)^{s/d} = \sum_{(h_1, h_2) \in \mathbb{N}_0^2} c_{s,d,(h_1, h_2)} b_1^{h_1} b_2^{h_2} y^{h_1 n_1 + h_2 n_2},$$

for certain universal coefficients $c_{s,d,(h_1, h_2)}$. Since $\tilde{h}(y)$ is a polynomial with $\deg \tilde{h} \leq m/d = n_2/d$, it follows that the only terms which may contribute to $\tilde{h}(y)$ are $\gamma_0 y^{m/d}$ and terms coming from $\gamma_{-1} \tilde{f}(y)^{1/d}$, for which $h_1 n_1 + h_2 n_2 \leq n_2/d$ holds. Since we assumed $2 \leq d$, it follows that $h_2 = 0$ and $h_1 \leq n_2/(2n_1) \leq 1224$. Taking also the terms with $h_1 = h_2 = 0$ and $\gamma_0 y^{m/d}$ into account, we conclude that \tilde{h} is a polynomial having no more than 1226 terms.

CASE 2. On the other hand, if $\{t_1, \dots, t_L, \tilde{h}(y)\}$ is linearly dependent over \mathbb{C} , then $\tilde{h}(y)$ may be written as the sum of at most $L \leq 17$ terms of the shape (3.1), where $\eta_p(y) = \eta_{p,1}(y)/\eta_{p,2}(y)$ and $\eta_{p,1}(y), \eta_{p,2}(y) \in \mathbb{C}[x]$ both have at most two terms and $k = h_1 \leq 2$ (this follows from $(1-s)m/d + h_1 n_{p+1} + \dots + n_{l-p} n_l = (1-s)m/d + h_1 n_l \leq 2n_l$). Recall that $s \in \{1, 0, \dots, -7\}$ and $d \leq 4$. It follows that $(s - kd)/e \in \{1, 0, -1, \dots, -15\}$. Since an n -th power of the sum of two terms can be written as the sum of $n+1$ terms, bringing all of the terms to common denominator $\eta_{p,1}(y)^{15} \eta_{p,2}(y)$, we can write $\tilde{h}(y)$ as a rational function with at most $2 \cdot 16 = 32$ terms in the denominator. In this notation we have

$$\left(\frac{\eta_{p,1}(y)}{\eta_{p,2}(y)} \right)^{(s-kd)/e} = \frac{n(y)}{\eta_{p,1}(y)^{15} \eta_{p,2}(y)},$$

where the numerator has the shape $n(y) = \eta_{p,1}(y)^{k_1} \eta_{p,2}(y)^{k_2}$ with $k_1 + k_2 = 16$. Therefore, the numerator of such a term has not more than $(k_1 + 1)(k_2 + 1) = (k_1 + 1)(17 - k_1)$ terms. This is maximal for $k_1 = k_2 = 8$, hence in the numerator there are at most 81 terms. Since $\tilde{h}(y)$ may be written as the sum of at most $L \leq 17$ such terms, and since this is also an upper bound in the first case, we may take $B_1(2) = 17 \cdot 81 = 1377$. \square

The obtained bound is still not very small. However, compared to the bound $B_1(2) = 2^{3 \cdot 2^{432}}$, this still gives a notable improvement.

REFERENCES

- [1] D. Dona, On lacunary polynomials and a generalization of Schinzel's conjecture, Master thesis, Concordia University/Université de Bordeaux, 2015.

- [2] C. Fuchs and L. Hajdu, *30 years of collaboration*, Period. Math. Hung. **74** (2017), 255–274.
- [3] C. Fuchs, V. Mantova and U. Zannier, *On fewnomials, integral points and a toric version of Bertini's theorem* J. Amer. Math. Soc. **31** (2018), 107–134.
- [4] C. Fuchs and A. Pethő, *Composite rational functions having a bounded number of zeros and poles*, Proc. Amer. Math. Soc. **139** (2011), 31–38.
- [5] C. Fuchs and U. Zannier, *Composite rational functions expressible with few terms*, J. Eur. Math. Soc. **14** (2012), 175–208.
- [6] A. Pethő and S. Tengely, *On composite rational functions*, in: Number theory, analysis, and combinatorics, De Gruyter Proc. Math., De Gruyter, Berlin, 2014, 241–259.
- [7] A. Schinzel, *On the number of terms of a power of a polynomial*, Acta Arith. **49** (1987), 55–70.
- [8] U. Zannier, *On the number of terms of a composite polynomial*, Acta Arith. **127** (2007), 157–167.
- [9] U. Zannier, *On composite lacunary polynomials and the proof of a conjecture of Schinzel*, Invent. Math. **174** (2008), 127–138.
- [10] U. Zannier, *Addendum to the paper: On the number of terms of a composite polynomial*, Acta Arith. **140** (2009), 93–99.

C. Karolus
Department of Mathematics
University of Salzburg
5020 Salzburg
Austria
E-mail: christina.karolus@sbg.ac.at

Received: 31.12.2018.

Revised: 1.4.2019.