

**RATIONAL SEQUENCES ON DIFFERENT MODELS OF
ELLIPTIC CURVES**GAMZE SAVAŞ ÇELİK, MOHAMMAD SADEK AND GÖKHAN SOYDAN
Bursa Uludağ University, Turkey and Sabancı University, Turkey

ABSTRACT. Given a set S of elements in a number field k , we discuss the existence of planar algebraic curves over k which possess rational points whose x -coordinates are exactly the elements of S . If the size $|S|$ of S is either 4, 5, or 6, we exhibit infinite families of (twisted) Edwards curves and (general) Huff curves for which the elements of S are realized as the x -coordinates of rational points on these curves. This generalizes earlier work on progressions of certain types on some algebraic curves.

1. INTRODUCTION

An algebraic (affine) plane curve C of degree d over some field k is defined by an equation of the form

$$\{(x, y) \in k^2 : f(x, y) = 0\}$$

where f is a polynomial of degree d . The algebraic affine plane curve C can also be extended to the projective plane by homogenising the polynomial f . If $P = (x, y)$, then we write $x = x(P)$ and $y = y(P)$.

Studying the set of k -rational points on C , $C(k)$, has been subject to extensive research in arithmetic geometry and number theory, especially when k is a number field. For example, if f is a polynomial of degree 2, then one knows that C is of genus 0, and so if C possesses one rational point then it contains infinitely many such points. If f is of degree 3, then C is a genus 1 curve if it is smooth. In this case, if $C(k)$ contains one rational point, then it is an elliptic curve, and according to Mordell-Weil Theorem, $C(k)$ is a finitely

2010 *Mathematics Subject Classification.* 11D25, 11G05, 14G05.

Key words and phrases. Elliptic curve, Edwards curve, Huff curve, rational sequence, rational point.

generated abelian group. In particular, $C(k)$ can be written as $T \times \mathbb{Z}^r$ where T is the subgroup of points of finite order, and $r \geq 0$ is the *rank* of C over k .

In enumerative geometry, one may pose the following question. Given a set of points S in k^2 , how many algebraic plane curves C of degree d satisfy that $S \subseteq C(k)$? It turns out that sometimes the answer is straightforward. For example, given 10 points in k^2 , in order for a cubic curve to pass through these points, a system of 10 linear equations will be obtained by substituting the points of S in

$$a_1x^3 + a_2x^2y + a_3x^2 + a_4xy^2 + a_5xy + a_6x + a_7y^3 + a_8y^2 + a_9y + a_{10} = 0$$

and solving for a_1, \dots, a_{10} . Therefore, there exists a nontrivial solution to the system if the determinant of the corresponding matrix of coefficients is zero, hence a cubic curve through the points of S . Thus, one needs linear algebra to check the existence of algebraic curves of a certain degree through various specified points in k^2 .

In this article, we address the following, relatively harder, question. Given $S \subset k$, are there algebraic curves C of degree d such that for every $x \in S$, $x = x(P)$ for some $P \in C(k)$? In other words, S constitutes the x -coordinates of a subset of $C(k)$. The latter question can be reformulated to involve y -coordinates instead of x -coordinates. It is obvious that linear algebra cannot be utilized to attack the problem as substituting with the x -values of S will not yield linear equations.

Given a set $S = \{x_1, x_2, \dots, x_n\} \subset k$, if (x_i, y_i) , $i = 1, \dots, n$, are k -rational points on an algebraic curve C , then these rational points are said to be an S -sequence of length n . In what follows, we summarize the current state of knowledge for different types of S .

We first describe the state-of-art when the elements of $S \subset \mathbb{Q}$ are chosen to form an *arithmetic progression*, Lee and Vélez ([10]) found infinitely many curves described by $y^2 = x^3 + a$ containing S -sequences of length 4. Bremner ([2]) showed that there are infinitely many elliptic curves with S -sequences of length 7 and 8. Campbell ([5]) gave a different method to produce infinite families of elliptic curves with S -sequences of length 7 and 8. In addition, he described a method for obtaining infinite families of quartic elliptic curves with S -sequences of length 9, and gave an example of a quartic elliptic curve with an S -sequence of length 12. Ulas ([17]) first described a construction method for an infinite family of quartic elliptic curves on which there exists an S -sequence of length 10. Secondly he showed that there is an infinite family of quartics containing S -sequences of length 12. Macleod ([11]) showed that simplifying Ulas' approach may provide a few examples of quartics with S -sequences of length 14. Ulas ([18]) found an infinite family of genus two curves described by $y^2 = f(x)$ where $\deg(f(x)) = 5$ possessing S -sequences of length 11. Alvarado ([1]) showed the existence of an infinite family of such curves with S -sequences of length 12. Moody ([12]) found an infinite number of

Edwards curves with an S -sequence of length 9. He also asked whether any such curve will allow an extension to an S -sequence of length 11. Bremner ([3]) showed that such curves do not exist. Also, Moody ([14]) found an infinite number of Huff curves with S -sequences of length 9, and Choudhry ([6]) extended Moody's result to find several Huff curves with S -sequences of length 11.

Now we consider the case when the elements of S form a *geometric progression*, Bremner and Ulas ([4]) obtained an infinite family of elliptic curves with S -sequences of length 4, and they also pointed out infinitely many elliptic curves with S -sequences of length 5. Ciss and Moody ([13]) found infinite families of twisted Edwards curves with S -sequences of length 5 and Edwards curves with S -sequences of length 4. When the elements of $S \subset \mathbb{Q}$ are *consecutive squares*, Kamel and Sadek ([9]) constructed infinitely many elliptic curves given by the equation $y^2 = ax^3 + bx + c$ with S -sequences of length 5. When the elements of $S \subset \mathbb{Q}$ are *consecutive cubes*, Çelik and Soydan ([7]) found infinitely many elliptic curves of the form $y^2 = ax^3 + bx + c$ with S -sequences of length 5.

In the present work, we consider the following families of elliptic curves due to the symmetry enjoyed by the equations defining them: (twisted) Edwards curves and (general) Huff curves. Given an arbitrary subset S of a number field k , we tackle the general question of the existence of infinitely many such curves with an S -sequence when there is no restriction on the elements of S . We provide explicit examples when the length of the S -sequence is 4, 5, or 6. This is achieved by studying the existence of rational points on certain quadratic and elliptic surfaces.

2. EDWARDS CURVES WITH S -SEQUENCES OF LENGTH 6

Throughout this work, k will be a number field unless otherwise stated.

An *Edwards curve* over k is defined by

$$(2.1) \quad E_d : x^2 + y^2 = 1 + dx^2y^2,$$

where d is a non-zero element in k . It is clear that the points $(x, y) = (-1, 0), (0, \pm 1), (1, 0) \in E_d(k)$. We show that given any set

$$S = \{s_{-1} = -1, s_0 = 0, s_1 = 1, s_2, s_3, s_4\} \subset k,$$

$s_i \neq s_j$ if $i \neq j$, there are infinitely many Edwards curves E_d that possess rational points whose x -coordinates are s_i , $-1 \leq i \leq 4$, i.e., the set S is realized as x -coordinates in $E_d(k)$. In other words, there are infinitely many Edwards curves that possess an S -sequence.

We start with assuming that s_2 is the x -coordinate of a point in $E_d(k)$, then one must have $y^2 = \frac{s_2^2 - 1}{s_2^2 d - 1}$, or $s_2^2 d - 1 = (s_2^2 - 1)p^2$ for some $p = 1/y \in k$.

Similarly, if s_3 is the x -coordinate of a point in $E_d(k)$, then $y^2 = \frac{s_3^2 - 1}{s_3^2 d - 1}$, or $s_3^2 d - 1 = (s_3^2 - 1)q^2$. So

$$d = \frac{(s_2^2 - 1)p^2 + 1}{s_2^2} = \frac{(s_3^2 - 1)q^2 + 1}{s_3^2}.$$

Thus we have the following quadratic curve

$$s_3^2 [(s_2^2 - 1)p^2 + 1] - s_2^2 [(s_3^2 - 1)q^2 + 1] = 0$$

on which we have the rational point $(p, q) = (1, 1)$. Parametrizing the rational points on the latter quadratic curve yields

$$p = \frac{2ts_2^2 - t^2s_2^2 - s_3^2 + s_2^2s_3^2 - 2ts_2^2s_3^2 + t^2s_2^2s_3^2}{-t^2s_2^2 + s_3^2 - s_2^2s_3^2 + t^2s_2^2s_3^2},$$

$$q = -\frac{(-1 + s_2^2)s_3^2 - 2t(-1 + s_2^2)s_3^2 + t^2s_2^2(-1 + s_3^2)}{-(-1 + s_2^2)s_3^2 + t^2s_2^2(-1 + s_3^2)}.$$

Therefore, fixing s_2 and s_3 in k , one sees that p and q lie in $k(t)$. Now we obtain the following result.

THEOREM 2.1. *Let $s_{-1} = -1$, $s_0 = 0$, $s_1 = 1$, s_2, s_3 and s_4 , $s_i \neq s_j$ if $i \neq j$, be a sequence in \mathbb{Z} such that*

$$h(s_2, s_3) = -3 + 4s_3^2 + s_2^4s_3^4 + s_2^2(4 - 6s_3^2) \neq 0$$

where either $g_1(s_2, s_3)/h(s_2, s_3)^2$ or $g_2(s_2, s_3)/h(s_2, s_3)^3$ are not integers, g_1 and g_2 are defined in (2.3). There are infinitely many Edwards curves described by

$$E_d : x^2 + y^2 = 1 + dx^2y^2, \quad d \in \mathbb{Q}$$

on which s_i , $-1 \leq i \leq 4$, are the x -coordinates of rational points in $E_d(\mathbb{Q})$. In other words, there are infinitely many Edwards curves that possess an S -sequence where $S = \{s_i : -1 \leq i \leq 4\}$.

PROOF. Substituting the value for p in $d = \frac{(s_2^2 - 1)p^2 + 1}{s_2^2}$ yields that

$$\begin{aligned} & (-t^2s_2^2 + s_3^2 - s_2^2s_3^2 + t^2s_2^2s_3^2)^2 d \\ &= (s_3^4 - 2s_2^2s_3^4 + s_2^4s_3^4) + (4s_3^2 - 8s_2^2s_3^2 + 4s_2^4s_3^2 - 4s_3^4 + 8s_2^2s_3^4 - 4s_2^4s_3^4)t \\ & \quad + (-4s_2^2 + 4s_2^4 - 4s_3^2 + 14s_2^2s_3^2 - 10s_2^4s_3^2 + 4s_3^4 - 10s_2^2s_3^4 + 6s_2^4s_3^4)t^2 \\ & \quad + (4s_2^2 - 4s_2^4 - 8s_2^2s_3^2 + 8s_2^4s_3^2 + 4s_2^2s_3^4 - 4s_2^4s_3^4)t^3 + (s_2^4 - 2s_2^2s_3^2 + s_2^4s_3^4)t^4. \end{aligned}$$

Thus, for fixed values of s_2 and s_3 , we have $d \in \mathbb{Q}(t)$.

Now we show the existence of infinitely many values of t such that s_4 is the x -coordinate of a rational point on E_d . In fact, we will show that t can be chosen to be the x -coordinate of a rational point on an elliptic curve with positive Mordell-Weil rank, hence the existence of infinitely many such

possible values for t . Forcing (s_4, r) to be a point in $E_d(\mathbb{Q})$ for some rational r yields that

$$(2.2) \quad r^2 = \frac{s_4^2 - 1}{s_4^2 d - 1} = (A_0 + A_1 t + A_2 t^2 + A_3 t^3 + A_4 t^4) / B(t)^2,$$

where $A_i \in \mathbb{Z}$ and $B(t) = -t^2 s_2^2 + t^2 s_2^2 s_3^2 + s_3^2 - s_2^2 s_3^2$. This implies that $A_0 + A_1 t + A_2 t^2 + A_3 t^3 + A_4 t^4$ must be a rational square. This yields the elliptic curve C defined by

$$z^2 = A_0 + A_1 t + A_2 t^2 + A_3 t^3 + A_4 t^4,$$

with the following rational point

$$(t, z) = (0, s_3^2(s_2^2 - 1)).$$

The latter elliptic curve is isomorphic to the elliptic curve described by the Weierstrass equation $E_{I,J} : y^2 = x^3 - 27Ix - 27J$ where

$$\begin{aligned} I &= 12A_0A_4 - 3A_1A_3 + A_2^2 \\ J &= 72A_0A_2A_4 + 9A_1A_2A_3 - 27A_1^2A_4 - 27A_0A_3^2 - 2A_2^3, \end{aligned}$$

see for example [16, §2]. The latter elliptic curve has the following rational point

$$P = (-12(-1 + s_2^2)(-1 + s_3^2)(-3 + s_2^2 + s_3^2), -216(-1 + s_2^2)^2(-1 + s_3^2)^2).$$

One notices that the coordinates of $3P$ are rational functions. Indeed,

$$(2.3) \quad 3P = \left(\frac{g_1(s_2, s_3)}{h(s_2, s_3)^2}, \frac{g_2(s_2, s_3)}{h(s_2, s_3)^3} \right), \quad \text{where } g_1, g_2 \in \mathbb{Q}[s_2, s_3]$$

and

$$h(s_2, s_3) = -3 + 4s_3^2 + s_2^4 s_3^4 + s_2^2(4 - 6s_3^2).$$

Hence, as long as $h(s_2, s_3) \neq 0$, and $g_1/h^2 \notin \mathbb{Z}$ or $g_2/h^3 \notin \mathbb{Z}$, one sees that $3P$ is a point of infinite order by virtue of Lutz-Nagell Theorem. Thus, P itself is a point of infinite order. It follows that $E_{I,J}$ is of positive Mordell-Weil rank. Since C is isomorphic to $E_{I,J}$, it follows that C is also of positive Mordell-Weil rank. Therefore, there are infinitely many rational points $(t, z) \in C(\mathbb{Q})$, each giving rise to a value for d , by substituting in (2.2), hence an Edwards curve E_d possessing the aforementioned rational points. That infinitely many of these curves are pairwise non-isomorphic over \mathbb{Q} follows, for instance, from [8, Proposition 6.1]. \square

3. TWISTED EDWARDS CURVES WITH S -SEQUENCES OF LENGTH 4

A *Twisted Edwards curve* over k is given by

$$(3.1) \quad E_{a,d} : ax^2 + y^2 = 1 + dx^2y^2,$$

where a and d are nonzero elements in k . Note that the point $(x, y) = (0, \pm 1) \in E_{a,d}(k)$. Given a set $\{u_0 = 0, u_1, u_2, u_3\} \subset k$, $u_i \neq u_j$ if $i \neq j$, we prove that

there are infinitely many twisted Edwards curves $E_{a,d}$ for which S is realized as the x -coordinates of rational points on $E_{a,d}$.

We begin by assuming that u_1 is the x -coordinate of a point in $E_{a,d}(k)$, then one must get $y^2 = \frac{au_1^2 - 1}{u_1^2 d - 1}$, or $u_1^2 d - 1 = (au_1^2 - 1)i^2$ for some $i \in k$.

Now, if u_2 is the x -coordinate of a point in $E_{a,d}(k)$, then $y^2 = \frac{au_2^2 - 1}{u_2^2 d - 1}$ or $u_2^2 d - 1 = (au_2^2 - 1)j^2$. So

$$d = \frac{(au_1^2 - 1)i^2 + 1}{u_1^2} = \frac{(au_2^2 - 1)j^2 + 1}{u_2^2}.$$

Hence we obtain the following quadratic surface

$$u_2^2 [(au_1^2 - 1)i^2 + 1] - u_1^2 [(au_2^2 - 1)j^2 + 1] = 0,$$

on which we have the rational point $(i, j) = (1, 1)$. Solving the above quadratic surface gives the following

$$i = \frac{-au_1^2 u_2^2 + u_2^2 + 2tau_1^2 u_2^2 - 2tu_1^2 - at^2 u_1^2 u_2^2 + u_1^2 t^2}{au_1^2 u_2^2 - u_2^2 - at^2 u_1^2 u_2^2 + u_1^2 t^2},$$

$$j = \frac{-2atu_1^2 u_2^2 + 2tu_2^2 + at^2 u_1^2 u_2^2 - u_1^2 t^2 + au_1^2 u_2^2 - u_2^2}{au_1^2 u_2^2 - u_2^2 - at^2 u_1^2 u_2^2 + u_1^2 t^2}.$$

Now we get the following result.

THEOREM 3.1. *Let $u_0 = 0$, u_1 , u_2 and u_3 , $u_i \neq u_j$ if $i \neq j$, be a sequence in \mathbb{Z} such that $h(u_1, u_2) \neq 0$, and either $g_1(s_2, s_3)/h(s_2, s_3)^2$ or $g_2(s_2, s_3)/h(s_2, s_3)^3$ are not integers, where h, g_1, g_2 are defined in (3.3). There are infinitely many twisted Edwards curves described by*

$$E_{a,d} : ax^2 + y^2 = 1 + dx^2 y^2, \quad d \in \mathbb{Q}, a \in \mathbb{Q}^\times \text{ is arbitrary}$$

on which u_i , $0 \leq i \leq 3$, are the x -coordinates of rational points in $E(\mathbb{Q})$. In other words, there are infinitely many twisted Edwards curves that possess an S -sequence where $S = \{u_i : 0 \leq i \leq 3\}$.

PROOF. Substituting the expression for i in $d = \frac{(au_1^2 - 1)i^2 + 1}{u_1^2}$ gives that

$$\begin{aligned} & (au_1^2 u_2^2 - u_2^2 - at^2 u_1^2 u_2^2 + u_1^2 t^2)^2 d \\ &= (u_1^4 a^3 u_2^4 - 2u_1^4 a^2 u_2^2 + u_1^4 a) t^4 + (-8au_1^2 u_2^2 + 4u_1^2 + 4u_1^2 a^2 u_2^4 - 4u_1^4 a \\ &\quad - 4u_1^4 a^3 u_2^4 + 8u_1^4 a^2 u_2^2) t^3 + (-4u_1^2 - 10u_1^2 a^2 u_2^4 + 14au_1^2 u_2^2 + 6u_1^4 a^3 u_2^4 \\ &\quad - 4u_2^2 - 10u_1^4 a^2 u_2^2 + 4u_1^4 a + 4au_2^4) t^2 + (4u_2^2 + 8u_1^2 a^2 u_2^4 - 8au_1^2 u_2^2 \\ &\quad + 4u_1^4 a^2 u_2^2 - 4au_2^4 - 4u_1^4 a^3 u_2^4) t + u_1^4 a^3 u_2^4 - 2u_1^2 a^2 u_2^4 + au_2^4. \end{aligned}$$

Then, assuming $(u_3, \ell) \in E(\mathbb{Q})$ yields

$$(3.2) \quad \ell^2 = \frac{au_3^2 - 1}{du_3^2 - 1} = (C_0 + C_1t + C_2t^2 + C_3t^3 + C_4t^4)/D(t)^2,$$

where $C_i \in \mathbb{Q}$ and $D(t) = au_1^2u_2^2 - u_2^2 - at^2u_1^2u_2^2 + u_1^2t^2$.

For the latter equation to be satisfied, one needs to find rational points on the elliptic curve C' defined by

$$z^2 = C_0 + C_1t + C_2t^2 + C_3t^3 + C_4t^4$$

that possesses the rational point

$$(t, z) = (0, u_2^2(au_1^2 - 1)).$$

The latter elliptic curve is isomorphic to the elliptic curve described by the Weierstrass equation $E_{I,J} : y^2 = x^3 - 27Ix - 27J$ where

$$I = 12C_0C_4 - 3C_1C_3 + C_2^2,$$

$$J = 72C_0C_2C_4 + 9C_1C_2C_3 - 27C_1^2C_4 - 27C_0C_3^2 - 2C_2^3,$$

see for example [16, §2]. The latter elliptic curve has the following rational point

$$Q =$$

$$(-12(-1 + au_2^2)(-1 + au_1^2)(-3 + au_2^2 + u_1^2), -216(-1 + au_2^2)^2(-1 + au_1^2)^2).$$

One notices that the coordinates of $3Q$ are rational functions. In fact,

$$3Q = \left(\frac{g_1(u_1, u_2)}{h(u_1, u_2)^2}, \frac{g_2(u_1, u_2)}{h(u_1, u_2)^3} \right), \quad \text{where } g_1, g_2 \in \mathbb{Q}[u_1, u_2]$$

and

$$(3.3) \quad \begin{aligned} h(u_1, u_2) = & -27 - 72u_1^2 + 36u_1^4 + 18u_1^2u_2^2 - 12u_1^4u_2^2 - 18u_2^4 + 12u_1^2u_2^4 \\ & + u_1^4u_2^4 - 2u_1^2u_2^6 + u_2^8 + a(36u_1^2 - 12u_1^4 - 24u_1^2(-3 + u_1^2) \\ & + 36u_2^2 + 72u_1^2u_2^2 - 24u_1^4u_2^2 - 12u_1^2u_2^4 + 4u_1^4u_2^4 \\ & - 4(-3 + u_1^2)u_2^6) + a^2(-144u_1^2u_2^2 + 36u_1^4u_2^2 + 18u_2^4 \\ & - 36u_1^2u_2^4 + 4u_1^4u_2^4 + 2u_1^2u_2^6 - 2u_2^8) + a^3(36u_1^2u_2^4 \\ & + 4(-3 + u_1^2)u_2^6) + a^4u_2^8. \end{aligned}$$

Therefore, as long as $h(u_1, u_2) \neq 0$ and $g_1/h^2 \notin \mathbb{Z}$ or $g_2/h^3 \notin \mathbb{Z}$, one sees that $E_{I,J}$ is of positive Mordell-Weil rank where the point Q is of infinite order. Since C' is isomorphic to $E_{I,J}$, it follows that C' is also of positive Mordell-Weil rank. Hence, there are infinitely many rational points $(t, z) \in C'(\mathbb{Q})$, each giving rise to a value for d , by substituting in (3.2), therefore a twisted Edwards curve $E_{a,d}$ possessing the aforementioned rational points. That infinitely many of these curves are pairwise non-isomorphic over \mathbb{Q} again follows from [8, Proposition 6.1]. \square

REMARK 3.2. Since $(0, -1), (0, 1)$ are rational points on any twisted Edwards curve, one can show that if $u_{-1} = -1, u_1 = 1, u_2, u_3$ and $u_4, u_i \neq u_j$ if $i \neq j$, is a sequence in \mathbb{Z} , there are infinitely many Edwards curves on which $u_i, i \in \{-1, 1, 2, 3, 4\}$, are the y -coordinates of rational points in $E_{a,d}(\mathbb{Q})$.

4. HUFF CURVES WITH S -SEQUENCES OF LENGTH 5

A Huff curve over a number field k is defined by

$$(4.1) \quad H_{a,b} : ax(y^2 - 1) = by(x^2 - 1),$$

with $a^2 \neq b^2$. Note that the points $(x, y) = (-1, \pm 1), (0, 0), (1, \pm 1)$ are in $H_{a,b}(k)$. We prove that given $s_{-1} = -1, s_0 = 0, s_1 = 1, s_2, s_3 \in k, s_i \neq s_j$ if $i \neq j$, there are infinitely many Huff curves on which these numbers are realized as the x -coordinates of rational points.

Assuming (s_2, p) and (s_3, q) are two points on $H_{a,b}$ yields

$$(4.2) \quad as_2(p^2 - 1) = bp(s_2^2 - 1),$$

and

$$(4.3) \quad as_3(q^2 - 1) = bq(s_3^2 - 1),$$

respectively. Using (4.2) and (4.3), one obtains

$$\frac{s_2(p^2 - 1)}{s_3(q^2 - 1)} = \frac{p(s_2^2 - 1)}{q(s_3^2 - 1)},$$

therefore, one needs to consider the curve

$$C' : Apq^2 - Ap - Bqp^2 + Bq = 0,$$

where $A = s_3s_2^2 - s_2$ and $B = s_2s_3^2 - s_2$. Dividing both sides of the above equality by q^3 gives

$$A\frac{p}{q} - A\frac{p}{q}\frac{1}{q^2} - B\left(\frac{p}{q}\right)^2 + B\frac{1}{q^2} = 0.$$

Substituting $x = \frac{p}{q}$ and $y = \frac{1}{q^2}$ in the above equation yields the following quadratic curve

$$Ax - Axy - Bx^2 + By = 0,$$

on which we have the rational point $(x, y) = (1, 1)$. Parametrizing the rational points on the latter quadratic curve gives

$$(4.4) \quad x = \frac{Bt - B}{At + B},$$

$$(4.5) \quad y = \frac{At(1-t) + B(1-t)^2}{At + B}.$$

Now we have the following result.

THEOREM 4.1. *Let $s_{-1} = -1, s_0 = 0, s_1 = 1, s_2, s_3, s_m \neq s_n$ if $m \neq n$, be a sequence in \mathbb{Z} such that*

$$h = -4 + A^2 - 3AB + B^2 \neq 0$$

where A and B are defined as above, and either g_1/h^2 or g_2/h^3 are not integers, where g_1, g_2 are defined in (4.6). There are infinitely many Huff curves described by

$$H_{a,b} : ax(y^2 - 1) = by(x^2 - 1), \quad a, b \in \mathbb{Q}, \quad a^2 \neq b^2$$

on which $s_m, -1 \leq m \leq 3$, are the x -coordinates of rational points in $H_{a,b}(\mathbb{Q})$. In other words, there are infinitely many Huff curves that possess an S -sequence where $S = \{s_i : -1 \leq i \leq 3\}$.

PROOF. Using the equalities (4.4) and (4.5), we obtain the following

$$p^2 = \frac{x^2}{y} = \frac{B^2(-1+t)}{(B(-1+t) - At)(B + At)},$$

$$q^2 = \frac{1}{y} = \frac{(B + At)}{(-1+t)(B(-1+t) - At)}.$$

In both cases we need $(B + At)(-1+t)(B(-1+t) - At)$ to be a square or in other words we need t to be the x -coordinate of a rational point on the elliptic curve C'' defined by

$$z^2 = (At + B)(t - 1)(t(B - A) - B),$$

with the following k -rational point $(t, z) = (0, B)$. The latter curve can be described by the following equation

$$Y^2 = X^3 + ((B - A)^2 - AB)X^2 - 2AB(B - A)^2X + A^2B^2(B - A)^2,$$

where $A(B - A)t = X$ and $A(B - A)z = Y$. This curve has the rational point

$$R = (X, Y) = (0, AB(B - A)).$$

Observing that

$$(4.6) \quad 3R = \left(\frac{g_1(A, B)}{h(A, B)^2}, \frac{g_2(A, B)}{h(A, B)^3} \right)$$

where $h(A, B) = -4 + A^2 - 3AB + B^2$, one concludes as in the proof of Theorem 2.1. \square

5. GENERAL HUFF CURVES WITH S -SEQUENCES OF LENGTH 4

A general Huff curve over a number field k is defined by

$$(5.1) \quad G_{a,b} : x(ay^2 - 1) = y(bx^2 - 1),$$

where $a, b \in k$ and $ab(a - b) \neq 0$. It is clear that the point $(x, y) = (0, 0) \in G_{a,b}(k)$. We show that given $u_0 = 0, u_1, u_2, u_3$ in $k, u_i \neq u_j$ if $i \neq j$, there

are infinitely many general Huff curves over which these points are realized as the x -coordinates of rational points.

We start by assuming that if u_1 is the x -coordinates of a point in $G_{a,b}(k)$, then one must have $\frac{ay^2 - 1}{y} = \frac{bu_1^2 - 1}{u_1}$ or $\frac{a - i^2}{i} = \frac{bu_1^2 - 1}{u_1}$ for some $i \in k$.

Similarly, if u_2 is the x -coordinate of a point in $G_{a,b}$, then $\frac{ay^2 - 1}{y} = \frac{bu_2^2 - 1}{u_2}$ or $\frac{a - j^2}{j} = \frac{bu_2^2 - 1}{u_2}$ for some $j \in k$. Thus, one obtains

$$a = \frac{(bu_1^2 - 1)i + u_1i^2}{u_1} = \frac{(bu_2^2 - 1)j + u_2j^2}{u_2},$$

which gives the following quadratic curve

$$S : Ai^2 + Bj^2 + Ciz + Djz = 0,$$

where $A = -u_1u_2$, $B = u_1u_2$, $C = -u_1^2u_2b + u_2$, $D = bu_1u_2^2 - u_1$. Then consider the line

$$mP + nQ = (np : nq : m + nr)$$

connecting the rational points $P = (i : j : z) = (0 : 0 : 1)$ and $Q = (p : q : r)$ lying on $S \subset \mathbb{P}^2$. The intersection of S and $mP + nQ$ yields the quadratic equation

$$n^2(Ap^2 + Bq^2 + Cpr + Dqr) + mn(Cp + Dq) = 0.$$

Using P and Q lying on S , one solves this quadratic equation and obtains formulae for the solution $(i : j : z)$ with the following parametrization:

$$\begin{aligned} i &= np = Cp^2 + Dpq, \\ j &= nq = Cpq + Dq^2, \\ z &= m + nr = -Ap^2 - Bq^2. \end{aligned}$$

Now we obtain the following result.

THEOREM 5.1. *Let $u_0 = 0$, u_1 , u_2 and u_3 , $u_i \neq u_j$ if $i \neq j$, be a sequence in k . There are infinitely many general Huff curves described by*

$$G_{a,b} : x(ay^2 - 1) = y(bx^2 - 1), \quad a, b \in k, \quad ab(a - b) \neq 0.$$

on which u_i , $0 \leq i \leq 3$, are the x -coordinates of rational points in $G_{a,b}(k)$. In other words, there are infinitely many general Huff curves that possess an S -sequence where $S = \{u_i : 0 \leq i \leq 3\}$.

PROOF. Substituting the value for i in $a = \frac{(bu_1^2 - 1)i + u_1i^2}{u_1}$ yields that

$$\begin{aligned} a &= u_2^2 (bu_1^2 - 1)^2 p^4 - 2u_1u_2 (bu_2^2 - 1) (bu_1^2 - 1) p^3 q \\ &\quad + u_1^2 (bu_2^2 - 1)^2 p^2 q^2 - \frac{u_2 (bu_1^2 - 1)^2}{u_1} p^2 + (bu_2^2 - 1) (bu_1^2 - 1) pq. \end{aligned}$$

Now we assume that $(u_3, \ell) \in G_{a,b}(k)$. This yields that

$$\begin{aligned} & pu_3 (bp^2u_1^3u_2 - bpqu_1^2u_2^2 - p^2u_1u_2 + pqu_1^2 - bu_1^2 + 1) \\ & (bpu_1^2u_2 - bqu_1u_2^2 - pu_2 + qu_1) \ell^2 - u_1 (bu_3^2 - 1) \ell - u_1u_3 = 0. \end{aligned}$$

This can be rewritten as

$$\begin{aligned} & Z^2(b^2p^4u_1^5u_2^2u_3 - 2bp^4u_1^3u_2^2u_3 - b^2p^2u_1^4u_2u_3 + p^4u_1u_2^2u_3 + 2bp^2u_1^2u_2u_3 \\ & - p^2u_2u_3) + qZ(-2b^2p^3u_1^4u_2^3u_3 + 2bp^3u_1^4u_2u_3 + 2bp^3u_1^2u_2^3u_3 + b^2pu_1^3u_2^2u_3 \\ & - 2p^3u_1^2u_2u_3 - bpu_1^3u_3 - bpu_1u_2^2u_3 + pu_1u_3) + q^2p^2u_1^3u_3(bu_2^2 - 1)^2 \\ & - TZu_1 (bu_3^2 - 1) - T^2u_1u_3 = 0, \end{aligned}$$

where $T = 1/\ell$. One sees that the rational point $P = (q : T : Z) = (1 : 0 : u_1(-1 + bu_2^2)/pu_2(-1 + bu_1^2))$ lies on the quadratic curve above, hence we may parametrize the rational points on the quadratic curve above. This is obtained by considering the intersection of the line $dP + eQ$ where $Q = (q_1 : q_2 : q_3)$ is a point on the quadratic curve. In fact, this yields that

$$\begin{aligned} d = & pu_2(bu_1^2 - 1)(q_3^2b^2p^4u_1^5u_2^2u_3 - 2q_3^2bp^4u_1^3u_2^2u_3 \\ & - q_3^2b^2p^2u_1^4u_2u_3 + q_3^2p^4u_1u_2^2u_3 + 2q_3^2bp^2u_1^2u_2u_3 \\ & - q_3^2p^2u_2u_3 - u_1q_2q_3bu_3^2 + u_1q_2q_3 + p^2u_1^3u_3q_1^2b^2u_2^4 \\ & - 2p^2u_1^3u_3q_1^2bu_2^2 + p^2u_1^3u_3q_1^2 - 2q_1q_3b^2p^3u_1^4u_2^3u_3 \\ & + 2q_1q_3bp^3u_1^4u_2u_3 + 2q_1q_3bp^3u_1^2u_2^3u_3 + q_1q_3b^2pu_1^3u_2^2u_3 \\ & - 2q_1q_3p^3u_1^2u_2u_3 - q_1q_3bpu_1^3u_3 - q_1q_3bpu_1u_2^2u_3 + q_1q_3pu_1u_3 \\ & - u_1u_3q_2^2), \\ e = & u_1(bu_2^2 - 1)(-pu_1^3u_3q_1b^2u_2^2 + p^2u_3q_3u_2b^2u_1^4 + pu_1u_3q_1bu_2^2 \\ & - 2p^2u_3q_3u_2bu_1^2 + pu_1^3u_3q_1b + u_1q_2bu_3^2 + p^2u_3q_3u_2 - u_1q_2 - pu_1u_3q_1). \end{aligned}$$

□

ACKNOWLEDGEMENTS.

We would like to thank the referees for carefully reading our manuscript and for giving such constructive comments which substantially helped improving the presentation of the paper.

REFERENCES

- [1] A. Alvarado, *An arithmetic progression on quintic curves*, J. Integer Seq. **12** (2009), Article 09.7.3., 6 pp.
- [2] A. Bremner, *On arithmetic progressions on elliptic curves*, Experiment Math. **8**, (1999), 409–413.
- [3] A. Bremner, *Arithmetic progressions on Edwards curves*, J. Integer Seq. **16** (2013), Article 13.8.5., 5 pp.

- [4] A. Bremner and M. Ulas, *Rational points in geometric progressions on certain hyper-elliptic curves*, Publ. Math. Debrecen **82** (2013), 669–683.
- [5] G. Campbell, *A note on arithmetic progressions on elliptic curves*, J. Integer Seq. **6** (2003), Article 03.1.3., 5 pp.
- [6] A. Choudhry, *Arithmetic progressions on Huff curves*, J. Integer Seq. **18** (2015), Article 15.5.2., 9 pp.
- [7] G. S. Çelik and G. Soydan, *Elliptic curves containing sequences of consecutive cubes*, Rocky Mountain J. Math. **48** (2018), 2163–2174.
- [8] H. Edwards, *A normal form for elliptic curves*, Bull. Amer. Math. Soc. (N.S.) **44** (2007), 393–422.
- [9] M. Kamel and M. Sadek, *On sequences of consecutive squares on elliptic curves*, Glas. Mat. Ser. III **52(72)** (2017), 45–52.
- [10] J.-B. Lee and W. Y. Vélez, *Integral solutions in arithmetic progression for $y^2 = x^3 + k$* , Period. Math. Hungar. **25** (1992), 31–49.
- [11] A. J. Macleod, *14-term arithmetic progressions on quartic elliptic curves*, J. Integer Seq. **9** (2006), Article 06.1.2., 4 pp.
- [12] D. Moody, *Arithmetic progressions on Huff curves*, Ann. Math. Inform. **38** (2011), 111–116.
- [13] A. A. Ciss and D. Moody, *Geometric progressions on elliptic curves*, Glas. Mat. Ser. III **52(72)** (2017), 1–10.
- [14] D. Moody, *Arithmetic progressions on Edwards curves*, J. Integer Seq. **14** (2011), Article 11.1.7., 4 pp.
- [15] J. H. Silverman, *The arithmetic of elliptic curves*, Springer, Dordrecht, 2009.
- [16] M. Stoll and J. E. Cremona, *Minimal models for 2-coverings of elliptic curves*, LMS J. Comput. Math. **5** (2002), 220–243.
- [17] M. Ulas, *A note on arithmetic progressions on quartic elliptic curves*, J. Integer Seq. **8** (2005), Article 05.3.1., 5 pp.
- [18] M. Ulas, *On arithmetic progressions on genus two curves*, Rocky Mountain J. Math. **39** (2009), 971–980.

G. S. Çelik
 Department of Mathematics
 Bursa Uludağ University
 16059 Bursa
 Turkey
E-mail: gamzesavascelik@gmail.com

M. Sadek
 Faculty of Engineering and Natural Sciences
 Sabancı University
 34956 Tuzla, İstanbul
 Turkey
E-mail: mmsadek@sabanciuniv.edu

G. Soydan
 Department of Mathematics
 Bursa Uludağ University
 16059 Bursa
 Turkey
E-mail: gsoydan@uludag.edu.tr

Received: 22.9.2018.

Revised: 5.3.2019.