



RESEARCH ARTICLE

JOHNSONGRASS MOSAIC VIRUS INFECTING SORGHUM IN BRAZIL

*¹Isabel Regina Prazeres de Souza, ¹Beatriz de Almeida Barros, ²André da Silva Xavier, ³Samanta Gabriela Medeiros Carvalho, ¹Elizabeth de Oliveira Sabato, ⁴Isabella Aparecida Maia Gonçalves, ¹Roberto Willians Noda and ¹José Avelino Santos Rodrigues

¹Embrapa Milho e Sorgo, Rod. MG 424 km 45, Sete Lagoas, MG, 35701-970, Brazil

²Post-doctoral researcher at Embrapa Milho e Sorgo, Rod. MG 424 km 45, Sete Lagoas, MG, 35701-970, Brazil

³MSc student at Departamento de Genética, Universidade Federal de Lavras, Lavras, MG, 37200-000, Brazil

⁴BS student in Biology from Centro Universitário de Sete Lagoas-UNIFEMM, Sete Lagoas, MG 35701-242, Brazil

ARTICLE INFO

Article History:

Received 20th September, 2017

Received in revised form

04th October, 2017

Accepted 11th November, 2017

Published online 31st December, 2017

Key words:

Potyvirus,
Capsid protein,
Mosaic,
RT-PCR,
Host range.

ABSTRACT

Sorghum bicolor (L.) Moench is cultivated in several tropical and subtropical regions in the world. Among the diseases, the mosaic caused by potyvirus is an important constraint for the agricultural production causing reduction in grain and forage sorghum production. In Brazil, only *Sugarcane mosaic virus* (SCMV) had previously been reported as the potyvirus species causing mosaic in sorghum and maize. A survey was carried out in sorghum plantations of the State of Minas Gerais, Brazil, during the 2014/2015 crop season for monitoring mosaic disease. Samples of sorghum plants expressing virus disease symptoms were collected for molecular analyzes. Molecular characterization of coat protein (CP) of the potyviruses naturally infecting sorghum, allowed us to identify the *Johnsongrass mosaic virus* (JGMV) as a new causal agent of mosaic disease in sorghum in Brazil. The sequences of the Brazilian JGMV sorghum-infecting (JGMV-Sr) isolates were deposited in the GenBank under the accession numbers KY952241, KY952242, and KY952243. Comparisons of the CP gene sequences of these Brazilian JGMV-Sr isolates revealed high nucleotide (nt) and amino acid (aa) sequence identities, ranging from 97.93 to 98.23%, and 99.12 to 99.20%, respectively, with the U07218.1 (JGMV-MDKS1) isolate. The Brazilian JGMV-Sr isolates were distinct from the Brazilian forage grasses-infecting (JGMV-Fg) isolates (KT833782 and KT289893). Transmission evaluations showed susceptibility of the teosinte, *Sorghum verticilliflorum* and *Sorghum bicolor* (L.) Moench, except line QL3. Maize and sugarcane genotypes were not infected by the Brazilian JGMV-Sr isolate. However, it is important to test more genotypes. This is the first report showing the identification and molecular characterization of the JGMV species naturally infecting sorghum at field conditions, expanding the knowledge about the dynamic and range of the mosaic causal agent for this crop in Brazil.

Copyright © 2017, Isabel Regina Prazeres de Souza et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Isabel Regina Prazeres de Souza, Beatriz de Almeida Barros, André da Silva Xavier et al. 2017. "Johnsongrass mosaic virus infecting sorghum in Brazil", *International Journal of Current Research*, 9, (12), 63415-63422.

INTRODUCTION

Sorghum bicolor (L.) Moench is cultivated in several tropical and subtropical regions in the world, and is the fifth major cereal in terms of production and acreage, after wheat, rice, maize and barley (FAO, 1999). Sorghum has water stress adaptive characteristics, which explains its cultivation in millions of hectares, in countries from Africa, Asia, Oceania and Americas (FAO, 1999). Among the diseases, the viruses are important constraint for the agricultural production worldwide, and the mosaic caused by potyviruses is widely

disseminated affecting *Poaceae* species. In susceptible sorghum cultivars, mosaic causes reduction in grain and forage production, and the symptoms can be expressed as typical mosaic or leaf necrosis depending on the genotype (Silva et al., 2012). Six species of potyviruses have been identified worldwide, causing mosaic symptoms in cultivated and weed grasses, including: *Sugarcane mosaic virus* (SCMV), *Sorghum mosaic virus* (SrMV), *Maize dwarf mosaic virus* (MDMV), *Johnsongrass mosaic virus* (JGMV) (Shukla et al., 1994), *Zea mosaic virus* (ZeMV) (Sheifers et al., 2000), and *Pennisetum mosaic virus* (PenMV) (Deng et al., 2008). In Brazil, mosaic in sorghum has as causal agent the SCMV species which strain constitutes a monophyletic group distinct from the others identified worldwide so far (Souza et al., 2012). Silva et al

*Corresponding author: Isabel Regina Prazeres de Souza, Embrapa Milho e Sorgo, Rod. MG 424 km 45, Sete Lagoas, MG, 35701-970, Brazil.

(2016) and Camelo-Garcia *et al.* (2016) identified isolates of the JGMV species infecting forage grasses, *Pennisetum purpureum* and *Panicum maximum* in Brazil, deposited in GenBank under numbers KT833782 and KT289893, respectively. However, none of these two isolates were identified naturally infecting maize or sorghum at field conditions. In addition, there is evidence of the presence of potyvirus species in maize whose identification is still necessary (Oliveira *et al.*, 2005). Maize aphid, *Rhopalosiphum maidis* (Fitch, 1856) (Hemiptera: Aphididae) is the most efficient vector in potyviruses transmission, and in Brazil is found mainly in regions where sorghum and maize are grown on the second harvesting (off-season), causing economic damages (Goussain, 2001). However, the green aphid, *Schizaphis graminum* (Rondani, 1852) (Hemiptera: Aphididae), considered an important pest of sorghum, also transmits the viruses (Cruz and Vendramin, 1995). The succession of grasses, maize and sorghum, and the increased incidence of vector in the off-season, are factors that contribute for the dissemination of the mosaic disease in both crops. Losses of up to 50% in production can be caused by this virosis (Waquil *et al.*, 1996). A survey was carried out in sorghum plantations of the State of Minas Gerais, Brazil, during the 2014/2015 crop season for monitoring mosaic disease. Samples of sorghum plants expressing virus disease symptoms were collected for molecular analyzes. Among the samples we identified JGMV isolates infecting grain and forage sorghums [*Sorghum bicolor* (L.) Moench], respectively, from Paracatu and Felixlândia municipalities. Here, we are reporting for the first time the occurrence of the JGMV species naturally infecting sorghum crops under field conditions in Brazil.

mosaic symptoms (i.e., small chlorotic areas interspersed with green areas (Figure 1 A-C) had their foliar tissue collected and the viruses identified through molecular analyzes. The molecular identification of the viral species had two steps, first the RNA was extracted from the leaf tissue using the RNeasy® Plant Mini Kit (Qiagen) and cDNA synthesized from 1.0 µg total RNA using oligo(dT)₁₈ and the SuperScript® III First-Strand Synthesis System (Invitrogen). In the first step, primers that amplify the conserved region of the CP of potyviruses were used PZEO1 x PZEO2 (Seifers *et al.*, 2000). Then, the positive samples were submitted to PCR reactions with primers specific for the region that encompasses the CP and a partial sequence of the nuclear inclusion protein (NIB) for all the six potyviruses from the mosaic complex (Table 1 from Souza and Barros 2016). The PCR reactions were also performed according to Souza and Barros, 2016. The primers set combinations PJG_F X PSR (Jiang and Zhou, 2002), and PJG_F (Jiang and Zhou, 2002) X PJG_R (design by the author Barros, B.A.) (Table 1) resulted in amplification of the JGMV in the samples with fragment sizes of the 813 bp and 414 bp, respectively. These primer sets amplify the N-terminal portion of the CP, allowing the identification of the JGMV isolates by sequencing (Shukla *et al.*, 1989). Amplicons were purified using ExoSAP-IT for PCR Product Cleanup (USB) and sequenced. The obtained sequences were analyzed using the Sequencher 1.4.1 software, performed by a two-step PCR. Sequence similarity searches against the NCBI sequence database (GenBank), allowed the identification of the JGMV species as the causal agent of the sorghum mosaic disease in Brazil. The JGMV isolate 7 infecting grain sorghum and the isolates 48 and 49 infecting forage sorghum were collected in




Figure 1. Mosaic symptoms in the leaves of grain sorghum (A), and forage sorghums (B and C) under field conditions, respectively, collected during the survey in Paracatu and Felixlândia municipalities of the Minas Gerais State (MG), Brazil

Table 1. Sequences of primers used in RT-PCR analysis

Potyvirus	Primer Identification	Sequence 5' – 3' ^a	Reference
SCMV, SrMV, MDMV and JGMV	PSR ^c	CAGCTGTGTGBCSTCTGTATT	JIANG & ZHOU (2002)
JGMV	PJG ^b -F	AAACCAGCTAGTGGTGAAGGC	JIANG & ZHOU (2002)
JGMV	PJG ^d -R	CACCAGACCATTAATCCGTTC	Design by author Barros, B. A.

^a – B= (G, C ou T); K = (G ou T); S = (G ou C); N = (A, G, C ou T); Y = (C ou T); R = (A ou G).

^b – Forward primer specific for determined potyviruses.

^c – Reverse primer common for all potyviruses.

^d – Reverse primer design by author Barros, B.A.

MATERIALS AND METHODS

Leaf sampling and virus identification

During the survey for common mosaic viral disease in sorghum crops carried out in the state of Minas Gerais (MG, Brazil) in the 2014/2015 crop season, plants expressing typical

the municipalities, respectively, Paracatu and Felixlândia, both in the state of Minas Gerais, Brazil. Sequences of the gene and the translated coat protein (CP) of these isolates were deposited in GenBank under the accession numbers KY952241 (isolate 7), KY952242 (isolate 48) and KY952243 (isolate 49).




Figure 2. Mosaic symptoms induced in leaves of the plant species mechanically inoculated with Brazilian JGMV-Sr isolate: Sorghum BRS 506 (A), *Euchlaena mexicana* (B), and *Sorghum verticilliflorum* (C)

Table 2. Pairwise percent coat protein nucleotide sequence identity (above diagonal) and amino acid (below diagonal) sequence identity among Brazilian JGMV-Sr isolates, Brazilian JGMV-Fg isolates and USA isolate of JGMV. The GenBank accession numbers are followed by the host species and country name or abbreviation

GenBankAccessions	U07218.1	KY952241	KY952242	KY952243	KT289893	KT833782
U07218.1 MDKSI-USA	-	98.23	97.93	98.01	79.18	78.82
KY952241-Sorghum_7-Brazil	99.20	-	98.82	98.74	77.36	76.43
KY952242-Sorghum_48-Brazil	99.12	100.00	-	99.71	79.09	78.50
KY952243-Sorghum_49-Brazil	99.16	100.00	100.00	-	78.12	77.14
KT289893-Panicum maximum -Brazil	81.52	81.20	84.51	83.12	-	95.87
KT833782-Pennisetum purpureum-Brazil	80.94	80.80	84.96	82.70	96.53	-

Mechanical transmission of the Brazilian JGMV-Sr isolate

The study was conducted at Embrapa Maize and Sorghum, experimental station, Sete Lagoas, Brazil, and carried out in a greenhouse with preventive insecticide applications to avoid the occurrence of insects. The experiment was conducted in pots filled with 25 kg of a Latosol, typical Brazilian Cerrado soil. Soil pH was corrected by liming and the fertilization was performed according to soil analysis. A rate equivalent to 400 kg ha⁻¹ of the formula 08-28-16 + 0.3% Zn and 40 kg ha⁻¹ of FTE BR12 was mixed to the soil in the pots. The experimental unit constituted of one pot containing five seedlings/pot. In the transmission test were evaluated *Sorghum bicolor* (L.) Moench: varieties BRS Ponta Negra and BRS 506, and the lines BRS 007B, ATF54, QL3, 9910032, 9929020, *Zea mays* L.: inbred lines L18, L19, L520, L3, and L2841 and a commercial hybrid; *Saccharum* spp: RB928064 and RB855536, besides *Sorghum verticilliflorum* and *Euchlaena mexicana*. Treatments consisted of the (i) inoculum of the Brazilian JGMV-Sr isolate, and (ii) a negative control for inoculation (mock treatment). The presence of the JGMV was confirmed molecularly and the inoculum was prepared using leaves of the symptomatic plants macerated in cooled phosphate buffer (10 mM, pH 7.0), at a ratio of 1:3 (weight/volume) (Souza *et al.*, 2008). Carborundum 600 mesh (Sigma-Aldrich) was added to the inoculum solution, which was kept on ice throughout the inoculation process. The first inoculation was performed in the middle to the basal part of the leaves when the seedlings had three to four leaves. The same mechanical procedure was adopted for the mock treatment using phosphate buffer (10 mM, pH 7.0) containing 600 mesh carborundum. Two inoculations were performed with a week interval. Phenotypic evaluations were initiated one week after the first inoculation and were repeated weekly along one month and five days period. Phenotypes were

evaluated based on mosaic symptoms and classified for the presence and absence (Figure 2).

Pairwise sequence alignment and phylogenetic analysis

In our analysis, we used a data set of 32 JGMV CP nucleotide sequences including isolates downloaded from NCBI database, and our three Brazilian JGMV-Sr isolates. Additionally, five other species as outgroups: *Sugarcane mosaic virus* (SCMV_KR108212.1), *Maize dwarf mosaic virus A* (MDMV-A_U07216.1), *Zea mosaic virus* (ZemV_AF228693.1), *Pennisetum mosaic virus* (PenMV_JX070151.1), and *Sorghum mosaic virus* (SrMV_KM025045.1). Apart from JGMV isolates described here, the sequences were downloaded from the NCBI database and comprised the complete ORF of the protein or were at least 700 base pairs (bp) in length. Alignments were obtained and conferred using ClustalOmega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>) and MUSCLE (executed in MEGA7 - Kumar, Stecher and Tamura, 2016). Pairwise comparisons were performed using SDT v. 1.2 (Muhire *et al.*, 2014). Phylogenetic reconstruction was carried out with MEGA7 (Kumar, Stecher, and Tamura, 2016) based on the Neighbor-Joining method (Saitou and Nei, 1987). The evolutionary distances were computed using the Kimura 2-parameter method (Kimura, 1980). The tree was generated using FigTree (<http://tree.bio.ed.ac.uk/software/figtree/>) and the percentage of replicate trees in the bootstrap test (1000 replicates) was shown next to the branches (Felsenstein, 1985).

RESULTS

Molecular characterization, sequencing and phylogenetic analysis of Brazilian JGMV-Sr isolates

Brazilian JGMV-Sr isolates showed the highest percentage of identities at coat protein nucleotide and amino acid sequences

with the JGMV-MDKS1, which values ranged from 97.93 to 98.23%, and 99.12 to 99.20%, respectively (Table 2). When compared with the Brazilian JGMV-Fg isolates (KT833782 and KT289893), these values ranged from 76.46 to 79.09%, and 80.80 to 84, 96%.

Alignments of the deduced amino acid (aa) sequences of the CP N-terminal showed the Brazilian JGMV-Sr isolates (KY952241, KY952242, and KY952243) and the JGMV-MDKS1 strain (U07218.1) with five aa residues less than Brazilian JGMV-Fg isolates (KT833782 and KT289893), residues 21, 32, 33, 45 and 46 (Figure 3). Also these two latter isolates showed several amino acids substitutions, as the MVWCIENG potyvirus conserved motif of the CP core was changed to MVWAIENG in relation to the Brazilian JGMV-Sr isolates, and JGMV-MDKS1 strain from USA.

other composed only by Brazilian JGMV-Fg isolates (KT833782, KT289893 and KC333416.1). Additionally, a third clade, grouped JGMV isolates from Australia.

Mechanical transmission of the Brazilian JGMV-Sr isolate

Among the sorghum genotypes, except the line QL3, the varieties BRS Ponta Negra and BRS 506, and lines BRS 007B, ATF54, 9910032, and 9929020 were infected and expressed the typical mosaic symptoms. Also, the *Poaceae* species *Euchlaena mexicana* and weed *Sorghum verticilliflorum* were infected and showed similar symptoms. All the five maize genotypes, inbred lines L18, L19, L520, L3, and L2841, and the two *Saccharum* spp. genotypes, varieties RB928064 and RB855536, were not infected. Variations in relation to mosaic symptoms expression were verified: (i) mosaic more confined to longitudinal sectors as observed at field (Figure 1A and C),




Figure 3. Multiple alignments of CP N-terminal amino acid sequences of JGMV strains from NCBI database and Brazilian JGMV-Sr isolates from this work. The accession numbers are followed by host plant species and country abbreviation, BR=Brazil and USA= United States of America. The conserved motifs (MVWCIENG, QMKAAA) of the potyviruses CP core are highlighted inside a box and the position with divergent amino acids is demarcated by black arrows

The phylogenetic tree (Figure 4) and color-coded pairwise identity matrix (Figure 5) showed the JGMV accessions from Brazil divided into two groups, one composed by the JGMV-Sr isolates (KY952241, KY952242 and KY952243) plus the JGMV isolates from USA (U07218.1 and U07217.1), and the

and greenhouse conditions, BRS 506 mechanically inoculated (Figure 1D), and (ii) Typical mosaic, chlorotic areas interspersed with green ones exhibited at field (Figure 1B) and greenhouse conditions, *Euchlaena mexicana* and weed *Sorghum verticilliflorum* mechanically inoculated (Figure 2B and C).




Figure 4. Phylogenetic tree derived from the CP nucleotide sequences of the 32 *Johnsongrass mosaic virus* (JGMV), plus five outgroups potyviruses. The tree was generated using MEGA 7 by the neighbor-joining method with 1,000 bootstrap replicates. Bootstrap values are shown at the internodes, and NCBI accession numbers are indicated. The Brazilian JGMV-Sr isolates from this study are in bold




Figure 5. Color-coded pairwise identity matrix based on the CP nucleotide coding sequences of Brazilian JGM-Sr isolates from this study, and 32 JGMV plus five outgroups potyviruses downloaded from GenBank. Each colored cell represents a percentage identity score between two sequences. A color key indicates the correspondence between pairwise identities and the colors displayed in the matrix

DISCUSSION

Distinct variations among the Brazilian JGMV-Sr isolates (GenBank KY952241, KY952242, and KY952243), and the Brazilian JGMV-Fg isolates, *Panicum maximum* (GenBank KT289893) and *Pennisetum purpureum* (GenBank KT833782), were demonstrated through DNA sequencing and alignment of the CP sequences, with average percentage identities of 77.77% and 82.88% at the nucleotide and correspondent amino acid sequences, respectively (Table 2 and Figure 4). The nucleotide identities of the Brazilian JGMV-Sr isolates meet the criteria for JGMV *Potyvirus* species classification due identity above the 76-77% for the CP (Adams et al., 2012). As mention by Shukla et al. (1991) the correct identification of a pathogen is the first step toward studies to allow its eventual control. The primers pair described by Camelo-Garcia et al. (2016) and Silva et al. (2016) to amplify the CP gene sequences of the Brazilian JGMV-Fg isolates from *Panicum maximum* and *Pennisetum purpureum*, respectively, does not produce amplicons using the Brazilian JGMV-Sr isolates. It is due variations caused by a mismatch at the 3' end of their forward primer (CAAGCCCCATACTTGTCGG), where only the underlined nucleotide sequence has complementary sequence to Brazilian JGMV-Sr isolates. In addition, the primers pair used by Lenardon and Giolliti (2004) to identify the JGMV in Argentina was not able to amplify the same fragment from samples infected with the Brazilian JGMV-Sr isolates. The highly conserved motif in the CPs potyviral, MVWCIENG (Pappu et al., 1993) was preserved in the Brazilian JGMV-Sr isolates, and "C" was replaced by "A", MVWAIENG, in the Brazilian JGMV-Fg isolates (Figure 3). The N-hypervariable region of the Brazilian JGMV-Sr isolates and JGMV-MDKS1 has a dinucleotide repeat (CA)₆, as part of the deduced ATHTQ peptide sequence of the CP, and is five amino acids shorter than Brazilian JGMV-Fg isolates. Zhao et al. (2011) found in potyviruses genome the predominance of the repeats AC/CA, AG/GA and AAG/GAA, and mentioned that microsatellites may have many important functions in terms of genomic organization and species evolution. A high mutation frequency of microsatellites allow to increase the diversity in pathogen populations (Pearson et al., 2005), and some of them may promote potyviruses species evolving to adapt to their hosts (Zhao, 2011). Majority of the CP aminoacid substitutions was verified in the amino terminal regions among the Brazilian JGMV-Sr and JGMV-Fg isolates. It was also observed by Shukla et al. (1991) when comparing U07218.1 (JGMV-MDKS1) and U07217.1 (JGMV-MDO) from USA with the JGMV-JG, a Johnsongrass-infecting isolate from Australia. Several differences in the CP nucleotide sequences among the Brazilian JGMV-Sr and JGMV-Fg isolates (Figure 3) grouped them in distinct monophyletic subgroups (Figure 4). The third and the largest cluster composed by JGMV isolates from Australia showed higher pairwise identities than the other two groups (Figure 4 and 5).

Through transmission test applying the Brazilian JGMV-Sr isolate, we were able to verify mosaic symptoms and infection in teosinte (*Euchlaena mexicana*), *Sorghum verticilliflorum*, and sorghum genotypes, except QL3, with the virus been detected by RT-PCR. All maize and the two sugarcane genotypes were not infected in the transmission evaluations. Persley et al (1981) found that most of the maize hybrids in Australia had at least good field JGMV resistance, besides a high correlation between ratings of resistance verified from

greenhouse and natural field infection. In addition, Camelo-Garcia et al. (2016) showed that the Brazilian JGMV-Fg isolate from *Panicum maximum* cv Mombaça, did not infected maize. Silva et al. (2016) showed through transmission evaluations that the Brazilian JGMV-CNPGL (KT833782) isolated from Elephant grass (*Pennisetum purpureum*) was capable to infect maize, suggesting a potential threat to this crop in Brazil. However, so far, this strain was not identified naturally infecting maize at field. Also, the KITC motif of HCPro involved in the interaction of the viral particle and the stylus (Blanc et al., 1998), was absent from the JGMV-CNPGL isolate (Silva et al., 2016). On the other hand the vegetative propagation of the Elefant grass by stem cuttings (Pereira et al., 2017), allows the viral transmission without the presence of the vector. Seifers et al. (2005) using transmission test demonstrated that JGMV-MDO (U07217.1), which also shares high identity with the Brazilian JGMV-Sr isolates, was capable to infect *Sorghum bicolor*, except line QL3 genotype, as found in this work. Originally, this JGMV-MDO strain from USA was described as MDMV oat-infecting strain (McDaniel and Gordon, 1989; McKern, 1990). Perennial wild *Sorghum verticilliflorum*, an important weed in many crop plantations in Brazil (Tessele et al., 2014), was susceptible to the Brazilian JGMV-Sr isolates identified in this research. It is an important factor in the intercropping periods allowing the survival of the JGMV between seasons acting as viral reservoirs. The most efficient way to control disease spreading is through the development of resistant cultivar. Studying JGMV resistance, Presley et al. (1981) observed ratios of maize segregating generations that were consistent with resistance being controlled by a single dominant gene, which is important to breeders toward introgression of disease resistance genes into susceptible cultivars.

Conclusion

The identification of *Johnsongrass mosaic virus* (JGMV) isolates naturally infecting sorghum (*Sorghum bicolor* (L.) Moench) at field level in Brazil is reported. These Brazilian JGMV-Sr isolates have high CP sequence identities with the USA JGMV strains, MDO_U07217.1 and MDKS1_U07218.1, and grouped with them in the phylogenetic tree. The transmission evaluations showed that *Sorghum bicolor*, except line QL3, teosinte and *Sorghum verticilliflorum* were infected and expressed mosaic symptoms. The sorghum line QL3 identified as a source of JGMV resistance could be used in breeding programs to develop new resistant cultivars. The Brazilian JGMV-Fg isolates previously identified infecting *Panicum maximum* (GenBank KT289893) and *Pennisetum purpureum* (GenBank KT833782), are different from that identified in this study, and so far, were not identified naturally infecting sorghum crops. Maize and sugarcane genotypes used in the transmission test were not infected by the Brazilian JGMV-Sr isolates described in this study. However more genotypes, and also more weed poaceas will be evaluated looking for host range of the Brazilian JGMV-Sr isolates.

Acknowledgments

The authors thank to FAPEMIG contract number 9761 CBB-RED-00005/14 and the Brazilian Agricultural Research Corporation (EMBRAPA) for the financial support provided to this research project. We also extend our thank to the technician Célio Ramos das Neves for his assistance in the lab and the greenhouse work.

REFERENCES

- Adams, M. J., Zerbini, F. M., French, R., Rabenstein, F., Stenger, D. C. and Valkonen, J. P. T. 2012. Family potyviridae. in *Virus Taxonomy: 9th report of the International Committee on the Taxonomy of Viruses*, ed. By A.M.Q. King, M.J. Adams, E.B. Carstens. San Diego, p. 1069–1089.
- Blanc S., Ammar E. D., Garcia-Lampasona, S., Dolja, V. V., Llave, C., Baker, J. and Pirone, T. P. 1998. Mutations in the potyvirus helper component protein: effects on interaction with virions and aphid stylets. *Journal of General Virology*, London, 79:3119–3122
- Camelo-García, V. M., Andrade, A. C. S., Geering, A. D. W., Kitajima, E. W. and Rezende, J. A. M. 2016. Genome organization and host range of a Brazilian isolate of johnsongrass mosaic virus. *Archives of Virology*, New York, v. 161, p. 1335-1341.
- Cruz, I. and Vendramim, J. D. 1995. Efeito de diferentes genótipos de sorgo resistentes no desempenho do pulgão-verde, *Schizaphis graminum* Rond. Anais da Sociedade Entomológica do Brasil, *Jaboticabal*, v. 24, p. 253-263.
- Deng, C. L., Wang, W. J., Wang, Z. Y., Jiang, X., Cao, Y., Zhou, T., Wang, F. R., Li, H. F. and Fan, Z. F. 2008. The genomic sequence and biological properties of Pennisetum mosaic virus, a novel monocot-infecting potyvirus. *Archives of Virology*, New York, v. 153, p. 921-927.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, Lancaster, v. 39, p. 783-791.
- Food and Agriculture Organization of United States. *SORGHUM: Post-harvest Operations*. 33 p. 1999.
- Goussain, M. M. Efeito da aplicação do silício em plantas de milho no desenvolvimento biológico da lagarta-do-cartucho /*Spodoptera frugiperda*/ (J. E. Smith, 1797) e do pulgão-da-folha /*Rhopalosiphum maidis*/ (Fitch, 1856) (Hemiptera: Aphididae). 2001. 64 f. Dissertação (Mestrado em Entomologia) - Universidade Federal de Lavras, Lavras, 2001.
- Jiang, J. X. and Zhou, X. P. 2002. Maize dwarf mosaic disease in different regions of China is caused by Sugarcane mosaic virus. *Archives of Virology*, New York, v. 147, p. 2437-2443.
- Kumar, S., Stecher, G. and Tamura, K. 2016. Mega7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, Chicago, v. 33, p. 1870-1874.
- Lenardon, S. L. 2004. Giolitti, F. Maize Argentina mosaic. In: Lapiere, L.; Signotet, P. A. *Viruses and virus disease of Poaceae (Gramineae)*. Montpellier, p. 635-636.
- McDaniel, L. L. and Gordon, D. T. 1989. Characterization of the oat-infecting strain of maize dwarf mosaic virus. *Phytopathology*, St Paul, v. 79, p. 113-120.
- McKern, N. M., Whittaker, L. A., Strike, P. M., Ford, R. E., Jensen, S. G. and Shukla, D. D. 1990. Coat protein properties indicate that maize dwarf mosaic virus-KS-1 is a strain of Johnsongrass mosaic virus. *Phytopathology*, St Paul, v. 80, p. 907-912.
- Muhire, B. M., Varsani, A. and Martin, D. P. 2014. SDT: a virus classification tool based on pairwise sequence alignment and identity calculation. *PLoS One*, San Francisco, v. 9, n. 9, p. 1-8.
- Oliveira, E., Fernandes, F. T. and Pinto, N. F. J. A. 2005. Doenças do milho: identificação e controle. Sete Lagoas: Embrapa Milho e Sorgo, 84 p.
- Pappu, S. S., Brand, R., Pappu, H. R., Rybicki, E. P., Gough, K. H., Frenkel, M. J. and Niblett, C. L. 1993. A polymerase chain reaction method adapted for selective amplification and cloning of 3' sequences of potyviral genomes: application to dasheen mosaic virus. *Journal of Virological Methods, Amsterdam*, v. 41, p. 9–20.
- Pearson, C. E., Edamura, K. N. and Cleary, J. D. 2005. Repeat instability: mechanisms of dynamic mutations. *Nature Reviews in Genetics*, London, v. 6, p. 729–742.
- Pereira, A. V., Ledo, F. J. S. and Machado, J. C. 2017. BRS Kurumi and BRS Capiaçú - New elephant grass cultivars for grazing and cut-and-carry system. *Crop Breeding and Applied Biotechnology*, Viçosa, v. 17, p. 59-62.
- Persley, D. M., Martin, I. F. and Greber, R. S. 1981. The resistance of maize inbred lines to sugarcane mosaic virus in Australia. *Australian Journal of Agricultural Research, Melbourne*, v. 32, p. 741 – 748.
- Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, Chicago, v. 4, p. 406-425.
- Seifers, D. L., Haber, S., Ens, W., She, Y. M., Standing, K. G. and Salomon, R. 2005. Characterization of a distinct Johnsongrass mosaic virus strain isolated from sorghum in Nigeria. *Archives of Virology*, New York, v. 150, p. 557-576.
- Seifers, D. L., Salomon, R., Marie-jeanne, V., Alliot, B., Signoret, P., Haber, S., Loboda, A., Ens, W., She, Y. M. and Standing, K. G. 2000. Characterization of a novel potyvirus isolated from maize in Israel. *Phytopathology*, Saint Paul, v. 90, p. 505-513.
- Shukla, D. D. and Ward, C. W. 1989. Identification and classification of potyviruses on the basis of coat protein sequence data and serology. *Archives of Virology*, New York, v. 106, p. 171-200.
- Shukla, D. D., Frenkel, M. J., Ward, C. W. Structure and function of the potyvirus genome with special reference to the coat protein coding region. *Canadian Journal of Plant Pathology, Burnaby*, v. 13, p. 178-191, 1991.
- Shukla, D. D., Ward, C. W. and Brunt, A. A. 1994. *The Potyviridae*. Cambridge: CAB.
- Silva, D. D., Cota, L. V. and Costa, R. V. 2012. Manejo de doenças. In: Sistema Embrapa de Produção Agroindustrial de Sorgo Sacarino para Bioetanol Sistema BRS1G – Tecnologia Qualidade Embrapa, ed. May, A.; Durães, F. O. M.; Filho, I. A. P.; Schaffert, R. E.; Parrella, R. A. C. Documento 139, Embrapa Milho e Sorgo, p. 68-84.
- Silva, K. N., Melo, F. L., Orílio, A. F., Nagata, T., Silva, M. S., Fernandes, C. D., Fragoço, R. R., Dessaune, S. N. and Resende, R. O. 2016. Biological and molecular characterization of a highly divergent johnsongrass mosaic virus isolate from Pennisetum purpureum. *Archives of Virology*, New York, v. 161, p. 1981-1986.
- Souza, I. R. P. and Barros, B. A. 2016. Molecular detection and identification of potyvirus species of the common-mosaic in maize and sorghum. In: Oliveira, C. M. de; Oliveira, E. de (Org.). *Diseases in maize: insect-vectors, mollicutes and virus*. Brasília, DF: Embrapa, p. 1-221.
- Souza, I. R. P. de, Giolitti, F., Carneiro, N. P., Lenardon, S. L., Oliveira, E. de, Gomes, E. A., Noda, R. W. and Souza, F. A. de. 2012. Sequence diversity in coat protein of SCMV infecting maize and sorghum in Brazil. *Revista Brasileira de Milho e Sorgo, Sete Lagoas*, v. 11, n. 2, p. 120-136.
- Souza, I. R. P., Schuelter, A. R., Guimarães, C. T., Chuster, I., Oliveira, E. de and Redinbaugh, M. 2008. Mapping QTL

- contributing to SCMV resistance in tropical maize. *Hereditas*, Lund, v. 145, p. 167-173.
- Tessele, A., Furtado, R. C. N., Moreno, G., Lorenzetti, J. B., Gris, D. J., Albretch, L. P. and Albretch, A. J. P. 2014. Sorghum spp.: situação como planta daninha e manejo no Brasil. *Journal of Agronomic Sciences*, Umuarama, v. 3, p.256-273.
- Waquil, I.M., Oliveira E., Pinto, N. F. J. A., Fernandes, F.T. and Corrêa, L. A. 1996. Efeito na produção e incidência de viroses em híbridos comerciais de milho. *Fitopatologia Brasileira*, Brasília, v. 21, p. 460-463.
- Zhao, X., Tan, Z., Feng, H., Yang, R., Li, M., Jiang, J., Shen, G. and Yu, R. 2011. Microsatellites in diferente Potyvirus genomes: Survey and analysis. *Gene*, Amsterdam, v. 488, p. 52-56.
