

## A review on the use of hormones in fish farming: Analytical methods to determine their residues

Celia A. Hoga<sup>a</sup>, Fernanda L. Almeida<sup>b</sup> and Felix G. R. Reyes<sup>a</sup>

<sup>a</sup>Department of Food Science, School of Food Engineering, University of Campinas – UNICAMP, Campinas, Brazil; <sup>b</sup>Brazilian Agricultural Research Corporation, Embrapa Amazônia Ocidental, Manaus, Brazil

### ABSTRACT

Hormones are used in fish farming to increase fish production when one sex of a species has the capacity to grow bigger and faster than the other sex. The technique to increase fish production based on sexual dimorphism mostly uses estrogens and androgens. These chemicals must be handled carefully to ensure environmental, biological and food safety, since they can contaminate the environment and promote changes in the endocrine system inducing adverse effects on the consumer health. Another important issue is the reliability of the analytical methods used to identify and/or quantify hormone residues in the meat of treated fish. Therefore, the aim of this review is to bring into view the use of hormones in fish farming, the possible impacts of this practice on humans and on the environment and to discuss the current methods of analysis for determining the hormone residues in food matrices, especially in fish.

### Una revisión sobre el uso de hormonas en la piscicultura. Métodos analíticos para determinar sus residuos

#### RESUMEN

Las hormonas se utilizan en la piscicultura para aumentar la producción de pescado cuando un sexo de una especie tiene la capacidad de crecer más y más rápido que el otro sexo. La técnica para aumentar la producción de pescado con base en el dimorfismo sexual utiliza, principalmente, estrógenos y andrógenos. Estas sustancias deben manipularse cuidadosamente para garantizar la seguridad ambiental, biológica y alimentaria, ya que pueden contaminar el medioambiente e inducir cambios en el sistema endocrino provocando efectos adversos en la salud del consumidor. Otro importante problema es la fiabilidad de los métodos analíticos utilizados para identificar y/o cuantificar los residuos de hormonas en la carne del pescado tratado. Así, el objetivo de esta revisión es presentar el uso de hormonas en piscicultura, los posibles impactos de esta práctica en seres humanos y en el medioambiente y discutir los métodos actuales de análisis para determinar los residuos de hormonas en varios alimentos, especialmente en el pescado.

### ARTICLE HISTORY


Received 5 February 2018  
Accepted 3 May 2018

### KEYWORDS

Fish farming; sexual reversion; sexual dimorphism; veterinary drugs; hormones; analytical methods; LC–MS/MS

### PALABRAS CLAVE

piscicultura; reversión sexual; dimorfismo sexual; medicamentos veterinarios; hormonas; métodos analíticos; LC–MS/MS



## 1. Introduction

The demand for fish has increased in recent years due to population growth and the constant search for a healthy

diet. According to the Brazilian Institute of Geography and Statistics (IBGE, 2015), in 2015, fish farming increased to 1.5% when compared to the previous year, on the total fish

production. The Brazilian aquaculture has been growing strongly in all regions of the country, mainly due to advances in the technical management of fish cultivation to cope with the increasing worldwide demand for fish as well as the economic and environmental sustainability of fish farming. In the Brazilian aquaculture scenario, tilapia (*Oreochromis niloticus*) and tambaqui (*Colossoma macropomum*) are the main farmed species.

Hormones are chemical messengers that are responsible for the communication between different types of cells that recognize their identity and function through receptors, which are protein structures specialized in molecular recognition. After the proximity and the hormone-receptor interaction, a series of biochemical reactions leading to specific biological responses occur (Reis-Filho, de Araújo, & Vieira, 2006).

Steroids are hormones produced from cholesterol and can be grouped into five subgroups based on their structural characteristics: estrogens, androgens, progestogens, glucocorticoids and mineral corticoids (Guedes-Alonso, Montesdeoca-Esponda, Sosa-Ferrera, & Santana-Rodríguez, 2014). The three former subgroups compose a group of sex steroids. The most important sex hormones used in fish farming are estrogens and androgens, which can be of natural origin (present in nature) or synthetic (synthesized or produced in laboratory).

Hormones in aquaculture are used for artificial reproduction and sex reversal. The first sustains the production chain with the constant production of seeds. The second is used when the growth rate and/or gain weight are different between the male and female. This difference between genders is very common in teleost fish and usually occurs during puberty (Taranger et al., 2010). Nevertheless, the use of hormonal products in fish farming can have harmful consequences, such as potential risks to human and environmental health related to hormone-dependent parameters. Moreover, its use outside of good animal husbandry practices may affect not only the fish farming production system itself but also the commercialization of this food commodity.

Currently, most issues related to food safety aim at residue control in foods due to the use of pesticides, veterinary drugs or accidents involving environmental contamination from chemicals. In relation to hormone residues in fish, Liu et al. (2011) studied the presence of steroidal hormones in crucian carp (*Carassius auratus*), carp (*Cyprinus carpio*) and silvery minnow (*Anabarilius alburnops*), which are fish species common in Dianchi Lake (China), as well as placed in the same position in the food chain. The authors conclude that steroidal hormones are likely ubiquitous contaminants in wild fish from that lake. The presence of steroid hormone residues was reported in fish tissue samples from two common fishes (*Boops boops* and *Sphoeroides marmoratus*) which are species that are at the bottom of the food chain and that were fished in the proximity of the outfall of the wastewater treatment plant of the city of Las Palmas de Gran Canaria (Spain) (Guedes-Alonso, Sosa-Ferrera, & Santana-Rodríguez, 2017). Steroid hormone residues were also reported in yellow croaker (*Larimichthys polyactis*) tissue samples collected from local supermarkets in Changchun (China) (Wang et al., 2012) and in tilapia (*Oreochromis mossambicus* and *Oreochromis niloticus*) purchased from supermarkets in Taipei City (Taiwan) (Chen, Wang, Gwo, & Chen, 2012). According to Epstein (1990 with Duarte, da Silva, & Meirelles, 2002), 3000 children in Puerto Rico had serious problems with premature sexual development and ovarian cysts due to ingestion of meat products with zeranol

residues (xenobiotic anabolic agent with estrogenic activity promoting growth). Thus, eating foods contaminated with steroids can lead to endocrine disorders and may even result in the development of cancer (Bergman, Heindel, Jobling, Kidd, & Zoeller, 2013). Nevertheless, the Joint FAO/WHO Expert Committee on Food Additives (JECFA) established an acceptable daily intake of 0–0.5  $\mu\text{g kg}^{-1}$  body weight (bw) for zeranol (JECFA, 1988), 0–0.02  $\mu\text{g kg}^{-1}$  bw for trenbolone acetate (JECFA, 1989), 0–0.05  $\mu\text{g kg}^{-1}$  bw for 17 $\beta$ -estradiol, 0–30  $\mu\text{g kg}^{-1}$  bw for progesterone, and 0–2  $\mu\text{g kg}^{-1}$  bw for testosterone (JECFA, 2000).

The use of hormones in food producing animals faces different legal regulations in different countries. The European Union prohibits the use of substances with hormonal or thyrostatic action and  $\beta$ -agonists in food producing animals, including aquaculture (EC, 1996, 2003, 2008). Other countries such as Canada, Australia, New Zealand, Argentina and the United States allow for the use of natural steroid hormones, such as testosterone, progesterone and 17 $\beta$ -estradiol, as well as the synthetics zeranol and trenbolone acetate. In those countries, only residues from synthetic compounds with maximum residue levels (MRLs) are controlled because they are considered to be the most potent endocrine compounds (Duarte et al., 2002). In Brazil, since 1991, the importation, production, marketing, and use of natural or artificial substances for animal growth purposes and/or weight gains have been prohibited. Some substances (i.e. testosterone, progesterone, estradiol, zeranol, trenbolone acetate), natural or synthetic, with estrogen or progestogen action are allowed for therapeutic purposes, estrous cycle synchronization, and preparation of donors and recipients for embryo transfer (MARA, 1991). Codex Alimentarius Commission (Codex) established that residues resulting from the use of 17 $\beta$ -estradiol, progesterone, and testosterone, as a production aid in cattle, in accordance with good animal husbandry practice are unlikely to pose a hazard to human health. Therefore, considering the establishment of an MRL (in muscle, liver, kidney, fat) is unnecessary. Thus, it is worth mentioning that the use of hormones for the purpose of sexual reversion is not established by the legislative framework of any country, nor in any recommendations of regional bodies (European Commission) nor international (JECFA or Codex) that deal with food safety.

In relation to the methods of analysis, the determination of hormone residues in foods of animal origin is difficult because of the complexity of the matrix and the low concentrations to be quantified (in the order of  $\text{ng g}^{-1}$ ) (Guedes-Alonso et al., 2014). In this regard, analytical methods for detecting hormone residues are generally immunoassays, such as radioimmunoassays (RIA) and enzyme-linked immunosorbent assays (ELISA) (Zanardi, 2007). For quantitative determination of hormone residues in fishery products, analytical methods using liquid chromatography (LC) coupled to a tandem mass spectrometry (LC-MS/MS) system is currently the recommended analytical technique, which has become popular and widely used in food analysis due to its high specificity and detectability in the determination of residue levels of contaminants in complex matrices (Reis-Filho et al., 2006; Zanardi, 2007). Nevertheless, the determination of hormone residues in fish matrices by LC-MS/MS has been seldom reported (Guedes-Alonso et al., 2017; Jakimska et al., 2013; Wang et al., 2012; Xu, Chu, Peng, Jin, & Wang, 2006).

The main goal of this article is to review the literature available on the use of hormones in fish farming, as well as the qualitative and quantitative analytical methods developed

for the determination of hormone residues in food from aquaculture. The aim is to provide advice to the readers on the advantages and disadvantages on the use of hormones in fish farming. Challenges and perspectives on future trends of analytical methodologies in determining hormones mainly in fish products are also discussed.

## 2. The use of hormones in fish farming

### 2.1. Sex reversal

Sex reversal can be achieved with hormonal treatment during sex differentiation. Fish sex may be separated into genotype, which is determined by the genes responsible for gonads formation, and phenotype, which is the appearance of the ovary or testis. Phenotype differentiation occurs naturally, generally earlier in female than in males, during the ontogeny of the fish larvae (Piferrer, 2001). This process is very complex, but it can be manipulated using androgen and estrogen hormones.

The use of hormones in fish farming for sex reversal aims at the production of monosex population to increase growth rate or weight gain. Commercially, it is advantageous to rear individuals of the most profitable gender, thereby achieving more uniform lots and controlling undesirable breeding (Singh, 2013; Taranger et al., 2010). There are two methods for producing monosex fish populations: (i) direct, by which the fish are treated with hormones for the purpose of developing the desired sex, and (ii) indirect, by which parents or breeders are treated with hormones to obtain neomale (XX♂), neofemale (XY♀, ZZ♀), or supermale (YY♂) populations, and with them getting the entire batch of larvae of the same sex (Piferrer, 2001).

Protocols for the production of monosex female stocks in 35 different species of teleosts using a variety of estrogens were reported by Piferrer (2001). Those protocols included the type of estrogen, route of administration, dose, duration of treatment and timing. The author emphasizes the convenience of using the natural estrogen 17β-estradiol rather than synthetic estrogens. Among androgens, 17α-methyl testosterone (synthetic) has been widely employed to reverse females into males, being commonly used in tilapia. Both steroids have the advantage of being easily metabolized after the period of hormonal treatment (Mainardes-Pinto, Fenerich-Verani, de Campos, & da Silva, 2000; Zanoni, Leal, Filho, de Oliveira, & Ribeiro, 2013). Table 1 shows hormones used for sex reversal and differentiation in different species of fish for both male and female induction.

The administration of hormones for sex reversal treatment can be done using systemic (direct injection and silastic implantation), immersion, or dietary supplementation (hormone incorporated in fish feed) (Pandian & Sheela, 1995). Commercially, the most successful treatments use immersion and diet, as both methods reach a large number of fish, while the systemic transfer method is expensive and requires technical ability to be applied on the fish. In the immersion technique, the dose administered not only affects the efficiency of hormonal treatments but also other parameters, such as the type of hormone, water temperature and exposure time. The addition of hormones in the feed is more efficient because it is easily controlled, and allows optimum steroid dose to induce the complete sex reversal of all individuals (Pandian & Sheela, 1995; Piferrer, 2001).

Most authors mention the use of 17β-estradiol, estradiol valerate, 17α-methyl testosterone, or 17α-methyl dihydrotestosterone (by immersion and diet technique) to produce an all-female population by the direct and indirect feminization. On the other hand, androgen is used to masculinize females, producing neomales (Piferrer, 2001). Females are produced due to the common problems associated with precocious male sexual maturity and the advantage associated with the fact that females grow faster than males. Several authors reported great results on the effects of hormones on the proportions of sexual types (de Bem, 2009; Haffray et al., 2009; Lin, Benfey, & Martin-Robichaud, 2012). Moreover, the authors who used the implant method to test different doses of the hormone, verify that all fishes died and showed liver injury at high doses of 17β-estradiol (higher than 8 mg kg<sup>-1</sup>) (Passini, 2013).

For production of an all-male population, tamoxifen (compound that blocks the estrogen receptor), fadrozole and letrozole (nonsteroidal aromatase inhibitor), 17α-methyl testosterone, 17α-ethynyl testosterone and 17α-ethynyl estradiol by direct and indirect methods have been used in species where males grow faster than females and reach a larger size (Garcia et al., 2013; Liao et al., 2014; Zanoni et al., 2013).

According to Pandian and Sheela (1995), the advantages of hormonal treatment ensure maximum growth, eliminate early maturation in males and allow broodstock management. The disadvantages of this technique are related to the possible presence of carcinogenic steroid residues that can affect consumer health. Hormonal induction of sex reversal can become a stressful process, resulting in low survival rates, delayed sexual maturity and reduction of fish fertility, while high doses can lead to sterility, paradoxical sexual reversal and growth suppression. On a large scale, sexual reversal may become a technique to pollute the environment because more than 99% of hormones are metabolized and released within hours or days into the water.

Treatments for sex reversal purposes are conducted at the initial stage of fish development, before or during the process of gonadal sexual differentiation, such as during the last few months before being marketed (Almeida, 2013; de Bem, 2009). The residue of 17β-estradiol, for instance, disappears in less than a month after the end of the treatment, meaning that there is no risk of meat contamination by hormones at the time of slaughter. According to Specker and Chandlee (2003), summer flounder (*Paralichthys dentatus*) larvae and juvenile whole-body estradiol levels were not significantly different compared to the untreated controls after 24 h. In addition to being a natural steroid, which is quickly metabolized and excreted, 17β-estradiol is also considered the hormone with the least impact on the environment (Hendry, Martin-Robichaud, & Benfey, 2003; Piferrer, 2001; Specker & Chandlee, 2003). While treating Rainbow trout (*Salmo gairdneri*), Adeel, Song, Wang, Francis, Yang, (2017) reported that steroidal estrogens decrease significantly in the aqueous phase, presenting short half-lives. In fact, the half-lives of estrogenic steroids (17β-estradiol, estradiol and ethinyl estradiol) were estimated in 2–6 days at 20°C, in water and sediment, with ethinyl estradiol being more stable than two other estrogens (Williams, Jürgens, & Johnson, 1999). Subsequently, Jürgens et al. (2002) reported that 17β-estradiol was much more resistant than ethinyl estradiol to the biodegradation in the water of English rivers.

**Table 1.** Androgens, estrogens and nonsteroidal agents used for sex reversal and differentiation in different species of fish.**Tabla 1.** Andr6genos, estr6genos y agentes no esteroides utilizados para la reversi3n sexual y la diferenciaci3n en diferentes especies de peces.

Hormone	Fish species	References	Region/country
17 $\beta$ -Estradiol	Rainbow trout ( <i>Salmogairdneri</i> )	Johnstone, Simpson, & Youngson, 1978	Aberdeen/Great Britain
	Atlantic salmon ( <i>Salmosalar</i> )	Johnstone et al., 1978	
	Atlantic halibut ( <i>Hippoglossus hippoglossus L.</i> )	Hendry et al., 2003	New Brunswick/Canada
	Summer flounder ( <i>Paralichthys dentatus</i> )	Specker & Chandlee, 2003	Narragansett/USA
	Bagrid catfish ( <i>Pseudo bagrus fulvidraco</i> )	Park, Kim, Cho, & Kim, 2004	Busan/South Korea
	Atlantic cod ( <i>Gadusmorhua</i> )	Lin et al., 2012	St. Andrews/Canada
17 $\alpha$ -Ethinylestradiol	Common snook ( <i>Centropomus undecimalis</i> )	Passini, 2013	Florian3polis/Brazil
	Tilapia ( <i>Oreochromis aureus</i> )	Melard, 1995	Tihange/Belgium
Estradiol valerate	Tetra ( <i>Astyanax altiparanae</i> )	de Bem, 2009	Rio Claro/Brazil
Tamoxifen	Bagrid catfish ( <i>Pseudo bagrus fulvidraco</i> )	Park et al., 2004	Busan/South Korea
Letrozole	Protogynous dusky grouper ( <i>Epinephelus marginatus</i> )	Garcia et al., 2013	S3o Paulo/Brazil
	Medaka fish ( <i>Oryzias latipes</i> )	Liao et al., 2014	Taipei/Taiwan
Fadrozole	Atlantic halibut ( <i>Hippoglossus hippoglossus L.</i> )	Babiak et al., 2012	Nordland/Norway
	Protogynous wrasse ( <i>Halichoeres trimaculatus</i> )	Nozu, Kojima, & Nakamura, 2009	Okinawa/Japan
17 $\alpha$ -Methyltestosterone	Brook trout ( <i>Salvelinus fontinalis</i> )	Haffray et al., 2009	Rennes/France
	Nile tilapia ( <i>Oreochromis niloticus</i> )	Curtis et al., 1991	Cowallis/USA
		Mainardes-Pinto et al., 2000	Nova Odessa/Brazil
		Toyama, Corrente, & Cyrino, 2000	Piracicaba/Brazil
		Hayashi, Boscolo, Soares, & Meurer, 2002	Maring3/Brazil
		Zanardi et al., 2011	Jaboticabal/Brazil
		El-Sayed, Abdel-Aziz, & Abdel-Ghani, 2012	Alexandria/Egypt
		Zanoni et al., 2013	Londrina/Brazil
		Tabata, Rigolino, & Tsukamoto, 1999	S3o Paulo/Brazil
		Johnstone, Macintosh, & Wright, 1983	Stirling/Great Britain
17 $\alpha$ -Methyldihydrotestosterone	Rainbow trout ( <i>Oncorhynchus mykiss</i> )	Johnstone et al., 1978; 1983	Aberdeen and Stirling,/ Great Britain
	Tilapia ( <i>Oreochromis Mossambicus</i> )		
	Rainbow trout ( <i>Salmo gairdneri</i> )		
17 $\alpha$ -Methyltestosterone	Atlantic salmon ( <i>Salmosalar</i> )	Johnstone et al., 1978	Aberdeen, /Great Britain
	Atlantic halibut ( <i>Hippoglossus hippoglossus L.</i> )	Hendry et al., 2003	New Brunswick/Canada
	Atlantic cod ( <i>Gadusmorhua</i> )	Lin et al., 2012	St. Andrews/Canada
17 $\alpha$ -Ethinyltestosterone	Tilapia hybrids ( <i>Oreochromis niloticus</i> x <i>O. aureus</i> )	Rothbard et al., 1990	Tel Aviv/Israel

## 2.2. Artificial reproduction

Another use of hormones in aquaculture is in seed production, i.e. artificial reproduction, which is important for manipulating the final maturation and ovulation in the gonad.

Artificial reproduction has endless applications, it can be controlled by environmental parameters for natural reproductive performance and by exogenous hormones to induce/advance or delay/arrest fish maturation, and in making spawning occur a few months earlier or later than normal. By advancing the period of reproduction, the fish farmers have a greater flexibility in the marketing of the larvae and young fish. By restricting spawning to a certain period, this maximizes time and resources, allowing for the higher throughput and turnover of nursery ponds. By synchronizing spawning males and females, the farmer obtains fingerlings in periods when profitability is higher (Mylonas, Fostier, & Zanuy, 2010; Venturieri & Bernardino, 1999). In addition, this technique is very important for the reproduction of fish that need external stimuli, since they are not suitable for production in captivity.

Hormonal techniques for fish breeding are based on intramuscular or intraperitoneal injection in broodstocks. The oldest and still the most used hormone is the crude extract of the pituitary gland (PE) of mature fish (carp and salmon). Another largely used hormone is the gonadotropin-releasing hormone (GnRH) of both mammals (mGnRH) and salmon (sGnRH). Gonadotropin-releasing hormone analogs (GnRH<sub>a</sub>), human gonadotropin (GtHs), dopamine antagonists (pimozide, domperidone, metoclopramide and reserpine) and synthetic hormones similar to gonadotropins (Ovopel) have been tested in several species to induce or block sexual maturation (Almeida, 2013; Ara3jo et al., 2014; Zohar & Mylonas, 2001). Zohar and Mylonas (2001) summarize the use of GnRH<sub>a</sub> and GtHs used to induce maturation and spawning in more than 30 species of fish.

The advantage of using dopamine antagonists is that they block the dopamine inhibitor mechanism, elevating GnRH secretion in the blood stream while inducing maturation. The most used and recommended dopamine antagonist is domperidone, as it is cheap, potent and does not cross the blood-brain barrier (Dasgupta, Sarkar, Sarangi, & Bhattacharya, 2009; Heyrati, Mostafavi, Toloei, & Dorafshan, 2007; Wang et al., 2009). When GnRH alone does not work properly, it is usually used in combination with dopamine antagonists (Almeida, 2013; Venturieri & Bernardino, 1999). The use of gonadotropin hormones to induce maturation and spawning in different fish species is shown in Table 2.

Regarding synthetic hormones, Ovopel is composed of a combination of the mGnRH<sub>a</sub> and dopamine antagonists (metoclopramide). It was used to induce the reproduction of wild mature asp (*Aspius aspius*) during out-of-season spawning (Targońska, Kucharczyk, Kujawa, Mamcarz, & 3arski, 2010), to induce spermiation of Amazon catfish (*Leirius marmoratus*) (Ara3jo et al., 2014) and to induce ovulation of pacu (*piaractus mesopotamicus*) (Santos, De Santos, Salom3o, & de Silva, 2012). Ovaprim is a synthetic hormone that combines sGnRH<sub>a</sub> and another dopamine antagonist (domperidone).

## 3. Impact of the hormone on the environment and consumer health

The 17 $\beta$ -estradiol residue in fish meat disappears in less than a month after finishing the sex reversal treatment, since this steroid is of natural origin, quickly metabolized and excreted. The residue of 17 $\alpha$ -methyl testosterone is also rapidly metabolized after a period of hormonal treatment (Piferrer, 2001). Thus, theoretically, it should not be found at a high level in fish meat if the treatment was done in a correct manner and over the correct period.



**Table 2.** Hormones used to induce maturation and spawning in different species of fish.**Tabla 2.** Hormonas usadas para inducir la maduración y el desove en diferentes especies de peces.

Hormone	Fish species	References	Region/country
Luteinizing hormone-releasing hormone analogous (LHRHa)	Milkfish ( <i>Chanos chanos Forsskal</i> )	Lee et al., 1986	Waimanalo/USA
	Black sea bass ( <i>Centropristis striata</i> )	Berlinsky, King, & Smith, 2005	Durham/USA
	Atlantic cod ( <i>Gadus morhua</i> )	Garber, Fordham, Symonds, Trippel, & Berlinsky, 2009	New Brunswick/Canada
Carp pituitary extract	Nase ( <i>Chondrostoma nasus</i> )	Szabó, Medgyasszay, & Horváth, 2002	Godollo/Hungary
	Kutum ( <i>Rutilus frisii kutum</i> )	Heyrati et al., 2007	Tehran/Iran
	Wild loach ( <i>Misgurnus anguillicaudatus</i> )	Wang et al., 2009	Hubei/China
	Dourado fish ( <i>Salminus brasiliensis</i> )	Flora, Maschke, Ferreira, & De A Pedron, 2010	Santa Maria/Brazil
	Carp ( <i>Cyprinus carpio carpio</i> )	Vazirzadeh et al., 2011	Karadj/Iran
	Amazon catfish ( <i>Leiarius marmoratus</i> )	Araújo et al., 2014	Cuiabá/Brazil
	Tambaqui ( <i>Colossoma macropomum</i> )	Filho, Brito, Cunha, Júnior, & Amorim, 2016	Recife/Brazil
Gonadotropin-releasing hormone analogous (GnRH <sub>a</sub> ) + dopamine antagonists	Nase ( <i>Chondrostoma nasus</i> )	Szabó et al., 2002	Godollo/Hungary
	Silver perch ( <i>Bidyanus bidyanus</i> )	Levavi-Sivan, Vaiman, Sachs, & Tzchori, 2004	Rehovot/Israel
	Kutum ( <i>Rutilus frisii kutum</i> )	Heyrati et al., 2007	Tehran/Iran
	Rohu ( <i>Labeo rohita</i> )	Dasgupta et al., 2009	Bhubaneswar/India
	Wild loach ( <i>Misgurnus anguillicaudatus</i> )	Wang et al., 2009	Hubei/China
Gonadotropin-releasing hormone analogous (GnRH <sub>a</sub> )	Carp ( <i>Cyprinus carpio carpio</i> )	Vazirzadeh et al., 2011	Karadj/Iran
	Nase ( <i>Chondrostoma nasus</i> )	Szabó et al., 2002	Godollo/Hungary
	Dusky grouper ( <i>Epinephelus marginatus</i> )	Marino et al., 2003	Rome/Italy
	Greater amberjack ( <i>Seriola dumerili</i> )	Mylonas, Papandroulakis, Smboukis, Papadaki, & Divanach, 2004	Crete/Greece
	Silver perch ( <i>Bidyanus bidyanus</i> )	Levavi-Sivan et al., 2004	Rehovot/Israel
	Rose snapper ( <i>Lutjanus guttatus</i> )	Ibarra-Castro & Duncan, 2007	Mazatlán/México
	Kutum ( <i>Rutilus frisii kutum</i> )	Heyrati et al., 2007	Tehran/Iran
	Atlantic cod ( <i>Gadus morhua</i> )	Garber et al., 2009	New Brunswick/Canada
	Wild loach ( <i>Misgurnus anguillicaudatus</i> )	Wang et al., 2009	Hubei/China
	Honeycomb grouper ( <i>Epinephelus merra</i> )	Kanemaru et al., 2012	Miyazaki/Japan
	Senegalese sole ( <i>Solea senegalensis</i> )	Rasines et al., 2013	Santander/Spain
	Meagre ( <i>Argyrosomus regius</i> )	Mylonas et al., 2013	Crete/Greece
	Longfin yellowtail ( <i>Seriola rivoliana</i> )	Fernández-Palacios, Schuchardt, Roo, Hernández-Cruz, & Izquierdo, 2015	Las Palmas/Spain
Synthetic hormone similar to gonadotropin (Ovaprim)	Freshwater rotifer ( <i>Brachionus calyciflorus</i> )	Sugumar & Munuswamy, 2006	Chennai/India
	Asp ( <i>Aspius aspius</i> )	Targońska et al., 2010	Olsztyn/Poland
	Pinfish ( <i>Lagodon rhomboids</i> )	DiMaggio, Broach, & Ohs, 2013	Gainesville/USA
Synthetic hormone similar to gonadotropin (Ovopel)	Asp ( <i>Aspius aspius</i> )	Targońska et al., 2010	Olsztyn/Poland
	Amazon catfish ( <i>Leiarius marmoratus</i> )	Araújo et al., 2014	Cuiabá/Brazil
	Pacu ( <i>Piaractus mesopotamicus</i> )	Santos et al., 2012	Presidente Prudente/Brazil
Human chorionic gonadotropin (hCG)	Silver perch ( <i>Bidyanus bidyanus</i> )	Levavi-Sivan et al., 2004	Rehovot/Israel
	Pinfish ( <i>Lagodon rhomboids</i> )	DiMaggio et al., 2013	Gainesville/USA

According to Reis-Filho et al. (2006) natural estrogens are continuously introduced into the environment by human excretion, which follows the sewage disposal system, even after entering the environment at low concentrations (nanograms per liter), making them very persistent. These estrogens introduced into the environment could affect wildlife and human health by disrupting their normal endocrine systems, such as the feminization effects of males, altering the offspring sex ratios, infertility, reduced fertility, inhibition of the development of sexual organs and sexual reversal.

The way to detect if the reproductive systems of aquatic organisms are affected by estrogen is by determining the level of vitellogenin (VTG) in the blood plasma of an organism. VTG is a protein that has an important role in the female reproductive system of oviparous vertebrates. It is synthesized in the liver, regulated by estrogen and transported via the blood to the ovaries, where it is incorporated into the developing eggs. If an increase is observed in the VTG plasma level of an organism, it means that it was exposed to substances with estrogenic activity. Although the concentrations of these hormones are very low (in the order of  $\text{ng L}^{-1}$  or  $\mu\text{g L}^{-1}$ ), they can be sufficient to induce VTG synthesis even in male fish (Bila & Dezotti, 2003, 2007).

The other way that hormones are introduced into the environment is by the incorrect or illegal discard of water containing residues of these compounds used to treat fish. The contamination could have originated by fish excretion, as well as from the medicated feed that remains unconsumed by fish. According to

some authors (Leonhardt, 1997; Specker & Chandlee, 2003), the liver metabolizes the hormones in water soluble compounds. Approximately 99% of the administered hormone through the diet during the sex reversal is metabolized and released into the water by bile and urine excretion within a few hours or days after treatment. Estrogens are then excreted mainly as inactive conjugates of sulfuric and glucuronic acids, which do not have direct biological activity, but can act as precursor hormone reservoirs that can be reconverted to free steroids with estrogenic activities by bacteria in the environment, such as the microorganisms in raw sewage and sewage treatment plants (Ying, Kookana, & Ru, 2002). In other words, the sex reversal technique can pollute the environment if the water in which fish were treated is not properly depurated.

Specific studies have demonstrated that the use of hormones does not result in residue accumulation in the tissues of the reversed fish. Rothbard et al. (1990) showed that in the muscle of tilapia treated for 11 weeks with a diet containing  $60 \text{ mg kg}^{-1}$  of  $17\alpha$ -ethynyl testosterone, a high concentration of the steroid was detected on day 0 ( $142.1 \text{ ng g}^{-1}$ ) and day 1 ( $168.6 \text{ ng g}^{-1}$ ) after hormonal exposure. The samples taken on days 3, 5 and 7 did not differ from the untreated controls ( $17\alpha$ -ethynyl testosterone concentrations were below the detectable level of  $50 \text{ ng g}^{-1}$ ). Curtis, Diren, Hurley, Seim, and Tubb (1991) fed Nile tilapia diets that contained  $30 \text{ mg kg}^{-1}$  of  $17\alpha$ -methyl testosterone for 30 days. The hormone concentration decreased logarithmically from day 1 to day 10 after treatment.

In humans, the exposure to hormones can cause endocrine disorders, such as early puberty in children, advances in bone age, as well as negative repercussions on growth, modification of sexual characteristics and cancer development (Bergman et al., 2013). These disorders especially occur in children because they are in the growing phase when puberty has not yet developed. Thus, Partsch & Sippell (2001 with Alves, Flores, Cerqueira, & Toralles, 2007) reported that precocious pseudo puberty could be induced in children by oral or dermal exposure to estrogen-containing foods or ointments, respectively, if they were chronically exposed to food (meat) derived from animals treated with high doses. In Milan, Bahrain, Puerto Rico and Jerusalem, there were reported cases of gynecomastia, larche, pubarche and precocious sexual development in children, which were induced by the consumption of poultry, meat, milk and cereals contaminated with estrogen, or its precursors, or by the contamination of drinking water with substances that had estrogenic activity (Alves et al., 2007).

#### 4. Methods of analysis for determination of hormone residues in several matrices

Determining hormone residues in foods from animal origin is a difficult task, first because of the matrix complexity, and second because of the low concentrations that need to be quantified, at the order of  $\mu\text{g kg}^{-1}$  (or  $\mu\text{g L}^{-1}$ ) and  $\text{ng kg}^{-1}$  (or  $\text{ng L}^{-1}$ ) (Guedes-Alonso et al., 2014). In most analyses involving actual samples, a substantial clean-up of the extract and analyte concentration during sample preparation step is necessary to isolate the target compounds from the matrix and to achieve the required limits for detection and quantification.

To determine the hormone residues in samples from the environment and in foods, analytical methods have been developed to identify and quantify these substances in simple and complex matrices. In this regard, according to Zanardi (2007), analytical methods for detecting hormone residues are, generally, immunoassays, such as RIA and ELISA, which are routinely used. High-performance thin layer chromatography and gas chromatography–MS (GC–MS) can also be an important tool for steroid determination. Nonetheless, for quantitative determination, analytical methods using LC coupled with several types of detectors are the most widespread techniques. For hormone determination, the most common detectors coupled to LC are electrochemical, ultraviolet diode array (DAD), fluorescence, UV-Vis, MS and tandem mass spectrometry (MS/MS) (Reis-Filho et al., 2006). Table 3 summarizes the analytical techniques reported for the determination of hormones in several matrices.

GC–MS permits the simultaneous determination of many analytes with a detection limit in the order of  $\text{ng kg}^{-1}$  or  $\text{ng L}^{-1}$ , but the sample preparation is laborious and time-consuming. To turn the molecules volatile, the most used derivation technique has been silylation and acetylation (Zhou & Zhang, 2011). Hartmann, Lacorn, and Steinhart (1998) investigated 12 steroids in beef, veal, milk, milk products, pork, meat products, poultry, eggs, fish, plants, yeast and fermented alcoholic beverages using GC–MS. For each matrix, different sample preparations were studied by evaluating the recovery of the spiked samples. The authors reported residue levels between 0.03 and  $1.41 \mu\text{g kg}^{-1}$  for 10 steroids, except estrogens that were not detectable. Long, Hong, and Jie (2007) reported a GC–MS method for the identification and quantitative determination

of  $17\beta$ -estradiol residues in the muscle of various fishery products. The authors reported extraction, clean-up and derivatization more able to enhance detection sensitivity. Great linearity and precision were obtained and the method was successfully applied to the analysis of samples from the retail market.

ELISA are simple and economical and are especially suitable for screening large numbers of hormonally active substances with great sensitivity. ELISA does not cause serious problems in discarding samples because it does not require the use of reagents that are dangerous to human and environmental health (Zhou & Zhang, 2011). Zanardi et al. (2011) analyzed tilapia carcass samples by ELISA using a commercial kit to determine testosterone at the residue concentration level of  $\text{ng g}^{-1}$ . The authors reported recovery values of 69.9%.

For the simultaneous separation of analytes, reversed-phase LC is performed (C18 columns). In general, for chromatographic separation, the mobile phases mainly consist of acetonitrile or methanol with water, and could contain a low level of an organic acid or base to help in the ionization of the compounds to be analyzed (Ardrey, 2003). Several spectroscopic techniques, such as ultraviolet (UV), fluorescence and diode-array (DAD), have been used in the LC determination of hormone residues. Nevertheless, the recommended and most used detectors are MS or MS/MS, which separate the analytes by the mass/charge ( $m/z$ ) of the ions. To generate charged ionic analytes, the most used interface ionization techniques are electrospray (ESI) and atmospheric pressure chemical ionization (APCI), in the both negative and positive ion mode (Zhou & Zhang, 2011). The use of ESI or APCI will depend on the analyte that will be analyzed.

The MS/MS detection system uses two stages of mass spectrometry (MS1 and MS2). The first is used to isolate the ion of interest, and the second is used to make the relationship of this ion with the others from which it may have been generated, or which it may generate on decomposition (Ardrey, 2003). Recently, the LC–MS/MS system has become more popular and widely used in food analysis because it has a high specificity and detectability in the determination of the residue levels of contaminants in complex matrices. However, there is the disadvantage that the LC–MS/MS systems are not accessible to all laboratories for routine analysis due to the high cost of the equipment and the requirement of an experienced operator.

Several studies have been reported in the literature that use LC–MS/MS to determine hormone residues in various matrices and to allow for the determination of a large number of compounds with a high precision and sensitivity (Guedes-Alonso et al., 2014, 2017). In this regard, a sample preparation step prior to the chromatographic separation is usually necessary and can include several steps, such as filtration, extraction, clean up, purification, evaporation, hydrolysis, and derivatization, among others (Reis-Filho et al., 2006). A summary of the advantages and disadvantages of the analytical technique used for determination of residues of hormones in several matrices is shown in Table 4.

Extracting hormones from water (including drinking water, ground water, surface water and effluents from sewage treatment plants, sewage, marine sediments, soil and biological sludge) is usually performed by solid phase extraction (SPE) disks or cartridges. Steroid hormones are excreted by humans and reach the aquatic environment through the sewage systems daily. Thus, several authors have stated that municipal wastewater is the main source of the contamination of aquatic

**Table 3.** Analytical techniques for the determination of hormone residues in several matrices.  
**Tabla 3.** Técnicas analíticas para la determinación de residuos de hormonas en varias matrices.

Analytical technique	Matrix	Analyte	Reference
LC/UV-Vis	Fishery products	ETES	Rothbard et al., 1990
	Animal foods	E1, E2, E3, EE	Hu et al., 2010
LC/fluorescence	Sediment	E2, E3, PRO, BOL, TES	Shi et al., 2011
	Water	METES	Falone, 2007
LC/DAD	Yogurt	METES	Falone, 2007
	Fishery products	E2, EE	Fernandes et al., 2011
LC-MS/MS	Water	E2, E3, PRO, BOL, TES	Shi et al., 2011
	Water	E2, E3, DES	Jiang et al., 2009
LC-MS/MS	Food of animal origin	E1, E2, E3, EE, MES, DES	Lopes et al., 2010
	Food of animal origin	E1, E2, E3, EE	de Alda & Barcelo, 2001
GC/MS	Food of animal origin	E1, E2, $\alpha$ E2, E3, EE, PRO, NORET, NORG, MES	Verbinen, Nunes, & Vieira, 2010
	Food of animal origin	E1, E2, E3, EE, DES, METES, TES, TBL, NAN, HEX, DIEN, STAN	Almeida & Nogueira, 2015
GC/MS	Food of animal origin	AP, BOL, CMA, FMT, MPA, MGA, MEBOL, METES, NE, NORG, NORTES, PRO, 16-OH-STAN, TES, TBL	Shao et al., 2005
	Food of animal origin	DES, E2, EE, BOL, NORTES, METES, TBL, TRIAM-ACE, DEXA, FMS, ZELZAL, ZEA, MGA, MLA, MPA	Xu et al., 2006
GC/MS	Food of animal origin	NAD, TBL, BOL, FMT, NAN, AED, MDROL, TES, DHEA, METES, AND, STAN, DHT, NET, 17-HPT, 21-HPT, NORG, MGA, MEPRO, CMA, PRO, MPA, TRIAM, PRD, PRNL, CORT, CORS, FMS, DEXA, FCN-ACE, MPRNL, TRIAM-ACE, FML, BDS, E1, E2, E3, EE, DES, HEX, DIEN, MEST, MEAND, DAN, MESOL, ALD, BEC, FLUA, PCL	Blasco, Poucke, & Peteghem, 2007
	Food of animal origin	CMA, MPA, MGA, PTS, NORG, METES, NAN, Z, E1, E2, $\alpha$ E2, E3, EE, EV, DES	Kaklamanos, Theodoridis, & Dabalís, 2009
GC/MS	Water	AED, TES, PRO, STAN, 17-HPT, PRNL, PRD, HCORT, MGA, MPA	Yang, Shao, Zhang, Wu, & Duan, 2009
	Water	E1, E2, EE	Wang et al., 2010
GC/MS	Fishery products	E1, E2, E3, EE, E1-3S, E1-3G, E2-3S, E2-3G, E2-17G, E2-3G17S, E2-3, 17DiS, E3-3S, E3-3G	Fan et al., 2014
	Fishery products	EPITES, NAN, METES, PTS, MEPRO, PRO, E1, E2, E3, EE	Ingrand, Herry, Beausse, & De Roubin, 2003
GC/MS	Fishery products	COR, E1, $\alpha$ E2, E3, MEPRO, PRO, 17-HPT, TES, NORTES	Isobe et al., 2003
	Fishery products	PRO, NORG, E1, E2, E3, EE, E1-3S	Xu et al., 2006
GC/MS	Foods, meats and vegetables	PRO, NORG, E1, E2, E3, EE, DES, TES, NAN, PRD, PRDNL, BOL, MGA, NORET, CORT	Wang et al., 2012
	Foods, meats and vegetables	P5, ADD, 17-HPT, DHEA, DHT, AND, $\alpha$ E2, E3	Jakimska et al., 2013
GC/MS	Foods, meats and vegetables	DES, Z	Guedes-Alonso et al., 2017
	Foods, meats and vegetables	DES, Z	Hartmann et al., 1998
RIA	Fishery products	DES, Z	Cardoso, Silva, & Santos, 1999
	Fishery products	DES, Z	Long et al., 2007
ELISA	Fishery products	DES, Z	Cardoso et al., 1999
	Fishery products	METES	Specker & Chandlee, 2003
ELISA	Fishery products	METES	Zanardi et al., 2011
	Fishery products	METES	Zanardi et al., 2011

**Abbreviations:** 16-OH-stan (16- $\beta$ -hydroxystanzolol); 17-HPT (17 $\alpha$ -hydroxyprogesterone); 21-HPT (21 $\alpha$ -hydroxyprogesterone); ADD (androstenedione); AED (4-androstene-3,17-dione); ALD (aldosterone); and (androsterone); AP (acetoxyprogesterone); BDS (budesonide); BEC (Beclomethasone); BOL (boldenone); CMA (chlormadinone acetate); COR (cortisol); CORT (cortisone); DAD (ultraviolet diode array); DAN (Danazol); DES (diethylstilbestrol); DEXA (dexamethasone); DHEA (dehydroepiandrosterone); DHT (dihydrotestosterone); DIEN (dienestrol); E1 (estrone); E1-3G (estrone-3-glucuronide); E1-3S (estrone-3-sulfate); E2 (17 $\beta$ -estradiol); E2-17G (estradiol-17-glucuronide); E2-3,17DiS (estradiol-3,17-disulfate); E2-3G (estradiol-3-glucuronide); E2-3S (estradiol-3-sulfate); E3 (estrone); E3-3G (estrone-3-glucuronide); E3-3S (estrone-3-sulfate); EE (17 $\alpha$ -ethynylestradiol); ELISA (enzyme-linked immunosorbent assay); EPITES (epitesterone); EV (estradiol valerate); FCN-ACE (flucinolone acetate); FML (florometholone); FMS (flumethasone); FMT (fluoymesterone); GC-MS (gas chromatography-mass spectrometry); HCORT (hydrocortisone); HEX (hexestrol); LC (Liquid chromatography); MDROL (mestrolone); MEAND (Methylandrosterone); MEBOL (methylboldenone); MEPRO (medroxyprogesterone); MES (mestranol); MESOL (Mestanolone); MEST (Mesterolone); METES (17 $\alpha$ -methyl-testosterone); MGA (megestrol acetate); MPA (medroxyprogesterone acetate); MLA (melengestrol acetate); MPRNL (methylprednisolone); M5/M5 (tandem mass spectrometry); NAD (19-nor-4-Androstene-3,17-dione); NAN (nandrolone); NE (nortestosterone); NET (19-norethandrone); NORET (19-norethisterone); NORG (levonorgestrel ou Norgestrel); NORTES (19-nortestosterone); P5 (pregnenolone); PCL (Clobetasol propionate); PRD (prednisone); PRNL (prednisolone); PRO (progesterone); PTS (testosterone 17-propionate); RIA (radioimmunoassay); STAN (stanzolol); TBL (trenbolone); TES (testosterone); TRIAM-ACE (triamcinolone); TRIAM-ACE (triamcinolone acetate); UV-Vis (ultraviolet-visible spectrophotometry); ZAL (zearalanol); ZEA (zearalanol);  $\alpha$ E2 (17 $\alpha$ -estradiol); Z (Zeranol).

**Table 4.** Advantages and disadvantages of the analytical technique used for determination of residues of hormones in several matrices.**Tabla 4.** Ventajas y desventajas de las técnicas analíticas utilizadas para la determinación de residuos de hormonas en diversas matrices.

Analytical technique	Advantages	Disadvantages	Reference
LC–UV/Vis	<ul style="list-style-type: none"> <li>● Relatively low cost analytical technique;</li> <li>● Easy to operate and interpret the data.</li> </ul>	<ul style="list-style-type: none"> <li>● Low sensitivity and, consequently, high LOQ. For low LOQ's values, a more elaborate sample preparation step is needed to clean-up the extract and to concentrate the analyte (e.g. using SPE cartridges, which are expensive and make the method more laborious);</li> <li>● The UV/Vis detector can detect single wavelength at one time.</li> <li>● The molecule to be analyzed need to present a chromophore in their structure.</li> </ul>	Rothbard et al., 1990 Hu et al., 2010 Shi et al., 2011 Falone, 2007 Fernandes et al., 2011
LC-fluorescence	<ul style="list-style-type: none"> <li>● Relatively low cost analytical technique;</li> <li>● High specificity and selectivity;</li> <li>● Easy to operate and interpret the data.</li> </ul>	<ul style="list-style-type: none"> <li>● To reach low LOQ's values, as well as LC-UV-Vis, a more elaborate sample preparation step is needed (e.g. using SPE cartridges);</li> <li>● The molecule to be analyzed needs to have fluorescence.</li> </ul>	Jiang et al., 2009 Lopes et al., 2010
LC-DAD	<ul style="list-style-type: none"> <li>● Relatively low cost analytical technique;</li> <li>● Easy to operate and interpret the data;</li> <li>● DAD can scan a wavelength range 190–800 nm region.</li> </ul>	<ul style="list-style-type: none"> <li>● As well as LC-UV-Vis, it features low sensitivity and, consequently, high LOQ. For low LOQ's values, a more elaborate sample preparation step is needed (e.g. using SPE cartridges);</li> <li>● The molecule to be analyzed need to present a chromophore in their structure.</li> </ul>	de Alda & Barcelo, 2001 Verbinnen et al., 2010 Almeida & Nogueira, 2015
LC–MS/MS	<ul style="list-style-type: none"> <li>● Ability of determining several analytes (multiresidue method), without the need for elaborated chromatographic separation;</li> <li>● High specificity, selectivity and sensitivity, allowing to achieve low LOQ's values;</li> <li>● No elaborate sample preparation is needed, making the sample preparation step simpler, faster and easier;</li> <li>● There is no need for derivatization steps;</li> <li>● It is possible to analyze both, unconjugated (i.e. free) and conjugates steroid hormones.</li> </ul>	<ul style="list-style-type: none"> <li>● It is an expensive analytical technique;</li> <li>● Usually there is the suppression or increase of signal caused by the competitive ionization between endogenous molecules present in the matrix to be evaluated and the target analyte, called matrix effect.</li> </ul>	Draisci, Palleschi, Ferretti, Lucentini, & Cammarata, 2000 Ferretti et al., 2008 Shao et al., 2005 Xu et al., 2006 Blasco et al., 2007 Kaklamanos et al., 2009 Yang et al., 2009 Wang et al., 2010 Fan et al., 2014 Ingrand et al., 2003 Isobe et al., 2003 Wang et al., 2012 Jakimska et al., 2013 Guedes-Alonso et al., 2017
GC–MS	<ul style="list-style-type: none"> <li>● High sensitivity and specificity, reaching low LOQ's values;</li> <li>● Good chromatographic resolution.</li> </ul>	<ul style="list-style-type: none"> <li>● There is a needed for derivatization of steroids (using silylation, acylation or alkylation), in order to reduce their polarity and thermal instability, which make the method tedious and time-consuming;</li> <li>● Limits the target compounds to unconjugated (i.e. free) estrogens;</li> <li>● Inability to determine some steroids such as estriol.</li> </ul>	Hartmann et al., 1998 Cardoso et al., 1999 Long et al., 2007
ELISA	<ul style="list-style-type: none"> <li>● High selectivity and specificity;</li> <li>● Available as commercial kits;</li> <li>● Easy to perform the assay.</li> </ul>	<ul style="list-style-type: none"> <li>● It is not possible to simultaneously determine multi-residues of steroids in complex matrices;</li> <li>● Although the kits are commercially available, they are expensive;</li> <li>● False positives and negatives are possible.</li> </ul>	Zanardi et al., 2011
RIA	<ul style="list-style-type: none"> <li>● High selectivity and specificity.</li> </ul>	<ul style="list-style-type: none"> <li>● Radiation hazard: need to use radiolabeled reagent;</li> <li>● Requires specially trained person to handle radioactive material;</li> <li>● Laboratories require special licenses to handle radioactive material.</li> </ul>	Specker & Chandlee, 2003 Cardoso et al., 1999

**Abbreviations:** LC–UV/Vis (liquid chromatography–ultraviolet/visible detector); LC-fluorescence (liquid chromatography–fluorescence detector); LC-DAD (liquid chromatography–diode array detector); LC–MS/MS (liquid chromatography–tandem mass spectrometry detector); GC–MS (gas chromatography–mass spectrometry detector); ELISA (enzyme-linked immunosorbent assay); RIA (radioimmunoassay); LOQ (limit of quantification); SPE (solid phase extraction).

environments (Alda, De, Díaz-Cruz, Petrovica, & Barcelo, 2003; Almeida & Nogueira, 2015; Guedes-Alonso et al., 2014).

Even though several articles have been published on the determination of steroid residues in matrices, such as water, sediments and food of animal origin, the determination of hormone residues in fish matrices have been seldom reported, even though fish farming is an activity of growing economic importance. In Table 5, analytical methods using LC and different detector systems to determine the hormone

residues in fishery products are reported. Analysis procedures, including extraction, recovery, detection and quantification limits, and their references for the determination of hormone residues in fishery samples are presented.

Rothbard et al. (1990) were one of the first groups to report hormone residue determination in fish (tilapia muscle). The authors used liquid-solid extraction and obtained low recovery (56.1%) in comparison to that reported later (80.0–94.1%) by Hu, Wang, Chen, Hu, and Li (2010), which used the same detection



Table 5. Analytical method conditions for determination of hormone residues in fishery products by liquid chromatography using different detectors.

Analyte	Matrix	ES	FM	DS	LOD	LOQ	Ref.
ETES	Tilapia muscle	HPLC Bondapak C18 column (7.8 × 300 mm; 10 µm)	H <sub>2</sub> O, ACN + 0.1% TFA 1 mL/min and 2 mL/min	UV-vis	50 ng/g	-	Rothbard et al., 1990
EPITES, NAN, METES, PTS, MEPRO, PRO, E1, E2, E3, EE	Fish muscle	Supelco Discovery C18 column (150 × 2.1 mm; 5 µm)	ACN, H <sub>2</sub> O 0.3 mL/min	MS/MS	0.06–0.22 ng/g	0.12–0.54 ng/g	Xu et al., 2006
E2, E3, DES	Tilapia and prawn	ODS C18 (VP-ODS, 150 × 4.6 mm, 5 µm)	H <sub>2</sub> O, MeOH 1.0 mL/min	FLD	0.023 mg/L	0.076 mg/L	Jiang et al., 2009
E1, E2, E3, EE	Fish and shrimp	Diamonsil C18 column (250 × 4.6 mm, 5 µm)	ACN, 0.12% acetate buffer 1.0 mL/min	UV-Vis	0.98–2.39 µg/L	-	Hu et al., 2010
COR, E1, αE2, E3, MEPRO, PRO, 17-HPT, TES, NORTES	Fish muscle	Zorbax Eclipse C18 column (150 × 4.6 mm, 3.5 µm)	ACN, H <sub>2</sub> O 0.8 mL/min	MS/MS	0.03–0.15 ng/g	0.11–0.47 ng/g	Wang et al., 2012
PRO, NORG, E1, E2, E3, EE, E1-3S	Fish muscle	Acquity UPLC BEH C18 column (50 × 2.1 mm, 1.7 µm)	MeOH, H <sub>2</sub> O (pH 9) 0.4 mL/min	MS/MS	0.002–3.09 ng/g	0.005–9.26 ng/g	Jakimska et al., 2013
PRO, NORG, E1, E2, E3, EE, DES, TES, NAN, PRD, PRDNL, BOL, MGA, NORET, COS	Fish muscle, skin, viscera	Acquity UPLC BEH C18 column (50 × 2.1 mm, 1.7 µm)	MeOH, H <sub>2</sub> O	MS/MS	0.14–49.0 ng/g	3.95 µg/g	Guedes-Alonso et al., 2017

Abbreviations: ES (stationary phase); MP (mobil phase); DS (detection system); LOD (limit of detection); LOQ (limit of quantification); FLD (fluorescence detector); 17-HPT (17α-hydroxyprogesterone); COR (corticosterone); DES (diethylstilbestrol); E1 (estrone); E1-3S (estrone-3-sulfate); E2 (17β-estradiol); E3 (estriol); EE (17α-ethynylestradiol); EPITES (epitesterone); ETES (17α-ethynyltestosterone); MEPRO (medroxyprogesterone); METES (17α-methyl-testosterone); NAN (nandrolone); NORG (norgestrel/levonorgestrel); NORTES (19-nortestosterone); PRO (progesterone); PTS (testosterone 17-propionate); TES (testosterone); αE2 (17α-estradiol); NORET (norethisterone); MGA (megestrol acetate); BOL (boldenone); COS (cortisone); PRD (prednisone); PRDNL (prednisolone).

technique to determine 4 estrogen residues in fish and shrimp. Hu et al. (2010) used molecularly imprinted polymer (MIP)-coated fiber for the extraction of estrogens in the fishery samples and demonstrated a higher extraction efficiency than the three commercial SPME fibers they had investigated (polydimethylsiloxane, polydimethylsiloxane/divinylbenzene and carbowax/template resin fiber). The MIP-coated SPME fiber was very stable, and could be repeatedly used more than 100 times for standard solutions.

Molecularly imprinted polymer (MIP) is a synthetic polymer with specific binding sites of complementary size, shape, and functional groups to the template molecule, which involves a retention mechanism based on molecular recognition. The advantages of MIP (stable, ease of preparation, low cost, and reusability) have led to its wide application in chromatography, catalysis, chemical sensors, and solid-phase extraction. Jiang et al. (2009) used MIP as an SPE sorbent (MISPE) to selectively extract 17β-estradiol from fish and prawn tissues prior to HPLC analysis because MIP showed affinity for this hormone in acetonitrile solution, which was confirmed by absorption experiments. The MISPE method could eliminate all matrix interference simultaneously, and showed good recoveries (78.3–84.5%).

Xu et al. (2006) using LC-MS/MS reported a recovery in the range of 64–75% and low quantitation limits (0.12–0.54 ng g<sup>-1</sup>) for 10 anabolic steroids. The authors describe a relatively simple sample preparation procedure using enzymolysis. Next, the sample was extracted with tert-butyl methyl ether and cleaned up through reverse solid-phase extraction.

Dynamic microwave-assisted extraction-salting-out liquid-liquid extraction (DMAE-SLLE) was an effective technique in reducing the sample preparation time and solvent consumption. DMAE has been developed to avoid the degradation or contamination of the analytes. SLLE is a classic homogeneous liquid-liquid extraction method using salt (ammonium sulfate, sodium chloride, ammonium acetate, sodium sulfate, and magnesium sulfate) to induce phase separation. Thus, Wang, Zhang, Pan, and Chen (2010) used DMAE-SLLE and ammonium acetate as a MS friendly salt and could obtain higher recoveries (75.3–95.4%).

Jakimska et al. (2013) selected three extraction methods for comparison based on their applicability in biota samples. The first extraction protocol consisted of pressurized liquid extraction (PLE) followed by gel permeation chromatography (GPC) clean-up. The second extraction method was a PLE extraction followed by a Florisil clean-up. The third approach was based on QuEChER approach, which involved two steps: liquid-liquid partitioning followed by the application of a specific salt, which was used for salting out water from the sample. The clean-up step with dispersive solid phase extraction (dSPE) was tested using four sorbent mixtures (MgSO<sub>4</sub> and PSA; MgSO<sub>4</sub>, PSA and C18; MgSO<sub>4</sub>, PSA, C18 and GPC; MgSO<sub>4</sub>, PSA and GPC). PLE with GPC clean-up was considered an inefficient method because only five out of the 19 compounds had recoveries higher than 40%, and PLE with Florisil clean-up allowed the extraction of most of the compounds; however, the estrone metabolite (estrone-3-sulfate) was still not extracted from the matrix. Therefore, the QuEChERS approach allowed for the simultaneous extraction of all target compounds, provided satisfactory recoveries and was chosen as the most efficient procedure for sample preparation. The best results were obtained from the combination of the extraction salt composed of sodium

acetate and MgSO<sub>4</sub>, and the dSPE sorbent composed of MgSO<sub>4</sub>, PSA and C18. This combination gave the most satisfactory results since this sorbent mixture is dedicated to samples with high lipid content.

## 5. Conclusions

The use of hormones in fish farming aiming sex reversal and artificial reproduction generally improve the profitability of the fish industry. However, it is recommended that the use of hormones for the purpose of sexual reversion should be established by the legislative framework of the countries that use this technique, in considering human and environmental safety. The most common hormones for sex reversal in fish are 17β-estradiol and 17α-methyl testosterone, which are natural steroids and therefore rapidly excreted soon after the period of the hormonal treatment. On the other hand, breeding induction treatments vary much more in different fish species. Different compounds have been tested over the years to achieve production of high-quality and high-quantity eggs and fingerlings, especially out of season. However, extreme care must be taken to produce the least possible environmental impact and assure food and human health safety, rendering sustainability in the fish sector.

LC coupled to MS in tandem is one of the best alternatives to determine hormonal residues regardless of the matrix. This allows qualitative analysis on a large number of compounds and quantitative analysis of the residues of these compounds with high precision and sensitivity over a shorter analysis time. Currently, the analytical methods intended for the determination of hormone residues in foods, and fish in particular, are focused on the quantitation of several analyte residues simultaneously (multi-residue determination).

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Acknowledgments

To the Brazilian National Council for Scientific and Technological Development for the researcher's scholarship (CNPq process number 306141/2017-5) and to Dr. Patrícia Aparecida de Campos Braga for her technical assistance in the preparation of the final version of the manuscript.

## Funding

This work was supported by São Paulo Research Foundation-Agilent Technologies (under Grant number 2013/50452-5); CAPES/PROEX (under Process number 3300301702P1); and Amazonas Research Foundation (under FAPEAM Grant number 015/2014 PAPAC).

## References

- Adeel, M., Song, X., Wang, Y., Francis, D., & Yang, Y. (2017). Environmental impact of estrogens on human, animal and plant life: A critical review. *Environment International*, 99, 107–119.
- Alda, M. J. L., de Díaz-Cruz, S., Petrovica, M., & Barcelo, D. (2003). Liquid chromatography–(tandem) mass spectrometry of selected emerging pollutants (steroid sex hormones, drugs and alkylphenolic surfactants) in the aquatic environment. *Journal of Chromatography A*, 1000, 503–526.
- Almeida, C., & Nogueira, J. M. F. (2015). Determination of steroid sex hormones in real matrices by bar adsorptive microextraction (BAμE). *Talanta*, 136, 145–154.
- Almeida, F. L. (2013). Endocrinologia aplicada na reprodução de peixes. *Revista Brasileira De Reprodução Animal*, 37(2), 174–180.
- Alves, C., Flores, L. C., Cerqueira, T. S., & Toralles, M. B. P. (2007). Exposição ambiental a interferentes endócrinos com atividade estrogênica e sua associação com distúrbios puberais em crianças [Environmental exposure to endocrine disruptors with estrogenic activity and the association with pubertal disorders in children]. *Cadernos De Saúde Pública*, 23(5), 1005–1014.
- Araújo, J. E. X. S., Streit, D. P., Jr, de Ribeiro, J. S. A., de Martins, E. F. F., de Souza, F. N. C. A. L., Ribeiro, R. P., ... Povh, J. A. (2014). Ovipel and carp pituitary extract as spawning inducers in males of the amazon catfish *Leicurus marmoratus* (Gill, 1970). *Brazilian Archives of Biology and Technology*, 57(6), 882–886.
- Ardrey, R. E. (2003). Liquid chromatography–mass spectrometry: An introduction. In D. J. Ando (Ed.), *Analytical techniques in the sciences*. Chichester, West Sussex, England: John Wiley & Sons, Ltd.
- Babiak, J., Babiak, I., Nes, S. V., Harboe, T., Haugen, T., & Norberg, B. (2012). Induced sex reversal using an aromatase inhibitor, Fadrozole, in Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 324–325, 276–280.
- Bergman, Å., Heindel, J. J., Jobling, S., Kidd, K. A., & Zoeller, R. T. (2013). *State of the science of endocrine disrupting chemicals*. Geneva, Switzerland: WHO Press, World Health Organization.
- Berlinsky, D. L., King, V. W., & Smith, T. I. J. (2005). The use of luteinizing hormone releasing hormone analogue for ovulation induction in black sea bass (*Centropristis striata*). *Aquaculture*, 250, 813–822.
- Bila, D. M., & Dezotti, M. (2003). Fármacos no meio ambiente [Pharmaceutical drugs in the environment]. *Química Nova*, 26(4), 523–530.
- Bila, D. M., & Dezotti, M. (2007). Desreguladores endócrinos no meio ambiente: Efeitos e consequências [Endocrine disruptors in the environment: Part 1 – Effects and consequences]. *Química Nova*, 30(3), 651–666.
- Blasco, C., Poucke, C. V., & Peteghem, C. V. (2007). Analysis of meat samples for anabolic steroids residues by liquid chromatography/tandem mass spectrometry. *Journal of Chromatography A*, 1154, 230–239.
- Cardoso, O. M. C., Silva, T. J. P., & Santos, W. L. M. (1999). Ocorrência de resíduos de dietilstilbestrol e zeranól em fígado de bovinos abatidos no Brasil [Occurrence of diethylstilbestrol and zeranól residues in liver of bovines slaughtered in Brazil]. *Ciência e Tecnologia de Alimentos*, 19(3), 1001–1014.
- Chen, W.-L., Wang, G.-S., Gwo, J.-C., & Chen, C.-Y. (2012). Ultra-high performance liquid chromatography/tandem mass spectrometry determination of feminizing chemicals in river water, sediment and tissue pretreated using disk-type solid-phase extraction and matrix solid-phase dispersion. *Talanta*, 89, 237–245.
- Curtis, L. R., Diren, F. T., Hurley, M. D., Seim, W. K., & Tubb, R. A. (1991). Disposition and elimination of 17-α-methyltestosterone in Nile tilapia. *Aquaculture*, 99(1), 193–201.
- Dasgupta, S., Sarkar, S. K., Sarangi, N., & Bhattacharya, S. (2009). Variation in spawning responses, egg and larvae productions from induced rohu (*Labeo rohita*) during pre-monsoon and monsoon seasons: Relationship with hormonal changes and oocyte responsiveness during final maturation. *Aquaculture*, 290, 320–326.
- de Alda, M. J. L., & Barcelo, D. (2001). Determination of steroid sex hormones and related synthetic compounds considered as endocrine disruptors in water by fully automated on-line solid-phase extraction–liquid chromatography–diode array detection. *Journal of Chromatography A*, 911, 203–210.
- de Bem, J. C. (2009). *Desenvolvimento gonadal inicial e reversão sexual em Astyanaxaltiparanae (Teleostei, characidae)* [Initial gonadal development and sexual reversal in *Astyanaxaltiparanae* (Teleostei, characidae)] (Master's thesis). Retrieved from <https://repositorio.unesp.br/handle/11449/87742>
- DiMaggio, M. A., Broach, J. S., & Ohs, C. L. (2013). Evaluation of Ovaprim and human chorionic gonadotropin doses on spawning induction and egg and larval quality of pinfish. *Lagodonrhomboides*. *Aquaculture*, 414–415, 9–18.
- Draisci, R., Palleschi, L., Ferretti, E., Lucentini, L., & Cammarata, P. (2000). Quantitation of anabolic hormones and their metabolites in bovine serum and urine by liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 870, 511–522.
- Duarte, K. M. R., da Silva, F. M. S. M., & Meirelles, C. F. (2002). Resíduos de anabolizantes na produção animal: Importância e métodos de detecção [Anabolic residues in livestock production: Relevancy and detection methods]. *Ciência Rural*, 32(4), 731–737.

- EC. (1996). Directive 96/22/EC of the Council of the European Union, of 29 April 1996, *Concerning the prohibition on the use in stockfarming of certain substances having a hormonal or thyrostatic action and of beta-agonists, and repealing Directives 81/602/EEC, 88/146/EEC and 88/299/EEC*. Official Journal of the European Communities of 23 May 1996, L 125/3-L 125/9. Retrieved from <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31996L0022&from=en>.
- EC. (2003). Directive 2003/74/EC of the European Parliament and of the Council of the European Union, of 22 September 2003, *Amending Council Directive 96/22/EC concerning the prohibition on the use in stockfarming of certain substances having a hormonal or thyrostatic action and of beta-agonists*. Official Journal of the European Communities of 14 October 2003, L 262/17- L 262/21. Retrieved from <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32003L0074&from=EN>
- EC. (2008). Directive 2008/97/EC of The European Parliament and of the Council of the European Union, of 19 November 2008, *Amending Council Directive 96/22/EC concerning the prohibition on the use in stockfarming of certain substances having a hormonal or thyrostatic action and of beta-agonists*. Official Journal of the European Communities of 28 November 2008, L 318/9- L 318/11. Retrieved from <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008L0097&rid=1>.
- El-Sayed, A. M., Abdel-Aziz, E. H., & Abdel-Ghani, H. M. (2012). Effects of phytoestrogens on sex reversal of Nile tilapia (*Oreochromis niloticus*) larvae fed diets treated with 17 $\alpha$ -methyltestosterone. *Aquaculture*, 360–361, 58–63.
- Falona, S. Z. (2007). *Desenvolvimento de métodos para a determinação do hormônio 17 $\alpha$ -metiltestosterona em amostras de água e de sedimentos de piscicultura: Ensaio ecotoxicológicos com cladóceros* [Development of methods for the determination of 17 $\alpha$ -metiltestosterona in water and sediments samples from pisciculture: Ecotoxicological assay with cladocera] (Doctoral dissertation). Retrieved from <http://www.teses.usp.br/teses/disponiveis/18/18139/tde-09042008-133314/pt-br.php>.
- Fan, Y., Yin, Y., Jiang, W., Chen, Y., Yang, J., Wu, J., & Xie, M. (2014). Simultaneous determination of ten steroid hormones in animal origin food by matrix solid-phase dispersion and liquid chromatography–electrospray tandem mass spectrometry. *Food Chemistry*, 142, 170–177.
- Fernandes, A. N., Giovanela, M., Almeida, C. A. P., Esteves, V. I., Sierra, M. M. D., & Grassi, M. T. (2011). Remoção dos hormônios 17 $\beta$ -estradiol e 17 $\alpha$ -etinilestradiol de soluções aquosas empregando turfas decompostas como material adsorvente [Removal of the hormones 17 $\beta$ -estradiol and 17 $\alpha$ -ethinylestradiol from aqueous solutions employing a decomposed peat as adsorbent material]. *Química Nova*, 34(9), 1526–1533.
- Fernández-Palacios, H., Schuchardt, D., Roo, J., Hernández-Cruz, C., & Izquierdo, M. (2015). Spawn quality and GnRH $\alpha$  induction efficiency in longfin yellowtail (*Seriolarivoltiana*) broodstock kept in captivity. *Aquaculture*, 435, 167–172.
- Ferretti, G., Ferranti, C., Crovella, T., Fiori, M., Civitareale, C., Marchiafava, C., ... Palleschi, L. (2008). Simultaneous analysis of 17 $\alpha$ -estradiol and 17 $\beta$ -estradiol in bovine serum by liquid chromatography–tandem mass spectrometry. *Journal of Chromatography B*, 871, 135–140.
- Filho, L. A. C., Brito, L. T., Cunha, K. L., Júnior, M. P., & Amorim, M. J. A. L. (2016). Técnica de Propagação Artificial em Tambaqui (*Colossoma macropomum*) (Cuvier, 1818) [Technique of Artificial Propagation in Tambaqui (*Colossoma macropomum*) (Cuvier, 1818)]. Retrieved from <https://www.researchgate.net/publication/228458020>
- Flora, M. A. D., Maschke, F., Ferreira, C. C., & De A Pedron, F. (2010). Biologia e cultivo do dourado (*Salminus brasiliensis*) [Biology and culture of dourado fish (*Salminus brasiliensis*)]. *Acta Veterinaria Brasilica*, 4(1), 7–14.
- Garber, A. F., Fordham, S. E., Symonds, J. E., Trippel, E. A., & Berlinsky, D. L. (2009). Hormonal induction of ovulation and spermiation in Atlantic cod (*Gadus morhua*). *Aquaculture*, 296, 179–183.
- Garcia, C. E. O., Araújo, B. C., Mello, P. H., Narcizo, A. M., Rodrigues-Filho, J. A., Medrado, A. T., ... Moreira, R. G. (2013). Involvement of pituitary gonadotropins, gonadal steroids and breeding season in sex change of protogynous dusky grouper, *Epinephelus marginatus* (Teleostei: Serranidae), induced by a non-steroidal aromatase inhibitor. *General and Comparative Endocrinology*, 192, 170–180.
- Guedes-Alonso, R., Montesdeoca-Esponda, S., Sosa-Ferrera, Z., & Santana-Rodríguez, J. J. (2014). Liquid chromatography methodologies for the determination of steroid hormones in aquatic environmental systems. *Trends in Environmental Analytical Chemistry*, 3–4, 14–27.
- Guedes-Alonso, R., Sosa-Ferrera, Z., & Santana-Rodríguez, J. J. (2017). Determination of steroid hormones in fish tissues by microwave-assisted extraction coupled to ultra-high performance liquid chromatography tandem mass spectrometry. *Food Chemistry*, 237, 1012–1020.
- Haffray, P., Petit, V., Guiguen, Y., Quillet, E., Rault, P., & Fostier, A. (2009). Successful production of monosex female brook trout *Salvelinus fontinalis* using gynogenetic sex reversed males by a combination of methyltestosterone immersion and oral treatments. *Aquaculture*, 290, 47–52.
- Hartmann, S., Lacorn, M., & Steinhart, H. (1998). Natural occurrence of steroid hormones in food. *Food Chemistry*, 62(1), 7–20.
- Hayashi, C., Boscolo, W. R., Soares, C. M., & Meurer, F. (2002). Exigência de proteína digestível para larvas de tilápia do Nilo (*Oreochromis niloticus*), durante a reversão sexual [Digestible protein requirement for Nile tilapia larvae (*Oreochromis niloticus*) during the sexual reversal] [suplemento]. *Revista Brasileira De Zootecnia*, 31(2), 823–828.
- Hendry, C. I., Martin-Robichaud, D. J., & Benfey, T. J. (2003). Hormonal sex reversal of Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 219, 769–781.
- Heyrati, F. P., Mostafavi, H., Toloei, H., & Dorafshan, S. (2007). Induced spawning of kutum, *Rutilus frisii kutum* (Kamenskii, 1901) using (D-Ala $_6$ , Pro $_9$ -NEt) GnRH $\alpha$  combined with domperidone. *Aquaculture*, 265, 288–293.
- Hu, Y., Wang, Y., Chen, X., Hu, Y., & Li, G. (2010). A novel molecularly imprinted solid-phase microextraction fiber coupled with high performance liquid chromatography for analysis of trace estrogens in fishery samples. *Talanta*, 80, 2099–2105.
- Ibarra-Castro, L., & Duncan, N. J. (2007). GnRH $\alpha$ -induced spawning of wild-caught spotted rose snapper *Lutjanus guttatus*. *Aquaculture*, 272, 737–746.
- IBGE. (2015). *Produção da Pecuária Municipal – 2015*. Rio de Janeiro, RJ: Instituto Brasileiro de Geografia e Estatística.
- Ingrand, V., Herry, G., Beausse, J., & De Roubin, M. R. (2003). Analysis of steroid hormones in effluents of wastewater treatment plants by liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 1020, 99–104.
- Isobe, T., Shiraiishi, H., Yasuda, M., Shinoda, A., Suzuki, H., & Morita, M. (2003). Determination of estrogens and their conjugates in water using solid-phase extraction followed by liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 984, 195–202.
- Jakimska, A., Huerta, B., Barganska, Z., Kot-Wasik, A., Rodríguez-Mozaz, S., & Barceló, D. (2013). Development of a liquid chromatography–tandem mass spectrometry procedure for determination of endocrine disrupting compounds in fish from Mediterranean rivers. *Journal of Chromatography A*, 1306, 44–58.
- JECFA. (1988). Joint FAO/WHO Expert Committee on Food Additives. *Evaluation of certain veterinary drug residues in food*. Geneva: World Health Organization. WHO Technical Report Series 763.
- JECFA. (1989). Joint FAO/WHO Expert Committee on Food Additives. *Evaluation of certain veterinary drug residues in food*. Geneva: World Health Organization. WHO Technical Report Series 788.
- JECFA. (2000). Joint FAO/WHO Expert Committee on Food Additives. *Evaluation of certain veterinary drug residues in food*. Geneva: World Health Organization. WHO Technical Report Series 893
- Jiang, T., Zhao, L., Chu, B., Feng, Q., Yan, W., & Lin, J. (2009). Molecularly imprinted solid-phase extraction for the selective determination of 17 $\beta$ -estradiol in fishery samples with high performance liquid chromatography. *Talanta*, 78, 442–447.
- Johnstone, R., Macintosh, D. J., & Wright, R. S. (1983). Elimination of orally administered 17 methyltestosterone by *Oreochromis mossambicus* (tilapia) and *Salmo gairdneri* (rainbow trout) juveniles. *Aquaculture*, 35, 249–257.
- Johnstone, R., Simpson, T. H., & Youngson, A. F. (1978). Sex reversal in salmonid culture. *Aquaculture*, 13, 115–134.
- Jürgens, M. D., Holthaus, K. I., Johnson, A. C., Smith, J. J., Hetheridge, M., & Williams, R. J. (2002). The potential for estradiol and ethinylestradiol degradation in English rivers. *Environmental Toxicology and Chemistry*, 21, 480–488.
- Kaklamanos, G., Theodoridis, G., & Dabalís, T. (2009). Determination of anabolic steroids in muscle tissue by liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 1216, 8072–8079.
- Kanemaru, T., Nakamura, M., Murata, R., Kuroki, K., Horie, H., Uchida, K., ... Kagawa, H. (2012). Induction of sexual maturation of the female honeycomb grouper, *Epinephelus merra*, in the non-breeding season



- by modulating environmental factors with GnRH analogue implantation. *Aquaculture*, 358–359, 85–91.
- Lee, C. S., Tamaru, C. S., Banno, J. E., Kelley, C. D., Bocek, A., & Wyban, J. A. (1986). Induced maturation and spawning of milkfish, *Chanos chanos* Forskal, by hormone implantation. *Aquaculture*, 52, 199–205.
- Leonhardt, J. H. (1997). *Efeito da reversão sexual em tilápia do Nilo, Oreochromis niloticus* (Linnaeus, 1757) [Effect of sexual reversal on Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1757)] (Doctoral dissertation). Retrieved from <https://repositorio.unesp.br/bitstream/handle/11449/144133/000105141.pdf?sequence=1>
- Levavi-Sivan, B., Vaiman, R., Sachs, O., & Tzchori, I. (2004). Spawning induction and hormonal levels during final oocyte maturation in the silver perch (*Bidyanus bidyanus*). *Aquaculture*, 229, 419–431.
- Liao, P., Chu, S., Tu, T., Wang, X., Lin, A. Y., & Chen, P. (2014). Persistent endocrine disruption effects in medaka fish with early life-stage exposure to a triazole-containing aromatase inhibitor (Ietrozole). *Journal of Hazardous Materials*, 277, 141–149.
- Lin, S., Benfey, T. J., & Martin-Robichaud, D. J. (2012). Hormonal sex reversal in Atlantic cod. *Gadus morhua*. *Aquaculture*, 364–365, 192–197.
- Liu, J., Wang, R., Huang, B., Lin, C., Wang, Y., & Pan, X. (2011). Distribution and bioaccumulation of steroidal and phenolic endocrine disrupting chemicals in wild fish species from Dianchi Lake, China. *Environmental Pollution*, 159, 2815–2822.
- Long, Z., Hong, L., & Jie, J. (2007). Determination of  $\beta$ -estradiol residues in fish/shellfish muscle by gas chromatography–mass spectrometry. *Chinese Journal of Analytical Chemistry*, 35(7), 983–987.
- Lopes, L. G., Marchi, M. R. R., Souza, J. B. G., Moura, J. A., Lorenzon, C. S., Cruz, C., & Amaral, L. A. (2010). Estrogênios em águas naturais e tratadas da região de Jaboticabal – São Paulo [Estrogens in natural and treated waters in the region of Jaboticabal – São Paulo, Brazil]. *Química Nova*, 33(3), 639–643.
- Mainardes-Pinto, C. S. R., Fenerich-Verani, N., de Campos, B. E. S., & da Silva, A. L. (2000). Masculinização da tilápia do Nilo, *Oreochromis niloticus*, utilizando diferentes rações e diferentes doses de 17  $\alpha$ -metiltestosterona [Masculinization of Nile tilapia, *Oreochromis niloticus*, using different diets and different doses of 17 $\alpha$ -methyltestosterone]. *Revista Brasileira De Zootecnia*, 29(3), 654–659.
- MARA. (1991). Ministério de Estado da Agricultura e Reforma Agrária [Brazil]. Portaria No. 51 de 24 de Maio 1991 [Brazilian Ministry of Agriculture and Agrarian Reform. Regulation No. 51 of 24 May 1991]. Published at *Diário Oficial da União* on 27 May 1991. Retrieved from [http://www3.servicos.ms.gov.br/iagro\\_ged/pdf/356\\_GED.pdf](http://www3.servicos.ms.gov.br/iagro_ged/pdf/356_GED.pdf).
- Marino, G., Panini, E., Longobardi, A., Mandich, A., Finoina, M. G., Zohar, Y., & Mylonas, C. C. (2003). Induction of ovulation in captive-reared dusky grouper, *Epinephelus marginatus* (Lowe, 1834), with a sustained-release GnRH implant. *Aquaculture*, 219, 841–858.
- Melard, C. (1995). Production of a high percentage of male offspring with 17 $\alpha$ -ethynylestradiol sex-reversed *Oreochromis aureus*. I. Estrogen sex-reversal and production of F2 pseudofemales. *Aquaculture*, 130, 25–34.
- Mylonas, C. C., Fostier, A., & Zanuy, S. (2010). Broodstock management and hormonal manipulations of fish reproduction. *General and Comparative Endocrinology*, 166, 516–534.
- Mylonas, C. C., Mitrizakis, N., Castaldo, C. A., Cerviño, C. P., Papadaki, M., & Sigelaki, I. (2013). Reproduction of hatchery-produced meagre *Argyrosus musregius* in captivity. II. Hormonal induction of spawning and monitoring of spawning kinetics, egg production and egg quality. *Aquaculture*, 414–415, 318–327.
- Mylonas, C. C., Papandroulakis, N., Smboukis, A., Papadaki, M., & Divanach, P. (2004). Induction of spawning of cultured greater amberjack (*Serioladu merili*) using GnRH implants. *Aquaculture*, 237, 141–154.
- Nozu, R., Kojima, Y., & Nakamura, M. (2009). Short term treatment with aromatase inhibitor induces sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *General and Comparative Endocrinology*, 161, 360–364.
- Pandian, T. J., & Sheela, S. G. (1995). Review: Hormonal induction of sex reversal in fish. *Aquaculture*, 138, 1–22.
- Park, I., Kim, J., Cho, S. H., & Kim, D. S. (2004). Sex differentiation and hormonal sex reversal in the bagrid catfish *Pseudobagrus fulvidraco* (Richardson). *Aquaculture*, 232, 183–193.
- Passini, G. (2013). *Indução da inversão sexual de machos do robalo-flecha, Centropomus undecimalis, com implantes do hormônio 17 $\beta$ -estradiol* [Induction of the sexual inversion of males of the shootworms, *Centropomus undecimalis*, with 17 $\beta$ -estradiol hormone implants] (Master's thesis). Retrieved from <https://repositorio.ufsc.br/bitstream/handle/123456789/106919/317597.pdf?sequence=1>
- Piferrer, F. (2001). Endocrine sex control strategies for the feminization of teleost fish. *Aquaculture*, 197, 229–281.
- Rasines, I., Gómez, M., Martín, I., Rodríguez, C., Mañanós, E., & Chereguini, O. (2013). Artificial fertilisation of cultured Senegalese sole (*Solea senegalensis*): Effects of the time of day of hormonal treatment on inducing ovulation. *Aquaculture*, 392–395, 94–97.
- Reis-Filho, R. W., de Araújo, J. C., & Vieira, E. M. (2006). Hormônios sexuais estrógenos: Contaminantes bioativos [Sexual estrogenic hormones: Bioactive contaminants]. *Química Nova*, 29(4), 817–822.
- Rothbard, S., Zohar, Y., Zmora, N., Sivan, B. L., Moav, B., & Yaron, Z. (1990). Clearance of 17 $\alpha$ -ethynyltestosterone from muscle of sex-inversed tilapia hybrids treated for growth enhancement with two doses of the androgen. *Aquaculture*, 89(3/4), 365–376.
- Santos, V. B., de Santos, R. S., Salomão, R. A. S., & de Silva, R. M. (2012). Reprodução induzida de pacu (*Piaractus Mesopotamicus*) com o uso de diferentes hormônios comerciais [Induced reproduction of pacu (*Piaractus Mesopotamicus*) with the use of different commercial hormones]. *Pesquisa & Tecnologia*, 9(1), 1–6.
- Shao, B., Zhao, R., Meng, J., Xue, Y., Wu, G., Hu, J., & Tu, X. (2005). Simultaneous determination of residual hormonal chemicals in meat, kidney, liver tissues and milk by liquid chromatography–tandem mass spectrometry. *Analytica Chimica Acta*, 548, 41–50.
- Shi, Y., Peng, D., Shi, C., Zhang, X., Xie, Y., & Lu, B. (2011). Selective determination of trace 17 $\beta$ -estradiol in dairy and meat samples by molecularly imprinted solid-phase extraction and HPLC. *Food Chemistry*, 126, 1916–1925.
- Singh, A. K. (2013). Introduction of modern endocrine techniques for the production of monosex population of fishes. *General and Comparative Endocrinology*, 181, 146–155.
- Specker, J. L., & Chandler, M. K. (2003). Methodology for estradiol treatment in marine larval and juvenile fish: Uptake and clearance in summer flounder. *Aquaculture*, 217, 663–672.
- Sugumar, V., & Munuswamy, N. (2006). Induction of population growth, mictic female production and body size by treatment of a synthetic GnRH analogue in the freshwater rotifer, *Brachionus calyciflorus* Pallas. *Aquaculture*, 258, 529–534.
- Szabó, T., Medgyasszay, C., & Horváth, L. (2002). Ovulation induction in nase (*Chondrostoma toxostoma*, Cyprinidae) using pituitary extract or GnRH analogue combined with domperidone. *Aquaculture*, 203, 389–395.
- Tabata, Y. A., Rigolino, M. G., & Tsukamoto, R. Y. (1999). Produção de lotes monossexos femininos triploides de truta arco-íris, *Oncorhynchus mykiss* (Pisces, Salmonidae). III – Crescimento até idade de primeira maturação sexual [Production of all female triploid rainbow trout, *Oncorhynchus mykiss* (Pisces, Salmonidae). III – Growth up to first sexual maturation]. *Boletim Do Instituto De Pesca*, 25, 67–76.
- Taranger, G. L., Carrillo, M., Schulz, R. W., Fontaine, P., Zanuy, S., Felip, A., & Hansen, T. (2010). Control of puberty in farmed fish. *General and Comparative Endocrinology*, 165, 483–515.
- Targońska, K., Kucharczyk, D., Kujawa, R., Mamcarz, A., & Źarski, D. (2010). Controlled reproduction of asp, *Aspius aspius* (L.) using luteinizing hormone releasing hormone (LHRH) analogues with dopamine inhibitors. *Aquaculture*, 306, 407–410.
- Toyama, G. N., Corrente, J. E., & Cyrino, J. E. P. (2000). Suplementação de vitamina C em rações para reversão sexual da tilápia do nilo [Vitamin C diet supplementation for sex reversal of the Nile tilapia]. *Scientia Agricola*, 57(2), 221–228.
- Vazirzadeh, A., Amiri, B. M., Yelghi, S., Hajimoradloo, A., Nematollahi, M. A., & Mylonas, C. C. (2011). Comparison of the effects of different methods of mammalian and salmon GnRH administration on spawning performance in wild-caught female carp (*Cyprinus carpio carpio*) from the Caspian Sea. *Aquaculture*, 320, 123–128.
- Venturieri, R., & Bernardino, G. (1999). Hormônios na reprodução artificial de peixes [Hormones in artificial fish breeding]. *Panorama Da Aquicultura*, 9(55), 39–48.
- Verbinnen, R. F., Nunes, G. S., & Vieira, E. M. (2010). Determinação de hormônios estrógenos em água potável usando CLAE-DAD [Determination of estrogens in drinking water using HPLC–DAD]. *Química Nova*, 33(9), 1837–1842.
- Wang, H., Zhou, X., Zhang, Y., Chen, H., Li, G., Xu, Y., & Ding, L. (2012). Dynamic microwave-assisted extraction coupled with salting-out liquid–liquid extraction for determination of steroid hormones in fish tissues. *Journal of Agricultural and Food Chemistry*, 60, 10343–10351.



- Wang, Q., Zhang, A., Pan, X., & Chen, L. (2010). Simultaneous determination of sex hormones in egg products by ZnCl<sub>2</sub> depositing lipid, solid-phase extraction and ultra performance liquid chromatography/electrospray ionization tandem mass spectrometry. *Analytica Chimica Acta*, 678, 108–116.
- Wang, Y., Hu, M., Wang, W., Liu, X., Cheung, S. G., Shin, P. K. S., & Song, L. (2009). Effects of GnRH<sub>a</sub> (D-Ala<sub>6</sub>, Pro<sub>9</sub>-NET) combined with domperidone on ovulation induction in wild loach *Misgurnus anguillicaudatus*. *Aquaculture*, 291, 136–139.
- Williams, R. J., Jürgens, M. D., & Johnson, A. C. (1999). Initial predictions of the concentrations and distribution of 17 $\beta$ -estradiol, oestrone and ethynyl oestradiol in 3 English rivers. *Water Research*, 33, 1663–1671.
- Xu, C. L., Chu, X. G., Peng, C. F., Jin, Z. Y., & Wang, L. Y. (2006). Development of a faster determination of 10 anabolic steroids residues in animal muscle tissues by liquid chromatography tandem mass spectrometry. *Journal of Pharmaceutical and Biomedical Analysis*, 41, 616–621.
- Yang, Y., Shao, B., Zhang, J., Wu, Y., & Duan, H. (2009). Determination of the residues of 50 anabolic hormones in muscle, milk and liver by very-high-pressure liquid chromatography–electrospray ionization tandem mass spectrometry. *Journal of Chromatography B*, 877, 489–496.
- Ying, G., Kookana, R. S., & Ru, Y. (2002). Occurrence and fate of hormone steroids in the environment. *Environment International*, 28, 545–551.
- Zanardi, M. F. (2007). *Determinação de resíduo hormonal na carcaça de tilápia do Nilo (Oreochromis niloticus) após reversão sexual* [Hormone residue determination in Nile tilapia carcass (*Oreochromis niloticus*) after sexual reversion] (Master's thesis). Retrieved from [https://repositorio.unesp.br/bitstream/handle/11449/86715/zanardi\\_mf\\_me\\_jabo.pdf?sequence=1](https://repositorio.unesp.br/bitstream/handle/11449/86715/zanardi_mf_me_jabo.pdf?sequence=1)
- Zanardi, M. F., Koberstein, T. C. R. D., Urbinati, E. C., Fagundes, M., Dos Santos, M. A., & Mataqueiro, M. I. (2011). Concentrações de hormônio na carcaça de tilápias-do-nilo e maturação precoce após reversão sexual [Hormone concentration in carcass of Nile tilapia submitted to early maturation after sexual reversion]. *Revista Brasileira De Zootecnia*, 40(1), 7–11.
- Zanoni, M. A., Leal, T. V., Filho, M. C., De Oliveira, C. A. L., & Ribeiro, R. P. (2013). Inversão sexual de alevinos de tilápias do Nilo (*Oreochromis niloticus*) variedade Supreme, submetidos a diferentes temperaturas durante fase de diferenciação sexual [Sex reversal of Nile tilapia (*Oreochromis niloticus*) Supreme variety, fingerlings submitted to different temperatures for sexual differentiation phase]. *Ciências Agrárias*, 34(1), 455–466.
- Zhou, J. L., & Zhang, Z. (2011). Analysis of Hormones in Food. In L. M. L. Nollet (Ed.), *Analysis of endocrine disrupting compounds in food*. Ames, IA: Wiley-Blackwell.
- Zohar, Y., & Mylonas, C. C. (2001). Endocrine manipulations of spawning in cultured fish: From hormones to genes. *Aquaculture*, 197, 99–136.