

Research Article

Cloning, Purification, and Partial Characterization of *Bacillus subtilis* Urate Oxidase Expressed in *Escherichia coli*

Pollyana Pfrimer,¹ Lidia Maria Pepe de Moraes,¹ Aleksandro Sobreira Galdino,¹
Loise Pedrosa Salles,¹ Viviane Castelo Branco Reis,¹ Janice Lisboa De Marco,¹
Maura Vianna Prates,² Carlos Bloch Jr.,² and Fernando Araripe Gonçalves Torres¹

¹Laboratório de Biologia Molecular, Universidade de Brasília, 70910-900 Brasília (DF), Brazil

²Laboratório de Espectrometria de Massa, EMBRAPA Recursos Genéticos e Biotecnologia, 70770-917 Brasília, (DF), Brazil

Correspondence should be addressed to Fernando Araripe Gonçalves Torres, ftorres@unb.br

Received 26 June 2009; Revised 14 October 2009; Accepted 17 November 2009

Academic Editor: Sherry Mowbray

Copyright © 2010 Pollyana Pfrimer et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Urate oxidase (EC 1.7.3.3) is an enzyme involved in purine metabolism which is used in the treatment of gout and as diagnostic reagent for detection of uric acid. In order to produce this enzyme in large quantities for biotechnological purposes, the gene coding for the *Bacillus subtilis* urate oxidase was cloned and heterologously expressed in *Escherichia coli*. Time course induction in *E. coli* showed an induced protein with an apparent molecular mass of ~60 kDa. Soluble recombinant enzyme was purified in a single-step procedure using Ni-NTA column. The enzyme was purified 2.1-fold with a yield of 56% compared to the crude extract. MALDI-TOF analysis revealed an ion with a mass of 58675 Da which is in agreement with the expected mass of the recombinant protein. The purified enzyme showed an optimal pH and temperature of 8.0 and 37°C, respectively, and retained 90% of its activity after 72 hours of incubation at -20°C and 4°C.

1. Introduction

Gout is a disease characterized by joint and kidney inflammation caused by high levels of uric acid in the bloodstream and tissues. Uric acid is water soluble and is normally eliminated by the kidneys in quantities between 600 and 700 mg/day when the organism receives a normal diet. Plasma concentrations of uric acid higher than 6 mg/dL and 7 mg/dL in women and men, respectively, cause hyperuricemia. This medical condition may be caused by a diet rich in proteins, fat, and alcohol as well as hereditary factors (inborn errors of purine-pyrimidine metabolism) [1]. There are other diseases associated to hyperuricemia such as type 2 diabetes which, in the presence of higher levels of uric acid, causes insulin absorption resistance in tissues, increase of nephropathy progression, and tumor lysis syndrome (TLS) [2–4].

Urate oxidase (uricase, EC 1.7.3.4) is an enzyme with copper bonds that catalyze the oxidative opening of the purine ring of uric acid to form allantoin which is 5–10 times more soluble than uric acid [5, 6]. This enzyme can be used therapeutically to reduce toxic urate accumulation.

Rasburicase (Fasturtec/Elitek) is a market name given to the *Aspergillus flavus* urate oxidase expressed in *Saccharomyces cerevisiae* which has been approved for clinical use. Studies showed that Rasburicase is more effective than other drugs in the treatment of hyperuricemia due to low incidence of hypersensitivity reaction [7, 8]. Urate oxidase is also used as a reagent in clinical diagnostic kits for enzymatic determination of uric acid. Native enzyme may be obtained from several microorganisms of the genus *Micrococcus*, *Brevibacterium*, *Streptomyces*, *Candida*, *Bacillus*, *Pseudomonas*, *Arthrobacter*, and *Aspergillus* [9–15]. Urate oxidase from thermophilic *Bacillus* sp. TB-90 has been extensively studied for diagnostic purposes since it exhibits high activity and thermostability in a wide range of pHs [16]. Purified recombinant *Bacillus subtilis* urate oxidase has been tested with success in a biochip system for diagnostic purposes [17]. In this case, the *pucLM* gene was cloned in-frame with the maltose-binding protein (MPB) coding sequence resulting in a large protein fusion (~98 kDa) which was not significantly overexpressed in *Escherichia coli*. This result prompted us to employ a different approach to improve the

expression of *B. subtilis* urate oxidase in *E. coli* by using a smaller 6x His tag which would also enable simple and fast purification of the recombinant enzyme for further biochemical characterization and large-scale production.

2. Materials and Methods

2.1. Strains and Reagents. *Bacillus subtilis subtilis* LMD 69.3 was used as a source of the urate oxidase gene. For heterologous expression, *E. coli* BL21 (DE3) pLysS and the expression vector pET21a (Novagen) were used. Ni Sepharose 6 Fast Flow (GE Healthcare) resin was used to purify recombinant urate oxidase. Restriction enzymes and the pGEM-T cloning vector were from Promega. Plasmid extraction was performed by the use of QIAprep Spin Mini Kit (Qiagen). T4 DNA ligase was obtained from New England Biolabs. *Taq* Platinum was purchased from Invitrogen. Monoclonal antipoly histidine AP antibody was purchased from Sigma. Other reagents were obtained from standard commercial sources.

2.2. DNA Manipulations. *B. subtilis* chromosomal DNA was extracted as described elsewhere [18]. The urate oxidase coding sequence (1485 pb) was obtained by PCR using primers PUCL5 (5'-CGGATCCATGTTTACAATGGATGACCTG-3') and PUCL3 (5'-GCTCGAGGGCTTTTCAGGCTCCGACAT-3') which were designed based on the *B. subtilis* *puclM* gene sequence (GeneBank gi: 32468813). Primer PUCL5 contained a *Bam*HI site (underlined) and the translation initiation codon (bold), whereas PUCL3 incorporated an engineered *Xho*I site (underlined). Amplification was performed using the following conditions: 30 cycles of 1 minute at 94°C, 45 seconds at 50°C, and 1.5 minutes at 72°C. The purified PCR product was ligated into pGEM-T resulting in plasmid pGEMURI. The sequence of the cloned gene was verified by automated DNA sequencing on the Genetic Analyzer Mega Bace 1000 (GE Healthcare) using the Mega Bace Dye Terminator kit. Plasmid pGEMURI was digested with *Bam*HI and *Xho*I and the *puclM* gene was ligated into pET21a digested with the same enzymes. Because primer PUCL3 lacks a stop codon, the urate oxidase gene was cloned in-frame with the 6x His tag coding sequence present in pET21a. The resulting plasmid, pETURI, was used to transform *E. coli* BL21 (DE3) pLysS.

2.3. Heterologous Expression. A single *E. coli* colony transformed with pETURI was inoculated in 6 mL Luria-Bertolin (LB) medium containing 100 µg/mL ampicillin and incubated at 37°C with shaking (200–300 rpm) overnight. Five milliliters of this preculture were transferred to 50 mL of LB medium in a 250 mL shake flask. The culture was grown under the same condition until OD₆₀₀ reached 0.6 when IPTG was added to a final concentration of 1 mM. One milliliter samples were collected at different intervals and cell pellet was resuspended in 50 µL of SDS-PAGE sample buffer and boiled for 5 minutes prior to gel electrophoresis analysis. Typically, 1–5 µg total protein were loaded in each lane.

2.4. Western Blotting. Protein samples were separated on 12% SDS-PAGE gels and electroblotted onto PVDF membrane. The membrane was blocked with 5% milk in 10 mM Tris-HCl with 150 mM NaCl (pH 8.0) and 0.1% Tween 20 (TBST) for 2 hours at room temperature. The membrane was then incubated for 2 hours at room temperature with monoclonal mouse antipoly histidine AP antibody diluted 1 : 1000 in TBST. After three washes with TBST, bands were visualized by using the NBT/BCIP detection method.

2.5. Recombinant Enzyme Purification. In order to purify and analyze the recombinant protein, one bacterial clone was induced with 1 mM IPTG in 100 mL LB during 3 hours. The cells were harvested by centrifugation at 5000 rpm for 20 minutes at 4°C. Cell pellet (1.5 g) was resuspended in 2 mL lysis buffer (10 mM imidazole, 50 mM NaH₂PO₄ [pH 8.0], 300 mM NaCl) and lysed by sonication (6 cycles of 10 seconds of pulses at 45% amplitude—59W). The suspension was centrifuged at 13000 rpm for 10 minutes and the supernatant loaded into an Ni-NTA column with a volume of 1 mL resin preequilibrated in lysis buffer. The column was washed 8 times with 1 mL wash buffer (20 mM imidazole, 50 mM NaH₂PO₄ [pH 8.0], 300 mM NaCl) and 4 times with 500 µL elution buffer (200 mM imidazole, 50 mM NaH₂PO₄ [pH 8.0], 300 mM NaCl). All fractions were stored at 4°C for further analysis.

2.6. MALDI TOF Mass Determinations. Molecular mass determinations were performed on an Ultraflex II spectrometer (Bruker Daltonics) using lyophilized sample dissolved in Milli-Q water and mixed separately with two different saturated matrices solutions, α -cyano-4-hydroxycinnamic acid, and 3,5-dimethoxy-4-hydroxycinnamic acid (sinapinic acid). MALDI spectra were acquired in linear mode with external calibration.

2.7. Urate Oxidase Assay. Urate oxidase degrades uric acid forming hydrogen peroxide and allantoin. Hydrogen peroxide reacts with 4-aminoantipyrine and 3,5-dichloro-2-hydroxybenzenesulfonic acid and forms quinoneimine, a red compound that is detected by an increase in absorbance at 555 nm. The enzymatic reaction was carried out at 37°C and contained 200 µL uric acid solution (10 mg in 100 mL of 10 mM Tris-HCl, pH 8.0), 50 µL distilled water, and 50 µL enzyme samples. After 15-minute reaction, the reagent solution (4 mM 4-aminoantipyrine, 2 mM 3,5-dichloro-2-hydroxybenzenesulfonic acid, 2 mM horseradish peroxidase, and 10 mM Tris-HCl buffer) was added, incubated at room temperature for 5 minutes, and measured at 555 nm. One unit of urate oxidase activity was defined as the amount of enzyme that catalyzes the transformation of 1 µmol of urate per minute at 37°C (pH 8.0). Urate oxidase activity was calculated using the math formula found at the Kikkoman Biochemicals website (<http://www.kikkoman.co.jp/bio/e/common/rinsyou.html>). Protein concentration was determined using the Bradford method [19].




FIGURE 1: Small-scale expression and time course study of urate oxidase expression in *E. coli*. Coomassie Blue stained 12% SDS-PAGE (a) and respective Western blotting analysis (b) of urate oxidase induction in *E. coli*. Lane M, protein BenchMarker (Invitrogen); Lane 1, supernatant of uninduced cells; Lanes 2–6, supernatant of induced cells after 15, 30, 60, 90, and 120 minutes, respectively. The arrow indicates the position of the band corresponding to recombinant urate oxidase (rUox).

2.8. Enzyme Characterization. Enzyme activity as a function of pH was tested with 50 mM sodium citrate for the followings values of pH: 2.5, 3.0, 4.5, 5.0; 50 mM sodium phosphate for pH values 6.0 and 7.0; 50 mM Tris-HCl for pH values 8.0, 9.0 and 10.0. The optimum temperature was evaluated by measuring urate oxidase activity for 30 minutes at different temperatures: 25, 37, 40, 60, and 80°C. The effect of temperature on enzyme stability was determined by measuring the residual activity after 0, 12, 24, 48 and 72 hours of preincubation in 10 mM Tris-HCl (pH 8.0) at –20, 4 and 37°C.

3. Results

A ~1.5 kb amplicon corresponding to the *B. subtilis pucLM* gene was successfully cloned, sequenced and transferred to the bacterial expression vector pET21a (data not shown). In order to analyze the kinetics of urate oxidase expression, a colony of *E. coli* cells harbouring pETURI was grown to mid-log and induced with IPTG for 0, 15, 30, 60, 90, and 120 minutes. Samples analyzed by SDS-PAGE revealed a ~60 kDa induction band (Figure 1(a)). The size of this band was consistent with the predicted mass of the recombinant enzyme. Western blot analysis showed that the induction band represented a protein which contained a 6x His tag (Figure 1(b)). Two smaller (<50 kDa) and less abundant bands were also observed in the Western blot (Figure 1(b)). Induction profile showed a progressive accumulation of the enzyme starting 15 minutes after addition of IPTG and reaching the highest peak 90 minutes later. After induction in 100 mL, the cleared cell lysate was chromatographed in a Ni-NTA resin column in order to purify recombinant urate oxidase in a single-step. Fractions collected in different steps of purification were analyzed by SDS-PAGE (Figure 2). In order to optimize the purification procedure of the enzyme different elution buffers containing increasing concentrations of imidazole (50, 100, 150, 200 mM) were tested (data not shown). After elution with 200 mM imidazole, a predominant band with an apparent molecular mass of ~60 kDa was observed (Figure 2, lane 6). The data presented in Table 1 shows that the enzyme was purified with a 2.1 fold increase in specific activity with a yield of ~58% as compared to the crude extract.




FIGURE 2: Recombinant enzyme purification. Fractions collected before and after Ni-NTA chromatography were visualized on 12% SDS-PAGE. Lane M, protein BenchMarker (Invitrogen); Lane 1, crude extract of uninduced *E. coli* cells; Lane 2, crude cell extract after induction for 3 hours; Lane 3, cellular extract (input); Lane 4, flow-through; Lane 5, washing step (20 mM imidazole); Lane 6, purified urate oxidase (~30.5 µg) after elution with 200 mM imidazole.

MALDI-TOF/MS analysis of the purified enzyme fraction revealed two major ions $M+H^+ = 58675$ Da and its double charged $M+2H^+ = 29338.1$ Da (Figure 3). The effects of pH and temperature on enzyme activity are shown in Figures 4(a) and 4(b), respectively. Purified urate oxidase showed maximum activity at pH 8.0 and 37°C (Figure 4(b)). As shown in Figure 5, the residual activity was practically constant at –20°C and 4°C after a 72 hours incubation period but was significantly affected by incubation at 37°C.

4. Discussion

Urate oxidase is an enzyme conserved in many species ranging from microorganisms to mammals; however, primates have lost this activity as consequence of evolutionary gene mutations [20, 21]. This enzyme catalyzes the oxidative reaction that converts urate to allantoin, a more soluble and easily excreted compound. Due to this property urate oxidase may be used for both therapeutical and diagnostic purposes [22].

TABLE 1: Summary of the purification steps.

Step	Total activity (IU)	Total protein (mg)	Specific activity (IU/mg)	Purification (fold)	Yield (%)
Crude Extract	54.0	3.0	18.0	1	100
Ni-NTA	31.2	0.8	39.0	2.1	56




FIGURE 3: Mass spectrometry. MALDI-TOF/MS spectrum of recombinant urate oxidase using sinapinic acid as matrix showed two related ions for the same polypeptide chain, $M+2H^+ = 29338.1$ Da and $M+H^+ = 58675$ Da.

Many sources of urate oxidase are available, but commercial production may be hampered by low productivity and difficulties in protein purification for clinical applications [22, 23]. DNA technology has been employed to overcome these difficulties by expressing urate oxidase genes in heterologous systems [24].

In *B. subtilis* the genes for purine degradation are clustered at positions 284 to 285° in the genome map [25, 26]. Mutational analysis of this region showed that inactivation of *puLM* resulted in a defect in the utilization of guanosine, hypoxanthine, and uric acid and was thus proposed to code for urate oxidase [26]. Although the protein coded from the *puLM* gene shows amino acid identity to other urate oxidases, this similarity is restricted to the C-terminal domain (amino acids 171–494) [26] with the highest identities being with *Bacillus* sp. TB-90 (66%), *Bacillus clausii* (63%), and *Bacillus fastidiosus* (55%). It has been proposed that the first 170 amino acids correspond to a peroxide reductase domain that could be involved in the removal of hydrogen peroxide, a product of the oxidation of uric acid to allantoin catalyzed by urate oxidase [26]. Unlike eukaryotic urate oxidases and similar to *Bacillus* sp. TB-90, the *B. subtilis* enzyme does not have the typical type 2 copper binding motif H-X-H-X-F [27].

In this work, the *puLM* was successfully expressed in *E. coli* in a soluble and active form. This result is particular important for the commercial production of this enzyme

since it eliminates the laborious denaturing/refolding steps often necessary to solubilize proteins present as inclusion bodies. Furthermore, the strategy used in this work involved the fusion of *B. subtilis* urate oxidase to a 6x His tag in order to allow simple one-step purification using Ni-NTA column. A derivation of this method using Ni ion-chelating magnetic beads to increase yield was reported for the purification of recombinant *Bacillus* sp. TB-90 urate oxidase [28]. The results presented in Table 1 show that the recombinant enzyme was purified almost to homogeneity and the yield was about 58% as compared to the crude extract. This yield was close to that reported for commercial Rasburicase (61%) and native *B. subtilis* urate oxidase (67%) [28, 29]. The smaller protein contaminants that copurified with the recombinant enzyme represent minor breakdown products because they were recognized by the antipoly histidine antibody (Figure 1(b)). The recombinant enzyme was submitted to biochemical characterization (molecular mass determination, thermostability, optimum temperature and pH) and in vitro enzymatic activity. The experimental molecular mass value of 58675 Da obtained by MALDI-TOF/MS for the recombinant enzyme is close to the theoretical values calculated for the predicted recombinant protein (58.9 kDa). The optimum temperature (37°C) and pH (8.0) were similar to that observed with *A. flavus* urate oxidase produced in *E. coli* [30] and the *B. subtilis* MPB-uricase fusion protein [17]. In enzymatic assays the *B. subtilis* urate oxidase exhibited a specific activity of 39 IU/mg protein, a value higher than those observed for other urate oxidase described in the literature under optimal conditions. For example, native *B. subtilis* urate oxidase specific activity was 5.29 IU/mg protein, the specific activity for the *A. flavus* urate oxidase expressed in *E. coli* was 26.96 IU/mg and the commercial Rasburicase was 26.98 IU/mg protein [30]. Thermostability is considered an important and useful criterion for industrial application of enzymes. The stability of the recombinant enzyme at the temperatures used to store commercial enzymatic diagnostic kits (−20°C and 4°C) was evaluated and the enzyme kept 80% of its residual activity after 72 hours of pre-incubation. However, at 37°C, 50% of the residual activity was lost after 12 hours which is in agreement with the Rasburicase half life of 19 hours reported in humans [30, 31]. It has been proposed that the human body temperature is responsible for the shorter circulating half-time exhibited by recombinant urate oxidase [30, 31].

The results presented in this work showed the feasibility of producing soluble recombinant urate oxidase at high levels. The purity of the enzyme together with its high activity and thermostability shows that the production of the *B. subtilis* urate oxidase in *E. coli* should be considered for the development of diagnostic kits and for clinical purposes.




FIGURE 4: Effect of pH and temperature on enzyme activity. Purified urate oxidase was incubated under different pHs (a) and temperatures (b) and assayed as described in Section 2.




FIGURE 5: Thermal stability. Purified enzyme was pre-incubated at -20°C , 4°C and 37°C for the indicated times, and assayed at 37°C . Residual activity corresponds to the percentage of activity determined for the nonincubated samples.

Acknowledgment

This work was supported by CNPq and FAP/DF.

References

- [1] H. R. Schumacher Jr. and L. X. Chen, "Newer therapeutic approaches: gout," *Rheumatic Disease Clinics of North America*, vol. 32, no. 1, pp. 235–244, 2006.
- [2] M. Modan, H. Halkin, A. Karasik, and A. Lusky, "Elevated serum uric acid—a facet of hyperinsulinaemia," *Diabetologia*, vol. 30, no. 9, pp. 713–718, 1987.
- [3] S. Bo, P. Cavallo-Perin, L. Gentile, E. Repetti, and G. Pagano, "Hypouricemia and hyperuricemia in type 2 diabetes: two different phenotypes," *European Journal of Clinical Investigation*, vol. 31, no. 4, pp. 318–321, 2001.
- [4] D.-H. Kang, T. Nakagawa, L. Feng, et al., "A role for uric acid in the progression of renal disease," *Journal of the American Society of Nephrology*, vol. 13, no. 12, pp. 2888–2897, 2002.
- [5] L. Fraisse, M. C. Bonnet, J. P. De Farcy, C. Agut, D. Dersigny, and A. Bayol, "A colorimetric 96-well microtiter plate assay for the determination of urate oxidase activity and its kinetic parameters," *Analytical Biochemistry*, vol. 309, no. 2, pp. 173–179, 2002.
- [6] Y. Lee, D. H. Lee, C. W. Kho, et al., "Transthyretin-related proteins function to facilitate the hydrolysis of 5-hydroxyisourate, the end product of the uricase reaction," *FEBS Letters*, vol. 579, no. 21, pp. 4769–4774, 2005.
- [7] C.-H. Pui, H. H. Mahmoud, J. M. Wiley, et al., "Recombinant urate oxidase for the prophylaxis or treatment of hyperuricemia in patients with leukemia or lymphoma," *Journal of Clinical Oncology*, vol. 19, no. 3, pp. 697–704, 2001.
- [8] B. Vogt, "Urate oxidase (rasburicase) for treatment of severe tophaceous gout," *Nephrology Dialysis Transplantation*, vol. 20, no. 2, pp. 431–433, 2005.
- [9] A. A. Ims̆enecki and L. S. Popova, "Decomposition of uric acid by aerobic bacteria isolated from soil," *Mikrobiologiya*, vol. 40, no. 2, pp. 269–274, 1971.
- [10] J. Kida and M. Kuniyama, "Studies on bacterial uricase—part I: isolation of uricase producing bacteria and some cultural conditions for production," *Journal of Fermentation Technology*, vol. 44, pp. 789–796, 1966.
- [11] W. Frank and G. E. Hahn, "Uricase chungen zum bakteriellen purin uber den abbau von amino-, oxy-, and methylpurinen durch *Pseudomonas aerogenosa* (*B. pyocyaneum*)," *Zeitschrift für Physiologische Chemie*, vol. 301, pp. 90–106, 1955.
- [12] K. Suzuki, S.-I. Sakasegawa, H. Misaki, and M. Sugiyama, "Molecular cloning and expression of uricase gene from *Arthrobacter globiformis* in *Escherichia coli* and characterization of the gene product," *Journal of Bioscience and Bioengineering*, vol. 98, no. 3, pp. 153–158, 2004.
- [13] Y. Watanabe, M. Yano, and J. Fukumoto, "Studies on the formation of uricase by *Streptomyces*—part I: the effect of purine bases on the induced formation of uricase by the cultured cells," *Agricultural & Biological Chemistry*, vol. 33, no. 2, pp. 1282–1290, 1969.
- [14] A. Tanaka, M. Yamamura, S. Kawamoto, and S. Fukui, "Production of uricase by *Candida tropicalis* using *n*-alkane as a substrate," *Applied and Environmental Microbiology*, vol. 34, no. 4, pp. 342–346, 1977.
- [15] G. P. A. Bongaerts and G. D. Vogels, "Uric acid degradation by *Bacillus fastidiosus* strains," *Journal of Bacteriology*, vol. 125, no. 2, pp. 689–697, 1976.
- [16] K. Yamamoto, Y. Kojima, T. Kikuchi, et al., "Nucleotide sequence of the uricase gene from *Bacillus* sp. TB-90," *Journal of Biochemistry*, vol. 119, no. 1, pp. 80–84, 1996.
- [17] S.-H. Huang, Y.-C. Shih, C.-Y. Wu, et al., "Detection of serum uric acid using the optical polymeric enzyme biochip system,"

- Biosensors and Bioelectronics*, vol. 19, no. 12, pp. 1627–1633, 2004.
- [18] J. Marmur and C. L. Schildkraut, “Growth of bacteria labelled with heavy isotopes for the isolation of nucleic acids,” *Nature*, vol. 189, no. 4765, pp. 636–638, 1961.
- [19] M. M. Bradford, “A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding,” *Analytical Biochemistry*, vol. 72, no. 1-2, pp. 248–254, 1976.
- [20] X. Wu, D. M. Muzny, C. C. Lee, and C. T. Caskey, “Two independent mutational events in the loss of urate oxidase during hominoid evolution,” *Journal of Molecular Evolution*, vol. 34, no. 1, pp. 78–84, 1992.
- [21] M. Oda, Y. Satta, O. Takenaka, and N. Takahata, “Loss of urate oxidase activity in hominoids and its evolutionary implications,” *Molecular Biology and Evolution*, vol. 19, no. 5, pp. 640–653, 2002.
- [22] S. Chohan and M. A. Becker, “Update on emerging urate-lowering therapies,” *Current Opinion in Rheumatology*, vol. 21, no. 2, pp. 143–149, 2009.
- [23] W. A. Lotfy, “Production of a thermostable uricase by a novel *Bacillus thermocatenulatus* strain,” *Bioresource Technology*, vol. 99, no. 4, pp. 699–702, 2008.
- [24] M. Hummel, D. Buchheidt, S. Reiter, J. Bergmann, K. Adam, and R. Hehlmann, “Recurrent chemotherapy-induced tumor lysis syndrome (TLS) with renal failure in a patient with chronic lymphocytic leukemia-successful treatment and prevention of TLS with low-dose rasburicase,” *European Journal of Haematology*, vol. 75, no. 6, pp. 518–521, 2005.
- [25] F. Kunst, N. Ogasawara, I. Moszer, et al., “The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*,” *Nature*, vol. 390, no. 6657, pp. 249–256, 1997.
- [26] A. C. Schultz, P. Nygaard, and H. H. Saxild, “Functional analysis of 14 genes that constitute the purine catabolic pathway in *Bacillus subtilis* and evidence for a novel regulon controlled by the PucR transcription activator,” *Journal of Bacteriology*, vol. 183, no. 11, pp. 3293–3302, 2001.
- [27] X. Wu, C. C. Lee, D. M. Muzny, and C. T. Caskey, “Urate oxidase: primary structure and evolutionary implications,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 86, no. 23, pp. 9412–9416, 1989.
- [28] Y. Nishiya, T. Hibi, and J.-I. Oda, “A purification method of the diagnostic enzyme *Bacillus* uricase using magnetic beads and non-specific protease,” *Protein Expression and Purification*, vol. 25, no. 3, pp. 426–429, 2002.
- [29] H. Y. Shin, H. J. Kang, E. S. Park, et al., “Recombinant urate oxidase (rasburicase) for the treatment of hyperuricemia in pediatric patients with hematologic malignancies: results of a compassionate prospective multicenter study in Korea,” *Pediatric Blood and Cancer*, vol. 46, no. 4, pp. 439–445, 2006.
- [30] J. Li, Z. Chen, L. Hou, et al., “High-level expression, purification, and characterization of non-tagged *Aspergillus flavus* urate oxidase in *Escherichia coli*,” *Protein Expression and Purification*, vol. 49, no. 1, pp. 55–59, 2006.
- [31] L. Cammalleri and M. Malaguarnera, “Rasburicase represents a new tool for hyperuricemia in tumor lysis syndrome and in gout,” *International Journal of Medical Sciences*, vol. 4, no. 2, pp. 83–93, 2007.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

