



Parasites in Loricariidae from Brazil: checklist and new records for fish from the Brazilian Amazon

William Felix Borges¹, Marcos Sidney Brito de Oliveira², Gracienhe Gomes Santos³ and Marcos Tavares-Dias^{2,4*}

¹Universidade do Estado do Amapá, Macapá, Amapá, Brazil. ²Universidade Federal do Amapá, Macapá, Amapá, Brazil. ³Faculdade de Macapá, Macapá, Amapá, Brazil. ⁴Empresa Brasileira de Pesquisa Agropecuária, Rod. Juscelino Kubitschek, Km 5, 2600, 68903-419, Macapá, Amapá, Brazil. *Author for correspondence. E-mail: marcos.tavares@embrapa.br

ABSTRACT. The aim of this study was to investigate the parasites fauna of *Ancistrus leucostictus*, *Hypostomus ventromaculatus*, *Ancistrus* sp. and *Hemiancistrus* sp. from the Igarapé Fortaleza River (Amapá State, Brazil), besides making a checklist of the parasite species in Loricariidae from Brazil. A total of 53 fishes were collected from November 2013 to August 2014. In the hosts, a total of 1,559 parasites of seven taxa were collected: *Unilatus unilatus*, *Trinigyryus mourei*, undetermined metacercariae, *Genarchella gemachella*, *Raphidascaris (Sprentascaris)* sp., *Gorythocephalus elongorchis* and *Proteocephalus* sp. Ectoparasite species were frequent in the examined Loricariidae species, which also had larval stages of endoparasites. The hosts with the highest sampled number, *H. ventromaculatus* and *Ancistrus* sp., had the highest parasite species richness. Loricariidae species from Brazil are parasitized by species of Protozoa, Monogenea, Nematoda, Digenea, Acantocephala, Cestoda, Crustacea and Hirudinea, but monogeneans, digeneans and nematodes were the predominant taxa.

Keywords: armored catfish ectoparasites; freshwater fish; Loricariidae.

Parasitos de Loricariidae do Brasil: checklist e novos registros para peixes da Amazônia brasileira

RESUMO. O objetivo deste estudo foi descrever a fauna parasitária de *Ancistrus leucostictus*, *Hypostomus ventromaculatus*, *Ancistrus* sp. e *Hemiancistrus* sp. da bacia Igarapé Fortaleza (Estado do Amapá, Brasil), além de fazer um checklist das espécies de parasitos em Loricariidae do Brasil. Foram coletados 53 peixes de novembro de 2013 a agosto de 2014. Nos hospedeiros foram coletados um total de 1.559 parasitos distribuídos em sete táxons: *Unilatus unilatus*, *Trinigyryus mourei*, metacercárias não identificadas, *Genarchella gemachella*, *Raphidascaris (Sprentascaris)* sp., *Gorythocephalus elongorchis* e *Proteocephalus* sp. Espécies de ectoparasitos foram frequentes nas espécies de Loricariidae examinadas, que também apresentaram estágios larvais de endoparasitos. Hospedeiros com maior número amostral, *H. ventromaculatus* e *Ancistrus* sp., apresentaram maior riqueza de espécies de parasitos. Espécies de Loricariidae do Brasil são parasitados por espécies de Protozoa, Monogenea, Nematoda, Digenea, Acantocephala, Cestoda, Crustacea e Hirudinea, mas monogeneas, digeneas e nematoides são os táxons predominantes.

Palavras-chave: cascudos; ectoparasitos; peixes de água doce; Loricariidae.

Introduction

Loricariidae is the most species-rich of freshwater fishes, with more than 915 valid species, distributed in 106 genera and six subfamilies (Lithogeneinae, Neoplecostominae, Ancistrinae, Hypoptopomatinae, Hypostominae and Loricariinae) and occurs in Costa Rica, Panama and South America (Alonso, Terán, Aguilera, & Mirande, 2016; Nelson, Grande, & Wilson, 2016). However, the highest occurrence of Loricariidae species is in the Amazon River basin system (Soares et al., 2011). Fish species of this family have size

ranging from 2.5 to 61.0 cm and detritivorous feeding habit, since they feed on debris, algae and invertebrates associated with the sediments of water bodies (Soares et al., 2011). Some species of Loricariidae such as *Ancistrus leucostictus* Günther, 1864; *Ancistrus* Rafinesque, 1815, *Hypostomus ventromaculatus* Boeseman, 1968 and *Hemiancistrus* Bleeker, 1862 are known for the Igarapé Fortaleza, a tributary of the Amazon River in the State of Amapá, eastern Amazon region (Brazil), the locality of this study. Loricariid species are fish that occupy a lower position in the food chain (Soares et al., 2011), thus they have a low parasitic fauna (Gonçalves, Oliveira, Santos, & Tavares-Dias, 2014).

Parasites play an important role in ecosystems, since they can regulate the abundance of host fish populations, destabilize food chains and alter the structure of communities of hosts (Luque & Poulin, 2007; Lagrue, Kelly, Hicks, & Poulin, 2011; Cardoso, Oliveira, Neves & Tavares-Dias, 2017; Baia, Florentino, Silva & Tavares-Dias, 2018). As parasites are components of most ecosystems, occurring in all food webs and at all trophic levels, several vertebrate and invertebrate species serve as hosts for one or more species of parasites (Lagrue et al., 2011; Cardoso et al., 2017; Baia et al., 2018). In wild fish populations, parasite communities differ in richness and diversity according to their behavior, diet of hosts and the parasite life cycle. The different species of fish living in sympatry may present a similar pattern of parasites when compared to species living in allopatry (Cardoso et al., 2017). Thus, this study compared the parasite fauna in *A. leucostictus*, *H. ventromaculatus*, *Ancistrus* sp. and *Hemiancistrus* sp. from the Igarapé Fortaleza River, and presented a checklist of the parasite species in Loricariidae from Brazil.

Material and methods

Study area, fish and locality of collection

The Igarapé Fortaleza basin, located in the municipalities of Macapá and Santana (AP), is formed by a main channel and an extensive area of floodplains that suffer periodic flooding and is strongly influenced by the high rainfall of the Amazon region and the daily tides of the Amazon River, thus providing shelter and feeding for different species of fish. The regional vegetation is composed of plants with characteristics of flooded forests and herbaceous fields, and several species of macrophytes (Thomaz, Costa-Neto, & Tostes, 2004; Tavares-Dias, Oliveira, Gonçalves, & Silva, 2014).

From November 2013 to August 2014, fifty three specimens of *A. leucostictus*, sp., *H. ventromaculatus*, *Ancistrus* sp. and *Hemiancistrus* sp. were collected in Igarapé Fortaleza River, municipality of Macapá, State of Amapá, Brazil (Figure 1). All fish were collected with gill nets of different mesh sizes, for parasitological analysis.

Collection procedures and analyses of parasites

All fish were weighed (g) and measured for total length (cm), and then necropsied for parasitological analysis. The mouth, opercula, gills and gastrointestinal tract were examined to collect the parasites (protozoans and metazoans). Gills were removed, fixed in formalin 5% and analyzed with the aid of a microscope. To quantify metazoan parasites, each viscera was dissected separately and washed with sodium chloride solution (0.85%) and examined under a stereomicroscope. Previously described techniques were used to collect, count, fix, preserve, and stain the parasites for identification (Eiras, Takemoto, & Pavanelli, 2006; Boeger & Viana, 2006). To analyze the parasites, the ecological terms used were those recommended by Bush, Lafferty, Lotz & Shostak (1997).


A review on the parasites of Loricariidae species in Brazil was performed by searching databases (SciELO, ISI, Scopus, Science Direct, Zoological Records, CAB Abstracts databases and Google Scholar), and available data regarding the parasitic fauna were added to Table 1.

Results

Species of Protozoa, Monogenea, Nematoda, Digenea, Acantocephala, Cestoda, Crustacea and Hirudinea were found parasitizing species of Loricariidae from Brazil, but the most frequent taxonomic groups are Monogenea, Digenea and Nematoda (Table 1). However, the greatest species richness is for Monogenea (Figure 2).

The weight, total length and number of examined hosts are listed in Table 2.

A total of 1,559 parasites were collected from 53 fish specimens, which infected gills and gut of hosts. *Unilatus unilatus* Mizelle et Kritsky, 1967 and *Trinigyryus mourei* Boeger and Bemont-Jégu, 1994 were the most frequent parasites, infecting four host species, followed by *Genarchella gernachella* Travassos 1928 and larvae of *Raphidascaris* (*Sprentascaris*) Peter and Cassone, 1984. Larvae of *Proteocephalus* Weinland, 1858 occurred only in one host species. *Hypostomus ventromaculatus* harbored the highest diversity of parasites, followed by *Ancistrus* sp. (Table 3).



Projection system: UTM
Datum horizontal: WGS 84

Legend

- ⊙ Macapá City
- Santana City
- Collection areas
- Drainage area
- Floodplain



Figure 1. Collection site of the Loricariidae species in Igarapé Fortaleza basin in the Amapá Satate (Brazil).

Table 1. Checklist of parasite species in Loricariidae fish from the Brazil.

Species of parasites	Species of hosts	Locality	References
<i>Spironucleus</i> sp.	<i>Harttia duriventris</i>	Igarapé Fortaleza River (AP)	Gonçalves et al. (2014)
<i>Ichthyophthirius multifiliis</i>		Igarapé Fortaleza River (AP)	Gonçalves et al. (2014)
	<i>Squaliforma emarginatas</i>	Igarapé Fortaleza River (AP)	Gonçalves et al. (2014)
	<i>Hypostomus iheringii</i>	Chavantes Reservoir (SP)	Zica et al. (2012)
	<i>Hypostomus regani</i>	Chavantes Reservoir (SP)	Zica et al. (2012)
<i>Unilatus unilatus</i>	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
	<i>Hypostomus strigaticeps</i>	Chavantes Reservoir (SP)	Zica et al. (2012)
	<i>Pterygoplichthys pardalis</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Unilatus irae</i>	<i>Leporacanthicus galaxias</i>	Guamá River (PA)	Branches & Domingues (2014)
<i>Unilatus</i> sp.	<i>Pterygoplichthys pardalis</i>	Manaus (AM)	Porto et al. (2012)
<i>Phanerothecium harrisii</i>	<i>Hypostomus</i> sp.	Tocantins River (TO)	Kritsky, Vianna, & Boeger (2007)
	<i>Hypostomus plecostomus</i>	Manaus (AM)	Kritsky & Boeger (1991)
<i>Phanerothecium spinatoides</i>	<i>Hypostomus</i> sp.	Tocantins River (TO)	Kritsky et al. (2007)
<i>Phanerothecium deiropedeum</i>		Capivari Dam (PR)	Kritsky et al. (2007)
<i>Phanerothecium spinatus</i>	<i>Hypostomus punctatus</i>	Guandu River (RJ)	Boeger, Kritsky, & Belmont-Jégu (1994)
<i>Phanerothecium spinulatum</i>	<i>Hypostomus regani</i>	Rio das Almas (SP)	Kritsky et al. (2007)
	<i>Hypostomus</i> sp.	Tocantins River (TO)	Kritsky et al. (2007)
<i>Phanerothecioides agostinhoi</i>	<i>Hypostomus affinis</i>	Guandu River (RJ)	Azevedo, Abdallah, & Luque (2010); Azevedo, Abdallah, & Luque (2011)
<i>Nothogrodactylus clavatus</i>		Manaus (AM)	Kritsky & Boeger (1991)
<i>Nothogrodactylus amazonicus</i>	<i>Ancistrus</i> sp.	Manaus (AM)	Kritsky & Boeger (1991)
<i>Nothogrodactylus plaesiophallus</i>		Manaus (AM)	Kritsky & Boeger (1991)
<i>Nothogrodactylus</i> sp.	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Onychogrodactylus hydaticus</i>	<i>Ancistrus multispinis</i>	Salto Morato River (PR)	Kritsky et al. (2007)
<i>Onychogrodactylus sudis</i>		Salto Morato River (PR)	Kritsky et al. (2007)
<i>Oogrodactylus farlowellae</i>	<i>Farlowella amazonum</i>	Manaus (AM)	Kritsky & Boeger (1991)
<i>Trinigyryus mourei</i>	<i>Squaliforma emarginatas</i>	Igarapé Fortaleza River (AP)	Gonçalves et al. (2014)
<i>Trinigyryus hypostomatis</i>	<i>Hypostomus affinis</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Hyperpletes malmbergi</i>		Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Aglaioyrodactylus salebrosus</i>		Patos River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus conei</i>	<i>Pareiorhaphis parmula</i>	Patos River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus ctenistus</i>		Patos River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus forficulatus</i>	<i>Kronichthys lacerta</i>	Salto Morato River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus pedunculatus</i>	<i>Hisonotus</i> sp.	Dois de Fevereiro River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus calamus</i>	<i>Schizolecis guntheri</i>	Várzea River (PR)	Kritsky et al. (2007)
<i>Aglaioyrodactylus forficuloides</i>		Cabral River (PR)	Kritsky et al. (2007)
<i>Paranaella luquei</i>	<i>Hypostomus</i> sp.	Paraná River (PR)	Kohn, Baptista-Farias, & Cohen (2000)
	<i>Rhinelepis aspera</i>	Paraná River (PR)	Kohn et al. (2000)
<i>Hyperpletes malmbergi</i>	<i>Rhineloricaria</i> sp.	Puraquequara River (AM)	Boeger et al. (1994)
<i>Demidospermus paranaensis</i>	<i>Loricariichthys platymetopon</i>	Paraná River (PR)	Ferrari-Hoeninghaus, Bellay, Takemoto, & Pavanelli (2010)
<i>Demidospermus spirophallus</i>	<i>Loricaria prolixa</i>	Batalha River	Pelegrini, Januário, Azevedo & Abdallah et al. (2018); Franceschini et al. (2018)
<i>Demidospermus anus</i>	<i>Loricariichthys platymetopon</i>	Paraná River	Franceschini et al. (2018)
<i>Demidospermus prolixus</i>	<i>Loricaria prolixa</i>	Sapucaí-Mirim River	Franceschini et al. (2018)
<i>Demidospermus rhinelepisii</i>	<i>Rhinelepis aspera</i>	Paraná River (PR)	Acosta, Scholz, Blasco-Costa & Silva (2018)
<i>Demidospermus</i> sp.	<i>Loricariichthys castaneus</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Heteropriapulius</i> sp.	<i>Pterygoplichthys pardalis</i>	Manaus (AM)	Porto et al. (2012)
<i>Paracapillaria piscicola</i>	<i>Hypostomus affinis</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
	<i>Hypostomus albopunctatus</i>	Paraná River (PR)	Moravec, Kohn, & Fernandes (1990)
	<i>Hypostomus commersoni</i>	Paraná River (PR)	Moravec et al. (1990)
	<i>Hypostomus derbyi</i>	Paraná River (PR)	Moravec et al. (1990)
	<i>Ancistrus cirrhosus</i>	Paraná River (PR)	Moravec et al. (1990)
	<i>Hypostomus</i> sp.	Paraná River (PR)	Kohn et al. (2011)
<i>Raphidascaris (Sprentascaris) hypostomi</i>	<i>Loricaria</i> sp.	Paraná River (PR)	Kohn et al. (2011)
	<i>Loricariichthys platymetopon</i>	Paraná River (PR)	Takemoto et al. (2009); Kohn et al. (2011)
	<i>Loricariichthys rostratus</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Pseudohemiodon laticeps</i>	Paraná River (PR)	Moravec et al. (1990); Kohn et al. (2011)
	<i>Loricariichthys</i> sp.	Paraná River (PR)	Kohn et al. (2011)


Species of parasites	Species of hosts	Locality	References
<i>Rhabdochona kidderi</i>	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Rhabdochona</i> sp.	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Oxyuricassis coronatus</i>	<i>Lasiancistrus saetiger</i>	Guamá River (PA)	Rodrigues, Santos, Furtado, & Melo (2017)
<i>Oxyuricassis hexaspinus</i>	<i>Lasiancistrus saetiger</i>	Guamá River (PA)	Rodrigues et al. (2017)
<i>Cucullanus (Cucullanus) pinnae</i>	<i>Loricaria</i> sp.	Paraná River (PR)	Kohn et al. (2011)
	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Ichthyouris brasiliensis</i>	<i>Megalancistrus parananus</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Hypostomus regani</i>	Paraná River (PR)	Kohn et al. (2011)
<i>Procamallanus (Procamallanus) annipetterae</i>	<i>Hypostomus</i> sp.	Paraná River (PR)	Kohn et al. (2011)
	<i>Megalancistrus parananus</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Squaliforma emarginatas</i>	Igarapé Fortaleza River (AP)	Gonçalves et al. (2014)
<i>Procamallanus (Spirocamallanus) inopinatus</i>	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Procamallanus (Spirocamallanus) rebecca</i>	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Parasynodontisia petterae</i>	<i>Rhinelepis strigosa</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Loricariichthys castaneus</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Contraecacum</i> sp.	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Hysterothylacium</i> sp.	<i>Loricariichthys</i> sp.	Paraná River (PR)	Kohn et al. (2011)
	<i>Hypostomus affinis</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
	<i>Hypostomus iheringii</i>	Chavantes Reservoir (SP)	Zica et al. (2011)
	<i>Hypostomus hermanni</i>	Chavantes Reservoir (SP)	Zica et al. (2011)
	<i>Hypostomus margaritifera</i>	Chavantes Reservoir (SP)	Zica et al. (2011)
		Paraná River (PR)	Yamada, Moreira, Ceschini, Takemoto, & Pavanelli (2008)
		Parapanema River (SP)	Zica et al. (2009)
	<i>Hypostomus regani</i>	Paraná River (PR)	Takemoto et al. (2009)
		Chavantes Reservoir (SP)	Zica et al. (2011)
		Chavantes Reservoir (SP)	Zica et al. (2011)
		Chavantes Reservoir (SP)	Zica et al. (2011)
	<i>Hypostomus</i> sp.	Chavantes Reservoir (SP)	Zica et al. (2011)
	<i>Hypostomus strigaticeps</i>	Chavantes Reservoir (SP)	Zica et al. (2011)
<i>Austrodiplostomum compactum</i>	<i>Potamohina pristigaster</i>	Solimões River (AM)	Vital, Morey, Pereira, & Malta (2016)
	<i>Loricariichthys castaneus</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
	<i>Potamohina latior</i>	Solimões River (AM)	Vital et al. (2016)
	<i>Pterygoplichthys pardalis</i>	Manaus (AM)	Porto et al. (2012)
	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Austrodiplostomum</i> sp.	<i>Loricariichthys castaneus</i>	Lages Reservoir (RJ)	Paraguassú & Luque (2007)
<i>Crassicutis intermedius</i>	<i>Hypostomus regani</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Hypostomus</i> sp.	Paraná River (PR)	Kohn et al. (2011)
<i>Crassicutis cichlasomae</i>	<i>Loricaria</i> sp.	Paraná River (PR)	Kohn et al. (2011)
<i>Clinostomum marginatum</i>		Paraná River (PR)	Eiras, Dias, Pavanelli, & Machado (1999); Takemoto et al. (2009)
<i>Clinostomum complanatum</i>		Paraná River (PR)	Ferrari-Hoeninghaus, Takemoto, & Pavanelli (2007); Dias et al. (2006)
<i>Crocodilicola pseudostoma</i>	<i>Loricariichthys platymetopon</i>	Paraná River (PR)	Ferrari-Hoeninghaus et al. (2007); Takemoto et al. (2009)
<i>Clinostomum detruncatum</i>	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Dadaytrema</i> sp.	<i>Megalancistrus parananus</i>	Paraná River (PR)	Kohn et al. (2011)
<i>Genarchella tropica</i>		Paraná River (PR)	Kohn et al. (2011)
<i>Genarchella genarchella</i>	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Saccocoeloides magnus</i>	<i>Megalancistrus parananus</i>	Paraná River (PR)	Kohn et al. (2011)
	<i>Pterygoplichthys pardalis</i>	Janauacá Lake (AM)	Thatcher & Varella (1981)
<i>Megacoelium spinicavum</i>	<i>Pterygoplichthys</i> sp.	Lago Janauacá River (AM)	Thatcher & Varella (1981)
	<i>Pterygoplichthys pardalis</i>	Manaus (AM)	Porto et al. (2012)
<i>Acanthostomum gnerii</i>		Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Dendrorchis</i> sp.	<i>Pterygoplichthys</i> sp.	Paraná River (PR)	Kohn et al. (2011)
<i>Gorytocephalus</i> sp.	<i>Hypostomus</i> sp.	Paraná River (PR)	Kohn et al. (2011)
	<i>Pterygoplichthys pardalis</i>	Manaus (AM)	Porto et al. (2012)
		Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Gorytocephalus elongorchi</i>	<i>Hypostomus carinatus</i>	Janauacá Lake (AM)	Thatcher (1979)
	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Proteocephalus</i> sp.		Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Riggia acuticaudata</i>	<i>Ancistrus</i> sp.	Manaus (AM)	Oda et al. (2015)
<i>Dolops longicauda</i>	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Ergasilus</i> sp.	<i>Hypostomus affinis</i>	Lages Reservoir (RJ)	Paraguassú & Luque (2007)
	<i>Loricariichthys castaneus</i>	Lages Reservoir (RJ)	Paraguassú & Luque (2007)
<i>Placobdella</i> sp.	<i>Hypostomus affinis</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Helobdella</i> sp.	<i>Loricariichthys castaneus</i>	Guandu River (RJ)	Azevedo et al. (2010); Azevedo et al. (2011)
<i>Oligobdella</i> sp.	<i>Loricaria prolixa</i>	Batalha River	Pelegri et al. (2018)
<i>Hirudinea</i> gen. sp.	<i>Peckoltia braueri</i>	Igarapé Fortaleza River (AP)	Cardoso et al. (2017)
<i>Hirudinea</i> Glossiphoniidae	<i>Hypostomus affinis</i>	Lages Reservoir (RJ)	Paraguassú & Luque (2007)
	<i>Loricariichthys castaneus</i>	Lages Reservoir (RJ)	Paraguassú & Luque (2007)

AM: Amazonas, AP: Amapá, PR: Paraná, PA: Pará, TO: Tocantins, SP: São Paulo, RJ: Rio de Janeiro.

Table 3. Parasitic indices four Loricariidae species from the Igarapé Fortaleza River, Amapá State, Brazil.

Species of parasites	<i>Unilatus unilatus</i> and <i>Trinigyryus mourei</i>	<i>Unilatus unilatus</i>	Undetermined metacercariae	<i>Genarchella gernachella</i>	<i>Raphidascaris (Sprentascaris sp.)</i>	<i>Gorythocephalus elongorchis</i>	<i>Proteocephalus sp.</i>
<i>Ancistrus leucostictus</i>	P (%)	75.0	0	0	25.0	25.0	0
	MI	43.0	0	0	34.0	24.0	0
	MA ± SD	32.3 ± 37.4	0	0	8.5 ± 17.0	6.0 ± 12.0	0
	Range	13.0-85.0	0	0	-	-	0
	TNP	129	0	0	34	24	0
	SI	Gills	Gills	Gills	Intestine	Intestine	Intestine
<i>Ancistrus sp.</i>	P (%)	64.3	0	0	15.0	5.0	0
	MI	141.4	0	0	7.3	1	0
	MA ± SD	90.9 ± 201.4	0	0	1.1 ± 3.9	0.1 ± 0.2	0
	Range	1.0-748.0	0	0	1.0 - 17.0	-	0
	TNP	1273	0	0	22	1	0
	SI	Gills	Gills	Gills	Intestine	Intestine	Intestine
<i>Hypostomus ventromaculatus</i>	P (%)	0	100	100	28.6	0	9.5
	MI	0	0.2	0.2	3.5	0	9.0
	MA ± SD	0	2.0 ± 0.0	2.0	1.0 ± 2.9	0	0.9 ± 3.7
	Range	0	-	-	1.0 - 13.0	0	1.0 - 17.0
	TNP	0	4	2	21	0	18
	SI	Gills	Gills	Gills	Intestine	Intestine	Intestine
<i>Hemiancistrus sp.</i>	P (%)	28.6	0	0	0	25.0	0
	MI	119.5	0	0	0	3.5	0
	MA ± SD	34.1 ± 89.9	0	0	0	9.0 ± 2.1	0
	Range	1.0-238.0	0	0	0	1.0-6.0	0
	TNP	239	0	0	0	7	0
	SI	Gills	Gills	Gills	Intestine	Intestine	Intestine

P: Prevalence, MI: Mean intensity, MA: Mean abundance SD: Standard deviation, TNP: Total number of parasites, SI: Site of infection.

**Figure 2.** Richness of the parasite taxa in Loricariidae species from the Brazil.**Table 2.** Body parameters of Loricariidae species from the Igarapé Fortaleza River, Amapá State (Brazil).

Species of hosts	N	Weight (g)	Length (cm)
<i>Ancistrus leucostictus</i>	4	15.1 ± 1.9	47.0 ± 17.8
<i>Ancistrus sp.</i>	20	15.0 ± 3.7	62.4 ± 53.0
<i>Hemiancistrus sp.</i>	8	12.4 ± 3.2	33.0 ± 37.1
<i>Hypostomus ventromaculatus</i>	21	17.2 ± 2.4	58.7 ± 19.1

Discussion

Some patterns of composition and structure of the parasite community in Loricariidae from Brazil were detected: (a) presence of ectoparasites (Protozoa, Monogenea, Crustacea and Hirudinea) and endoparasites (Nematoda, Digenea, Acanthocephala and Cestoda); (b) dominance of *Austrodiplostomum compactum* Lutz, 1928; (c)

infracommunities of ectoparasites richer and more diverse than endoparasites; (d) the presence of endoparasites at the larval stage with high prevalence and low abundance; (e) overdispersion of parasites and (f) interspecific associations of parasite infracommunities (Table 1). There are few studies on parasites in Loricariidae species in the Igarapé Fortaleza basin (Gonçalves et al., 2014; Cardoso et al., 2017). However, this was the first study for *A. leucostictus*, *H. ventromaculatus*, *Ancistrus sp.* and *Hemiancistrus sp.*

In wild fish populations, the parasitic fauna depend on internal and external factors such as: seasonal variations, habitat, limnological characteristics, water body depth, local biota, biological and physiological characteristics of the hosts, trophic level of those population and geography of region (Dogiel, 1961; Lagrue et al., 2011; Cardoso et al., 2017; Baia et al., 2018). The parasites, including monogeneans, can then be positively or negatively influenced by environmental stressors, mainly by anthropic actions that favor the eutrophication of water bodies, altering the parasites abundance in polluted environments (Takemoto et al., 2009).

Monogenea has life cycle in a single host and, in general, of a same family of fish, such as Loricariidae, which are commonly parasitized by species of *Demidospermus*, *Paranaella*, *Trinigyryus* and *Unilatus* (Boeger & Vianna, 2006; Braga, Araújo, & Boeger, 2014). In general, monogeneans are found in low prevalence and abundance in wild fish

populations, when the environmental characteristics do not facilitate the reproduction of these ectoparasites (Tavares-Dias, Rigôr, Pinheiro, Oliveira, & Marinho, 2013). *Unilatus unilatus* and *T. mourei* have been reported infecting diverse Loricariidae species from Brazil (Table 1), and now also *A. leucostictus*, *Ancistrus* sp., *H. ventromaculatus* and *Hemiancistrus* sp., which are new hosts for such monogeneans.

Digeneans are parasites of different host species and have mollusks as primary intermediate hosts, and fish can be intermediate, paratenic or definitive hosts (Ferrari-Hoeninghaus et al., 2007; Cardoso et al., 2017). *Genarchella germachella* does not have host specificity, since it infects several Characiformes and Siluriformes fish from Brazil, which can be definitive hosts for this digenean specie (Cardoso et al., 2017). Undetermined metacercariae and *G. germachella* were found in *A. leucostictus*, *Ancistrus* sp. and *H. ventromaculatus*, but the greatest prevalence occurred in intestine of *H. ventromaculatus*. This was the first report of *G. genarchella* for *Ancistrus* sp., *A. leucostictus* and *H. ventromaculatus*.

Species of *Raphidascaris* (*Sprentascaris*) are nematodes with little known life cycle, but require fish and aquatic crustaceans as intermediate hosts; in addition amphibians, aquatic insects, zooplankton, etc. which are less likely to be intermediate or paratenic hosts (Moravec, 1998). For *A. leucostictus*, *Ancistrus* sp. and *Hemiancistrus* sp., there were varied infection levels of *Raphidascaris* (*Sprentascaris*) sp. *Raphidascaris* (*S.*) *hipostomi* Peter and Cassone, 1984 and *Raphidascaris* (*S.*) *mahnerti* Peter and Cassone, 1984 (Table 1) are known infecting Loricariidae species, but this was the first record of *Raphidascaris* (*Sprentascaris*) sp. for *A. leucostictus*, *Ancistrus* sp. and *Hemiancistrus* sp.

Acanthocephalans are endoparasites of fish, birds and mammal (Cardoso et al., 2017). *Gorytocephalus elongorchis* Thatcher (1979), a parasite of Loricariidae species (Table 1), was found only in *H. ventromaculatus*, which are new hosts for this acanthocephalan. Species of *Proteocephalus* are common cestodes of Siluriformes, which consume planktonic crustaceans (Copepoda and Cyclopoida) that serve as intermediate hosts. Thus, some species of Siluriformes can be definitive hosts, with direct infection after consuming microcrustaceans (Scholz, 1999; Cardoso et al., 2017). Plerocercoids of *Proteocephalus* sp. were found only in *H. ventromaculatus*, which can be secondary intermediate host for this endoparasite. In addition, *H. ventromaculatus* is a new host for *Proteocephalus* sp.

Conclusion

Ectoparasites were predominant in the examined Loricariidae species, which also had species of endoparasites at the larval stage. The hosts with the highest number of collected samples (*H. ventromaculatus* and *Ancistrus* sp.) had a higher richness of parasite species. The species of Loricariidae from Brazil present ecto- and endoparasites, but the predominance is of ectoparasites. Species of monogeneans, digeneans and nematodes are the most diversified parasites in Brazilian Loricariidae.

Acknowledgements

Dr. M. Tavares-Dias receives a Research Fellowship (# 303013/2015-0) from the *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq, Brazil).

References

- Acosta, A. A., Scholz, T., Blasco-Costa, I., Alves, P. V., & Silva, R. J. (2018). A new genus and two new species of dactylogyrid monogeneans from gills of Neotropical catfishes (Siluriformes: Doradidae and Loricariidae). *Parasitology International*, 67(1), 4-12. doi: 10.1016/j.parint.2017.09.012
- Alonso, F., Terán, G. E., Aguilera, G., & Mirande, J. M. (2016). First record of *Hypostomus boulengeri* (Siluriformes: Loricariidae) from Bermejo River basin. *Revista del Museo Argentino Ciencias Naturales*, 18(1), 85-88.
- Azevedo, R. K., Abdallah, V. D., & Luque, J. L. (2010). Acanthocephala, Annelida, Arthropoda, Myxozoa, Nematoda and Platyhelminthes parasites of fishes from the Guandu River, Rio de Janeiro, Brazil. *Check List*, 6(4), 659-667. doi: 10.15560/6.4.659
- Azevedo, R. K., Abdallah, V. D., & Luque, J. L. (2011). Biodiversity of fish parasites from Guandu River, southeastern Brazil: an ecological approach. *Neotropical Helminthology*, 5(2), 185-199.
- Baia, R. R. J., Florentino, A. C., Silva, L. M. A., & Tavares-Dias, M. (2018). Patterns of the parasite communities in a fish assemblage of a river in the Brazilian Amazon region. *Acta Parasitology*, 63(2), 304-316. doi: 10.1515/ap-2018-0035
- Boeger, W. A., & Viana, R. T. (2006). Monogenoidea. In: Thatcher V. E. (ed.). *Amazon fish parasites* (p. 42-116). Sofia, BUL: Pensoft Publishers.
- Boeger, W. A., Kritsky D. C., & Belmont-Jégu, E. (1994). Neotropical Monogenoidea. 20. Two new species of oviparous Gyrodactylidea (Polyonchoinea) from loricariid catfishes (Siluriformes) in Brazil and the phylogenetic status of Oeogyrodactylidae Harris, 1983. *Journal of Helminthological Society of Washington*, 61(1), 34-44.

- Branches, B., & Domingues, M. V. (2014). A new species of *Unilatus* (Platyhelminthes: Monogeneoidea) from the gills of *Leporacanthicus galaxias* Isbrücker et Nijssen (Siluriformes: Loricariidae) from Brazil. *Acta Parasitologica*, 59(1), 91-97. doi: 10.2478/s11686-014-0213-7.
- Braga, M. P., Araújo, S. B. L., & Boeger, W. A. (2014). Patterns of interaction between Neotropical freshwater fishes and their gill Monogeneoidea (Platyhelminthes). *Parasitology Research*, 113(2), 481-490. doi: 10.1007/s00436-013-3677-8
- Bush, A. O., Lafferty, K. D., Lotz, J. M., & Shostak, W. (1997). Parasitology meets ecology on its own terms: Margolis et al. Revisited. *Journal of Parasitology*, 83(4), 575-583. doi: 10.2307/3284227.
- Cardoso, A. C. F., Oliveira, M. S. B., Neves, L. R., & Tavares-Dias, M. (2017). Metazoan fauna parasitizing *Peckoltia braueri* and *Pterygoplichthys pardalis* (Loricariidae) catfishes from the northeastern Brazilian Amazon. *Acta Amazonica*, 47(2), 147-154. doi: 10.1590/1809-4392201603232.
- Dias, M. L. G. G., Mente-Vera, C. V. Eiras, J. C., Machado, M. H., Souza G. T. R., Pavanelli, G. C. (2006). Ecology of *Clinostomum complanatum* Rudolphi, 1814 (Trematoda: Clinostomidae) infecting fish from the floodplain of the high Paraná River, Brazil. *Parasitology Research*, 99, 675-681. doi: 10.1007/s00436-006-0205-0
- Dogiel, V. A. (1961). Ecology of the parasites of freshwater fishes. In V. A. Dogiel, G. K. Petrushevsky, and Y. I. Polyanski (Eds.). *Parasitology of fishes* (p. 1-47). Leningrad, RUS: University Press, Russian Soviet Federative Socialist Republic.
- Eiras, J. C., Dias, M. L. G. G., Pavanelli G. C., & Machado M. H. (1999). Histological studies on the effects of *Clinostomum marginatum* (Digenea, Clinostomidae) in its second intermediate host *Loricariichthys platymetopon* (Osteichthyes, Loricariidae) of the upper Paraná River, Brazil. *Acta Scientiarum, Biological Sciences*, 21(2), 237-241. doi: 10.4025/actascibiolsci.v21i0.4416
- Eiras, J. C., Takemoto, R. M., & Pavanelli, G.C. (2006). *Métodos de estudos e técnicas laboratoriais em parasitologia de peixes*. Maringá, PR: Eduem.
- Ferrari-Hoeninghaus, A. P., Takemoto, R. M., & Pavanelli, G. C. (2007). Digenetic trematode parasites of *Loricariichthys platymetopon* (Loricariidae, Siluriformes) of the upper Paraná river floodplain, Brazil. *Acta Scientiarum. Biological Sciences*, 29(3), 327-329. doi: 10.4025/actascibiolsci.v29i3.508
- Ferrari-Hoeninghaus, A. P., Bellay, S., Takemoto, R. M., & Pavanelli, G. C. (2010). A new species of *Demidosperrmus* Suriano, 1983 (Monogenea, Dactylogyridae) parasitic on *Loricariichthys platymetopon* Isbrücker et Nijssen (Loricariidae, Siluriformes) from the upper Paraná River floodplain, Brazil. *Acta Parasitologica*, 55(1), 16-19. doi:10.2478/s11686-010-0007-5
- Franceschini, L., Zago, A. C., Müller, M. I., Francisco, C. J., Takemoto, R. M., & Silva, R. J. (2018). Morphology and molecular characterization of *Demidosperrmus spirophallus* n. sp., *D. prolixus* n. sp. (Monogenea: Dactylogyridae) and a redescription of *D. anus* in siluriform catfish from Brazil. *Journal of Helminthology*, 92(2), 228-243. doi:10.1017/S0022149X17000256
- Gonçalves, R. A., Oliveira, M. S. B., Santos, E. F., & Tavares-Dias, M. (2014). Aspectos ecológicos da comunidade de parasitos em duas espécies de Loricariidae da bacia Igarapé Fortaleza, estado do Amapá, Brasil. *Biota Amazonia*, 4(1), 15-21. doi: 10.18561/2179-5746/biotaamazonia.v4n1p15-21
- Kohn, A., Baptista-Farias, M. F. D., & Cohen S. C. (2000). *Paranaella luquei* gen. et sp. n. (Monogenea: Microcotylidae), a new parasite of Brazilian catfishes. *Folia Parasitologica*, 47(4), 279-283. doi: 10.14411/fp.2000.048
- Kohn, A., Moravec, F., Cohen, S. C., Canzi, C., Takemoto, R. M., & Fernandes, B. M. M. (2011). Helminths of freshwater fishes in the reservoir of the Hydroelectric Power Station of Itaipu, Paraná, Brazil. *Check List*, 7(5), 681-690. doi: 10.15560/7.5.681
- Kritsky, D. C., & Boeger, W. A. (1991). Neotropical Monogenea. 16. New species of oviparous Gyrodactylidae with proposal of *Nothogyrodactylus* gen. n. (Oogyrodactylidae). *Journal of the Helminthological Society of Washington*, 58(1), 7-15.
- Kritsky, D. C., Vianna, R. T., & Boeger, W. A. (2007). Neotropical Monogeneoidea. 50. Oviparous gyrodactylids from loricariid and pimelodid catfishes in Brazil, with the proposal of *Phanerothecioides* n. g., *Onychogyrodactylus* n. g. and *Aglaigyrodactylus* n. g. (Polyonchoinea: Gyrodactylidae). *Systematic Parasitology*, 66(1), 1-34. doi: 10.1007/s11230-006-9053-7.
- Lagrange, C., Kelly, D. W., Hicks, A., & Poulin, R. (2011). Factors influencing infection patterns of trophically transmitted parasites among a fish community: host diet, host-parasite compatibility or both? *Journal of Fish Biology*, 79(2), 466-485. doi: 10.1111/j.1095-8649.2011.03041.x
- Luque, J. L., & Poulin R. (2007). Metazoan parasite species richness in Neotropical fishes: hotspots and the geography of biodiversity. *Parasitology*, 134(6), 865-878. doi:10.1017/S0031182007002272
- Moravec, F., Kohn, A., Fernandes, B. M. M. (1990). First record of *Raphidascaris (Sprentascaris) hypostomi* (Petter et Cassone, 1984) comb. n. and *R. (S.) mahnerti* (Petter et Cassone, 1984) comb. n. (Nematoda: Anisakidae) from Brazil with remarks on the taxonomic status of the genus *Sprentascaris* Petter et Cassone, 1984. *Folia Parasitologica*, 37(2), 131-140.
- Moravec, F. (1998). *Nematodes of freshwater fish of the Neotropical region*. Praha, CZ: Academia of Sciences of the Czech Republic.
- Nelson, J. S., Grande, T. C., & Wilson, M. V. H. (2016). *Fishes of the world* (5th) New Jersey, EUA: John Wiley & Sons, Inc., Hoboken.
- Oda, F. H., Graça, R. J., Tencatt, L. F. C., Tavares, L. E. R., Froehlich O., & Takemoto R. M. (2015). The

- poorly known *Riggia acuticaudata* (Crustacea: Isopoda) parasitizing *Ancistrus* sp. (Siluriformes: Loricariidae) from the Paraguay River basin, Brazil, with comments on its reproductive biology. *Comparative Parasitology*, 82(1), 25-28. doi: 10.1654/4738.1
- Paraguassú, A. R., & Luque J. L. (2007). Metazoários parasitos de seis espécies de peixes do reservatório de Lajes, Estado do Rio de Janeiro, Brasil. *Revista Brasileira de Parasitologia Veterinária*, 16(3), 121-128. doi: 10.1590/S1984-29612007000300002.
- Pelegrini, L. S., Januário, F. F., Azevedo, R. K., & Abdallah, V. D. (2018). Biodiversity and ecology of the parasitic infracommunities of *Loricaria prolixa* (Siluriformes: Loricariidae) from the Tietê-Batalha Basin, SP, Brazil. *Acta Scientiarum. Biological Sciences*, 40(1), e36294. doi: 10.4025/actasciobiolsci.v40i1.36294.
- Porto, D. B., Vital, J. F., Santos, A. K. S., Morais, A. M., Varella, A. M. B., Malta, J. C. O. (2012). Metazoários parasitos de *Pterygoplichthys pardalis* (Castelnau, 1855) (Siluriformes: Loricariidae) da Amazônia central, Brasil. *Revista Brasileira de Zootecnia*, 14 (1-3): 35-40.
- Rodrigues, A. R. O., Santos, J. N., Furtado, A. P., & Melo, F. T. V. (2017). *Oxyuricassis coronatus* n. gen. n. sp. and *O. hexaspinatus* n. sp. (Oxyurida: Pharyngodonidae): parasites of *Lasiancistrus saetiger* (Siluriformes: Loricariidae) in freshwater rivers of the Brazilian Amazon. *Journal of Parasitology*, 103(4), 390-398. doi: 10.1645/16-138.
- Scholz, T. (1999). Life cycles of species of *Proteocephalus*, parasites of fishes in the Palearctic region: a review. *Journal of Helminthology*, 73(1), 1-19. doi: 10.1017/S0022149X99000013
- Soares, M. G. M., Costa E. L., Siqueira-Souza, F. K., Anjos, H. D. B., Yamamoto K. C., & Freitas, C. E. C. (2011). *Peixes de lagos do médio Rio Solimões* (2a ed.). Manaus, AM: Instituto Piatam.
- Takemoto, R. M., Pavanelli, G. C., Lizama, M. A. P., Lacerda, A. C. F., Yamada, F. H., Moreira, L. H. A., ... Bellay, S. (2009). Diversity of parasites of fish from the upper Paraná River floodplain, Brazil. *Brazilian Journal of Biology*, 69(2), 691-705. doi: 10.1590/S1519-69842009000300023
- Tavares-Dias, M., Rigôr, L. N., Pinheiro, D. A., Oliveira, M. S. B., & Marinho, R. G. B. (2013). Parasites in *Curimata cyprinoides* (Characiformes: Curimatidae) from eastern Amazon, Brazil. *Acta Scientiarum. Biological Sciences*, 35(4), 595-601. doi: 10.4025/actasciobiolsci.v35i4.19649
- Tavares-Dias, M., Oliveira, M. S. B., Gonçalves, R., Silva, L. M. A. (2014). Ecology and seasonal variation of parasites in wild *Aequidens tetramerus*, a Cichlidae from the Amazon. *Acta Parasitologica*, 59(1), 158-164. doi: 10.2478/s11686-014-0225-3
- Thatcher, V. E. (1979). Uma nova espécie de *Gorytocephalus* Nickol e Thatcher, 1971 (Acanthocephala: Neoechinorhynchidae) do acari bodó (Pisces: Loricariidae) da Amazônia, Brasil. *Acta Amazonica*, 9(1), 199-202. doi: 10.1590/1809-43921979091199
- Thatcher, V. E., & Varella, A. B. (1981). Duas novas espécies de *Megacoelium* Szidat, 1954 (Trematoda: Heploporidae), parasitas estomacais de peixes da Amazônia Brasileira, com uma definição do gênero. *Acta Amazonica*, 11(2), 285-289. doi: 10.1590/1809-43921981112285
- Thomaz, D. O., Costa-Neto, S. V., Tostes L. C. L. (2004). Inventário florístico das ressacas das bacias do Igarapé do Lago. In Takiyama, L. R., & Silva, A. Q. *Diagnostico de ressacas do estado do Amapá: bacias do Igarapé da Fortaleza e do Rio Curiaú* (p. 13-32). Macapá, AP: IEPA.
- Vital, J. F., Morey, G. A. M., Pereira, N. B., & Malta, J. C. O. (2016). Metacercárias de *Austrodiplostomum compactum* (Lutz, 1928) em peixes de lagos de várzea da Amazônia Brasileira. *Folia Amazonica*, 25(2), 153-158. doi: 10.24841/fa.v25i2.399
- Yamada, F. H., Moreira, L. H. A., Ceschini, T. L., Takemoto, R. M., Pavanelli, G. C. (2008). Novas ocorrências de metacercária de *Austrodiplostomum compactum* (Lutz, 1928) (Platyhelminthes: Digenea) parasito de olhos de peixes da bacia do Rio Paraná. *Revista Brasileira de Parasitologia Veterinária*, 17(3), 163-166. doi:10.1590/S1984-29612008000300010.
- Zica, E. O. P., Santos, K. R., Ramos, I. P., Zanatta, A. S., Carvalho, E. D., & Silva, R. J. (2009). First case of an infection of the metacercariae of *Austrodiplostomum compactum* (Lutz, 1928) (Digenea, Diplostomidae) in *Hypostomus regani* (Ihering, 1905) (Siluriformes: Loricariidae). *Pan-American Journal of Aquatic Sciences*, 4(1), 35-38.
- Zica, E. O. P., Brandão, H., Zawadzki, C. H., Nobile, A. B., Carvalho E. D., & Silva, R. J. (2011). The occurrence of *Austrodiplostomum compactum* (Lutz, 1928) (Digenea: Diplostomidae) metacercariae in the eyes of loricariid fish (Siluriformes: Osteichthyes: Loricariidae) from Brazil. *Journal of Helminthology*, 85(1), 73-79. doi: 10.1017/S0022149X10000271.
- Zica, E. O. P., Abdallah, V. D., Azevedo R. K., Wunderlich A. C., Carvalho E. D., & Silva R. J. (2012). *Unilatus unilatus* Mizelle & Kritsky, 1967 (Monogenea, Ancyrocephalinae) in *Hypostomus* spp. (Siluriformes, Loricariidae) from the Chavantes reservoir, São Paulo State, Brazil. *Helminthologia*, 49(2), 87-91.

Received on November 28, 2017.

Accepted on May 7, 2018.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.