

## University of Wollongong Research Online

---

Faculty of Science, Medicine and Health - Papers:  
Part B

Faculty of Science, Medicine and Health

---

2019

# Sperm collection and storage for the sustainable management of amphibian biodiversity

Robert Browne  
*Sustainable America*

Aimee J. Silla  
*University of Wollongong, asilla@uow.edu.au*

Rose Upton  
*University of Newcastle*

Gina Della Togna  
*Smithsonian Conservation Biology Institute*

Ruth Marcec-Greaves  
*Detroit Zoological Society*

*See next page for additional authors*

---

### Publication Details

Browne, R. K., Silla, A. J., Upton, R., Della-Togna, G., Marcec-Greaves, R., Shishova, N. V., Uteshev, V. K., Proano, B., Perez, O. D., Mansour, N., Kaurova, S. A., Gakhova, E. N., Cosson, J., Dyzuba, B., Kramarova, L. I., McGinnity, D., Gonzalez, M., Clulow, J. & Clulow, S. (2019). Sperm collection and storage for the sustainable management of amphibian biodiversity. *Theriogenology*, 133, 187-200.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# Sperm collection and storage for the sustainable management of amphibian biodiversity

## Abstract

Current rates of biodiversity loss pose an unprecedented challenge to the conservation community, particularly with amphibians and freshwater fish as the most threatened vertebrates. An increasing number of environmental challenges, including habitat loss, pathogens, and global warming, demand a global response toward the sustainable management of ecosystems and their biodiversity. Conservation Breeding Programs (CBPs) are needed for the sustainable management of amphibian species threatened with extinction. CBPs support species survival while increasing public awareness and political influence. Current CBPs only cater for 10% of the almost 500 amphibian species in need. However, the use of sperm storage to increase efficiency and reliability, along with an increased number of CBPs, offer the potential to significantly reduce species loss. The establishment and refinement of techniques over the last two decades, for the collection and storage of amphibian spermatozoa, gives confidence for their use in CBPs and other biotechnical applications. Cryopreserved spermatozoa has produced breeding pairs of frogs and salamanders and the stage is set for Lifecycle Proof of Concept Programs that use cryopreserved sperm in CBPs along with repopulation, supplementation, and translocation programs. The application of cryopreserved sperm in CBPs, is complimentary to but separate from archival gene banking and general cell and tissue storage. However, where appropriate amphibian sperm banking should be integrated into other global biobanking projects, especially those for fish, and those that include the use of cryopreserved material for genomics and other research. Research over a broader range of amphibian species, and more uniformity in experimental methodology, is needed to inform both theory and application. Genomics is revolutionising our understanding of biological processes and increasingly guiding species conservation through the identification of evolutionary significant units as the conservation focus, and through revealing the intimate relationship between evolutionary history and sperm physiology that ultimately affects the amenability of sperm to refrigerated or frozen storage. In the present review we provide a nascent phylogenetic framework for integration with other research lines to further the potential of amphibian sperm banking.

## Publication Details

Browne, R. K., Silla, A. J., Upton, R., Della-Togna, G., Marcec-Greaves, R., Shishova, N. V., Uteshev, V. K., Proano, B., Perez, O. D., Mansour, N., Kaurova, S. A., Gakhova, E. N., Cosson, J., Dyzuba, B., Kramarova, L. I., McGinnity, D., Gonzalez, M., Clulow, J. & Clulow, S. (2019). Sperm collection and storage for the sustainable management of amphibian biodiversity. *Theriogenology*, 133 187-200.

## Authors

Robert Browne, Aimee J. Silla, Rose Upton, Gina Della Togna, Ruth Marcec-Greaves, Natalia Shishova, Victor Uteshev, Belin Proano, Oscar D. Perez, Nabil Mansour, Svetlana Kaurova, Edith Gakhova, Jacky Cosson, Borys Dyzuba, Ludmila Kramarova, Dale McGinnity, Manuel Gonzalez, John Clulow, and Simon Clulow

# Sperm collection and storage for the sustainable management of amphibian biodiversity

Robert K. Browne <sup>a\*</sup>, Aimee.J. Silla <sup>b</sup>, Rose Upton <sup>c</sup>, Gina Della-Togna <sup>d,e</sup>, Ruth Marcec-Greaves <sup>f</sup>, Natalia V. Shishova <sup>g</sup>, Victor K. Uteshev <sup>g</sup>, Belin Proaño <sup>h</sup>, Oscar D. Pérez <sup>h</sup>, Nabil Mansour <sup>i</sup>, Svetlana A. Kaurova <sup>g</sup>, Edith N. Gakhova <sup>g</sup>, Jacky Cosson <sup>j</sup>, Borys Dyzyuba <sup>j</sup>, Ludmila I. Kramarova <sup>k</sup>, Dale McGinnity <sup>l</sup>, Manuel Gonzalez <sup>m</sup>, John Clulow <sup>c</sup>, Simon Clulow <sup>c,n</sup>

a \*Corresponding author. Sustainability America, Sarteneja, Corozal District, Belize. robert.browne@gmail.com.

b School of Earth, Atmospheric and Life Sciences, University of Wollongong NSW 2522.

c School of Environmental and Life Sciences, University of Newcastle, Callaghan Drive, Callaghan, NSW 2308, Australia.

d Smithsonian Tropical Research Institute, Panama Amphibian Rescue and Conservation Project, Panama City, Panama.

e Universidad Interamericana de Panamá, Dirección de Investigación, Sede Central, Panama.

f National Amphibian Conservation Center Detroit Zoological Society, Detroit, USA.

g Institute of Cell Biophysics, Russian Academy of Sciences, Pushchino, Moscow Region, 142290, Russia.

h Escuela de Ciencias Biológicas, Pontificia Universidad Católica del Ecuador, Ecuador.

i Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt.

j University of South Bohemia in Ceske Budejovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, 38925 Vodnany, Czech Republic.

k Institute of Theoretical and Experimental Biophysics, Russian Academy of Sciences, Pushchino, Moscow Region, 142290, Russia.

l Nashville Zoo at Grassmere, Nashville, TN 37211, USA

m Departamento de Producción Animal, Universidad de Guadalajara, , Guadalajara, Jalisco, Mexico.

n Department of Biological Sciences, Macquarie University, Sydney NSW 2109 Australia.

Corresponding author: R. Browne, PhD, robert.browne@gmail.com.

**Abstract**

37 Current rates of biodiversity loss pose an unprecedented challenge to the conservation community,  
38 particularly with amphibians and freshwater fish as the most threatened vertebrates. An increasing  
39 number of environmental challenges, including habitat loss, pathogens, and global warming, demand a  
40 global response toward the sustainable management of ecosystems and their biodiversity. Conservation  
41 breeding programs (CBPs) are needed for the sustainable management of amphibian species threatened  
42 with extinction. CBPs support species survival while increasing public awareness and political  
43 influence. Current CBPs only cater for 10% of the almost 500 amphibian species in need. However, the  
44 use of sperm storage to increase efficiency and reliability, along with an increased number of CBPs,  
45 offer the potential to significantly reduce species loss. The establishment and refinement of techniques  
46 for the collection and storage of amphibian spermatozoa, over the last two decades, gives confidence for  
47 their use in CBPs and other biotechnical applications. Cryopreserved spermatozoa has produced  
48 breeding pairs of frogs and salamanders and the stage is set for lifecycle proof of concept studies that  
49 use cryopreserved sperm in CBPs along with repopulation, supplementation, and translocation  
50 programs. The application of cryopreserved sperm in CBPs, is complimentary to but separate from  
51 archival gene banking and general cell and tissue storage. However, where possible amphibian sperm  
52 banking should be standardised and integrated into other global biobanking projects, especially those for  
53 fish, and those that include the use of cryopreserved material for genomics and other research. Research  
54 over a broader range of amphibian species, and more uniformity in experimental methodology, is needed  
55 to inform both theory and application. Genomics is revolutionising our understanding of biological  
56 processes and increasingly guiding species conservation through the identification of evolutionary  
57 significant units as the conservation focus, and through revealing the intimate relationship between  
58 evolutionary history and sperm physiology that ultimately affects the amenability of sperm to  
59 refrigerated or frozen storage. In the present review we provide a nascent phylogenetic framework for  
60 integration with other research lines to further the potential of amphibian sperm banking.

61

62

## 1. Introduction

Conservation Breeding Programs (CBPs) are required for the sustainable management of amphibian species threatened with extinction. The use of cryopreserved spermatozoa in CBPs perpetuates male genetic variation, lowers costs, increases biosecurity, reduces the number of required captive individuals, enables the fertilization of a single female's spawn with spermatozoa from many genetically diverse males, and reduces the need for animal transport [1,2,3]. Sperm banks for fishes exist globally for projects ranging from the perpetuation of zebra-fish cell lines [4] to maintaining genetic variation in sport fishing based CBPs and in aquaculture programs [5]. However, the cryopreserved spermatozoa of amphibians has only been practically applied to maintain *Xenopus* transgenic lines for biotechnological research [6]. Techniques for the post-thaw recovery of cryopreserved fishes [7,8] and amphibians [9] oocytes or embryos have not succeeded. However, primordial germ cells from cryopreserved fish embryos have been transplanted to amphibian embryos and then have developed into the gonads of fertile adults [10]. Similar technology for amphibians offers the greatest current potential for the cryopreserved storage of female germplasm. In any case, live females are needed to provide oocytes for *in vitro* fertilization with stored spermatozoa, or to supply larvae for primordial germ cell transplantation [9].

The Amphibian Ark [AArk, [11]) was established as a zoo based organisation to support amphibian CBPs. The AArk Species Conservation Assessments [12] recommends that of the ~570 amphibian species requiring CBPs, that ~500 species or 90% need the support of gene banks including the use of cryopreserved spermatozoa ([Supplementary Table 1](#)).

The loss of genetic variation in CBPs can result in poor reproduction, health, and survival [13,14]. Even with large founder populations, genetic selection can occur in a few generations for rapid growth, early maturity, amenability to husbandry or ease of reproduction in captivity [13,15]. Selection toward domestication can be reduced but not eliminated by strict studbook management. However, even strict studbook management is subject to loss of broodstock and difficulties in transporting broodstock between breeding groups. The optimal approach is to use cryopreserved spermatozoa to reduce the effective number of male generations to one, and to provide an easy means of transport [13].

95 The natural genetic variation of species can be recovered through the use of cryopreserved  
 96 spermatozoa using oocytes from highly domestic strains (Fig. 1). Consequently, domestic strains of  
 97 species without studbook requirement, both in aquaculture and in private keeper's collections, could  
 98 contribute female brood stock to CBPs if adequate stocks of cryopreserved sperm were available to  
 99 restore the species genetic variation. In the case of *Andrias davidianus*, where genomics recently  
 100 revealed the taxon consists of 5 component species now mainly as aquaculture hybrids [16], and with all  
 101 5 species are functionally extinct in nature [17], these species could be re-established with the use of  
 102 stored sperm.

103

104 Insert Fig. 1.

105

106 The spermatozoa of fishes and amphibians can remain viable for days to weeks during refrigerated  
 107 storage at ~4°C, or indefinitely when cryopreserved in liquid nitrogen at -196°C (Supplementary Tables  
 108 2,3. [16,18]). Post-thaw, motile spermatozoa can be used for artificial fertilization [2] and immotile  
 109 spermatozoa for intracytoplasmic injection into the oocyte (ICSI; [19]). In amphibians, post-thaw  
 110 spermatozoa from Anurans (frogs and toads) has resulted in reproducing pairs of *Xenopus* [6] and  
 111 sexually mature males of tree frogs [20], and in Caudata reproducing pairs of \*Salamanders [21], and  
 112 almost mature Cryptobranchids (Giant salamanders, Dale McGinnity, personal communication).

113

114 \*Urodeles include all extinct and extant salamanders. The Caudata, Fig. 2. have three main lineages; the Cryptobranchidae, Sirenidae, and  
 115 "other salamanders". In this review for grammatical simplicity where appropriate we simply use the term "Salamander" rather than "other  
 116 salamander).

117

118 We use studies of freshwater fishes spermatozoa to provide the closest phylogenetic, morphological, and  
 119 physiological comparisons to amphibian spermatozoa. The parameters used to assess the quality of  
 120 amphibian spermatozoa include the percentage with flagella movement (percent activation) and velocity  
 121 (percentage motility), and the velocity and longevity of motile spermatozoa [22]. Membrane integrity,  
 122 spermatozoa concentration, DNA integrity, and acrosome integrity, and relationship between these and  
 123 fertilisation and larval growth to adults are also assessed [23].

124

## 125 **2. Amphibian phylogenetics, fertilisation history, and reproductive strategies**

126

127 phylogenetic patterns in the amenability of amphibian species spermatozoa, to the physiological and  
 128 morphological stressors during storage, could facilitate the development of storage techniques and in  
 129 general inform amphibian spermatology [2,24,25]. Most Anurans externally fertilise through  
 130 spermatozoa shed in spermic urine. Spermatozoa is released in semen in the Cryptobranchidae  
 131 (Cryptobranchids and Hydronobids) and Sirenidae. Fertilization is internal in all Salamanders through  
 132 the deposition of spermatophores by males that are then picked up by the cloaca of females and all  
 133 species of Caecilians are internal fertilizers [26].

134

135 Of amphibians, the Anurans have the most complicated evolutionary history of fertilisation:  
 136 primordially with external fertilisation, then internal fertilization as Lissamphibians, external  
 137 fertilization as Batrachians, a 40 million year period of internal fertilization, then a reversion to external  
 138 fertilisation from 275 mya to the present [27]. One primitive Anuran retains internal fertilisation [25,28],  
 139 and less than 15 known species have reverted to internal fertilisation [28]. The Caudata have a less  
 140 complicated evolutionary history than the Anurans with two families the Sirenidae and the  
 141 Cryptobranchidae retaining the ancestral Batrachian external fertilisation, and Salamanders readopting  
 142 internal fertilisation. Caecilians are internal fertilisers ([27] Fig. 2).

143

144 Insert Fig. 2.

145

146 The three sub-orders of Anurans are the Archaeobatrachia with 4 families and ~27 species, the  
 147 Mesobatrachia with 6 families and ~168 species, and Neobatrachia with 21 families and the ~5000  
 148 species. In Anurans, the spermatozoa of 30 species has been cryopreserved: species in seven  
 149 Neobatrachia families, two species in one Mesobatrachia family, the Pipidae, which includes *Xenopus*.  
 150 In anurans, post-thaw assessment of spermatozoa viability as defined by live/dead stains was reported  
 151 for 12 species, motility for 16 species, and life stages to first cleavage for 4 species, larval development  
 152 for 7 species, and development to adults for 2 species. In the Caudata the spermatozoa of 5 species have  
 153 been cryopreserved, 3 salamanders and 2 cryptobranchids. Post-thaw assessment of spermatozoa  
 154 viability in spermatophores was reported for one species, motility for two species, development to late  
 155 juvenile/adults with one species, and to fertile adults in one species (Supplementary Table 3).

156

### 157 3. Sperm collection and sperm concentration

158

159 Sperm can be collected as testicular macerates or suspensions from any sexually mature male  
160 amphibian. Anuran spermatozoa can also be collected through hormonal induction, either as spermic  
161 urine through abdominal massage [23] or through cannulation of the cloaca [29,30,31,32,33,34].  
162 Hormonal induction causes internally fertilising salamanders to deposit spermatophores [35] or to  
163 express sperm in cloacal fluid (Fig. 3,4 [36,37,38]).

164

165 Insert Fig. 3.

166

167 Both phylogeny and the environment, especially climate, influence the reproductive strategies of  
168 amphibian species and their amenability to hormonal induction [2,22,40]. Hormones can be  
169 administered safely and efficiently by injection even with small frogs [31,32,33], and generally most  
170 species are amenable to hormonal induction of sperm release with gonadotropin releasing hormones  
171 (GnRH) or human chorionic gonadotropin (hCG [2]). GnRH is generally more effective at inducing  
172 spermiation than hCG across a wide range of species, however, there are a number of species, mainly  
173 from the Bufonidae and Limnodynastidae families, where hCG elicits a stronger response [22,39,41].  
174 Inter-taxon variation also occurs between closely related species, where due to its fertilisation strategy a  
175 single species from a family otherwise amenable to hormonal induction responded poorly both hCG and  
176 GnRH [2,29]. This and other exceptions may elucidate the specific evolutionary drivers behind  
177 reproductive strategies [2].

178

179 Hormonal induction depends on the presence of mature spermatozoa in the testes [24]. Seasonal  
180 quiescence in spermatozoa maturation can be circumvented through the use of both priming, where sub-  
181 inducing doses of hormones, along with dopamine antagonists [42,43], are administered days before the  
182 final inducing dose [22,24]. Hormone administration generally induces Anuran spermic urine over  
183 periods between 2 -12 h with clear peaks in spermatozoa concentration between 3 and 7 h  
184 (Supplementary Table 2. [22,23,24,33,41,44,45]).

185

186 The most reliable collection technique for large quantities of mature Anuran spermatozoa is  
187 through the maceration of the testes to produce spermatozoa suspensions [6,18]). The concentration of  
188 spermatozoa in testicular macerates at  $\sim 10^{8-9}/\text{ml}$  is generally one to three magnitudes higher than in



189 spermic urine (Supplementary Table 2). The high concentration, volume and quality of testicular  
190 spermatozoa has resulted in its use in most studies of Anuran spermatozoa cryopreservation until  
191 recently, and in the only two studies resulting in mature reproducing pairs [6] or sexually mature males  
192 [20]. As well, spermatozoa in suspension from testes have higher refrigerated storage potentials than  
193 those stored in intact testes, spermic urine, or in semen (Supplementary Table 2. [6,46]).

194

195 Caudata spermatozoa in high concentrations and volumes in semen is easy to collect from  
196 seasonally mature or hormonally induced Cryptobranchidae [47,48], and as hormonally induced  
197 spermatozoa in cloacal fluid, even from small salamanders approximately 8 g in weight, making the  
198 collection of spermatozoa from testes unnecessary unless from recently dead individuals (Ruth Marcec,  
199 personal communication). The semen of the cryptobranchid, *A. davidianus*, is collected at concentrations  
200 of  $\sim 11 \times 10^6/\text{mL}$  and up to 12mL/kg of male weight [47,48] with adult males weighing up to 50 kg [49],  
201 but in lower absolute and comparative volumes of 2 mL in *Cryptobranchus* (the North American giant  
202 salamander clade, Dale McGinnity, personal communication). The comparative differences in sperm  
203 numbers between amphibian species may generally relate to different levels of sperm competition [45],  
204 including internal competition in Salamanders where the number of spermatozoa per spermatophore  
205 varies by more than three magnitudes (Supplementary Table 2. [47]).

206

207 Insert Fig. 4.

208

209 Processing during cryopreservation, thawing, and recovery can reduce the concentration of  
210 spermatozoa to less than <15% of the original [50]. Consequently, in samples derived from spermic  
211 urine the post-thaw spermatozoa concentrations could be lower than the fertilization optimum, and even  
212 lower than the fertilisation threshold. In Anurans, fertilization rates decline in a sigmoidal curve from  
213 the optimum spermatozoa concentration, and concentration three magnitudes lower than the optimum  
214 provided only 30% fertilization in one species [51] no fertilisation in another [52]. The optimal  
215 spermatozoa concentrations for fertilization may depend on other factors in addition to spermatozoa  
216 motility. These include chemo-attractants found in the oocyte gel, oocyte size, or in terrestrial-breeding  
217 and foam-nesting Anurans the direct deposition of spermatozoa onto oocytes [2].

218

219 Centrifugation is used to concentrate spermatozoa from spermic urine, to remove protein and  
 220 lipid components from fresh spermatozoa suspensions [44], and post-thaw to remove cryodiluents [20].  
 221 The use of centrifugation should be minimised as spermatozoa motility can be reduced by up to 50%  
 222 through morphological damage [53]. Therefore, in cases where there are surplus amphibians in breeding  
 223 programs [6], or males can be taken from wild populations [20], the collection of high numbers and  
 224 concentrations of spermatozoa directly through testes maceration may be the preferred option.

225  
 226 In some fish species the contamination of semen with urine decreases spermatozoa viability  
 227 [54,55], and in others urine is needed for spermatozoa maturation [56]. Anuran spermatozoa from  
 228 spermic urine exhibits slightly lower viability than testicular spermatozoa possibly as a consequence of  
 229 activation in the lower osmolality of urine in comparison to the isotonic testicular environment [45].  
 230 Nevertheless, spermatozoa stored in spermic urine at room temperature have been reported to retain  
 231 similar high levels of motility to testicular sperm for up to 45 m [34]. The effect of urine contamination  
 232 on Caudata spermatozoa in semen, or when sampled in milt is unknown, but urine contamination of  
 233 fishes semen negatively affects spermatozoon metabolism [54,55].

234

#### 235 **4. Effects of environmental factors on sperm motility activation and fertilization rate**

236

237 The major environmental factors affecting spermatozoa motility in externally fertilizing amphibians and  
 238 fishes are media osmolality, ionic composition, pH, and temperature [25,45]. The motility of  
 239 spermatozoa in some freshwater fishes is highly influenced by the extracellular concentration of  
 240 electrolytes [57,58]. The motility of spermatozoa is activated when sperm transition from the high  
 241 osmolality of the testes to the low osmolality of the freshwater environment, and with tested amphibians  
 242 is the major factor controlling activation [25,59,60]. With Anuran spermatozoa an osmolality of ~250  
 243 mOsmolkg<sup>-1</sup> prevents activation [25], and dependent on the species osmolalities below 70 mOsmolkg<sup>-1</sup>  
 244 [60] to 105 mOsmolkg<sup>-1</sup> [61] promote activation. Inter-specific variation in the optimal osmolality for  
 245 fertilization in Anurans was shown where in one species fertility was maintained up to 40 mM [60], in  
 246 another a steady decline in fertility occurred as osmolalities increased to more than 7 mOsmolkg<sup>-1</sup> [61],  
 247 and even intra-specific variation was shown in the optimum osmolality in one species [62].

248

249 Anuran spermatozoa show the longest period of sperm motility of all amphibians [25,61] with an  
 250 average period of motility of 1 h [25]. However, at extremes the spermatozoa of *Xenopus* only maintains

251 motility for 2 min [63,64] and in one species motility is extended to 7 h [61]. The spermatozoa of  
252 Cryptobranchidae and some freshwater fish stays motile for up to 10 min, but with most freshwater fish  
253 motility only lasts for seconds to a few min [25]. Higher osmolalities increase the longevity of  
254 spermatozoa possibly from less energy being partitioned from motility to maintaining osmotic  
255 equilibration [25,65].

256

257 The longevity of both fish [65,66,67] and amphibian spermatozoa [68] depends on specific  
258 metabolic pathways and the availability of energy substrates [69,70]. Adenosine triphosphate (ATP),  
259 adenosine diphosphate, and creatine phosphate [71,72] provide energy for flagella motion and maintain  
260 ionic and water balance across plasma membranes [57]. In many fish species increased spermatozoa  
261 velocity also positively corresponds with ATP levels [70]. Species variability and specificity of energy  
262 metabolism has been shown for fishes [72], however, species specificity has not been shown in the few  
263 studies of amphibians. In the Bufonid, *Anaxyrus fowleri*, ATP/adenosine monophosphate over a wide  
264 range of concentrations did not affect spermatozoon velocity or longevity (Robert Browne pers.  
265 communication).

266

#### 267 4.1 Diluents

268

269 Diluents are formulated to simulate the cellular concentrations of ions but may also contain organic  
270 supplements (Supplementary Table 3 [73,74]). Diluents approximating 220 mOsmolk<sup>-1</sup> are generally  
271 used as cryodiluents [37,38,73,74], to deactivate spermatozoon motility [74,75], and at low  
272 concentrations are used for post-thaw sperm equilibration, activation and fertilization [75]. The ionic  
273 composition of diluents and their osmolality for fish are similar to those of blood plasma, and generally  
274 include Na<sup>+</sup>, Ca<sup>2+</sup>, K<sup>+</sup>, Cl<sup>-</sup>, and H<sub>2</sub>CO<sub>3</sub><sup>-</sup> ions. Other ions that may be formulated in diluents are Mg<sup>2+</sup>,  
275 SO<sub>4</sub><sup>-</sup>, and increased relative concentrations of K<sup>+</sup> or Ca<sup>2+</sup> or H<sub>2</sub>CO<sub>3</sub><sup>-</sup>, however, their benefits are species  
276 specific in fish [57].

277

278 The formulation of diluents for amphibians were originally taken from physiological salines used  
279 for general biology [75], and now include formulations used for cell culture, and for the storage or  
280 fertilisation of fish and mammalian spermatozoa [1,2]. However, the K<sup>+</sup> concentration in most of these  
281 diluents is only ~2 mM, whereas, K<sup>+</sup> concentrations in anuran testicular plasma is ~70 mM in *Xenopus*

282 and ~40 mM in *Bufo* [76]. A low K<sup>+</sup> concentration in diluents can produce membrane damage, therefore,  
283 a greater knowledge of the role of K<sup>+</sup> and other ions in diluents for amphibian spermatozoa would be  
284 beneficial.

285  
286 With some fish species the pH of diluents is a major factor affecting spermatozoon motility,  
287 where a pH similar to or higher than that of seminal plasma promotes the activation and longevity of  
288 motility [58,77]. Because of few studies, the role of pH in spermatozoa motility in amphibians is not  
289 clear. Studies in Anurans show a higher pH 7.1-7.8 in spermic urine than that of urine (pH 6.7-6.8) [34].  
290 In the Caudata, with *A. davidianus* an artificially high pH 7.0-7.5 [47,48] increased refrigerated storage  
291 life and low pH inhibited flagella movement, and in contrast the semen of *Cryptobranchus* had a low pH  
292 6.4 (unpublished), and where the highest motility of *Axolotl* spermatozoa was found at pH 10.0-12.0  
293 (Nabil Mansour personal communication).

294  
295 Diluents are commonly mixed 1:1 by testis weight or semen volume to create sperm suspensions  
296 [50]. If spermatozoon suspensions are prepared from non-ionic diluents they will have only 50% of the  
297 original concentration of the various ions in testicular tissue, spermic urine, or semen. If sperm  
298 suspensions are prepared from ionic diluents the final concentration of ions in the suspensions will  
299 depend on the initial concentration of the diluent and the tissues or semen's ionic concentration.

300

#### 301 4.2 Stimulants and antioxidants

302

303 The stimulants caffeine and/or theophylline have been shown to increase the activation of Anuran  
304 spermatozoa [61,78]. The beneficial effects of these compounds on motility are likely the result of  
305 phosphodiesterase inhibition increasing levels of cyclic adenosine monophosphate, however, they may  
306 also influence spermatozoa motility and longevity as antioxidants by acting as antioxidants, and by  
307 suppressing reactive oxygen species formation within the sperm suspension [61]. The effects of other  
308 antioxidant compounds on Anuran spermatozoa have also been investigated, however, Vitamin C  
309 supplementation was detrimental and Vitamin E supplementation had no effect [79]. Pentoxifylline did  
310 not increase the motility of fresh *Xenopus* spermatozoa [64].

311

### 312 **5. Sperm processing and refrigerated storage**

313

314 Refrigeration at 4°C extends the storage life of spermatozoa by lowering metabolic rates, and most  
315 amphibian [53,80,81] and fish [82] spermatozoa largely tolerate refrigeration temperatures to above  
316 freezing. Anuran spermatozoa has retained moderate motility (~10-20%) after refrigerated storage of  
317 testis in carcasses for ~7 d [83], excised testes for ~14 d [80,83], testicular macerates for between ~14-  
318 21 d [80], spermic urine for ~7 d [40], and in the semen of a Cryptobranchid for ~4.5 d [48], and in  
319 hormonally induced cloacal fluid from a Salamander for ~3 d (Supplementary Table 2 [38]).

320

### 321 5.1 Oxygenation

322

323 Anuran spermatozoa uses both aerobic and anaerobic metabolism [67,84,85], whereas fish spermatozoa  
324 can only use aerobic metabolism [68,86]. Nevertheless, oxygenation extends the refrigerated storage life  
325 of spermatozoa in many fish species [87], however, in some species it decreases storage life and in  
326 others has no effect [86]. Increased oxygen concentration was shown to increase the refrigerated storage  
327 life of the spermatozoa of two anurans [85,88]. Oxidative damage to fish spermatozoa may be limited by  
328 components of seminal fluid [89], and this may also be the case with cryptobranchid spermatozoa in  
329 semen.

330

### 331 5.2 Processing Osmolalities

332

333 Spermatozoa are affected by varying osmolalities at various stages of processing, storage, activation,  
334 and fertilisation [1]. Spermatozoa suspensions whether created from testicular macerates [6,20],  
335 Cryptobranchid semen [47], salamander spermatophores [35] or cloacal fluid [37,39], or Anuran  
336 spermic urine [73], have high osmolalities. The greatest changes in osmolality between spermatozoa and  
337 diluents occur with the processing of spermic urine for refrigerated or cryopreserved storage, and after  
338 both refrigerated and cryopreserved storage when spermatozoa are equilibrated to recover motility [90].

339

### 340 5.3 Antibiotics and light

341

342 The ability of antibiotics to increase the storage life of refrigerated spermatozoa is a balance between  
343 their efficacy and their toxicity with contrasting results in different species and studies. In fish,

344 antibiotics have been shown to improve the storage of spermatozoa in several species [91,92]. In  
345 contrast, the use of antibiotics (penicillin–streptomycin and gentamicin) in diluents reduced motility  
346 during the refrigerated storage of Anuran spermatozoa from spermic urine [85] and spermatozoa  
347 suspensions from testicular macerates [88]. Gentamicin in concentrations of up to 4 mg mL<sup>-1</sup> did not  
348 affect the refrigerated storage life of spermatozoa in Anuran testicular macerates or in spermic urine, but  
349 did reduce bacterial contamination [46]. While there is currently no evidence that antibiotics increase  
350 the storage life of amphibian spermatozoa, antibiotics may inhibit bacteria and reduce the risk of  
351 pathogen transmission through stored samples [46]. In *A. davidianus* has strong light has been shown to  
352 decrease the longevity of fresh sperm [48].

353

## 354 **6. Sperm cryopreservation**

355

356 In response to the amphibian extinction crisis, the cryopreservation of testicular anuran spermatozoa with  
357 subsequent fertilization was achieved in the late-1990's by independent research teams in the Russian  
358 Federation [93,94] and Australia [52]. Early studies compared a wide range of penetrating  
359 cryoprotectants between species, diluents, and processing methods [52,74,95,96,97,98,99,100,101]  
360 focussing on dimethyl sulphoxide (DMSO [52,97]), glycerol [52,97], and methanol [97]. By 2010,  
361 DMSO was the most consistent penetrating cryoprotectant in achieving post-thaw motile and fertile  
362 spermatozoa. To extend the collection of spermatozoa to non-lethal techniques, research then focussed  
363 on techniques for the cryopreservation of hormonally induced spermatozoa in spermic urine. Motility  
364 and fertilization with post-thaw spermatozoa from spermic urine was first achieved in 2011, with the  
365 novel penetrating cryoprotectant dimethyl formamide (DMFA) giving greater recovery than DMSO  
366 [73]. DMFA was then successfully used with four other phylogenetically diverse Anuran species,  
367 *Pelophylax lessonae* [40], *Anaxyrus b. boreas* and *Lithobates sevosa* [100] and *Atelopus zeteki* [101].

368

369 Insert Fig. 5.

370

371 The preparation of cryosuspensions involves the mixing of spermatozoa samples with  
372 cryodiluents (Fig. 4). Compounds in cryodiluents act in synergy to protect spermatozoa during the rigors  
373 of freezing and thawing. Cryodiluents are formulated from penetrating cryoprotectants and non-  
374 penetrating cryoprotectants, and with amphibian spermatozoa now favour ionic/saccharide non-

375 penetrating cryoprotectants. Both sucrose [40,64] and trehalose [100,101] as saccharide non-penetrating  
376 cryoprotectants have provided high post-thaw recovery.

377

378 With fish spermatozoa, non-penetrating cryoprotectants including proteins, lipoproteins and  
379 lipids have increased plasma membrane resistance to osmotic stress along with post-thaw recovery  
380 [102,103,104]. With amphibians spermatozoa the inclusion of protein/lipids in cryodiluents provided  
381 high post-thaw recovery of viability in an Anuran [64], and in Caudata motility in *Cryptobranchus* (Dale  
382 McGinnity, unpublished data), and increased fertility in a Salamander [37]. A negative effect of a buffer  
383 was shown through a higher post-thaw recovery of the motility of Bufonid spermatozoa using DMSO  
384 alone than with the addition of HEPES buffer [40], while the addition of TRIS buffer did not affect the  
385 recovery of *Xenopus* spermatozoa [6].

386

387 Concentrations of DMSO or DMFA between 5-10% (v/v) in cryosuspensions have proven the  
388 most successful for the cryopreservation of amphibians spermatozoa [6,20,40,52,73] with up to 15%  
389 concentration in a Litorid frog [105]. However, high concentrations of up to 15% DMSO have generally  
390 proven effective with fish. Glycerol [106], trehalose alone [107], and propylene glycol [108] have  
391 proven more effective than DMSO in some fish species and may be suitable for some amphibian  
392 species. In some Anurans, even low concentrations of DMSO reduce hatch rates [66], and with  
393 spermatozoa from spermic urine reduce fertility and larval survival [83]. In contrast, with *Xenopus*  
394 spermatozoa DMSO proved less toxic than glycerol [6]. In one species post-thaw motility was more  
395 highly correlated to fertilisation with glycerol in contrast to DMSO [105].

396

397 Cryosuspensions are generally refrigerated for ~10 min before freezing to equilibrate  
398 spermatozoa to penetrating cryoprotectants in a low temperature environment that reduces  
399 cryoprotectant toxicity [44,73,101]. The penetration rates of penetrating cryoprotectants vary widely,  
400 where DMSO reaches equilibrium with fishes spermatozoa within 10 s [109], but with the penetration  
401 rate of glycerol being much lower [100,111]. Longer equilibration periods may benefit some Anuran  
402 cryopreservation protocols and did not affect post-thaw recovery after 20 min [6], with some sperm  
403 recovering motility and fertility after 6 d of refrigerated storage [112]. Consequently, to enable more  
404 flexibility in the timing of techniques, and to facilitate the use of some slowly penetrating  
405 cryoprotectants, the equilibration period of refrigerated cryosuspensions may extend to 20 min or more.

406

## 407 6.1 Cooling rates and cryopreservation

408

409 Samples can be frozen in the field using dry ice or through suspension into LN<sub>2</sub> vapour, and in facilities  
410 also using -80°C or programmable freezers. We categorise freezing rates as very slow (10°C/min), slow  
411 (30°C/min), moderate (110°C/min), fast (300°C/min), very fast (1200°C/min) from a broad canvassing  
412 of the studies in Supplementary Table 2. Stepped freezing rates for amphibians spermatozoa have  
413 achieved high post-thaw recovery with testicular spermatozoa [20,99] and with spermatozoa from  
414 spermic urine (Supplementary Table 2. [73,100]). The cryopreservation of some fishes [113] and  
415 amphibians [52] spermatozoa is affected by changes in cooling rates as low as 5°C/min, and in these  
416 cases the use of programmable freezers may be necessary.

417

418 The spermatozoa of several Caudata families have proven amenable to cryopreservation at slow  
419 to fast cooling rates. With *Axolotl* spermatophores, cooling rates between ~10°C/min and ~300°C/min  
420 did not affect viability [35], and high post-thaw recovery was shown with Salamander sperm using  
421 stepped freezing [37]. The sperm of *Cryptobranchus* proved amenable to cryopreservation with the slow  
422 lowering of straws into a LN<sub>2</sub> vapour (Dale McGinnity personal communication), but using a similar  
423 freezing method with *Andrias* only recovered <10% motility [47]. In fish optimal cooling rates are  
424 membrane lipid dependent [114,115] and this is expected to be the case with amphibian sperm.

425

426 **Sperm processing, activation, and fertilisation**

427

428 High levels of motility and fertility are recovered from cryopreserved amphibian sperm when thawed in  
429 a wide range of conditions from air at room temperature, to unheated tap water, to immersion into 37°C  
430 water baths (Supplementary Table 2, [6,100,101]). The first post-thaw recovery of hormonally induced  
431 Anuran sperm was achieved through a four-step osmotic equilibration process at 4°C [83], and with  
432 another Anuran the percentage activation, velocity, morphology, longevity, and DNA integrity were  
433 higher when spermatozoa were held at 4°C during the post-thaw processing. Therefore, once thawed as  
434 shown by the last remaining ice just thawing, cryosuspensions should be held as close to 4°C as  
435 possible. The period that post-thaw spermatozoa can be stored and maintain viability is unknown for  
436 most species, but with *Xenopus* last 20 min without any effect on fertilisation or embryo survival rates  
437 [6,44].



438

439 The highest fertilization rates are achieved through high spermatozoa per an oocyte ratios. The  
440 general practice of the mixing of cryodiluents with sperm samples at a 1:1 ratio reduces spermatozoa  
441 concentration by 50%. For activation, these suspensions are then generally mixed at a ratio of 1:2 or  
442 more with water resulting in a further reduction of spermatozoa concentration to ~20% or less of the  
443 original sample [51,73]. This lowering of spermatozoa concentrations is particularly significant with low  
444 concentrations of hormonally induced spermatozoa and where fertilisation requires high sperm  
445 concentrations [73], and also where a significant percentage of sperm lyse or do not recover motility  
446 [53]. For the highest fertilisation rates, a process known as dry fertilisation is used where sperm  
447 suspensions are deposited directly onto oocytes and then after 5 to 10 min the oocytes are flooded with  
448 fresh water [52,75].

449

## 450 **7. Morphological integrity of sperm**

451

452 Many morphological deformations can be found in post-thaw amphibian spermatozoa, such as swelling  
453 or rupture of the plasma membrane, loss of the nuclear envelope, fracture of the perforatorium and  
454 axoneme, degeneration of the undulating membrane and disappearance of the mitochondrial ridge  
455 [48,115]. Morphological damage may be associated with impacts on post-thaw activation mechanisms  
456 where fish [116,117] or Anuran [53,112] spermatozoa are intact but unable to activate. The positive  
457 correspondence between high post-thaw plasma membrane integrity and fertilization rates was shown  
458 with *Silurana tropicalis* in contrast to low membrane integrity and fertilisation rates in *Xenopus laevis*  
459 [6,115].

460

### 461 **Sperm DNA fragmentation (SDF)**

462

463 The main objective of spermatozoa storage is to provide unfragmented and viable genetic material to the  
464 oocyte upon fertilisation. Sperm DNA fragmentation (SDF) is a highly dynamic process that continues  
465 from spermatozoa collection until fertilization. Evolutionary history, morphology and physiology of  
466 spermatozoon, and DNA-protein interactions, affect SDF during refrigerated storage, cryopreservation,  
467 and post-thaw activation [118].

468

469 Sperm DNA fragmentation interferes with syngamy and embryonic development in fish [118].  
 470 However, to date there are only seven research publications of SDF in amphibians; *R. temporaria* sperm  
 471 stored in refrigerated carcasses [83], refrigerated storage in testes or macerates in *X. laevis* and *S.*  
 472 *tropicalis* [6,115,119], and the fresh hormonally induced and cryopreserved spermatozoa of *A. zeteki*  
 473 [44,101] and *Epidalea calamata* [120]. Sperm DNA fragmentation increased and fertilization rates  
 474 decreased during refrigerated storage of anuran sperm in carcasses [83], in spermic urine [45,101], and  
 475 in post-thaw spermatozoa [101,115]. However, SDF was not a predictor of survival rates from first  
 476 cleavage oocytes [6,83]. Sperm DNA fragmentation in post-thaw spermatozoa was higher in seasonally  
 477 collected spermatozoa than in unseasonal spermatozoa, but also did not correspond to reduced embryo  
 478 survival [120].

479  
 480 **9. Phylogenetic patterns in sperm induction and amenability to storage**

481  
 482 Phylogeny and environment interact to mold the reproductive strategies of amphibians [2,24,33]. Most  
 483 studies of amphibian spermatozoa collection and storage are on Anurans from regions in the temperate  
 484 zones of Australia, or the cool to warm temperate zones of Europe and North America (Supplementary  
 485 Table 1, 2). Southern Australia, has a cool to warm climate with stochastic seasonal rainfall and has  
 486 more studies of Anurans than any other region. In this climate Anurans generally reproduce over  
 487 extended seasonal periods (Supplementary Table 3. [22]). Recent developments of cryopreservation  
 488 techniques for fish spermatozoa have also focused on species from temperate climates and with seasonal  
 489 reproduction, including numerous studies in Brazil [104].

490  
 491 More studies over a wider range of families, and species within families, are needed to reveal  
 492 phylogenetic patterns in species amenability to spermatozoa cryopreservation Spermatozoa  
 493 cryopreservation has been trialed in 2 Bufonid species and 6 Ranid from the cool to warm temperate  
 494 zones of Eurasia and North America; and in 3 Bufonid species, 2 Hylid and 1 Eleutherodactylid from the  
 495 tropical and subtropical zones of South and Central America (Supplementary Table 3). Although  
 496 Bufonidae and Hylidae are sister clades, there were different responses to similar protocols, showing  
 497 high recovery for Bufonid spermatozoa [44] and low recovery for Hylid spermatozoa (Belin Proaño and  
 498 Oscar D. Pérez, personal communication). In contrast, the spermatozoa of both Ranids and Bufonids,  
 499 which diverged ~170mya [121], are amenable to cryopreservation. However, Pelodryadid sperm showed  
 500 greater amenability to cryopreservation than Myobatrachid sperm [98], where Pelodryadids diverged

501 from the Myobatrachids ~140 mya [122,123], and with no difference between spermatozoa from two  
502 Myobatrachid subfamilies that diverged 70 mya [99,123].

503

504 The Pipidae genera *Silurana* and *Xenopus* diverged only ~20-40 mya [124], however, post-thaw  
505 *S. tropicalis* sperm retains higher motility, membrane and DNA integrity than *X. laevis* sperm [15]. The  
506 two species of Pipidae (sub-order Mesobatrachia) in which cryopreservation has been trialed, are  
507 phylogenetically distant from other trialed anurans which are all from the Neobatrachidae, and in  
508 contrast to the spermatozoa of Neobatrachia the spermatozoa of both Pipidae species successfully  
509 cryopreserves in an ionic/sucrose diluent alone [6].

510

511 Insert Fig. 6.

512

513 A possible relationship between amphibians climatic range and the amenability of Anuran  
514 spermatozoa to cryopreservation was shown where the spermatozoa of freeze-tolerant wood frogs *Rana*  
515 *sylvatica* had a much higher post-thaw recovery than the more temperate climate leopard frogs *R.*  
516 *pipiens* and American toads *A. americanus* (Fig. 6, [125]) but this concept remains to be investigated  
517 over a wider range of species. Caudate spermatozoa from two distantly related families has been  
518 successfully cryopreserved in the Cryptobranchidae (Dale McGinnity personal communication, [46])  
519 and the Ambystomatida [35,36]. Phylogenetic patterns of spermatozoa cryopreservation still need to be  
520 established in the remaining eight Caudate families.

521

## 522 **10. Evaluation of techniques and their standardization**

523

524 The development of techniques for the collection, processing, and storage of amphibians spermatozoa  
525 depend on assessing spermatozoa quality through standardised metrics. In fishes, percentage motility  
526 and velocity [82], and plasma membrane integrity [115,126], have mainly been used as metrics of  
527 spermatozoa quality, with far fewer studies extending to fertility and development [127]. Research on  
528 Anurans spermatozoa has generally used percentage motility and velocity as a metric [61,87,128],  
529 though a number of studies have used live/dead (viability) stains [27,29,31], and to a lesser extent  
530 fertility and development depending on the study goals and the availability of oocytes [34,73,83,100].  
531 With Anurans, two recent Proof of Concept Studies used cryopreserved spermatozoa to produce mature

532 adults [6,20]. The post-thaw recovery of Caudate spermatozoa from semen has been assessed by fertility  
533 and development to mature adults (Dale McGinnity personal communication, [37]), and from  
534 spermatophores by live/dead stains [35]. However, vital stains may not always be reliable when  
535 assessing the membrane integrity of spermatozoa in spermatophores (Manuel Gonzalez pers.  
536 communication. Fig. 7.).

537  
538 Insert Fig. 7.

539  
540  
541 To further the development of both research and practical application, at each stage of processing the  
542 quality of spermatozoa should be assessed by standardized methods for percentage of activation,  
543 velocity, period of motility, concentration and volume. Any procedures to induce spermatozoa or  
544 oocytes should be recorded including the body length, weight, age, and reproductive condition of males,  
545 and testes weight when collecting testicular spermatozoa. Spermatozoa suspensions can be measured for  
546 pH, osmolality, and ionic composition. If a study includes spermatozoa morphology, where possible the  
547 cataloguing of images from both stained slides and electron microscopy should be undertaken.

548  
549 Some studies have shown an unexpected lowering of spermatozoa concentration during processing and  
550 storage [24] and a better understanding of the extent and nature of these lysed and missing spermatozoa  
551 is needed. Fertilization techniques should be quantified in terms of the number of spermatozoa used for  
552 specific numbers of oocytes (e.g. sperm-to-oocyte ratio), the associated water volumes and  
553 concentrations, gamete holding times, and protocols used for activation of the gametes.

554  
555 The percentage activation, motility, and also the velocity of spermatozoa can be assessed by observers  
556 using phase contrast microscopes, or more accurately percentage motility and velocity by Computer  
557 Assisted Sperm Analysis (CASA). CASA objectively measures the percentage of motile spermatozoa  
558 and their various types of velocity [82] and analyses the resultant data with a sophisticated statistical  
559 programs. In almost all cases, CASA systems rely on head movement of spermatozoa, and free software  
560 developed for fish spermatozoa is also suitable for amphibian spermatozoa as a plug-in of ImageJ  
561 software [129].

562

## 563 **11. Application of amphibian sperm banks**

564

565 Amphibians produced by *in vitro* fertilisation have been released in supplementation programs (Robert  
566 Browne, personal communication), and refrigerated spermatozoa transported between facilities to  
567 successfully fertilise oocytes [130], but no amphibians from cryopreserved spermatozoa have been  
568 released. In contrast, fish aquaculture has a long history with the use of *in vitro* fertilization since 600  
569 BC in mainland China (Fan-Li The Art of Fish-Breeding 600 BC), and has been widely used globally  
570 for the restocking of fish since the mid-20<sup>th</sup> century [131].

571

572 Although the spermatozoa of 200 aquaculture and 60 threatened species has been cryopreserved  
573 [132] it has only had limited use in practice especially for threatened populations [5,133]. For example,  
574 cryopreserved spermatozoa have been used in a CBP for marble trout (*Salmo marmoratus*) to maintain  
575 pure strains [5], and pallid sturgeon (*Scaphirhynchus albus*) juveniles included in general releases  
576 (William Wayman, personal communication).

577

578 The concept of sperm banks to support CBPs for select species is separate from archival genetic  
579 resource banking (AGRB) for the widest range of possible species for taxonomic and other purposes.  
580 The banking of amphibian spermatozoa requires cryopreservation, whereas, AGRB for taxonomic  
581 purposes requires the storage of samples, preferably including the whole specimen, at room temperature.  
582 In contrast to the indefinite storage period of AGRB, the storage period and the use of cryopreserved  
583 spermatozoa must be defined within a Sustainable Management Plan. Limited resources must be focused  
584 on species where the programs goal is the reestablishment of genetically varied populations in nature,  
585 with these examples then extending to the broader amphibian CBP community. Not all CBPs will  
586 require the use of cryopreserved spermatozoa [104] including those where genetically varied populations  
587 may be rebuilt without the use of cryopreserved spermatozoa. At an extreme, access to cryopreserved  
588 spermatozoa alone is useless if females are not available.

589

590 There are many fish sperm banks in Europe, Brazil, Mexico, USA, and Canada that are mostly  
591 dedicated to commercial fish, but between them and others globally house sperm samples from hundreds  
592 of threatened species. Some of these collections include spermatozoa from species that were common at  
593 the time of collection but are now endangered or extinct (see review [104]). Sperm banking should be  
594 based on forming links and partnerships, between the target CBP and other participating entities

595 including civil and governmental institutions, within a framework of overlapping and shared interests.  
596 The establishment of communication networks and information portals, along with standardization of  
597 terminologies and lexicon provide for efficient communication [104].

598  
599 Future expansion of the use of cryopreserved spermatozoa for aquatic species in aquaculture and  
600 will mostly be based on advances in high-throughput cryopreservation and commercial-scale application  
601 [134,135]). With amphibians the optimal cryopreservation protocol will vary dependent on the species,  
602 however, as the field develops greater standardization of protocols for at least each species will be  
603 advantageous [104,134], through increasing offspring production, and minimizing variability and the  
604 waste of samples (Fig. 8).

605

606 Insert Fig. 8.

607

608 Lifecycle Proof of Concept studies (Fig. 8. [18]) should now be integrated within select CBPs;  
609 based on the species conservation status, CBP facilities ability to complete the lifecycle, and the  
610 potential for release into their natural environment. Complete integration requires three stages, the  
611 development of appropriate technologies, funding for the establishment and maintenance of sperm  
612 banks, and the integration of sperm banks into CBPs.

613

614 The minimum number of males needed for CBP to maintain 90% of a species genetic variation  
615 in a 55 year CBP is 75 males, with numbers dramatically increasing with shorter generation times and  
616 lower longevity [136]. The cost of sperm banking depends on the scale where the minimum of one  
617 cryostorage container can house many hundreds of samples, and as storage capacity increase the storage  
618 costs per sample lowers and costs for other capital items such as a microscope stay static. The location  
619 of facilities in the low income regions of the highest amphibian biodiversity will generally lower labour  
620 costs [104]. The estimated total costs for sperm processing, cryopreservation, and storage of each  
621 sample in the USA, based on Caffey and Tiersch [135], is ~\$5.00 USD for the first year and less for  
622 subsequent years. The initial part of this cost for processing will be species specific, but in any case the  
623 cost of sperm banking is one to two orders of magnitude less than keeping live males and more secure.

624

625 The development of techniques for the cryopreservation of amphibian sperm have almost  
626 exclusively been in moderate to high income industrialised countries, except recently for three anuran  
627 species in Ecuador (Belin Proaño and Oscar D. Pérez, personal communication, [44]). Yet most  
628 amphibian species, except SE North America and eastern Australia, are found in the low to moderate  
629 income countries of Central and South America, SE Asia, New Guinea, Africa and Madagascar (Fig. 9).  
630 Most currently threatened amphibians come from Central and South America (Fig 9. [137]).

631

632 Insert Fig. 9.

633

634 Sperm banks and CBPs are ideally be located within species range, where males can be sampled  
635 opportunistically, cumbersome legislation regarding spermatozoa transport between facilities does not  
636 apply, and the CBP relates to the general sustainable management of the regional environment.  
637 However, the number of institutionally supported CBPs in or out of range countries can only support  
638 about 10% of species in need of conservation actions. Private keepers CBPs in or out of range offer an  
639 opportunity to prevent the extinction of the 90% of neglected amphibian species, along with supporting  
640 in range CBPs, habitat protection and restoration, and increasing public perception and political  
641 influence for the sustainable management of amphibian biodiversity [138].

642

## 643 **12. Conclusion and future directions**

644

645 There is an increasing need for the storage of spermatozoa in the sustainable management of amphibian  
646 biodiversity. Techniques for the use of cryopreserved spermatozoa are developed and there is no  
647 technical reason that sperm banking cannot be implemented for many species programs. Considering  
648 the depth of the amphibian conservation crisis it is imperative to develop "Proof of Concept Projects" for  
649 the use of cryopreserved spermatozoa in CBPs, and that broadly engage the global amphibian  
650 conservation community and promote the sustainable management of the environment in general.

651

652 Techniques for the collection and refrigerated storage of amphibian spermatozoa are well  
653 advanced in the Anura and Caudata, but neglected in the Sirenidae and Caecilians. Sperm banking must  
654 embrace the diversity of reproductive modes in amphibians, and further develop techniques to optimize  
655 the cryopreservation of their spermatozoa. Patterns between species phylogeny, their evolutionary

656 history, and reproductive modes will lead to more generalized concepts regarding the cryopreservation  
 657 of amphibian spermatozoa. This progress will be furthered by a greater understanding of the critical  
 658 components of protocols, and a greater standardization of methods to enable more meaningful  
 659 comparisons between studies and to focus on critical points in the cryopreservation process.

660

#### 661 **Acknowledgements:**

662

663 R. B. is thankful to Nashville Zoo, USA. B. D. and J. C. to the Ministry of Education, Youth and Sports  
 664 of the Czech Republic—projects CENAKVA (No. CZ.1.05/2.1.00/01.0024), CENAKVA II (No.  
 665 LO1205 under the NPU I program) CZ.02.1.01./0.0/0.0/16\_025/0007370 Reproductive and genetic  
 666 procedures for preserving fish biodiversity and aquaculture and by the Grant Agency of the University  
 667 of South Bohemia in Ceske Budejovice (125/2016/Z and 114/2013/Z), and by the Czech Science  
 668 Foundation (No. P502/15-12034S). G. Della Togna is thankful to the Secretaría Nacional de Ciencia y  
 669 Tecnología de Panamá (SENACYT) and the Sistema Nacional de Investigación de Panamá (SNI).

670

#### 671 **References**

- 672 [1] Clulow J, Clulow S. Cryopreservation and other assisted reproductive technologies for the  
 673 conservation of threatened amphibians and reptiles: bringing the ARTs up to speed. *Reprod Fert*  
 674 *Devel* 2016;28(8):1116-1132.
- 675 [2] Silla AJ, Byrne PG. The role of reproductive technologies in amphibian conservation breeding  
 676 programs. *Ann Rev Anim Biosci* 2019;7:10-21.
- 677 [3] Ananjeva NB, Uteshev VK, Orlov NL, Ryabov SA, Gakhova EN, Kaurova SA, Kramarova L,  
 678 Shishova NV, Browne RK. Comparison of the modern reproductive technologies for  
 679 amphibians and reptiles. *Russian J Herpet* 2017;24(4):275-290.
- 680 [4] Carmichael C, Westerfield M, Varga ZM. Cryopreservation and in vitro fertilization at the Zebrafish  
 681 International Resource Center. *Methods Mol Biol* 2009;546:45-65.
- 682 [5] Crivelli AJ, Poizat G, Berrebi P, Jesensek D, Rubin JF. Conservation biology applied to fish: The  
 683 example of a project for rehabilitating the marble trout (*Salmo marmoratus*) in Slovenia. *Cybiurn*  
 684 2000;24:211-230.



- 685 [6] Pearl E, Morrow S, Noble A, Lerebours A, Marko Horb M, Guille M. An optimized method for  
686 cryogenic storage of *Xenopus* sperm to maximize the effectiveness of research using genetically  
687 altered frogs. *Theriogenology* 2017;92:149-155.
- 688 [7] Khosla K, Wang Y, Hagedorn M, Qin Z, Bischof J. Gold nanorod induced warming of embryos  
689 from the cryogenic state enhances viability. *ACS Nano* 2017;11:7869-7878
- 690 [8] Hagedorn M, Kleinhans F, Wildt D, Rall W. Chill sensitivity and cryoprotectant permeability of  
691 dechorionated zebrafish embryos, *Brachydanio Rerio*. *Cryobiology* 1997;34:251-263.
- 692 [9] Clulow J, Trudeau V, and Kouba A. Amphibian declines in the twenty-first century: why we need  
693 assisted reproductive technologies. In: Holt WV, Brown JL, Comizzoli P, editors. *Reproductive*  
694 *Sciences in Animal Conservation*, Vol. 753. Springer: New York; 2014, pp. 275–316.
- 695 [10] Higaki S, Eto Y, Kawakami Y, Yamaha E, Kagawa N, Kuwayama M, Nagano M, Katagiri S,  
696 Takahashi, Y. Production of fertile zebrafish (*Danio rerio*) possessing germ cells (gametes)  
697 originated from primordial germ cells recovered from vitrified embryos. *Reproduction*  
698 2010;139:733-740.
- 699 [11] AArk. 2018a. Amphibian Ark – about us, <http://www.amphibianark.org/about-us/>; [accessed 20<sup>th</sup>  
700 November 2018]
- 701 [12] AArk. 2018b. Amphibian Ark Conservation Assessments,  
702 <http://www.amphibianark.org/conservation-programs/assessment-results/>; [accessed 20<sup>th</sup>  
703 November 2018]
- 704 [13] Williams SE, Hoffman EA. Minimizing genetic adaptation in captive breeding programs: A review.  
705 *Biol Conserv* 2009;142:2388-2400.
- 706 [14] Baker A. Animal ambassadors: an analysis of the effectiveness and conservation impact of ex situ  
707 breeding efforts. In: Zimmerman A, Hatchwell M, West C, Lattis R, editors. *Zoos in the 21st*  
708 *Century Catalysts for Conservation*. Cambridge, UK, 2007, p. 373.
- 709 [15] Frankham R. Genetic adaptation to captivity in species conservation programs. *Mol Ecol*  
710 2008;17:325-333.
- 711 [16] Yan F, Lu J, Zhang B, Yuan Z, Zhao H, Huang S, Wei F, Wu M, Xiao H, Liang Z, Jin J, Wu S, Xu  
712 C, Tapley B, Turvey ST, Papenfuss TJ, Cunningham AA, Murphy RW, Zhang Y, Che J. 2018.  
713 The Chinese giant salamander exemplifies the hidden extinction of cryptic species. *Curr Biol*  
714 2018;28:R592–R594.

- 715 [17] Turvey ST, Chen S, Tapley B, Wei G, Xie F, Yan F, Yang j, Liang Z, Tian H, Wu M, Okada S,  
716 Wang J, Lu J, Zhou F, Papworth SK, Redbond J, Brown T, Che J, Cunningham AA. Imminent  
717 extinction in the wild of the world's largest amphibian. *Curr Biology* 2018;28:590–592.
- 718 [18] Browne RK, Li H, Robinson H, Uteshev VK, Shishova NR, McGinnity D, Nofs S, Figiel CR,  
719 Mansour N, Lloyd R, Agnew D, Carleton C, Gakhova EN. Reptile and amphibian conservation  
720 through gene banking and other reproduction technologies. *Russian J Herpetol* 2011;18(3):165-  
721 174.
- 722 [19] Poleo GA. 2002. Intracytoplasmic sperm injection (ICSI) in fishes. PhD Dissertation. The School of  
723 Forestry, Wildlife, and Fisheries, Graduate Faculty of the Louisiana State University and  
724 Agricultural and Mechanical College, USA.
- 725 [20] Upton R, Clulow S, Mahony MJ, Clulow J. Generation of a sexually mature individual of the  
726 Eastern dwarf tree frog, *Litoria fallax*, from cryopreserved testicular macerates: proof of capacity  
727 of cryopreserved sperm derived offspring to complete development. *Conserv Physiol*  
728 2018;6(1):coy043; doi:10.1093/conphys/coy043.
- 729 [21] Vance CK, Julien A, Counsell K, Marcec R, Agcanas L, Tucker A, Kouba AJ. Amphibian ART  
730 over generations: frozen sperm offspring produce viable F2 generation. *Abstracts / Cryobiology*  
731 2018;85:178.
- 732 [22] Browne RK, Seratt J, Vance, C, Kouba, A. Hormonal induction with priming and *In vitro*  
733 fertilization increases egg numbers and quality in the Wyoming toad (*Bufo baxteri*). *Reprod Biol*  
734 *Endocrinol* 2006;4:34.
- 735 [23] Costanzo JP, Mugnano JA, Wehrheim HM. Osmotic and freezing tolerance in spermatozoa of  
736 freeze-tolerant and -intolerant frogs. *Am J Physiol* 1998;44:713-719.
- 737 [24] Clulow J, Pomeroy M, Herbert D, Upton R, Calatayud N, Clulow S, Mahony MJ, Trudeau VL.  
738 Differential success in obtaining gametes between male and female Australian temperate frogs  
739 by hormonal induction: A review. *Gen Comp Endocrin* 2018;doi:  
740 <https://doi.org/10.1016/j.ygcen.2018.05.032>
- 741 [25] Browne RK, Kaurova SA, Uteshev VK, Shishova NV, Kramarova L., McGinnity D, Figiel CR,  
742 Mansour N, Agnew D, Wu M, Gakhova EN, Dzyuba B, Cosson J. Sperm motility of externally  
743 fertilizing fish and amphibians. *Theriogenology* 2015;83(1):1-13.
- 744 [26] Kuehnelt S, Kupfer A. 2012. Sperm storage in caecilian amphibians. *Frontiers Zool* 2012;9:12.

- 745 [27] Kühnel S, Reinhard S, Kupfer A. Evolutionary reproductive morphology of amphibians: an  
746 overview. *Bonn Zool Bull* 2010;57(2):119-126.
- 747 [28] Iskandar DT, Evans BJ, McGuire JA. A novel reproductive mode in frogs: A new species of fanged  
748 frog with internal fertilization and birth of tadpoles. *PLoS ONE* 2014;9(12): e115884.
- 749 [29] Byrne PG, Silla, AJ. Hormonal induction of gamete release, and in-vitro fertilisation, in the  
750 critically endangered Southern Corroboree Frog, *Pseudophryne corroboree*. *Reprod Biol*  
751 *Endocrin* 2010; 30(10):1352-1358.
- 752 [30] Lipke C, Meinecke-Tillmann S, Meinecke B. Induced spermiation and sperm morphology in the  
753 green poison frog, *Dendrobates auratus*. *Salamandra* 2009;45(2):65-74.
- 754 [31] Silla AJ. Effects of luteinizing hormone-releasing hormone and argininevasotocin on the sperm-  
755 release response of Günther's Toadlet, *Pseudophryne guentheri*. *Reprod Biol Endocrinol*  
756 2010;8:139. doi: 10.1186/1477-7827-8-139
- 757 [32] Silla AJ. Effect of priming injections of luteinizing hormone-releasing hormone on spermiation and  
758 ovulation in Günther's toadlet, *Pseudophryne guentheri*. *Reprod Biol Endocrinol* 2011;9:68.  
759 doi.org/10.1186/1477-7827-9-68
- 760 [33] Silla AJ, Roberts JD. Investigating patterns in the spermiation response of eight Australian frogs  
761 administered human chorionic gonadotropin (hCG) and luteinizing hormone-releasing hormone  
762 (LHRHa). *Gen Comp Endocrinol* 2012;179:128-136.
- 763 [34] Della Togna G, Trudeau VL, Gratwicke B, Evans M, Augustine L, Chia H, Bronikowski EJ,  
764 Murphy JB, Comizzoli P. Effects of hormonal stimulation on the concentration and quality of  
765 excreted spermatozoa in the critically endangered Panamanian golden frog (*Atelopus zeteki*).  
766 *Theriogenology*. 2017;91:27-35.
- 767 [35] Figiel CR Jr. Cryopreservation of sperm from the axolotl *Ambystoma mexicanum*: Implications for  
768 conservation. *Herpet Conserv Biol* 2013;8(3):748-755.
- 769 [36] Mansour N, Lahnsteiner F, Patzner RA. Collection of gametes from live axolotl, *Ambystoma*  
770 *mexicanum*, and standardization of *in vitro* fertilization. *Theriogenology* 2011;75(2):354-361.
- 771 [37] Marcec R, Langhorne C, Vance C, Kouba A, Willard S. Cryopreservation of spermic milt in the  
772 model species *Ambystoma tigrinum* (Tiger salamander) for application in endangered  
773 salamanders. *Cryobiology* 2014;69(3):349-530.

- 774 [38] Uteshev VK, Kourova SA, Shishova NV, Stolyarov SD, Browne RK, Gakhova EN. *In vitro*  
775 fertilisation with hormonally induced sperm and eggs from sharp-ribbed newts *Pleurodeles waltl*.  
776 Russian J Herpet 2015;22(1):35-40.
- 777 [39] Uteshev VK, Shishova N V, Kaurova S A, Browne RK, Gakhova EN. Hormonal induction of  
778 spermatozoa from amphibians with *Rana temporaria* and *Bufo bufo* as anuran models. *Reprod*  
779 *Fertil Dev* 2012;24(4):599-607.
- 780 [40] Uteshev VK, Shishova N, Kaurova S, Manokhin A, Gakhova E. Collection and cryopreservation  
781 of hormonally induced sperm of pool frog (*Pelophylax lessonae*). *Russian J Herpet*  
782 2013;20(2):105-109.
- 783 [41] Silla AJ, McFadden M and Byrne PG. Hormone-induced sperm-release in the critically endangered  
784 Booroolong frog (*Litoria booroolongensis*): effects of gonadotropin-releasing hormone and  
785 human chorionic gonadotropin. *Conserv Physiol* 2019;7(1): coy080
- 786 [42] Browne RK, Seratt J, Li H, Kouba A. Progesterone improves the number and quality of hormonally  
787 induced fowler toad (*Bufo fowleri*) oocytes. *Reprod Biol Endocrin* 2006;4:3.
- 788 [43] Trudeau VL, Somoza GM, Natale GS, Pauli B, Wignall J, Jackman P, Doe K, Schueler FW.  
789 Hormonal induction of spawning in 4 species of frogs by coinjection with a gonadotropin  
790 releasing hormone agonist and a dopamine antagonist. *Reprod Biol Endocrinol* 2010;8:36.
- 791 [44] Proaño B, Pérez OD. *In vitro* fertilizations with cryopreserved sperm of *Rhinella marina* (Anura:  
792 Bufonidae) in Ecuador. *Amphib Rept Conserv* 2017;11(2):1–6.
- 793 [45] Della Togna G, Gratwicke B, Evens M, Augustine L, Chai H, Bronikowski E, Murphy JB,  
794 Comizzoli P. Influence of extracellular environment on the motility and structural properties of  
795 spermatozoa collected from hormonally stimulated Panamanian Golden Frog (*Atelopus zeteki*).  
796 *Theriogenology* 2018;108:153-160.
- 797 [46] Peng L, Yue YM, Liu Y. Effect of cryopreservation and short-term storage of Chinese giant  
798 salamander sperm. *Acta Hydrobiologica Sinica* 2011;35:325-332.
- 799 [47] Zhao H, Zhang H, Wang Q, Chen J. Preliminary study on conservation of *Andrias davidianus*  
800 sperm *In vitro*. *J Hydroecol* 2012;33(1):149-151.
- 801 [48] Keogh LM, Byrne PG, Silla AJ. The effect of gentramycin on sperm motility and bacterial  
802 abundance during chilled sperm storage in the Booroolong frog. *Gen Comp Endocrinol*  
803 2016;243:51-59.
- 804 [49] Browne RK, Li H, Wang Z, Okada S, Hime P, McMillan A, Wu M, Diaz R, McGinnity D, Briggler

- 805 JT. The giant salamanders (Cryptobranchidae): Part B. Biogeography, ecology and reproduction.  
806 Amphib Rept Conserv 2013;5(4):30-50.
- 807 [50] Doyle JM, McCormick CR, DeWoody JA. The quantification of spermatozoa by real-time  
808 quantitative PCR, spectrophotometry, and spermatophore capsizes. Mol Ecol Resour  
809 2011;11:101-106.
- 810 [51] Edwards DL, Mahony MJ, Clulow J. 2004. Effect of sperm concentration, medium osmolality and  
811 oocyte storage on artificial fertilisation success in a myobatrachid frog (*Limnodynastes*  
812 *tasmaniensis*). Reprod Fertil Devel 16:347-354.
- 813 [52] Browne RK, Clulow J, Mahony M, Clark A. Successful recovery of motility and fertility of  
814 cryopreserved cane toad (*Bufo marinus*) sperm. Cryobiology 1998;37:339-345
- 815 [53] Fitzsimmons C, Mahony MJ, Clulow J. Cold shock during rapid cooling of sperm from a tropical  
816 anuran, the cane toad, *Bufo marinus*. Reprod Fert Develop 2009;21(9):103-103.
- 817 [54] Dreanno C, Suquet M, Desbruyères E, Cosson J, Le Delliou H, Billard R. Effect of urine on semen  
818 quality in turbot (*Scophthalmus maximus*). Aquaculture 1998;169:247-262.
- 819 [55] Perchec G, Cosson J, André F, Billard R. Degradation of the quality of carp sperm by urine  
820 contamination during stripping. Aquaculture 1995;129:133-137
- 821 [56] Dzyuba B, Cosson J, Boryshpolets S, Bondarenko O, Prokopchuk G, Gazo I, Dzyuba V, Rodina M,  
822 Linhart O. *In vitro* maturation of spermatozoa in *Acipenser ruthenus*. Reprod. Biol 2014;14:160-  
823 163.
- 824 [57] Cosson J. 2008. The motility apparatus of fish spermatozoa. In: Alavi, S.M.H., Cosson, J.J.,  
825 Coward, K. & Rafiee, G., (Eds). *Fish Spermatology* (pp. 281-316). Alpha Science: Oxford.
- 826 [58] Alavi SM, Cosson J. Sperm motility in fishes. (II) Effects of ions and osmolality: A review. Cell  
827 Biol Internat 2006;30(1):1-14.
- 828 [59] Rugh, R. Induced breeding. In Experimental Embryology. Techniques and Procedures, Burgess  
829 Publishing Company, Minneapolis 1962;1:91-103.
- 830 [60] Clark AK. The effects of ionicity in diluents on the motility of cane toad (*Bufo marinus*) sperm. B.  
831 Sc. (Hons) thesis. University of Newcastle, Australia; 1997.
- 832 [61] Silla AJ, Keogh LM, Byrne PG. Sperm motility activation in the critically endangered Booroolong  
833 frog: the effect of medium osmolality and phosphodiesterase inhibitors. Reprod Fertility  
834 Develop 2017;29(11):2277-2283.

- 835 [62] Byrne PG, Dunne C, Munn AJ, Silla AJ. Environmental osmolality influences sperm motility  
836 activation in an anuran amphibian. *J Evol Biol* 2015;28:521-534.
- 837 [63] Wolf DP, Hedrick JL. A molecular approach to fertilization 11. Viability and artificial fertilization  
838 of *Xenopus laevis* gametes. *Develop Biol* 1971;25:360-376.
- 839 [64] Sargent MG, Mohun TJ. Cryopreservation of sperm of *Xenopus laevis* and *Xenopus tropicalis*.  
840 *Genesis* 2005;41(1):41-46.
- 841 [65] Billard R, Cosson MP. The energetics of teleost sperm motility. In: Gascon C, editor. Controls of  
842 sperm motility: Biological and Clinical Aspects. CRC Press;1990, p. 155-173.
- 843 [66] Burness G, Casselman SJ, Schulte-Hostedde A I, Moyes CD, Montgomerie R. Sperm swimming  
844 speed and energetics vary with sperm competition risk in bluegill (*Lepomis macrochirus*). *Behav*  
845 *Ecol Sociobiol* 2004;56(1):65-70.
- 846 [67] Cosson J. 2010. Frenetic activation of fish spermatozoa flagella entails short-term motility,  
847 portending their precocious decadence. *J Fish Biol* 2010;76:240-279.
- 848 [68] Del Rio AG. Metabolism of amphibian spermatozoa in relation to their motility. *Experimentia*  
849 1979;35:185-186.
- 850 [69] Cosson J. ATP, the sperm movement energizer. In: Kuester E, Traugott G, editors. Adenosine  
851 Triphosphate Chemical Properties, Biosynthesis and Functions in Cells. Nova Science Pub Inc;  
852 UK ed. Edition,2013, pp. 1-46.
- 853 [70] Dzyuba V, Dzyuba B, Cosson J, Rodina M. Enzyme activity in energy supply of spermatozoon  
854 motility in two taxonomically distant fish species (sterlet *Acipenser ruthenus*, Acipenseriformes  
855 and common carp *Cyprinus carpio*, Cypriniformes). *Theriogenology* 2016b;85(4):567-574.
- 856 [71] Fedorov P, Dzyuba B, Fedorova G, Grabic R, Cosson J, Rodina. M. Quantification of adenosine  
857 triphosphate, adenosine diphosphate, and creatine phosphate in sterlet spermatozoa during  
858 maturation. *J Anim Sci* 2015;93(11):5214–5221.
- 859 [72] Biegniewska A. Ziętara M S, Rurangwa E, Ollevie, ., Swierczynski J, Skorkowski EF. Some  
860 differences between carp (*Cyprinus carpio*) and African catfish (*Clarias gariepinus*)  
861 spermatozoa motility. *J App Ichthyol* 2012;26(5):674–677.
- 862 [73] Browne RK, Mahony M, Clulow J. A comparison of sucrose, saline, and saline with egg-yolk  
863 diluents on the cryopreservation of cane toad (*Bufo marinus*) sperm. *Cryobiology*  
864 2002d;44(3):251-257.

- 865 [74] Hollinger TG, Corton GL. Artificial fertilization of gametes from the South African clawed frog,  
866 *Xenopus laevis*. Gamete Res 1980;3:45-57.
- 867 [75] Shishova NR, Uteshev VK, Kaurova SA, Browne RK, Gakhova EN. Cryopreservation of  
868 hormonally induced sperm for the conservation of threatened amphibians with *Rana temporaria*  
869 as a model research species. Theriogenology 2011;75:220-232
- 870 [76] Inoda T, Morisawa M. Effect of osmolality on the initiation of sperm motility in *Xenopus laevis*.  
871 Comp Biochem Physiol Part A: Physiol 1987;88:539-542.
- 872 [77] Baynes SM, Scott AP, Dawson AP. Rainbow trout, *Salmo gairdnerii* Richardson, spermatozoa:  
873 effects of cations and pH on motility. J Fish Biol 1981;19:259–267.
- 874 [78] Fitzsimmons C, McLaughlin EA, Mahony MJ, Clulow J. Optimisation of handling, activation and  
875 assessment procedures for *Bufo marinus* spermatozoa. Reprod Fertil Dev 2007;19(4):594-601.
- 876 [79] Keogh LM, Byrne PG, Silla AJ. The effect of antioxidants on sperm motility activation in the  
877 Booroolong frog. Anim Reprod Sci 2017;183:126-131.
- 878 [80] Browne RK, Clulow J, Mahony M. Short-term storage of cane toad (*Bufo marinus*) gametes.  
879 Reproduction 2001;121:167-173.
- 880 [81] Unger S, Mathis A, Wilkinson R. A comparison of sperm health in declining and stable populations  
881 of Hellbenders (*Cryptobranchus alleganiensis alleganiensis* and *C.a. bishopi*). Am. Midl. Nat.  
882 2013;170:382-392
- 883 [82] Fauvel C, Suquet M, Cosson J. Evaluation and determinism of fish sperm quality. J. Appl Ichthyol  
884 2010;26:636-643.
- 885 [83] Shishova NR, Uteshev VK, Sirota NP, Kuznetsova EA, Kaurova SA., Browne RK, Gakhova EN.  
886 The quality and fertility of sperm collected from European common frog (*Rana temporaria*)  
887 carcasses refrigerated for up to 7 days. Zoo Biol 2013;32(4):400-406.
- 888 [84] Bernardini, G., Belgiojoso P, Camatini M. 1988b. *Xenopus* spermatozoon: Is there any correlation  
889 between motility and oxygen consumption? Gamete Res 21(4):403-408.
- 890 [85] Germano M, Arregui L, Kouba AJ. Effects of aeration and antibiotics on short-term storage of  
891 Fowler's toad (*Bufo fowleri*) sperm. Aquaculture 2013;396-399:20-24.
- 892 [86] Kowalski R, Cejko BI, Irnazarow I, Szczepkowski M, Dobosz S, Glogowski J. Short-term storage  
893 of diluted fish sperm in air versus oxygen. Turkish J Fisher Aquat Sci 2014;14:831-834.
- 894 [87] Dzyuba V, Cosson J, Dzyuba B, Yamaner G, Rodina M, Linhart O. The antioxidant system of  
895 seminal fluid during in vitro

- 896 [88] Silla AJ, Keogh LM, Byrne PG. Antibiotics and oxygen availability affect the short-term storage of  
897 spermatozoa from the critically endangered Booroolong frog, *Litoria booroolongensis*. *Reprod*  
898 *Fertil Dev* 2015;27:1147–1153.
- 899 [89] Dzyuba V, Dzyuba B, Cosson J, Boryshpolets S, Yamaner G, Kholodniy V, Rodina M. The  
900 antioxidant system of sterlet seminal fluid in testes and Wolffian ducts. *Fish Physiol Biochem*  
901 2014;40(6)-1731-1739.
- 902 [90] Gwo JC. Methods for sperm collection. In: Cabrita E, Robles V, Herraez P, editors. *Methods in*  
903 *Reproductive Aquaculture: Marine and Freshwater Species*. CRC Press; 2008, p. 81-92.
- 904 [91] Segovia M, Jenkins JA, Paniagua-Chavez C, Tiersch TR. Flow cytometric evaluation of antibiotic  
905 effects on viability and mitochondrial function of refrigerated spermatozoa of Nile tilapia.  
906 *Theriogenology* 2000;53:1489-1499.
- 907 [92] Rahimi R, Hajirezaee S, Shalvei F, & Kaboutari Katadj J. Antibiotics, penicillin and streptomycin  
908 improve semen quality indices of endangered Caspian brown trout, *Salmo trutta caspius*  
909 (Kessler, 1870) during in vitro short-term storage. *Aquacult Res* 2015;47(11):3662-3666.
- 910 [93] Kaurova SA, Chekurova NR, Melnikova EV, Uteshev VK, Gakhova EN. Cryopreservation of frog  
911 *Rana temporaria* sperm without loss of fertilizing capacity. Presented at the Proceedings of XIV  
912 Working Meeting of Genetic Resource Conservation, Pushchino. 1996, p. 106–108.
- 913 [94] Kaurova SA, Uteshev VK, Chekurova NR, Gakhova EN. Cryopreservation of testis of frog *Rana*  
914 *temporaria*. *Infusionsther Transfusionsmed* 1997;24(5):78–79.
- 915 [95] Hopkins BK, Herr C. Cryopreservation of frog (*Rana pipiens*) sperm collected by non-lethal  
916 methods. *Canadian J Zool* 2004;82:555-563.
- 917 [96] Michael SF, Jones C. Cryopreservation of spermatozoa of the terrestrial Puerto Rican frog,  
918 *Eleutherodactylus coqui*. *Cryobiology* 2004;48-90-94
- 919 [97] Mugnano JA, Costanzo JP, Beesley SG, Lee RE. Evaluation of glycerol and dimethyl sulfoxide for  
920 the cryopreservation of spermatozoa from the wood frog (*Rana sylvatica*). *Cryo-Letters*  
921 1998;19:249-254.
- 922 [98] Mansour N, Lahnsteiner F, Robert A, Patzer RA. Motility and cryopreservation of spermatozoa of  
923 European common frog, *Rana temporaria*. *Theriogenology* 2010;74:724-732.
- 924 [99] Browne RK, Clulow J, Manony M. The short-term storage and cryopreservation of spermatozoa  
925 from hylid and myobatrachid frogs. *Cryo Letters* 2002;23:129-136.



- 926 [100] Langhorne CJ, Calatayud NE, Kouba AJ, Feugang JM, Vance CK, Willard ST. Cryoconservation:  
 927 Successful sperm cryopreservation and developmental outcomes using endangered North  
 928 American amphibians. *Cryobiology* 2013;67(3):405.
- 929 [101] Della Togna NG. Structural and functional characterization of the Panamanian golden frog  
 930 (*Atelopus zeteki*) spermatozoa - impact of medium osmolality and cryopreservation on motility  
 931 and cell viability. **University of Maryland, College Park, ProQuest Dissertations Publishing,**  
 932 **2015. 10011589**
- 933 [102] Kopeika E, Kopeika J. In: Alavi SMH, Cosson J, Coward K, Rafiee G, editors. Variability of  
 934 sperm quality after cryopreservation in fish: *Fish Spermatology*. Oxford: Alpha Science  
 935 International; 2008, p. 347-396.
- 936 [103] Suquet M, Dreanno C, Fauvel C, Cosson J, Billard R. Cryopreservation of sperm in marine fish.  
 937 *Aquacult Res* 2000;31(3):231-343.
- 938 [104] Martínez-Páramo S, Horváth A, Labbé C, Zhang T, Robles V, Herráez P, Suquet M, Adams S,  
 939 Viveiros A, Tiersch TR, Cabrita C. Cryobanking of aquatic species. *Aquaculture* 2017;472:156-  
 940 177.
- 941 [105] Upton R, Clulow S, Seeto Rebecca, Wong L, Mahony M, Clulow J. Successful sperm  
 942 cryopreservation and generated offspring of the endangered frog, *Litoria aurea*. *Abstracts /*  
 943 *Cryobiology* 2018;85:148-149.
- 944 [106] Lim HK, Le MH. Evaluation of extenders and cryoprotectants on motility and morphology of  
 945 longtooth grouper (*Epinephelus bruneus*) sperm. *Theriogenology* 2013;79:867-871.
- 946 [107] Kiriya A, Gallardo WG, Bart AN. Successful hybridization of groupers (*Epinephelus coioides* x  
 947 *Epinephelus lanceolatus*) using cryopreserved sperm. *Aquaculture* 2011;320:106-112.
- 948 [108] Liu QH, Ma DY, Xu SH, Xiao ZZ, Xiao YS, Song ZC, Li J. Summer flounder (*Paralichthys*  
 949 *dentatus*) sperm cryopreservation and application in interspecific hybridization with olive  
 950 flounder (*P. olivaceus*). *Theriogenology* 2015;83:703-710.
- 951 [109] Prokopchuk G, Dzyuba B, Rodina M, Cosson J. Control of sturgeon sperm motility: Antagonism  
 952 between K(+) ions concentration and osmolality. *Anim Reprod Sci* 2016;164:82-89.
- 953 [110] Bickis IJ, Kazaks K, Finn JJ, Henderson IWD. Permeation kinetics of glycerol and dimethyl  
 954 sulfoxide in Novikoff hepatoma ascites cells \**Cryobiology* 1967;4(1):1-10.

- 955 [111] Gilmore JA, Liu J, Woods EJ, Peter AT, Critser JK. Cryoprotective agent and temperature effects  
 956 on human sperm membrane permeabilities: convergence of theoretical and empirical approaches  
 957 for optimal cryopreservation methods. *Human Reprod* 2000;15(2):335-343.
- 958 [112] Browne RK, Davis J, Pomeroy M, Clulow J. Storage of cane toad (*Bufo marinus*) sperm for 6  
 959 days at 0°C with subsequent cryopreservation. *Reprod Fert Develop* 2002b;14:267-273.
- 960 [113] Yang H, Norris M, Winn R, Tiersch TR. Evaluation of cryoprotectant and cooling rate for sperm  
 961 cryopreservation in the euryhaline fish medaka *Oryzias latipes*. *Cryobiology* 2010;61:211-219.
- 962 [114] Horokhovatskyi Y, Samples S, Dzyuba B, Cosson J, Linhart O, Rodina M, Fedorov P, Blecha M.  
 963 2015. Cryoresistance in relation with lipid composition in sperm of common carp *Cyprinus*  
 964 *carpio*. (2015) Int. Meeting on Cryobiologies, Ostrava (Cz. Rep.). *Cryobiology* 71:537e573.
- 965 [115] Morrow S, Gosálvez J, López-Fernández C, Arroyo F, Holt WV, Guille MJ. Effects of freezing  
 966 and activation on membrane quality and DNA damage in *Xenopus tropicalis* and *Xenopus laevis*  
 967 spermatozoa. *Reprod Fertil Develop* 2014;29(8):1556-1566.
- 968 [116] Billard R, Cosson J, Noveiri SB, Pourkazemi M. Cryopreservation and short-term storage of  
 969 sturgeon sperm, a review. *Aquaculture* 2004;236(1-4):1-9 .
- 970 [117] Riley KL, Chesney EJ, Tiersch TR. Field collection, handling, and refrigerated storage of sperm of  
 971 red snapper and gray snapper. *North American J Aquacult* 2008;70:356-364.
- 972 [118] Gosálvez J, Holt WV, Johnston SD. Sperm DNA fragmentation and its role in wildlife  
 973 conservation. *Adv Exp Med Biol.* 2014;753:357-384.
- 974 [119] Pollock K, Gosálvez J, López-Fernández J, Johnston S. Amphibian sperm chromatin structure and  
 975 function and its relevance to sperm preservation. *J Herpet* 2018;52(4):487-492.
- 976 [120] Arregui L, Boveda P, Gosálvez J, Kouba AJ. Seasonal effect of sperm cryopreservation in  
 977 *Epidalea calamata* (Bufonidae). *Abstracts / Cryobiology* 2018;85:177.
- 978 [121] Brelsford A, Stock M, Betto-Collard C, Dubey D, Dufresnes C, Jourdan-Pineau H, Rodrigues N,  
 979 Savary R, Sermier R, Perrin N. Homologous sex chromosomes in three deeply divergent anuran  
 980 species. *Evolution* 2013;67-8:2434–2440.
- 981 [122] Duellman WE, Marion AB, Hedges SB. Phylogenetics, classification, and biogeography of the  
 982 treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 2016;4104:101-109.
- 983 [123] Zhang P, Liang D, Mao R.L, Hillis DM, Wake DB, Cannatella DC. Efficient sequencing of anuran  
 984 mtDNAs and a mitogenomic exploration of the phylogeny and evolution of frogs. *Mol. Biol.*  
 985 *Evol* 2013;30:1899-1915.

- 986 [124] Evens BJ. Genome evolution and speciation genetics of clawed frogs (*Xenopus* and *Silurana*).  
987       Frontiers Biosci 2008;4687-4706.
- 988 [125] Beesley SG, Costanzo JP, Lee RE. Cryopreservation of spermatozoa from freeze-tolerant and  
989       intolerant anurans. Cryobiology 1998;37:155-162.
- 990 [126] Flajshans M, Cosson J, Rodina M, Linhart O. The application of image cytometry for viability  
991       assessment of dual fluorescent stained spermatozoa of fish. A technical note. Cell Biol Intern  
992       2004;28/12:955-959.
- 993 [127] Linhart O, Rodina M, Cosson J. Cryopreservation of sperm in common carp *Cyprinus carpio*:  
994       sperm motility and hatching success of embryos. Cryobiology 2000;41:241-250.
- 995 [128] Silla AJ. Artificial fertilisation in a terrestrial toadlet *Pseudophryne guentheri*: effect of medium  
996       osmolality, sperm concentration and gamete storage. Reprod Fert Develop 2013;25(8):1134-  
997       1141
- 998 [129] Wilson-Leedy & Ingermann, 2015 Computer Assisted Sperm Analyzer.  
999       <https://imagej.nih.gov/ij/plugins/casa.html> [Accessed 9th December 2018]
- 1000 [130] Kouba A, Willis E, Vance C, Hasenstab S, Reichling S, Krebs J, Linhoff L, Snoza M, C.  
1001       Langhorne C, Germano J. Development of assisted reproduction technologies for the  
1002       Endangered Mississippi gopher frog (*Rana sevosa*) and sperm transfer for in vitro fertilization  
1003       Reprod Fert Devel 2011;24(1):170-170.
- 1004 [131] Cabrita E, Robles V, Herraes P. 2009. Methods in Reproductive Aquaculture: Marine and  
1005       Freshwater Species. CRC Press; 2008, p. 456.
- 1006 [132] Tiersch TR, Green CC. Cryopreservation in Aquatic Species, 2nd Edition. World Aquaculture  
1007       Society, Baton Rouge, Louisiana; 2011, p. 1-14.
- 1008 [133] Horvath A, Wayman WR, Dean JC, Urbanyi B, Tiersch TR, Mims SD, Johnson D, Jenkins JA.  
1009       Viability and fertilizing capacity of cryopreserved sperm from three North American acipenseriform  
1010       species: a retrospective study. J Appl Ichthyol 2012;24:443-449.
- 1011 [134] Hu E, Tiersch TR. Development of high-throughput cryopreservation for aquatic species. In:  
1012       Tiersch TR, Green CC, editors. Cryopreservation in Aquatic Species, 2nd Ed. World  
1013       Aquaculture Society, Baton Rouge, Louisiana; 2011, p. 995-1003.
- 1014 [135] Caffey RH, Tiersch TR. Economics and marketing of cryopreserved fish sperm. In In: Tiersch TR,  
1015       Green CC, editors. Cryopreservation in Aquatic Species, 2nd Ed. World Aquaculture Society,  
1016       Baton Rouge, Louisiana; 2011, p. 896-916

- 1017 [136] Schad K. Amphibian Population Management Guidelines. Amphibian Ark Amphibian Population  
1018 Management Workshop; December 10-11; San Diego, CA, USA. Amphibian Ark, 2007. 31 p.
- 1019 [137] Nori J, Villalobos F, Loyola R. Global priority areas for amphibian research. *J Biogeog*  
1020 2018;2588-2594
- 1021 [138] Browne RK, Janzen P, Bagaturov MF, van Houte DK. Amphibian Keeper Conservation Breeding  
1022 Programs. *J Zool Res* 2018;2(1):29-46.
- 1023
- 1024
- 1025

1026 **Legends for Figures**

1027

1028 Fig. 1. The rapid effects of selection in captive breeding are shown in these colour varieties of the  
1029 Chinese giant salamander (*Andrias davidianus*) which now has more than  $12 \times 10^6$  individuals in  
1030 aquaculture, mostly hybrids between 5 recently revealed cryptic species. *Image Robert Browne.*

1031

1032 Fig. 2. The evolutionary relationships between major amphibian clades, orders, families and species  
1033 with respect to major parameters informing an understanding of sperm storage and cryopreservation.  
1034 Studies may have included more than one species, and the assessment methods are for all studies of each  
1035 species (see Supplementary Table 2,3 for more detail). Blue = External fertilisation, Red = Internal  
1036 fertilisation.

1037

1038 Fig. 3. Hormonal stimulation and sperm collection in the Panamanian Golden Frog (*Atelopus zeteki*). A.  
1039 Intraperitoneal injection of GnRH; B. Spermic urine collection by abdominal massage; and C. Spermic  
1040 urine collection by gentle insertion of a catheter in the cloacae. *Image Gina Della Togna.*

1041

1042 Fig. 4. A. Collecting semen from a cryptobranchid, *Cryptobranchus alleganiensis*, through abdominal  
1043 massage at Nashville Zoo, USA. *Image Robert Browne*; and B) hormonally induced tiger salamander,  
1044 *Amystoma tigrinum*, semen from the cloaca at National Amphibian Conservation Center Detroit  
1045 Zoological Society, Detroit, USA. *Image Ruth Marcec.*

1046

1047 Fig. 5. A flow diagram of the different stages in the collection, preparation of cryosuspensions,  
1048 acclimation, and freezing of amphibian sperm as reported across various amphibian studies  
1049 (Supplementary Table 2,3). Temperatures in °C. *Neurergus kaiseri*. *Image Richard Bartlett*  
1050 [http://news.mongabay.com/2010/0208-hance\\_luristannewt.html](http://news.mongabay.com/2010/0208-hance_luristannewt.html)

1051

1052 Fig. 6. The phylogenetic relationship between four anuran families, with the Pipidae diverging from the  
1053 others ~210 mya, the Ranidae from the Hylidae and Bufonidae ~170 mya, and the Hylidae and  
1054 Bufonidae diverging ~70 mya (Adapted from Brelsford et al. [119]).

1055

1056 Fig. 7. A. *Ambystoma mexicanum* sperm stained with eosin nigrosin, 40× magnification. B. *Ambystoma*  
1057 spermatophore stained with Trypan Blue. Glycoproteins on the spermatophore highly stain but the  
1058 sperm package is intact to the vital stain and the enclosed sperm do not stain, and C. Hoesch staining is  
1059 used to test acrosomal integrity. *Image Manuel Gonzalez.*

1060

1061 Fig. 8. Flow chart of suggested Proof of Concept study to complete the life cycle of a threatened species  
1062 using cryopreserved sperm in a CBP. Warm temperatures in brown, and cool and colder temperatures in  
1063 blue. *Image Robert Browne.*

1064

1065 Fig. 9. Global diversity of amphibian species, the distribution of threatened amphibians, average income,  
1066 and the locations of research for the collection and storage of amphibian sperm for the sustainable  
1067 management of amphibian biodiversity. Anurans (black circles) and Caudata (Yellow circles). Both  
1068 Anura and Caudata are found in Australia, but only Anura are native. The size of circles roughly  
1069 approximates research on sperm storage.

1070

1071

1072

1073