

DISEASE NOTE

FIRST REPORT OF BACTERIAL LEAF SPOT CAUSED BY *PSEUDOMONAS SYRINGAE* pv. *PORRI* ON ONION IN IRAN

**Y. Moradi Amirabad, G. Khodakaramian
and H. Ramezani Zalandi**

*Department of Plant Protection, College of Agriculture,
Bu-Ali Sina University, Hamadan, Iran*

In October 2016, bacterial leaf spots on onion (*Allium cepa* L.) were observed in fields with rainy irrigation of Eghlid, Fars province in Iran with the rate of occurrence varying from 90 to 100% of the plants. Symptoms on leaves comprised leaf blight and water soaked, irregular, white and brown spots on the leaf surface. Bacterial isolates were obtained on nutrient agar (NA) from leaf spot and blight lesions that were surface disinfected in 70% ethanol for 45 s. The isolates were fluorescent on King's B agar, gram-negative and strictly aerobic. All isolates belonged to *P. syringae* (LOPAT) group Ia (+, -, -, +) (Lelliott *et al.*, 1996). Isolates were deposited in the Culture Collection of the Bu-Ali Sina University of Iran. DNA was extracted from the representative isolate (YMAO17) by the alkali lysis method and the 16S rRNA, *gyrB* and *rpoD* regions were partially sequenced using fD1/rP2 (Weisburg *et al.*, 1991), *gyrB*-Fpsc/*gyrB*-Rpsc, and *rpoD*-Fpc/*rpoD*-Rpsc (Sarkar and Guttman, 2004) primers, respectively. PCR product were sequenced and their sequences were aligned and compared with those deposited in GenBank. Sequences of the 16S rRNA (Accession No. KY684039), *gyrB* (KY684037) and *rpoD* (KY684038) of a representative isolate showed 100% identity with those of *Pseudomonas syringae* pv. *porri*. To confirm Koch's postulates, leaf spot symptoms were reproduced on onion leaves inoculated with bacterial suspensions (1×10^6 CFU/ml) and the same bacterial isolate was reisolated from artificially infected leaves as confirmed by LOPAT. To our knowledge, this is the first report of bacterial leaf spot of onion caused by *P. syringae* pv. *porri* in Iran.

Lelliott R.A., Billing E., Hayward A.C., 1966. A determinative scheme for the fluorescent plant pathogenic Pseudomonads. *Journal of Applied Microbiology* **29**: 470-489.

Sarkar S.F., Guttman D.S., 2004. Evolution of the core genome of *Pseudomonas syringae*, a highly clonal, endemic plant pathogen. *Applied and Environmental Microbiology* **70**: 1999-2012.

Weisburg W.G., Barns S.M., Pelletier D.A., Lane D.J., 1991. 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* **173**: 697-703.

Corresponding author: G. Khodakaramian
E-mail: khodakaramian@yahoo.com

Received March 16, 2017
Accepted March 31, 2017

DISEASE NOTE

FIRST REPORT OF *RASPBERRY BUSHY DWARF VIRUS* INFECTING RASPBERRY IN ARGENTINA

**A. Dal Zotto¹, A. Cardozo², D. Cabrera Mederos^{1,3},
C. Nome¹, F. Giolitti¹ and C. Cobelo²**

¹*Instituto de Patología Vegetal (IPAVE), Centro de Investigaciones Agropecuarias (CIAP), Instituto Nacional de Tecnología Agropecuaria (INTA), Av. 11 de Septiembre 4755, X5020ICA Córdoba, Argentina*

²*Agencia de Extensión Rural (AER) El Bolsón, INTA, Río Negro, Argentina*

³*Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Godoy Cruz 2290, C1425FBQ Buenos Aires, Argentina*

In 2014-2015, raspberry (*Rubus idaeus*) cv. Autumn bliss plants with dwarfism and foliar chlorosis were observed in commercial fields in parallel 42° in the Patagonia region of Argentina. Symptoms were similar to those observed in raspberry infected by *Raspberry bushy dwarf virus* (RBDV) in the USA (Ellis *et al.*, 2005; Martin *et al.*, 2013). A survey was conducted to determine if RBDV was also present in Argentinian raspberry fields. Transmission electron microscope observations of leaf dip preparations revealed the presence of spherical particles of 30nm, resembling those of virus members of the genus *Ideaovirus*. RBDV was detected in 88 of the 130 plants (68%) tested using double-antibody sandwich ELISA with a specific polyclonal antiserum (Bioreba AG, Switzerland). RNA was extracted with Spectrum™ Plant Total RNA Kit (Sigma-Aldrich, USA) and tested by RT-PCR using a pair of specific primers to the capsid protein gene (Wang *et al.*, 2008). Expected fragments of 825 bp were obtained and custom-sequenced in both directions (Macrogen, Korea). Nucleotide BLAST analysis of the complete sequence obtained (GenBank accession No. KY308191) showed 97% identity at the nucleotide level with virus isolates from Slovenia, Belarus, Switzerland and Japan (EU796088, FR687356, FR687358, AB948216, respectively). To our knowledge, this is the first report of RBDV infecting raspberry in Argentina. Further research is needed to determine the distribution of this virus in the country and its effects on raspberry plant production.

This work was supported in part by the PNFRU 1105073 and PNPV 1135022 projects of INTA. The authors thank Aylene Gonzalez for the collaboration in raspberry sampling.

Ellis M., Martin R.R., Wright S.R., 2005. First report of *Raspberry bushy dwarf virus* in Ohio. *Plant Health Progress* PHP-2005-0510-01-HN.

Martin R., MacFarlane S., Sabanadzovic S., Quito D., Poudel B., Tzanetakis L., 2013. Viruses and Virus Diseases of Rubus. *Plant Disease* **97**: 168-182.

Wang Q., Cuellar W., Rajamäki M.L., Hirata Y., Valkonen J.P., 2008. Combined thermotherapy and cryotherapy for efficient virus eradication: relation of virus distribution, subcellular changes, cell survival and viral RNA degradation in shoot tips. *Molecular Plant Pathology* **9**: 237-250.

Corresponding author: A. Dal Zotto
E-mail: dalzotto.angelica@inta.gob.ar

Received March 4, 2017
Accepted April 4, 2017