



Molecular Confirmation of the Relationship between *Candida guilliermondii* Fks1p Naturally Occurring Amino Acid Substitutions and Its Intrinsic Reduced Echinocandin Susceptibility

Catiana Dudiuk,^{a,b} Daiana Macedo,^a Florencia Leonardelli,^{a,b} Laura Theill,^a Matias S. Cabeza,^{a,b} Soledad Gamarra,^a Guillermo Garcia-Effron^{a,b}

Laboratorio de Micología y Diagnóstico Molecular, Cátedra de Parasitología y Micología, Facultad de Bioquímica y Ciencias Biológicas, Universidad Nacional del Litoral, Santa Fe, Argentina^a; Consejo Nacional de Investigaciones Científicas y Tecnológicas, Centro Científico Tecnológico Santa Fe, Santa Fe, Argentina^b

ABSTRACT *Candida guilliermondii* shows intrinsic reduced echinocandin susceptibility. It harbors two polymorphisms (L633M and T634A) in the Fks1p hot spot 1 region. Our objective was to confirm that the reduced echinocandin susceptibility of *C. guilliermondii* is due to those naturally occurring substitutions. We constructed a *Saccharomyces cerevisiae* mutant in which a region of the *FKS1* gene (including hot spot 1) was replaced with that from *C. guilliermondii*. The chimeric mutants showed 32-fold increases in echinocandin MIC values, confirming the hypothesis.

KEYWORDS *Candida guilliermondii*, FKS mutations, echinocandin resistance, molecular mechanism

Echinocandins are the treatment of choice for candidemia, since most *Candida* spp. are susceptible to these drugs (1–4). However, some highly prevalent *Candida* spp., such as species of the *Candida parapsilosis* complex and *Candida guilliermondii*, show natural reduced echinocandin susceptibility (RES) (5–7). Echinocandin drugs interact with the Fksp subunits of the β -1,3-glucan synthase complex, inhibiting the biosynthesis of the principal cell wall glucan. Clinical resistance was linked to hot spot mutations in Fksp (5). It was molecularly confirmed that the *C. parapsilosis sensu lato* RES phenotype is due to a naturally occurring substitution (P660A) in the hot spot 1 region of Fks1p (8). *C. guilliermondii* harbors two polymorphisms (L633M and T634A) in the same Fks1p region; it was assumed that the RES in this species is attributable to these amino acid substitutions, but the hypothesis was never confirmed (5).

The objective of this work was to verify that the intrinsic RES of *C. guilliermondii* is due to the naturally occurring amino acid changes in the hot spot 1 region of Fks1p. To reach this goal, we constructed a *Saccharomyces cerevisiae* strain with a hybrid *FKS1* gene. A region of the *FKS1* gene that includes hot spot 1 was replaced with that from *C. guilliermondii*. We then evaluated the echinocandin susceptibility.

S. cerevisiae BY4742 was used as the parental strain to obtain *S. cerevisiae* LMDM 537 (BY4742 *FKS1* deletant in which codons 453 to 649 were replaced with *URA3*, i.e., *fks1* Δ 453–649::*URA3*) (8). *C. guilliermondii* ATCC 6260 (*Meyerozyma guilliermondii*) was used to obtain the 695-nucleotide (nt) region of the *FKS1* gene, which was fused to generate the chimeric *FKS1* gene (*S. cerevisiae*-*C. guilliermondii*). DNA was extracted by phenol-based extraction (9). Caspofungin (CSF) and anidulafungin (ANF) susceptibility tests were performed following CLSI guidelines (10–12), with modifications. For *S.*

Received 14 December 2016 Returned for modification 25 January 2017 Accepted 18 February 2017

Accepted manuscript posted online 27 February 2017

Citation Dudiuk C, Macedo D, Leonardelli F, Theill L, Cabeza MS, Gamarra S, Garcia-Effron G. 2017. Molecular confirmation of the relationship between *Candida guilliermondii* Fks1p naturally occurring amino acid substitutions and its intrinsic reduced echinocandin susceptibility. Antimicrob Agents Chemother 61:e02644-16. <https://doi.org/10.1128/AAC.02644-16>.

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Guillermo Garcia-Effron, ggarcia@unl.edu.ar.




FIG 1 (A) Clustal alignment of the deduced Fks1p sequences of *S. cerevisiae* BY4742, *S. cerevisiae* harboring the chimeric *FKS1*, and *C. guilliermondii* ATCC 6260. Black lines show where the *C. guilliermondii* *FKS1* fragment was fused with the *S. cerevisiae* gene. The black box shows the *FKS1* hot spot 1 region. (B) PCR confirmation of the *FKS1* reconstitution. Lane 1, markers; lanes 2 to 7, 872-nt PCR fragment of the chimeric *FKS1* gene obtained by using the *FKS1*-305F and *Cg/h-R* primers and DNA isolated from uracil-auxotrophic and CSF- and FK506-resistant *S. cerevisiae* transformants. (C, D, and E) Diffusion susceptibility testing using caspofungin disks, following CLSI guidelines (11), for *S. cerevisiae* BY4742 (C), *S. cerevisiae* harboring the chimeric *FKS1* (D), and *C. guilliermondii* ATCC 6260 (E).

cerevisiae susceptibility evaluations, RPMI 1640 broth and Mueller-Hinton 2% glucose-methyl blue agar were replaced by YPD broth (1% yeast extract, 2% peptone, 2% glucose) and YPD agar (YPD broth plus 1.5% agar), respectively. Also, the incubation temperature was changed to 30°C. These modifications were necessary since *S. cerevisiae* strains did not grow properly with the CLSI-proposed culture media and temperature (8).

A two-step method for PCR-based *FKS1* mutagenesis was employed to generate a *S. cerevisiae* *FKS1* chimeric mutant (8, 13, 14). Briefly, the first step consisted of partial deletion of the *FKS1* gene of *S. cerevisiae* BY4742 using an *URA3* cassette. The deletion included the Fks1p hot spot 1 region (amino acid residues 453 to 649). This disruption leads to FK506 (tacrolimus) and echinocandin hypersensitivity (8, 15). The transformation was performed and the deletion was confirmed using methodologies described previously (8, 16). The second step of the method was the replacement of the partially deleted *S. cerevisiae* *FKS1* gene (*fks1*Δ453–649::*URA3*) by a construction that included a 695-nt portion of the *C. guilliermondii* *FKS1* gene (nt 1591 to nt 2286 of the sequence reported in GenBank under accession number [XM_001487741.1](http://www.ncbi.nlm.nih.gov/nuccore/XM_001487741.1)) surrounded by *S. cerevisiae* *FKS1* regions from nt 1123 to nt 1611 and from nt 2308 to nt 2548 (*S. cerevisiae* *FKS1* sequence reported in GenBank under accession number [U12893.1](http://www.ncbi.nlm.nih.gov/nuccore/U12893.1)). This construction was designed to yield, after homologous recombination, a functional *FKS1* gene encoding a chimeric Fks1p (*S. cerevisiae* *fks1*Δ537–762::*C. guilliermondii* *fks1* 531–762) (Fig. 1). The transformation vector was obtained by four PCRs, including amplification of the *C. guilliermondii* *FKS1* fragment (695 nt), the 488-nt portion of the 5' region of *S. cerevisiae* *FKS1*, and the 240 nt of the 3' fragment of *S. cerevisiae* *FKS1* using the primer pairs Fus5-*FKS1*-Sc-CguiF/Fus3-*FKS1*-Sc-CguiR, Fus5-*FKS1*-Sc-CguiR/*FKS1*-375, and Fus3-*FKS1*-Sc-CguiF/LMDM85, respectively. The last PCR was designed to fuse the three fragments using *FKS1*-375 and LMDM85 primers (see Table 1 for primer details). PCRs were performed in a 25-μl volume, following the manufacturer's instructions for the Pegasus DNA polymerase (PBL, Buenos Aires, Argentina), in an Applied

TABLE 1 Primers used in this study

Primer	Target organism	Orientation (5'→3')	Sequence (5'→3')	Purpose	Reference
FKS1-453-URA3F	<i>S. cerevisiae</i>	Sense	AAAGAGACCCGCTACTTGGTTACATTTGGTCACCAACTTCAGAGTG CACCATACCACAGCT	Generation of <i>S. cerevisiae</i> FKS1-deleted mutant	8
FKS1-649-URA3R	<i>S. cerevisiae</i>	Antisense	GTATTCACCTGTACACCTCATTGCAGTGGTGGACAAAAATTGGTAT TTCACACCCGATAGG	Generation of <i>S. cerevisiae</i> FKS1-deleted mutant	8
URA3IR	<i>S. cerevisiae</i>	Antisense	TGCCTTTAGCGGCTTAACTG	Confirmation of <i>S. cerevisiae</i> FKS1 deletion	8
FKS1-375	<i>S. cerevisiae</i>	Sense	GGTCGTTTTGTCAAGCGTGA	Confirmation of <i>S. cerevisiae</i> FKS1 deletion, chimeric FKS1 construction, and <i>S. cerevisiae</i> FKS1 reconstruction	8
FKS1-707R	<i>S. cerevisiae</i>	Antisense	ATTTCCCAACAGAGAAAAATGG	<i>S. cerevisiae</i> FKS1 reconstruction	8
LMDM85	<i>S. cerevisiae</i>	Antisense	CGTAGTGAGGAGTCAATACTGTG	Chimeric FKS1 construction	This study
Fus5-FKS1-Sc-CguiF	<i>S. cerevisiae</i> and <i>C. guilliermondii</i>	Sense	CGTAGATTCTGGTTTTATGCATCATCTTCGTGGTTAACTTGGCCCC	Chimeric FKS1 construction	This study
Fus5-FKS1-Sc-CguiR	<i>S. cerevisiae</i> and <i>C. guilliermondii</i>	Antisense	GGGGCCCAAGTTAACCCACCGAAGATGATGCATAAAAACCAGAAATCTACG	Chimeric FKS1 construction	This study
Fus3-FKS1-Sc-CguiF	<i>S. cerevisiae</i> and <i>C. guilliermondii</i>	Sense	GTACAGAGAACATTTGTTGGCTATTGACCATGTACAAAAAATTACTATATC	Chimeric FKS1 construction	This study
Fus3-FKS1-Sc-CguiR	<i>S. cerevisiae</i> and <i>C. guilliermondii</i>	Antisense	GATATAGTAATTTTTGTACATGGTCAATAGCCAACAATAATGTTCTCTGTAC	Chimeric FKS1 construction	This study
FKS1-305F	<i>S. cerevisiae</i>	Sense	CCCTGGAAAGAGTTCGTCATATC	Confirmation of FKS1 chimeric reconstitution	This study
Cg/h-R	<i>C. guilliermondii</i>	Antisense	CAAAACCCACCAAGGCATAAC	Confirmation of FKS1 chimeric reconstitution	This study

Biosystems thermocycler (Tecolab-AB, Buenos Aires, Argentina). The construction was cloned into a PGEM-T Easy vector (Promega; Biodynamics, Buenos Aires, Argentina) to yield the plasmid LMDM-P113. *S. cerevisiae* LMDM 537 (BY4742 *fks1*Δ453–649::*URA3*) was transformed with 1 μg of the LMDM-P113 plasmid, following the protocol (lithium acetate) described by Geitz et al. (17). Transformants were selected on FK506 plates and screened for the loss of *URA3* as described previously (8). The RES phenotype was screened for using YPD agar with 0.25 μg/ml CSF. Seven chimeric mutants showing the expected phenotype, i.e., uracil auxotrophic, FK506 resistant, and able to grow in the presence of 0.25 μg/ml CSF, were obtained. The *FKS1* reconstitution was assessed in all seven mutants by PCR using the primers FKS1-305F and Cg/h-R. These primers hybridize to the *S. cerevisiae FKS1* gene (around 900 nt upstream of the start codon) and *C. guilliermondii FKS* (near hot spot 1). The expected PCR band of 872 nt would be obtained only if the chimeric *FKS1* gene was present (Fig. 1). The *FKS1* reconstitution was confirmed by sequencing. A revertant control strain (LMDM 537R) was obtained by transforming a 1,000-bp PCR fragment into the *S. cerevisiae* LMDM 537 strain (17). This fragment was amplified using *S. cerevisiae* BY4742 DNA and the primers FKS1-375 and FKS1-707R (Table 1). The amplified fragment included the *S. cerevisiae FKS1* hot spot 1 region. Homologous recombination produced a strain with a functional *FKS1* that was FK506 resistant.

Parental and revertant *S. cerevisiae* strains (BY4742 and LMDM 537R, respectively) showed very low CSF and ANF MIC values (0.008 μg/ml), while the seven chimeric strains showed 32-fold higher MICs for both echinocandins (0.25 μg/ml). These values mimicked the *C. guilliermondii* ATCC 6260 RES phenotype (MICs of 0.25 μg/ml).

C. guilliermondii causes 1.4% of all *Candida* infections worldwide but is more prevalent in South America (>4%) (6, 18–20). Despite the RES phenotype, most of these infections are treated with echinocandins, since this yeast is also less susceptible to fluconazole than other *Candida* spp. (21). Most of these infections respond well to echinocandin therapy. However, *C. guilliermondii* breakthrough infections during caspofungin therapy have been reported (22). Under such circumstances, there is a concern that the use of echinocandins would contribute to an increased prevalence of this species.

It is well known that echinocandin clinical resistance is linked to *FKS* hot spot substitutions (5). The 32-fold increases in echinocandins MICs observed for the seven chimeric *S. cerevisiae* mutants support the assumption that L633M and T634A polymorphisms at *C. guilliermondii Fks1p* hot spot 1 are responsible for the observed RES in this species.

ACKNOWLEDGMENTS

This study was supported in part by a CONICET grant (grant PIP 2011-331) and an Universidad Nacional del Litoral CAI+D grant to G.G.-E. C.D., M.S.C., and F.L. have fellowships from CONICET, and D.M. has a fellowship from FONCYT.

REFERENCES

- Cornely OA, Vazquez J, De Waele J, Betts R, Rotstein C, Nucci M, Pappas PG, Ullmann AJ. 2014. Efficacy of micafungin in invasive candidiasis caused by common *Candida* species with special emphasis on non-*albicans* *Candida* species. *Mycoses* 57:79–89. <https://doi.org/10.1111/myc.12104>.
- Reboli AC, Rotstein C, Pappas PG, Chapman SW, Kett DH, Kumar D, Betts R, Wible M, Goldstein BP, Schranz J, Krause DS, Walsh TJ. 2007. Anidulafungin versus fluconazole for invasive candidiasis. *N Engl J Med* 356:2472–2482. <https://doi.org/10.1056/NEJMoa066906>.
- Reboli AC, Shorr AF, Rotstein C, Pappas PG, Kett DH, Schlamm HT, Reisman AL, Biswas P, Walsh TJ. 2011. Anidulafungin compared with fluconazole for treatment of candidemia and other forms of invasive candidiasis caused by *Candida albicans*: a multivariate analysis of factors associated with improved outcome. *BMC Infect Dis* 11:261. <https://doi.org/10.1186/1471-2334-11-261>.
- Vazquez J, Reboli AC, Pappas PG, Patterson TF, Reinhardt J, Chin-Hong P, Tobin E, Kett DH, Biswas P, Swanson R. 2014. Evaluation of an early step-down strategy from intravenous anidulafungin to oral azole therapy for the treatment of candidemia and other forms of invasive candidiasis: results from an open-label trial. *BMC Infect Dis* 14:97. <https://doi.org/10.1186/1471-2334-14-97>.
- Perlin DS. 2015. Echinocandin resistance in *Candida*. *Clin Infect Dis* 61(Suppl 6):S612–S617. <https://doi.org/10.1093/cid/civ791>.
- Pfaller MA, Boyken L, Hollis RJ, Kroeger J, Messer SA, Tendolkar S, Diekema DJ. 2008. In vitro susceptibility of invasive isolates of *Candida* spp. to anidulafungin, caspofungin, and micafungin: six years of global surveillance. *J Clin Microbiol* 46:150–156. <https://doi.org/10.1128/JCM.01901-07>.
- Pfaller MA, Boyken L, Hollis RJ, Kroeger J, Messer SA, Tendolkar S, Jones RN, Turnidge J, Diekema DJ. 2010. Wild-type MIC distributions and epidemiological cutoff values for the echinocandins and *Candida* spp. *J Clin Microbiol* 48:52–56. <https://doi.org/10.1128/JCM.01590-09>.

8. Garcia-Effron G, Katiyar SK, Park S, Edlind TD, Perlin DS. 2008. A naturally occurring proline-to-alanine amino acid change in Fks1p in *Candida parapsilosis*, *Candida orthopsilosis*, and *Candida metapsilosis* accounts for reduced echinocandin susceptibility. *Antimicrob Agents Chemother* 52:2305–2312. <https://doi.org/10.1128/AAC.00262-08>.
9. Sambrook J, Fritsch EF, Maniatis T. 1998. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
10. Clinical and Laboratory Standards Institute. 2008. Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard—3rd ed. CLSI document M27-A3. Clinical and Laboratory Standards Institute, Wayne, PA.
11. Clinical and Laboratory Standards Institute. 2009. Method for antifungal disk diffusion susceptibility testing of yeast; approved guideline—2nd ed. CLSI document M44-A2. Clinical and Laboratory Standards Institute, Wayne, PA.
12. Clinical and Laboratory Standards Institute. 2012. Reference method for broth dilution antifungal susceptibility testing of yeast; 4th informational supplement. CLSI document M27-S4. Clinical and Laboratory Standards Institute, Wayne, PA.
13. Johnson ME, Katiyar SK, Edlind TD. 2011. New Fks hot spot for acquired echinocandin resistance in *Saccharomyces cerevisiae* and its contribution to intrinsic resistance of *Scedosporium* species. *Antimicrob Agents Chemother* 55:3774–3781. <https://doi.org/10.1128/AAC.01811-10>.
14. Katiyar SK, Edlind TD. 2009. Role for Fks1 in the intrinsic echinocandin resistance of *Fusarium solani* as evidenced by hybrid expression in *Saccharomyces cerevisiae*. *Antimicrob Agents Chemother* 53:1772–1778. <https://doi.org/10.1128/AAC.00020-09>.
15. Douglas CM, Foor F, Marrinan JA, Morin N, Nielsen JB, Dahl AM, Mazur P, Baginsky W, Li W, el-Sherbeini M. 1994. The *Saccharomyces cerevisiae* FKS1 (ETG1) gene encodes an integral membrane protein which is a subunit of 1,3-beta-D-glucan synthase. *Proc Natl Acad Sci U S A* 91:12907–12911. <https://doi.org/10.1073/pnas.91.26.12907>.
16. Edlind TD, Henry KW, Vermitsky JP, Edlind MP, Raj S, Katiyar SK. 2005. Promoter-dependent disruption of genes: simple, rapid, and specific PCR-based method with application to three different yeast. *Curr Genet* 48:117–125. <https://doi.org/10.1007/s00294-005-0008-3>.
17. Gietz RD, Schiestl RH, Willems AR, Woods RA. 1995. Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. *Yeast* 11:355–360. <https://doi.org/10.1002/yea.320110408>.
18. Cordoba S, Vivot W, Bosco-Borgeat ME, Taverna C, Szusz W, Murisengo O, Isla G, Davel G. 2011. Species distribution and susceptibility profile of yeasts isolated from blood cultures: results of a multicenter active laboratory-based surveillance study in Argentina. *Rev Argent Microbiol* 43:176–185.
19. Pfaller M, Neofytos D, Diekema D, Azie N, Meier-Kriesche HU, Quan SP, Horn D. 2012. Epidemiology and outcomes of candidemia in 3648 patients: data from the Prospective Antifungal Therapy (PATH Alliance®) registry, 2004–2008. *Diagn Microbiol Infect Dis* 74:323–331. <https://doi.org/10.1016/j.diagmicrobio.2012.10.003>.
20. Pfaller MA, Diekema DJ, Mendez M, Kibbler C, Erzsebet P, Chang SC, Gibbs DL, Newell VA. 2006. *Candida guilliermondii*, an opportunistic fungal pathogen with decreased susceptibility to fluconazole: geographic and temporal trends from the ARTEMIS DISK antifungal surveillance program. *J Clin Microbiol* 44:3551–3556. <https://doi.org/10.1128/JCM.00865-06>.
21. Pappas PG, Kauffman CA, Andes D, Benjamin DK, Jr, Calandra TF, Edwards JE, Jr, Filler SG, Fisher JF, Kullberg BJ, Ostrosky-Zeichner L, Reboli AC, Rex JH, Walsh TJ, Sobel JD. 2009. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. *Clin Infect Dis* 48:503–535. <https://doi.org/10.1086/596757>.
22. Kabbara N, Lacroix C, Peffault de Latour R, Socie G, Ghannoum M, Ribaud P. 2008. Breakthrough *C. parapsilosis* and *C. guilliermondii* blood stream infections in allogeneic hematopoietic stem cell transplant recipients receiving long-term caspofungin therapy. *Haematologica* 93:639–640. <https://doi.org/10.3324/haematol.11149>.