

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2016

CSF sTREM2: Marking the tipping point between preclinical AD and dementia?

Suzanne E. Schindler

Washington University School of Medicine in St. Louis

David M. Holtzman

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Schindler, Suzanne E. and Holtzman, David M., "CSF sTREM2: Marking the tipping point between preclinical AD and dementia?." *EMBO Molecular Medicine*.8,5. 437-438. (2016).
https://digitalcommons.wustl.edu/open_access_pubs/6623

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.



CSF sTREM2: marking the tipping point between preclinical AD and dementia?

Suzanne E Schindler^{1,2,3} & David M Holtzman^{1,2,3}

Biomarkers for Alzheimer's disease (AD) have improved our understanding of the temporal sequence of biological events that lead to AD dementia (Jack *et al*, 2013). AD is characterized neuropathologically by amyloid plaques comprised of the amyloid- β peptide and neurofibrillary tangles comprised of tau. Brain amyloid deposition, as evidenced by a decline in amyloid- β peptide 42 (A β 42) in the cerebrospinal fluid (CSF) or by binding of amyloid PET ligands, is thought to be a key initiating event in AD and begins many years prior to the onset of dementia. A rise in CSF tau and phosphorylated tau in the setting of A β deposition appears to reflect neurodegeneration and also begins years prior to the onset of dementia but after A β deposition has begun to accumulate. Individuals with "preclinical AD," that is, normal cognition but abnormal AD biomarkers, have a much higher risk for developing AD dementia but may remain cognitively normal for years (Vos *et al*, 2013). While deposition of amyloid and formation of tau tangles are necessary for AD to occur, it is likely that additional events involving inflammation or other processes contribute to crossing the tipping point from preclinical AD to AD dementia. Current efforts are aimed at defining the biomarker(s) that best predict the transition from cognitive normality to abnormality. A biomarker that is closely associated with the onset of cognitive decline could help us to understand the biological events that connect amyloid deposition and tangle formation to cognitive decline and could have significant

practical value in AD diagnosis and clinical trial design.

See also: M Suárez-Calvet *et al* (May 2016)

AD researchers have long suspected that inflammation plays an important role in the biology of AD, but genetic evidence was largely lacking until 2013, when two studies reported that rare, missense mutations in TREM2 (triggering receptor expressed on myeloid cells 2) increased risk for AD by about twofold to fourfold (Guerreiro *et al*, 2013; Jonsson *et al*, 2013). Since then the biology of TREM2, and the role of TREM2 in AD, has been a very active area of investigation (for review see Ulrich & Holtzman, 2016). TREM2 is a receptor that is expressed on phagocytic cells, including microglia, and can activate a number of different signaling cascades that modulate inflammatory cytokines and other processes. Mice that express mutant human APP and develop amyloid deposition have upregulation in TREM2 in regions with amyloid deposition (Frank *et al*, 2008). One report suggested that loss-of-function mutations in TREM2 decrease the phagocytic activity of microglia and reduce A β clearance (Kleinberger *et al*, 2014). Further, mice with brain amyloid deposition and TREM2 haploinsufficiency have decreased clustering of microglia around amyloid plaques (Ulrich *et al*, 2014). Although the details of how TREM2 modulates risk for AD remain to be clarified, it appears that TREM2 helps to mediate a protective inflammatory response to amyloid pathology. Thus, TREM2 may be a

link between amyloid pathology and complex downstream events including inflammation.

In their paper, Suárez-Calvet *et al* (2016) examined CSF levels of the soluble ectodomain of TREM2 (sTREM2) in individuals representing the spectrum of AD, from cognitively normal with no biomarker evidence of AD pathology to preclinical AD to mild cognitive impairment (MCI) to AD dementia. Interestingly, they found that levels of CSF sTREM2 were lowest in controls and preclinical AD, peaked in MCI-AD, and then declined slightly in AD dementia. The authors also found correlations between CSF sTREM2 and CSF A β 42, tau, and ptau. CSF sTREM2 levels correlated better with CSF tau and ptau levels than A β 42 levels, suggesting that elevated sTREM2 levels occur later in the course of the AD process. Two other groups also recently reported that CSF sTREM2 levels are elevated in AD cases compared to cognitively normal controls and that sTREM2 levels are correlated with CSF tau and ptau (Hesgrave *et al*, 2016; Piccio *et al*, 2016). However, these groups did not examine how sTREM2 levels changed across the AD spectrum. Although the findings of Suárez-Calvet *et al* (2016) are cross-sectional in nature and must be replicated in cohorts sampled and followed longitudinally, they show that CSF sTREM2 levels may increase at the onset of cognitive decline (see Fig 1).

If CSF sTREM2 levels are reproducibly associated with the onset of cognitive decline, this biomarker could have significant value in AD diagnostics and drug trials. Current clinical CSF diagnostics

1 Department of Neurology, Washington University School of Medicine, St. Louis, MO, USA. E-mail: holtzman@neuro.wustl.edu

2 Knight Alzheimer's Disease Research Center, Washington University School of Medicine, St. Louis, MO, USA

3 Hope Center for Neurological Disorders, Washington University School of Medicine, St. Louis, MO, USA

DOI 10.15252/emmm.201606245 | Published online 14 March 2016




Figure 1. Possible temporal ordering of biomarkers along the AD time course incorporating CSF sTREM2.

evaluate levels of A β 42, tau, and ptau. The combination of low CSF A β 42 and high tau/ptau is consistent with the presence of AD pathology, but it is not specific—it may occur either in patients with dementia due to AD or preclinical AD. Addition of sTREM2 to CSF testing may improve our ability to determine whether mild cognitive problems are due to AD or other etiologies. CSF sTREM2 levels could also be helpful in AD drug trials. Biomarkers are being used as endpoints as well as enrollment criteria in some clinical trials. If CSF sTREM2 is reliably associated with the onset of cognitive decline, reduced sTREM2 levels in drug-treated individuals with preclinical AD could indicate a favorable drug effect. Additionally, drugs for AD may be most effective when given at a particular stage of the AD process. Individuals who have abnormal biomarkers for amyloid, tau/ptau but normal CSF sTREM2 may be at an earlier stage of the AD process than individuals with abnormalities in all three biomarkers. If CSF sTREM2 further refines our ability to stage AD, this would allow us to evaluate which drugs are most appropriate for patients at various stages of the AD process. If additional studies confirm the results of Suárez-Calvet *et al* (2016), CSF sTREM2 levels may be an important biomarker that

represents some of the biological events that connect amyloid deposition and neurofibrillary tangle formation to cognitive decline. A biomarker that marks the transition from preclinical AD to dementia could have a major impact on our ability to understand, diagnose, and treat this formidable disease.

References

Frank S, Burbach GJ, Bonin M, Walter M, Streit W, Bechmann I, Deller T (2008) TREM2 is upregulated in amyloid plaque-associated microglia in aged APP23 transgenic mice. *Glia* 56: 1438–1447

Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogava E, Majounie E, Cruchaga C, Sassi C, Kauwe JS, Younkin S *et al* (2013) TREM2 variants in Alzheimer's disease. *N Eng J Med* 368: 117–127

Heslegrave A, Heywood W, Paterson R, Magdalino N, Svensson J, Johansson P, Ohrfelt A, Blennow K, Hardy J, Schott J *et al* (2016) Increased cerebrospinal fluid soluble TREM2 concentration in Alzheimer's disease. *Mol Neurodegener* 11: 3

Jack CR Jr, Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD *et al* (2013) Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of

dynamic biomarkers. *Lancet Neurol* 12: 207–216

Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, Bjornsson S, Huttenlocher J, Levey AI, Lah JJ *et al* (2013) Variant of TREM2 associated with the risk of Alzheimer's disease. *N Eng J Med* 368: 107–116

Kleinberger G, Yamanishi Y, Suarez-Calvet M, Czirri E, Lohmann E, Cuyvers E, Struyfs H, Pettkus N, Weninger-Weinzierl A, Mazaheri F *et al* (2014) TREM2 mutations implicated in neurodegeneration impair cell surface transport and phagocytosis. *Sci Transl Med* 6: 243ra286.

Piccio L, Deming Y, Del-Aguila JL, Ghezzi L, Holtzman DM, Fagan AM, Fenoglio C, Galimberti D, Borroni B, Cruchaga C (2016) Cerebrospinal fluid soluble TREM2 is higher in Alzheimer's disease and associated with mutation status. *Acta Neuropathol* doi:10.1007/s00401-016-1533-5

Suárez-Calvet M, Kleinberger G, Araque Caballero MA, Brendel M, Rominger A, Alcolea D, Fortea J, Lleó A, Blesa R, Gispert JD *et al* (2016) sTREM2 cerebrospinal fluid levels are a potential biomarker for microglia activity in early-stage Alzheimer's disease and associate with neuronal injury markers. *EMBO Mol Med* 8: 466–476

Ulrich JD, Finn MB, Wang Y, Shen A, Mahan TE, Jiang H, Stewart FR, Piccio L, Colonna M, Holtzman DM (2014) Altered microglial response to Abeta plaques in APPPS1-21 mice heterozygous for TREM2. *Mol Neurodegener* 9: 20

Ulrich JD, Holtzman DM (2016) TREM2 function in Alzheimer's disease and neurodegeneration. *ACS Chem Neurosci*. doi:10.1021/acscchemneuro.5b00313

Vos SJ, Xiong C, Visser PJ, Jasielec MS, Hassenstab J, Grant EA, Cairns NJ, Morris JC, Holtzman DM, Fagan AM (2013) Preclinical Alzheimer's disease and its outcome: a longitudinal cohort study. *Lancet Neurol* 12: 957–965



License: This is an open access article under the terms of the Creative Commons Attribution 4.0 License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.