

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2017

Declines in prevalence of adolescent substance use disorders and delinquent behaviors in the USA: A unitary trend?

R. A. Gruzca

Washington University School of Medicine in St. Louis

R. F. Krueger

University of Minnesota

Arpana Agrawal

Washington University School of Medicine in St. Louis

A. D. Plunk

Eastern Virginia Medical School

M. J. Krauss

Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Gruzca, R. A.; Krueger, R. F.; Agrawal, Arpana; Plunk, A. D.; Krauss, M. J.; Bongu, J.; Cavazos-Rehg, P. A.; and Bierut, L. J., "Declines in prevalence of adolescent substance use disorders and delinquent behaviors in the USA: A unitary trend?" *Psychological Medicine*, 1-12. (2017).

https://digitalcommons.wustl.edu/open_access_pubs/6363

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Authors

R. A. Grucza, R. F. Krueger, Arpana Agrawal, A. D. Plunk, M. J. Krauss, J. Bongu, P. A. Cavazos-Rehg, and L. J. Bierut

Declines in prevalence of adolescent substance use disorders and delinquent behaviors in the USA: a unitary trend?

R. A. Gruzca^{1*}, R. F. Krueger², Arpana Agrawal¹, A. D. Plunk³, M. J. Krauss¹, J. Bongu¹,
P. A. Cavazos-Rehg¹ and L. J. Bierut¹

¹Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA

²Department of Psychology, University of Minnesota, Minneapolis, MN, USA

³Department of Pediatrics, Eastern Virginia Medical School, Norfolk, VA, USA

Background. Downward trends in a number of adolescent risk behaviors including violence, crime, and drug use have been observed in the USA in recent years. It is unknown whether these are separate trends or whether they might relate to a general reduction in propensity to engage in such behaviors. Our objectives were to quantify trends in substance use disorders (SUDs) and delinquent behaviors over the 2003–2014 period and to determine whether they might reflect a single trend in an Externalizing-like trait.

Methods. We analyzed data from 12 to 17 year old participants from the National Survey on Drug Use and Health, a representative survey of the household dwelling population of the USA, across the 2003–2014 period ($N=210\,599$). Outcomes included past-year prevalence of six categories of substance use disorder and six categories of delinquent behavior.

Results. Trend analysis suggested a net decline of 49% in mean number of SUDs and a 34% decline in delinquent behaviors over the 12-year period. Item Response Theory models were consistent with the interpretation that declines in each set of outcomes could be attributed to changes in mean levels of a latent, Externalizing-like trait.

Conclusions. Our findings suggest that declines in SUDs and some delinquent behaviors reflect a single trend related to an Externalizing-like trait. Identifying the factors contributing to this trend may facilitate continued improvement across a spectrum of adolescent risk behaviors.

Received 26 April 2017; Revised 5 September 2017; Accepted 12 September 2017

Key words: Epidemiology, juvenile delinquency, substance use disorder, structural models, trends.

Introduction

Downward trends in a number of adolescent health risk behaviors have been observed over the past 15 or more years in the USA. For example, arrest rates for both assault and theft dropped by 75% between 1992 and 2010, and this trend is consistent with those based on results from crime victimization surveys (Robers *et al.* 2010; White & Lauritsen, 2012; Finkelhor *et al.* 2014; Child Trends Data Bank, 2015; Morgan *et al.* 2015; Office of Juvenile Justice & Delinquency Prevention, 2015). Self-reported survey measurements also indicate declines in problem behaviors including bullying and fighting, binge drinking, cigarette smoking, use of most classes of illicit drugs, and early sexual

involvement (Abma *et al.* 2010; Finkelhor, 2013; Johnston *et al.* 2013; Perlus *et al.* 2014).

The phenomenon of reduced rates for a broad array of risk behaviors raises an important question: Have these changes resulted from separate trends across multiple domains of behavior, or are they better described by a single trend that involves predisposition to risk-taking behaviors more generally? The answer to this question has important implications. Separate trends suggest behavior-specific causes. For example, state policies implemented since the late 1990s may have led to a reduction in bullying and other types of violence (Hatzenbuehler *et al.* 2015), but there is no *a priori* reason to think that these policies would have direct effects on substance use behaviors or on non-violent crime. Similarly, policies adopted to restrict access to alcohol and tobacco by minors at the state and municipal levels in recent years have likely had their intended effects (Gruenewald, 2011; Farrelly *et al.* 2013; Gruzca *et al.* 2013). But any ‘spillover’ effects on violent behaviors

* Address for correspondence: R. A. Gruzca, Ph.D. Department of Psychiatry, Washington University School of Medicine, 660 South Euclid Avenue, Box 8134, St. Louis, Missouri 63110, USA.
(Email: gruczar@psychiatry.wustl.edu)

and other crime would presumably be indirect and smaller in magnitude.

In contrast, if we were to discover that these individual trends were manifestations of a more far-reaching underlying trend, we would think differently about the potential causes. Rather than asking why teenagers are smoking less or drinking less, etc., we might ask why adolescents seem less disposed toward risk behaviors more generally. Several lines of research suggest that much of variation in proclivity to engage in different problem behaviors stems from an underlying latent characteristic. For example, results from developmental studies based on Problem Behavior Theory suggested that adolescent risk behaviors including substance use, precocious sexual involvement, and delinquency is linked to an underlying behavioral syndrome that was subsequently labeled 'risk behavior syndrome' (Jessor & Jessor, 1977; Donovan & Jessor, 1985; Jessor, 1991). More recently, psychopathology-focused studies have suggested that conduct disorder, alcohol and drug use disorders, and impulsivity share common etiologies and represent a core 'externalizing' or 'disinhibition' factor (Young *et al.* 2000; McGue *et al.* 2001; Krueger *et al.* 2002; Kendler *et al.* 2003; Hicks *et al.* 2004; Dick *et al.* 2005; Krueger & South, 2009). These earlier lines of research characterized variation in risk behaviors within cohorts, but no research to date has examined whether the population-level mean values of these traits might change over time.

Much of the research on externalizing and related constructs has emerged from the behavior genetics literature and these latent factors have been shown to be highly heritable. However, this does not mean that externalizing-like traits are unmodifiable by the environment. Biometrical modeling studies suggest significant influence for environmental factors that are shared by siblings and those that are unique to the individual environment (see Burt, 2009 for a review and meta-analysis). There are also examples of specific environmental factors that may influence externalizing. For example, Hicks *et al.* (2009) showed that the heritability of externalizing was modified by the environment such that heritability was higher in the presence of multiple risk factors such as antisocial peer affiliations and stressful life events. This suggests an important role for the environment in modulating risk for multiple externalizing outcomes. Relatedly, Verona & Sachs-Ericsson (2005) showed that the transmission of externalizing from parent to offspring was mediated by physical and sexual abuse, again suggesting this highly heritable trait is substantially modifiable by the environment. Notably, several indicators suggest declines in child abuse and neglect in recent decades, including physical and sexual abuse (Board on Children, Youth, and Families, 2012). This provides

us with at least one example of a societal-level environmental change that could influence risk for multiple externalizing outcomes.

Given the possibility that environmental change can lead to reductions in multiple adverse outcomes, it is essential to know the degree to which observed reductions in externalizing (or 'problem') behaviors reflect a single, multi-faceted trend in these behaviors as a behavioral syndrome as opposed to multiple, separate but concurrent trends. Distinguishing between these possibilities requires multivariate analysis. It is not sufficient to merely examine whether these declines are occurring in parallel. Rather, we need to know whether a reduction in risk for any one outcome for a given individual corresponds to reductions in risk of comparable magnitude for other outcomes for that same individual.

Addressing these questions requires historical data, and no single series of US youth behavioral health surveys has measured all behaviors of interest. Therefore, this report focuses outcomes related to externalizing that have been regularly assessed in the National Survey on Drug Use and Health (NSDUH). Externalizing outcomes include substance use disorders (SUDs) and disruptive behaviors (Krueger, 1999; Krueger *et al.* 2002). The NSDUH is annually administered to a cross-section of the population (i.e. a new sample every year) and is the only US national survey that regularly assesses SUDs among adolescents. The NSDUH does not formally assess disruptive behavior disorders, but queries several delinquent behaviors that partially overlap with conduct disorder (Lahey, 2008). Our first objective was to describe trends in all outcomes using conventional univariate methods. We then employed Item Response Theory models (IRT) to first determine whether trends in various SUDs among adolescents could be attributed to a single trend in a latent trait or factor conferring liability to all SUD outcomes, then whether trends in different delinquent behaviors could similarly be attributed to a trend in an underlying latent trait for delinquency, and finally whether trends in both SUDs and delinquent behaviors were consistent with a trend in a single Externalizing-like trait. Our analyses were motivated by our previous work in which we suggested that trends in delinquency were related to the recent decline in the prevalence of marijuana use disorder among adolescents (Grucza *et al.* 2016). However, formal modeling of the co-occurrence between delinquent behaviors and marijuana use disorder was beyond the scope of that work. (Nor did that work examine other SUDs). The work described here constitutes the first comprehensive and multivariate examination of recent trends in adolescent SUDs and delinquent behaviors. Further, despite several decades of research indicating that a full understanding of

these behaviors requires multivariate analysis, we are unaware of prior studies that have used such methods to examine population-level trends in any set of adolescent risk behaviors.

Methods

Survey overview and sample

We utilized data from the adolescent sample (ages 12–17) of the NSDUH, a yearly survey of the non-institutionalized, civilian population of the USA, overseen by the Substance Abuse and Mental Health Services Administration (SAMHSA, 2012). The NSDUH utilizes household-based multistage probability sampling from all 50 states and the District of Columbia, and includes those living in group-quarters. Consistent sampling and recruitment methods have been employed since 2002, rendering the data comparable from year-to-year on most measures. Interviews are conducted in dwelling units; behavioral questions are administered by audio-computer assisted self-interview to maximize privacy and confidentiality. Detailed methods are available through SAMHSA (Substance Abuse and Mental Health Services Administration, 2015). Because of slight changes to items assessing delinquency in 2003, our analyses cover the period 2003 through 2014, the most recent year for which data was available. Weighted adolescent response rates for that period range 80–87% (SAMHSA, 2014). Public use NSDUH files were obtained from the Interuniversity Consortium for Social and Political Research (ICPSR, 2016). Annual sample sizes ranged from 13 409 to 18 518. After removing 2144 subjects with missing data, the final combined sample size was 210 599.

Outcome measures

Main outcomes were measures of past-year delinquent behaviors and SUDs. Frequencies of engaging in six delinquent behaviors were assessed: participation in a serious fight, involvement in a group fight, attacking a person with intent to injure, stealing an item worth \$50 or more, selling drugs, and handgun carrying. NSDUH SUD assessments for alcohol and eight classes of drugs—including prescription drugs used non-medically—are based on DSM-IV abuse and dependence criteria covering the past 12 months. We analyzed six SUD outcomes related to alcohol, nicotine, marijuana, prescription opioids, other non-prescription illicit drugs, and other (non-opioid) prescription drugs. The two ‘other’ categories were created because several of the specific SUD diagnoses were rare. The NSDUH does not assess DSM-IV nicotine dependence, but includes both the Fagerström test for Nicotine Dependence and the Nicotine

Dependence Syndrome Scale (Heatherton *et al.* 1991). Participants who met criteria using either of those measures were counted as nicotine dependent. For other substances, SUD was defined as meeting criteria for DSM-IV abuse or dependence.

Our primary outcomes were the mean counts of (i) past-year delinquent behaviors reported and (ii) SUD categories for which past-year diagnostic criteria were met. In order to derive summary statistics to describe the overall trends in these variables for the entire observation period—as opposed to year-to-year differences that might fluctuate over time—we used regression methods to model each variable as a function of year (described below). This also allowed for adjustment for demographic covariates that might also change over time. We also examined trends in individual delinquent behaviors and SUDs.

Demographic variables

Sex, age, race/ethnicity, population density (urban/rural status) and poverty status were used as stratification variables in descriptive analyses and covariates in adjusted trend analyses. Age was categorized into three groups: 12–13, 14–15 and 16–17. Race/ethnicity was recoded into six groups: White, Black, Hispanic, Asian, multiple reported races, and other. The population density variable was recoded to indicate whether or not the respondent lived in a core-based statistical area (CBSA) with a population of 10 000 or more, or whether they lived outside of a CBSA (labeled ‘non-rural’ and ‘rural’, respectively). A poverty variable, derived from the ratio of total family income to the federal poverty level, included the following categories: family incomes below the federal poverty threshold (FPT), below 200% FPT, and equal to or above 200% FPT.

Statistical analysis

Stata version 14 was used for descriptive statistics and regression analyses. For dichotomous outcomes, we modeled each dependent variable as a function of year using log-binomial regression. The exponent of the regression coefficient yields the risk ratio (RR) associated with year. For example an RR of 0.9 would correspond to a 10% reduction in risk per year. We report the average annual relative change in prevalence, which is equivalent to the average annual relative change in risk, calculated as $-100 \times (1-RR)$. For count variables (number of delinquent behaviors and number of SUDs), we proceeded in a similar manner except using negative binomial regression. In this case, the exponent of the regression coefficient yields the ‘rate ratio’ associated with year, which can be interpreted similarly; i.e. a ratio of 0.9 would mean a reduction in count of 10% per year. As with the dichotomous outcomes, we report the

average annual change, or $-100 \times (1 - \text{Rate Ratio})$. To account for the complex design of the surveys, all analyses were conducted using Stata algorithms that incorporate survey design variables and utilize robust variance estimation procedures.

Structural analyses: IRT modeling

IRT analyses were conducted to examine whether changes in each set of outcomes could be attributed to changes in underlying latent traits, which we label DQ (delinquency) and SUD, respectively. Development of 2-parameter logistic (2PL) IRT models for single-factor DQ and SUD models are described in the Supplementary Material; 2PL models yield estimates of item discrimination coefficients (a) and item difficulty (b). For the two single-factor models, we analyzed six delinquent behaviors and six SUDs, respectively, as indicators of the underlying unidimensional factor. We then examined measurement invariance (MI) of each unidimensional model across survey years. This was done by estimating a series of models: (1) a model in which the discrimination coefficients and difficulty parameters were allowed to vary across years—typically called a configural model; factor means and variances are fixed at 0 and 1, respectively, for model identification. (2) A model in which discrimination coefficients and difficulty parameters were held constant across years, but factor means and variances were estimated separately for each year—typically called a scalar model. (3) Model 2, with variances held constant (at 1) for each year but factor means estimated separately for each year. (4) Model 2 with means held constant and variances estimated separately for each year and (5) Model 2 with both factor means and variances held constant across years. Superiority of the scalar model (2) over the configural model (1) is evidence of MI and justifies constraining of the discrimination coefficients and item difficulty parameters to be constant over time, indicating that the relations between the manifest indicators and the latent factor means remain constant over time, such that changes in indicator values can be interpreted as changes in the distribution of the underlying latent traits rather than temporal differences in model properties (e.g., Eaton *et al.* 2012). Superiority of Model 3 would further indicate that the variance of the underlying factor remained constant and that changes in indicator values reflect changes in the mean levels of the latent trait. Models 4 and 5 were estimated to rule-out alternative hypotheses that changes stemmed from changes in variance only, or that the distribution of the latent factor remained relatively constant over time.

We compared models using the Bayesian Information Criterion (BIC; Schwarz, 1978). The BIC is derived from maximum likelihood estimation and is an indicator of

the likelihood that the model is correct based on goodness of fit and model parsimony (lower values indicate preferred models). To ensure our results were not dependent on a particular estimator or fit statistic, we also report the comparative fit index (CFI), which is a parsimony-weighted fit index derived from weighted-least squares estimation. The CFI ranges from 0 to 1, with higher values suggesting a better model (Cheung & Rensvold, 2002). All multivariate analyses were conducted in MPlus utilizing either the 'MLR' estimator (maximum-likelihood with robust standard errors) or the 'WLSMV' (weighted least squares with mean and variance adjusted chi-square statistic) estimator (Muthén & Muthén, 1998).

After establishing that the single-factor models for both DQ and SUD exhibited strong MI across years and that the variance of each factor was constant, we turned to the question of whether changes in mean values for all six SUDs and all six delinquent behaviors might be related to changes in mean values of a single higher-order externalizing factor (EXT). Ideally, we would estimate a hierarchical model, in which DQ and SUD are sub-factors of EXT, but this model is under-identified (see online Supplemental Material, Part II). Therefore, we modeled all indicators as a function of EXT and estimated the series of five invariance-testing models described above. Although this model exhibited a suboptimal fit compared with the single-factor models, we justified the use of a single-factor model by estimating a model in which the two latent DQ and SUD factors were correlated with each other and demonstrating that this correlation coefficient was relatively high and constant over time (see online Supplemental Material, Part II).

Results

Table 1 reports the number of participants in each demographic group and the mean numbers of delinquent behaviors and SUDs per 100 participants, overall and for each demographic group. The means of these quantities for the first and last year of the observation period are listed in the bottom two rows of the table, and a decline in each is readily apparent. Across the 2003–2014 period, the estimated mean number of delinquent behaviors per 100 persons was 52.5 (95% CI 52.0–53.1) and the mean number of SUDs was 13.5 per 100 (95% CI 13.2–13.8). Because our primary interest was in trends over time, we do not discuss the demographic distribution of these variables further, but interested readers may refer to Table 1 for details.

Results of epidemiological trend analyses are shown in Figs 1 and 2, and Table 2. Panels A and B of Fig. 1 plot the yearly prevalence estimates for each of the six delinquent behaviors and for each of the six SUD categories, respectively, while Fig. 2 shows the mean

Table 1. Estimates for mean number of six delinquent behaviors and mean number six categories of SUDs per 100 persons, by demographic group, for the 2003–2014 NSDUH adolescent samples

	Unweighted n	Weighted % of Sample	Mean delinquent behaviors (95% CI) [Weighted] ^a		Mean SUDs (95% CI) [Weighted] ^a	
Full sample	210 599	100.0	52.5	(52.0–53.1)	13.5	(13.2–13.8)
Sex						
Males	107 232	51.1	63.0	(62.2–63.8)	13.1	(12.7–13.5)
Females	103 367	48.9	41.7	(41.0–42.3)	13.9	(13.5–14.4)
Age						
12–13	66 984	32.1	48.0	(47.2–48.8)	2.7	(2.5–2.8)
14–15	71 207	34.0	55.3	(54.3–56.3)	12.2	(11.7–12.6)
16–17	72 408	33.9	54.0	(53.0–55.1)	25.2	(24.6–25.8)
Race/ethnicity						
White	125 312	58.4	47.5	(46.9–48.2)	15.7	(15.3–15.9)
Black	28 808	14.7	70.5	(69.0–72.1)	7.7	(7.2–8.4)
Hispanic	36 775	19.5	56.7	(55.2–58.1)	12.2	(11.4–12.9)
Asian	6687	4.3	31.4	(29.2–33.5)	4.2	(3.4–5.1)
Multiple Race	8854	2.2	63.2	(58.9–67.5)	16.1	(14.3–18.1)
Other ^b	4163	0.9	70.2	(64.7–75.8)	9.8	(8.9–10.7)
Population density						
Non-rural	191 289	91.0	52.2	(51.6–52.7)	13.2	(12.9–13.5)
Rural	19 310	9.0	56.3	(54.5–58.1)	17.2	(16.2–18.3)
Poverty						
Below FPT	40 692	19.5	67.0	(65.6–68.4)	14.5	(13.7–15.3)
1–2× FPT	47 799	22.0	59.2	(58.1–60.3)	14.4	(13.8–15.1)
>2× FPT	124 252	58.5	45.2	(44.6–45.9)	12.8	(12.5–13.2)
Year						
2003	18 067	8.36	62.6	(60.6–64.7)	17.1	(16.1–18.2)
2014	13 409	8.28	40.6	(39.0–42.2)	8.8	(8.0–9.6)

^a Per 100 people; range = 0–600.

^b Includes Native American, Native Hawaiian and other Pacific Islanders; included in main analyses, but not in stratified trend analyses due to small sample size.

FPT, federal poverty threshold.

number of delinquent behaviors and the mean number of SUDs per 100 persons. [Table 2](#) lists the average annual changes in each outcome; i.e. the average annual relative change in prevalence for dichotomous variables and the average annual relative change in means for count variables. These are derived from regression estimates of risk ratios and rate ratios, respectively, describing the association between the outcome variable and year. These parameters are related to the slopes of the trend lines in [Figs 1](#) and [2](#). The first column of [Table 2](#) lists the unadjusted estimates of average annual relative change while the second column lists the same parameters adjusted for demographic variables. (Although some trends deviated from the log-linear form, we opted not to introduce quadratic or higher order terms into the trend models so that we could summarize and compare trend magnitudes for all outcomes using the annual average percentage change.)

The top half of [Table 2](#) shows that, with the exception of handgun carrying, the prevalence of each delinquent

behavior underwent a significant decrease, with unadjusted rates of decline ranging from 3.0% to 5.0% annually; adjustment for demographics had little impact on these estimates. Based on these rates, overall declines in the prevalence of each behavior (except for handgun carrying) for the 2003–2014 period ranged from 29 to 44%. The annual average decline in the mean number of delinquent behaviors was 3.7%, which corresponds to an overall decline of 34%. The bottom half of [Table 2](#) summarizes changes in the past-year prevalence of the six SUD categories; all SUDs underwent significant and substantial decreases in prevalence, with unadjusted average annual reductions ranging from 2.5% for marijuana use disorder to 8.0% for nicotine dependence. These changes correspond to overall declines ranging from 25% to 60% for the 2003–2014 period. The average annual decline in mean number of SUDs per 100 participants was 6.0%, which corresponds to a net decline of 49%.

Results of analyses of structural relationships among individual delinquent behaviors and individual SUDs




Fig. 1. Prevalence by year for each of six delinquent behaviors (a), and each of six SUD categories (b). Trend lines represent fits to log-binomial models of each variable as a function of year. Error bars represent 95% confidence intervals. Y-axes are logarithmically scaled.




Fig. 2. Mean by year for an average number of delinquent behaviors and SUDs per person. Trend lines represent fits to log-binomial model. Error bars represent 95% confidence intervals. Y-axis is logarithmically scaled.

are described in the Supplementary Material and below. Development of the one-factor models for DQ and SUD are provided in the Supplementary Material, Part I with results in online Supplementary Tables S1–S3. Model fit statistics (BIC and CFI) from MI analyses of the single-factor models are shown in the first two sections of Table 3. In both cases, the models with parameters constrained to be equivalent across years (scalar models) yielded lower BIC and higher CFI values than the unconstrained models, and further constraining the variance to be constant across years resulted in further improvements in those parameters. These results indicate that both the DQ and SUD measurement models exhibit MI across survey years and that there was little change in the variance of the corresponding latent traits over time. Finally, we estimated the model in which all 12 behaviors linked by a single factor labeled EXT. This model also exhibited strong MI and constant factor variance across years. Final models are diagrammed in Fig. 3. Online Supplementary Material Part III describes estimation of the mean values the latent DQ, SUD, and EXT factors; these estimates are plotted in online Supplementary Fig. S2. The plot illustrates that mean values of all three factors declined by about 0.3–0.4 standard deviations during the period under study.

Discussion

Over the years 2003–2014, we estimate a 34% decline in the number of delinquent behaviors reported by

Table 2. Average annual relative change in the prevalence of each of six delinquent behaviors, each of six substance use disorders, and the mean numbers of each, as estimated from linear trend models

	Average annual reduction, % (95% CI)	
	Unadjusted ^a	Adjusted ^b
Delinquent behaviors		
Serious fight	-3.0 (-3.3 to -2.7)	-3.2 (-2.9 to -3.5)
Group fight	-5.0 (-5.4 to -4.5)	-5.2 (-4.8 to -5.7)
Attack to injure	-4.5 (-5.1 to -3.9)	-4.8 (-4.2 to -5.4)
Stealing item >\$50	-4.2 (-4.9 to -3.5)	-5.0 (-4.3 to -5.7)
Selling drugs	-4.0 (-4.8 to -3.2)	-4.7 (-3.8 to -5.6)
Hand gun carrying	+0.3 (-0.5 to 1.1)	+0.7 (-0.1 to 1.5)
Mean # of behaviors	-3.7 (-4.0 to -3.4)	-4.0 (-3.7 to -4.3)
Substance use disorders		
Alcohol	-6.7 (-6.1 to -7.4)	-6.9 (-6.2 to -7.5)
Nicotine	-8.0 (-7.2 to -8.8)	-7.8 (-7.1 to -8.6)
Marijuana	-2.5 (-1.6 to -3.4)	-3.3 (-2.4 to -4.2)
Analgesics	-6.1 (-4.4 to -7.8)	-6.5 (-4.7 to -8.2)
Other non-prescription Drugs	-7.0 (-5.5 to -8.4)	-7.6 (-6.1 to -9.1)
Other prescription drugs	-4.3 (-2.0 to -6.5)	-4.7 (-2.3 to -7.0)
Mean # of SUDs	-6.0 (-5.4 to -6.5)	-6.3 (-5.8 to -6.9)

^a Unadjusted analyses include year as a predictor variable, with no covariates.

^b Adjusted analyses include demographic covariates (age, sex, race/ethnicity, rural status, and household income), with categories defined as shown in Table 1.

Table 3. Comparisons of models to assess measurement invariance of the unidimensional DQ, SUD, and EXT models

	# Parameters	BIC	Difference	CFI	Difference
Delinquency model					
Unconstrained (ref)	144	1 660 689	-	0.954	-
Constrain a, b	34	1 660 051	-638	0.965	0.011
Constrain a, b, σ^2	23	1 659 969	-720	0.973	0.019
Constrain a, b, \bar{x}	23	1 660 397	-292	0.952	-0.002
Constrain a, b, \bar{x}, σ^2	12	1 661 023	+334	0.961	0.007
SUD model					
Unconstrained (ref)	144	1 268 727	-	0.985	-
Constrain a, b	34	1 268 001	-726	0.984	-0.001
Constrain a, b, σ^2	23	1 267 905	-822	0.987	0.002
Constrain a, b, \bar{x}	23	1 268 126	-601	0.970	-0.015
Constrain a, b, \bar{x}, σ^2	12	1 268 529	-198	0.974	-0.011
Externalizing model					
Unconstrained (ref)	288	1 874 060	-	0.916	-
Constrain a, b	46	1 872 550	-1510	0.938	0.022
Constrain a, b, σ^2	35	1 872 456	-1604	0.946	0.030
Constrain a, b, \bar{x}	35	1 873 226	-834	0.929	0.013
Constrain a, b, \bar{x}, σ^2	24	1 873 745	-315	0.939	0.023

Notes: a , item discrimination coefficient; b , threshold, \bar{x} , latent factor mean; σ^2 , latent factor variance.

12–17 year olds, and a 49% decline in the number of SUDs. Results of multivariate modeling analyses are consistent with the interpretation that these declines

reflect a trend in an underlying Externalizing-like trait rather than multiple trends in specific behaviors or specific types of SUD. In the first stage of modeling,




Fig. 3. Structural (IRT) models for (a) Delinquency, (b) Substance use disorders, (c) A unidimensional externalizing model in which all SUD and DQ indicators are joined to a single factor.

single-factor models for DQ and SUD both exhibited consistent measurement properties from year to year, supporting the interpretation that prevalence declines in each set of outcomes can be attributed to declines in mean levels of the hypothesized underlying traits. We then demonstrated that the latent DQ and SUD

traits were strongly correlated with each other ($R = 0.74$), and that the magnitude of that correlation was invariant over time. This justified estimation of a single-factor externalizing model summarizing both delinquent behaviors and SUDs. This model also exhibited consistent measurement properties and

constant variance over time suggesting that declines in all manifest variables are largely due to declines in mean levels of an Externalizing-like trait.

We do not discount the possibility that outcome-specific factors—such as alcohol or tobacco policies—may have had some influence on trends in individual outcomes. In fact, this may be particularly true in the case of gun-carrying, which is the only delinquent behavior that did not undergo an appreciable or even statistically significant decline—an observation that is consistent with the findings of at least one other report (Webster *et al.* 2014). What this means in terms of manifest behavior is that the prevalence of gun-carrying in the context of other delinquent behaviors likely declined whereas the prevalence of gun-carrying as a standalone behavior likely increased. This is explored further in the online Supplementary Material Part IV and online Supplementary Fig. S3. Nonetheless, our modeling results suggest that the bulk of change in prevalence in SUDs and delinquent behaviors can be attributed to a trend in a common factor.

The main implication of our findings is that the primary causal factors for the trends we observed probably influence individual characteristics, such as disinhibition or risk-preferences, rather than impacting risk for specific outcomes. Given the fairly sharp decline in the prevalence of behaviors we examined, the potential causes are likely to be environmental factors that have undergone relatively rapid changes in recent years. There are a large number of such factors, but at least two have been nominated by other investigators as potential causes of reductions in delinquency and other risk behaviors that could be considered as part of the externalizing spectrum. Several investigators have suggested that reductions in childhood lead exposure may be linked to declining rates of delinquency, unwed pregnancy, low IQ, violent crime, and other problems (e.g. Nevin, 2000; Dietrich *et al.* 2001; Stretesky & Lynch, 2001; Reyes, 2007; Lane *et al.* 2008). Environmental lead levels dropped precipitously during the 1970s and 1980s and the drop in preschool blood lead levels continued even as the rate of decline in environmental lead asymptotically slowed (Nevin, 2007; Centers for Disease Control & Prevention, 2013). More recently, Finkelhor & Johnson (2015) proposed that the decline in juvenile delinquency may be related to increased rates of psychotropic medication utilization in pediatric populations. This seems particularly plausible in the case of stimulant medications, which are known to reduce aggression in school-aged children and criminal behavior among adults (Hinshaw *et al.* 1989; Hinshaw, 1991; Sinzig *et al.* 2007; Patti & Vanderschuren, 2008; Lichtenstein *et al.* 2012). Use of these medications was relatively rare prior to the mid-1990s but prescribing rates have increased

dramatically since then (Diller, 1996; Zuvekas & Vitiello, 2012; Olsson *et al.* 2015). We also noted the declining rates of child maltreatment in recent years (see the section Introduction). Maltreatment has similarly been linked to multiple externalizing-related outcomes and so this is another factor that could be contributing to the trends observed here (Teicher *et al.* 2003; Hicks *et al.* 2009; Heim *et al.* 2010). These factors do not comprise an exhaustive list of candidates and the trends we observe are likely to be multi-causal in nature. But our results underscore the need to consider a broad spectrum of behaviors in examining potential causes for these trends and to utilize multivariate approaches when possible.

Some caveats and limitations to our findings must be enumerated. As noted earlier, there have been reductions in several domains of adolescent risk behaviors over the past 15 or more years, including specific crimes and sexual-risk taking. Our interpretation that trends are a result of a trend in an underlying trait is limited to trends in the outcomes and the time-period studied here. A final noteworthy limitation is that we cannot evaluate or rule out the possibility of causal relationships among our outcomes; for example, alcohol use is likely to influence violent behavior, notwithstanding the likelihood of those behaviors sharing common etiologies. Finally, standard limitations associated with observational studies and self-reported outcomes should be kept in mind.

While the full explanation for recent trends in SUDs and delinquent behaviors is almost certainly multicausal, the analyses presented here provides some clues into their etiology by showing that they appear to reflect an overall trend in an Externalizing-like trait. Future research should characterize this phenomenon in greater detail by investigating what other behaviors have been influenced by this trend and identifying which segments of the population have been most affected. This could provide further clues into the contributing causes to this phenomenon, elucidation of which would be invaluable toward facilitating continuation of these improvements in adolescent health as far into the future and across as much of the adolescent population as possible.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717002999>

Acknowledgements

This work was supported by grants from the National Institute on Drug Abuse: DA23668, DA042195, DA32573 and DA04041 and DA031288. The funding

agency had no role in the design, conduct, collection, management, analysis or interpretation of data, or the preparation, review or approval of this paper. NSDUH data were obtained from the Interuniversity Consortium for Social and Political Research. We are grateful to Dr. David Finkelhor, University of New Hampshire, and Dr. John Constantino, Washington University School of Medicine, for helpful discussion in the early stages of drafting this manuscript.

References

- Abma JC, Martinez GM, Copen CE** (2010). Teenagers in the United States: sexual activity, contraceptive use, and childbearing. *Vital Health Statistics* **23**, 1–47.
- Board on Children, Youth, and Families** (2012). *Social Trends and Child Maltreatment Trends in Child Maltreatment Research, Policy, and Practice for the Next Decade*. National Academies Press: Washington, DC.
- Burt SA** (2009). Rethinking environmental contributions to child and adolescent psychopathology: a meta-analysis of shared environmental influences. *Psychological Bulletin* **135**, 608–637. doi:10.1037/a0015702
- Centers for Disease Control and Prevention (CDC)** (2013). Blood lead levels in children aged 1-5 years - United States, 1999–2010. *MMWR Morbidity and Mortality Weekly Report* **62**, 245–248.
- Cheung GW, Rensvold RB** (2002). Evaluating Goodness-of-Fit indexes for testing measurement invariance. *Structural Equation Modeling Multidisciplinary Journal* **9**, 233–255. doi:10.1207/S15328007SEM0902_5
- Child Trends Data Bank** (2015). Teen Homicide, Suicide, and Firearm Deaths. [Online] (<http://www.childtrends.org/?indicators=lead-poisoning>). Accessed 5 03 2016.
- Dick DM, Viken RJ, Kaprio J, Pulkkinen L, Rose RJ** (2005). Understanding the Co-variation among childhood externalizing symptoms: genetic and environmental influences on conduct disorder, attention deficit hyperactivity disorder, and oppositional defiant disorder symptoms. *Journal of Abnormal Child Psychology* **33**, 219–229. doi:10.1007/s10802-005-1829-8
- Dietrich KN, Douglas RM, Succop PA, Berge OG, Bornschein RL** (2001). Early exposure to lead and juvenile delinquency. *Neurotoxicology and Teratology* **23**, 511–518. doi:10.1016/S0892-0362(01)00184-2
- Diller LH** (1996). The run on ritalin: attention deficit disorder and stimulant treatment in the 1990s. *Hastings Center Reports* **26**, 12–18. doi:10.2307/3528571
- Donovan JE, Jessor R** (1985). Structure of problem behavior in adolescence and young adulthood. *Journal of Consulting and Clinical Psychology* **53**, 890–904. doi:10.1037/0022-006X.53.6.890
- Eaton NR, Keyes KM, Krueger RF, Balsis S, Skodol AE, Markon KE, Grant BF, Hasin DS** (2012). An invariant dimensional liability model of gender differences in mental disorder prevalence: evidence from a national sample. *Journal Abnormal Psychology* **121**, 282–288. doi: 10.1037/a0024780
- Farrelly MC, Loomis BR, Han B, Gfroerer J, Kuiper N, Couzens GL, Dube S, Caraball RS** (2013). A comprehensive examination of the influence of state tobacco control programs and policies on youth smoking. *American Journal of Public Health* **103**, 549–555. doi:10.2105/AJPH.2012.300948
- Finkelhor D** (2013). *Trends in Bullying and Peer Victimization*. University of New Hampshire: Crimes against Children Research Center: Durham.
- Finkelhor D, Johnson M**, (2015). Has psychiatric medication reduced crime and delinquency? *Trauma Violence & Abuse* **18**, 1–9. doi:10.1177/1524838015620817.
- Finkelhor D, Shattuck A, Turner HA, Hamby S** (2014). Trends in children’s exposure to violence, 2003 to 2011. *JAMA Pediatrics* **168**, 540–546.
- Grucza RA, Agrawal A, Krauss MJ** (2016). Declining prevalence of marijuana Use disorders among adolescents in the United States, 2002–2013. *Journal of the American Academy of Child and Adolescent Psychiatry* **55**, 487–494. doi:10.1016/j.jaac.2016.04.002
- Grucza RA, Plunk AD, Hipp PR, Cavazos-Rehg P, Krauss MJ, Brownson RC, Bierut LJ** (2013). Long-term effects of laws governing youth access to tobacco. *American Journal of Public Health* **103**, 1493–1499. doi:10.2105/AJPH.2012.301123
- Gruenewald PJ** (2011). Regulating availability: how access to alcohol affects drinking and problems in youth and adults. *Alcohol Research and Health* **34**, 248–256. doi:SPS-AR&H-39
- Hatzenbuehler ML, Schwab-Reese L, Ranapurwala SI, Hertz MF, Ramirez MR** (2015). Associations between antibullying policies and bullying in 25 states. *JAMA Pediatrics* **169**, e152411. doi:10.1001/jamapediatrics.2015.2411
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO** (1991). The Fagerstrom test for nicotine dependence: a revision of the fagerstrom tolerance questionnaire. *Addiction* **86**, 1119–1127.
- Heim C, Shugart M, Craighead WE, Nemeroff CB** (2010). Neurobiological and psychiatric consequences of child abuse and neglect. *Developmental Psychobiology* **52**, 671–690. doi:10.1002/dev.20494
- Hicks BM, Krueger RF, Iacono WG, McGue M, Patrick CJ** (2004). Family transmission and heritability of externalizing disorders: a twin-family study. *Archives of General Psychiatry* **61**, 922–928. doi:10.1001/archpsyc.61.9.922
- Hicks BM, South SC, DiRago AC, Iacono WG, McGue M** (2009). Environmental adversity and increasing genetic risk for externalizing disorders. *Archives of General Psychiatry* **61**, 922–928. doi:10.1001/archgenpsychiatry.2008.554
- Hinshaw SP** (1991). Stimulant medication and the treatment of aggression in children with attentional deficits. *Journal of Clinical Child Psychology* **20**, 301–312. doi:10.1207/s15374424jccp2003_9
- Hinshaw SP, Buhrmester D, Heller T** (1989). Anger control in response to verbal provocation: effects of stimulant medication for boys with ADHD. *Journal of Abnormal Child Psychology* **17**, 393–407. doi:10.1007/BF00915034
- ICPSR** (2016). National Survey on Drug Use and Health (NSDUH) Series. [Online] (<http://www.icpsr.umich.edu/icpsrweb/ICPSR/series/64>). Published 2015. Accessed 1 August 2016.

- Jessor R (1991). Risk behavior in adolescence: a psychosocial framework for understanding and action. *Journal of Adolescent Health* **12**, 587–605.
- Jessor R, Jessor S (1977). *Problem Behavior and Psychosocial Development: A Longitudinal Study of Youth*. Academic Press: New York.
- Johnston L, O'Malley PM, Miech RA, Bachman JG, Schulenberg JE, (2013). *Monitoring the Future: National Results on Adolescent Drug Use: 2013. Overview of Key Findings on Adolescent Drug Use*. University of Michigan, Institute for Social Research: Michigan.
- Kendler KS, Prescott CA, Myers J, Neale MC (2003). The structure of genetic and environmental risk factors for common psychiatric and substance Use disorders in Men and women. *Archives of General Psychiatry* **60**, 929–937. doi:10.1001/archpsyc.60.9.929
- Krueger RF (1999). The structure of common mental disorders. *Archives of General Psychiatry* **56**, 921–926. doi: 10.1001/archpsyc.56.10.921
- Krueger RF, Hicks BM, Patrick CJ, Carlson SR, Iacono WG, McGue M (2002). Etiologic connections among substance dependence, antisocial behavior and personality: modeling the externalizing spectrum. *Journal of Abnormal Psychology* **111**, 411–424. doi:10.1037/0021-843X.111.3.411
- Krueger RF, South SC (2009). Externalizing disorders: Cluster 5 of the proposed meta-structure for DSM-V and ICD-11. *Psychological Medicine* **39**, 2061–2070. doi:10.1017/S0033291709990328
- Lahey BB (2008). Oppositional defiant disorder, conduct disorder, and juvenile delinquency. In: *Child and Adolescent Psychopathology* (ed. T. Beauchaine and S. P. Hinshaw), pp. 335–369. Wiley: New York.
- Lane SD, Webster NJ, Levandowski BA, Rubinstein RA, Keefe RH, Wojtowycz MA, Cibula DA, Kingdon JEF, Aubry RH, (2008). Environmental injustice: childhood lead poisoning, teen pregnancy, and tobacco. *Journal of Adolescent Health* **42**, 43–49. doi:10.1016/j.jadohealth.2007.06.017
- Lichtenstein P, Halldner L, Zetterqvist J, Sjölander A, Serlachius E, Fazel S, Långström N, Larsson H (2012). Medication for attention deficit-hyperactivity disorder and criminality. *The New England Journal of Medicine* **367**, 2006–2014. doi:10.1056/NEJMoa1203241
- McGue M, Iacono WG, Legrand LN, Elkins I (2001). Origins and consequences of age at first drink. II. Familial risk and heritability. *Alcoholism: Clinical and Experimental Research* **25**, 1166–1173. doi:dx.Doi.org/10.1111/j.1530-0277.2001.tb02331.x.
- Morgan R, Musu-Gillette L, Robers S, Zhang A (2015). *Indicators of School Crime and Safety: 2014 (No. NCJ 248036)*. Bureau of Justice Statistics: Washington, DC.
- Muthén L, Muthén B (1998). *Mplus User's Guide*. 7th edn. Muthén & Muthén: Los Angeles, CA.
- Nevin R (2000). How lead exposure relates to temporal changes in IQ, violent crime, and unwed pregnancy. *Environmental Research* **83**, 1–22. doi:10.1016/j.envres.2007.02.008
- Nevin R (2007). Understanding international crime trends: the legacy of preschool lead exposure. *Environmental Research* **104**, 315–336. doi:10.1006/enrs.1999.4045
- Office of Juvenile Justice and Delinquency Prevention OJJDP, (2015). OJJDP Statistical Briefing Book. Online. [Online] (http://www.ojjdp.gov/ojstatbb/crime/JAR_Display.asp?ID=qa05206) Accessed 11 March 2016.
- Olsson M, Druss BG, Marcus SC (2015). Trends in mental health care among children and adolescents. *The New England Journal of Medicine* **372**, 2029–2038. doi:10.1056/NEJMsa1413512
- Pattij T, Vanderschuren L (2008). The neuropharmacology of impulsive behaviour. *Trends in Pharmacological Sciences*, **29**, 192–199. doi:10.1016/j.tips.2008.01.002
- Perlus J, Brooks-Russell A, Wang J, Iannotti R (2014). Trends in bullying, physical fighting, and weapon carrying among 6th- through 10th-grade students from 1998 to 2010: findings from a national study. *American Journal of Public Health* **104**, 1100–1106. doi:10.2105/AJPH.2013.301761
- Reyes JW (2007). Environmental policy as social policy? The impact of childhood lead exposure on crime. *The B E Journal of Economic Analysis & Policy* **7**, 51. doi:10.2202/1935-1682.1796
- Robers S, Zhang J, Truman J (2010). *Indicators of School Crime and Safety: 2010 (NCES 2011-002/ NCJ 230812)*. National Center for Education Statistics, U.S. Department of Education, and Bureau of Justice Statistics, Office of Justice Programs, U.S. Department of Justice: Washington, DC.
- Schwarz G (1978). Estimating the dimension of a model. *Annals of Statistic* **6**, 461–464.
- Sinzig J, Döpfner M, Lehmkuhl G, German Methylphenidate Study Group, Uebel H, Schmeck K, Poustka F, Gerber WD, Günter M, Knölker U, Gehrke M, Hässler F, Resch F, Brünger M, Ose C, Fischer R (2007). Long-acting methylphenidate has an effect on aggressive behavior in children with attention-deficit/hyperactivity disorder. *Journal of Child and Adolescent Psychopharmacology* **17**, 421–432. doi:10.1089/cap.2007.0011
- Stretesky PB, Lynch MJ (2001). The relationship between lead exposure and homicide. *Archives of Pediatrics and Adolescent Medicine Journal* **155**, 579–582. doi:10.1001/archpedi.155.5.579
- Substance Abuse and Mental Health Services Administration. Results from the 2012 National Survey on Drug Use, Health: Summary of National Findings (No. NSDUH Series H-46, HHS Publication No. (SMA) 13–4795). Rock Hill MD. 2014.
- Substance Abuse and Mental Health Services Administration (2014). National Survey on Drug Use and Health Methodological Summary and Definitions. <https://www.samhsa.gov/data/sites/default/files/NSDUH-MethodSummDefs2014/NSDUH-MethodSummDefs2014.htm>. Accessed October 4, 2017.
- Substance Abuse and Mental Health Services Administration (2015). Methodological Resource Books. Population Data/NSDUH. <http://www.samhsa.gov/data/population-data-nsduh/reports?tab=39>. Accessed August 24, 2016.

- Teicher MH, Andersen SL, Polcari A, Anderson CM, Navalta CP, Kim DM** (2003). The neurobiological consequences of early stress and childhood maltreatment. *Neuroscience & Biobehavioral Reviews* **27**, 33–44. doi:10.1016/S0149-7634(03)00007-1
- Verona E, Sachs-Ericsson N** (2005). The intergenerational transmission of externalizing behaviors in adult participants: the mediating role of childhood abuse. *Journal of Consulting and Clinical Psychology* **27**, 1135–1145. doi:10.1037/0022-006X.73.6.1135
- Webster DW, Meyers JS, Buggs MS** (2014). Youth acquisition and carrying of firearms in the United States: Patterns, consequences, and strategies for prevention. Presented at the Proceedings of Means of Violence Workshop, Forum of Global Violence Prevention. Washington, DC, Institutes of Medicine of the National Academies.
- White N, Lauritsen JL** (2012). Violent crime against youth, 1994–2010. Off. Justice Programs Bur. Justice Stat. NJC 240106.
- Young SE, Stallings MC, Corley RP, Krauter KS, Hewitt JK** (2000). Genetic and environmental influences on behavioral disinhibition. *American Journal of Medical Genetics* **96**, 684–695. doi:10.1002/1096-8628(20001009)96:5<684::AID-AJMG16>3.0.CO;2-G
- Zuvekas SH, Vitiello B** (2012). Stimulant medication use in children: a 12-year perspective. *American Journal of Psychiatry* **169**, 160–166. doi:10.1176/appi.ajp.2011.11030387